

Spatial and Seasonal Thermal Microclimate Variability in Reef Fisheries Zones on the Southwestern Coast of Madagascar

Andrew R. Villeneuve¹, Merrill Baker-Médard², Bemahafaly Randriamanantsoa³, Easton R.
White¹

¹Department of Biological Sciences, University of New Hampshire, NH, USA

²Department of Environmental Studies, Middlebury College, Middlebury, VT, USA

³Wildlife Conservation Society Madagascar Program, Soavimbahoaka, Antananarivo 101,
Madagascar

ORCID: Andrew Robert Villeneuve: 0000-0001-7303-5931; Merrill Baker-Médard: 0000-0002-
3724-8320; Easton Radley White: 0000-0002-0768-9555

Keywords: marine microclimate, marine heatwaves, fisheries climate sensitivity, Western Indian
Ocean, reef flats, gleaning

Author Contributions

Conceptualization: Andrew Robert Villeneuve, Easton Radley White; Methodology: Andrew
Robert Villeneuve, Merrill Baker-Médard, Bemahafaly Randriamanantsoa, Easton Radley
White; Formal Analysis: Andrew Robert Villeneuve; Writing - first draft: Andrew Robert
Villeneuve; Writing - review and editing: Andrew Robert Villeneuve, Merrill Baker-Médard,
Bemahafaly Randriamanantsoa, Easton Radley White; Funding Acquisition: Merrill Baker-
Médard, Easton Radley White; Project Administration: Merrill Baker-Médard, Bemahafaly
Randriamanantsoa, Easton Radley White

Statements and Declarations

The authors declare no competing interests.

Abstract

1. Climate change poses an understudied threat to the coastal reef systems of the Western Indian Ocean (WIO), a biodiversity hotspot critical for the maintenance of subsistence fisheries. The southwestern coast of Madagascar is especially sensitive to disturbances due to the high level of socioecological dependence on functioning reefs.
2. Fishing in this region is spatially structured. For example, gleaning (*pêche-à-pied*) is undertaken predominantly by women and children walking in intertidal reef flat and crest habitats during low tide, while boat-based fishing occurs across a range of reef geomorphic zones. These zones may experience different microclimates due to differences in depth, water residence time, and oceanic exposure.
3. To characterize the thermal microclimates of these zones, we deployed temperature loggers for over a year across reef geomorphic zones on Salary Reef, Madagascar. We further examined modeled tide elevation data over the deployment period and the 18.6-year tidal cycle.
4. Subtidal loggers showed similar mean temperature but diverged up to 27% in cumulative hours above 30 °C. Compared with gridded temperature datasets, logger observations revealed higher extremes and lower minima, underscoring the added fidelity of in-situ measurements.
5. The intertidal logger recorded extreme thermal stress (up to 48.8 °C) in gleaning habitats, with maximum heat events occurring in transitional “shoulder” seasons rather than peak summer. Tidal analyses showed that midday low tides in these transitional periods drive extreme intertidal warming.

6. Fisheries in Madagascar will likely experience uneven and seasonally distinct thermal risks. Climate adaptation must consider how microclimate variability mediates socioecological vulnerability in coastal oceans.

Résumé

1. Le changement climatique représente une menace encore peu étudiée pour les systèmes récifaux côtiers de l'océan Indien occidental (OIO), un hotspot de biodiversité essentiel au maintien des pêches de subsistance. La côte sud-ouest de Madagascar est particulièrement sensible aux perturbations en raison de la forte dépendance socio-écologique à des récifs fonctionnels.
2. Les pratiques de pêche dans cette région sont spatialement structurées. Par exemple, la pêche à pied (*gleaning*), pratiquée principalement par les femmes et les enfants, se déroule dans les habitats intertidaux de platiers et de crêtes récifales à marée basse, tandis que la pêche embarquée couvre un ensemble de zones géomorphologiques récifales. Ces zones peuvent présenter des microclimats distincts en fonction de la profondeur, du temps de résidence de l'eau et de leur exposition à l'océan ouvert.
3. Afin de caractériser ces microclimats thermiques, nous avons déployé des enregistreurs de température pendant plus d'un an dans différentes zones géomorphologiques du récif de Salary, à Madagascar. Nous avons également examiné les données de hauteur de marée modélisées sur la période de déploiement ainsi que sur un cycle tidal de 18,6 ans.
4. Les enregistreurs subtidaux ont montré des températures moyennes similaires, mais jusqu'à 27% de différence dans les heures cumulées au-dessus de 30 °C. Comparées aux jeux de données de température interpolées, les observations des enregistreurs ont

révélé des extrêmes plus élevés et des minima plus bas, soulignant la plus grande fidélité des mesures *in situ*.

5. L'enregistreur intertidal a mis en évidence un stress thermique extrême (jusqu'à 48,8 °C) dans les habitats exploités par la pêche à pied, les événements maximaux de chaleur survenant durant les saisons de transition plutôt qu'au cœur de l'été austral. Les analyses tidales ont montré que les marées basses de milieu de journée, propres à ces périodes de transition, entraînent un réchauffement intertidal extrême.
6. Les pêcheries de Madagascar connaîtront probablement des risques thermiques inégaux et distincts selon les saisons. Les stratégies d'adaptation au climat doivent donc intégrer la variabilité microclimatique qui module la vulnérabilité socio-écologique des océans côtiers.

83 Introduction

84 Chronic warming trends and acute heatwaves, already predicted to accelerate under climate
85 change in the tropical Western Indian Ocean (WIO) (McClanahan et al. 2009; Harrington and
86 Otto 2020; Mawren et al. 2022; Thorai et al. 2022), are predicted to increase temperature trends
87 within reef systems. Due to complex interactions between bathymetry and water movements in
88 the near-shore zone of coastal reefs, distinct microclimates may arise and experience climate
89 change differently from the open ocean. Since human activity is greatly concentrated in these
90 coastal zones, understanding the current and future microclimatic trends of reefs can greatly
91 improve conservation and management decision-making (Woodson et al. 2019). Coastal
92 socioeconomic systems in the WIO are strongly linked to the ecological functioning of
93 ecosystems, and are thus predicted to be at a higher risk of disruption compared to areas of
94 more rapid warming in developed countries (Cinner et al. 2012; Le Manach et al. 2012; Cullen-
95 Unsworth et al. 2014).

96 Within the WIO, southwestern Madagascar is a region particularly vulnerable to the social
97 and ecological impacts of climate change and heatwaves. Reefs in the region provide critical
98 fisheries and ecosystem services for people in nearby cities and villages, where livelihoods and
99 income are highly dependent on the reef function in proportion to the rest of the WIO (Allison et
100 al. 2009; Harris et al. 2010; Cinner et al. 2012; Gough et al. 2020; Ranaivomanana et al. 2023).
101 However, compound stressors from climate change, elevated fishery pressure, decreasing
102 water quality, enhanced cyclone activity, changing land use patterns, and increasing
103 populations due to coastal migration have placed profound stress on these reefs (Bruggemann
104 et al. 2012; Andréfouët et al. 2013; Carter et al. 2022). While much of the conservation focus on
105 the fringing reef system of southwestern Madagascar has been on unsustainable harvest and
106 adherence to marine protected area (MPA) policies, there has been relatively little work

examining the sensitivity of this marine system to climate change. Nonetheless, these reefs are critically imperiled by thermal stress (Ateweberhan and McClanahan 2010; D'Agata and Maina 2022; Obura et al. 2022), which contributed to historic wide-scale coral bleaching in the region in 1998 and 2016 (Obura et al. 2022; Randrianarivo et al. 2022).

However, assessments of seawater temperature on reefs in southwestern Madagascar have almost exclusively been made using remote-sensed data products (McClanahan et al. 2009; Harris et al. 2010), which masks the information from coastal reef ecosystems in which most artisanal fisheries operate. Internal waves, water mass pooling, and other hydrodynamic phenomena can result in subtidal heterogeneity on reef systems, further complicating assessments of thermal stress on complex coastal reef systems (Leichter et al. 2005; Davis et al. 2011; Woodson et al. 2019; Reid et al. 2020; Brown et al. 2022). Reef flat and shallow lagoon environments are expected to experience higher average and maximum temperatures than deep lagoons and backreefs/forereefs, mainly due to the mixing of cooler offshore waters from greater depths with warmed shallow waters (Reid et al. 2020; Kekuwa et al. 2021; Brown et al. 2022). Therefore, fisheries that are spatially segregated across depth and distance offshore may end up fishing in different thermal microclimates.

In addition to national and multinational industrial fishing, a large proportion of fisheries in the WIO are artisanal (catches sold at the market) and subsistence (catches consumed at home), and provide much of the coastal population's nutrition and livelihoods (Le Manach et al. 2012; White et al. 2022). In the WIO, men and women tend to fish in different habitats using distinct gear, with men typically using boats to access offshore sites and employing nets and lines, but also fishing on foot. Women, by contrast, fish on foot in nearshore areas, harvesting benthic invertebrates with hand-held tools (Westerman and Benbow 2013; Fröcklin et al. 2014; Baker-Médard and Kroger 2024). This type of fishing by foot, or gleaning, is the predominant fishing method for women and low-income households (Cinner and Bodin 2010; Fröcklin et al.

2014)Fortuna-Jackson *forthcoming*(Cinner and Bodin 2010; Fröcklin et al. 2014). Gleaning, (fishing on foot) tends to be restricted to coastal areas accessible by foot, such as mangroves, intertidal flats, seagrass beds, and reef crests, and occurs primarily during calmer seasons (de la Torre-Castro et al. 2017; Lemahieu et al. 2018; Grantham et al. 2021; Tilley et al. 2021; Wallner-Hahn et al. 2022). While men engage primarily in boat-based fishing, some—often those with fewer economic resources—practice gleaning, which does not require a boat or substantial capital investment. Similar species can be found in both deep and shallow habitats, the species composition and abundance within catches vary dramatically between habitats (Fröcklin et al. 2014; Jankowski et al. 2015; Medina-Valmaseda et al. 2020; Stiepani et al. 2023).

Given the asymmetric impact of climate change and heatwaves (marine and atmospheric) on shallow water coastal ecosystems, it is possible that species and populations targeted by women and the least wealthy of a given fishing community will be impacted first and to a greater magnitude than fisheries dominated by slightly wealthier men in deeper waters. This is especially significant given the importance of gleaned organisms for nutrients in the local food supply. Further, it's likely that pressure will increase on gleaning fisheries with climate change, drought, and overfishing, which are transitions from boat-based fisheries and inland agriculture to gleaning (Willer et al. 2023).

Complicating any assessment of thermal stress on this reef system is the additional interaction of the macrotidal semidiurnal tide system in southwestern Madagascar, with a tidal range of around three meters (Chevalier et al. 2015; Moustapha et al. 2021). These tides expose reef flats and even the reef crest at periodic intervals, providing opportunities for gleaning to occur. Tidal emersion on reefs, when timed with solar exposure, can result in devastating bleaching in corals and mass mortality in reef invertebrates, including lucrative fished species such as sea cucumbers (Glynn 1968; Yamaguchi 1975; Anthony and Kerswell

2007; Buckee et al. 2020, 2022; Mejía-Rentería et al. 2020; Schoepf et al. 2020). Given the additional factors of tidal cycles, reef hydrodynamics, and substrate albedo, satellite-derived temperature measurements are inappropriate to describe the thermal environment in which both reefs and fisheries coexist in southwestern Madagascar.

Analyses of climate sensitivity of Malagasy reefs and fisheries have historically been limited to satellite-derived sea surface temperature products, potentially ignoring microclimatic complexities on reefs that experience a spatial segregation of fishery types. Here, we describe the thermal environment of Salary Reef, southwestern Madagascar, across three reef geomorphic zones (backreef, lagoon, and flat) that encompass subtidal and intertidal sites by deploying long-term temperature loggers. We hypothesize maximum temperatures and variability to be highest at intertidal reef flat sites, and amongst subtidal sites, to be highest in backreef and shallow lagoon environments. These zones encompass fishing sites regularly used separately by boat-based and gleaning fishers, thereby describing the thermal environment used by fisheries-targeted species. We further modeled the 18-year tidal cycle (Denny and Paine 1998) of Salary Reef to assess reef flat immersion timing on a yearly, seasonal, and daily basis.

Methods

Study Site

The study was conducted on Salary Reef (22.53 S, 43.25 E; Figure 2), a fringing reef system located in southwestern Madagascar, approximately 100 km north of the region's major city, Toliara. This fringing reef system is part of "Mikea Coast", and is distinct from the major, better-studied barrier reef off the coast of Toliara, Grand Récif de Toliara (GRT), because it receives little riverine inputs. The GRT has long been known to be influenced by the hydrology

of the Onilahy and Fiherenana rivers, and excess sedimentation from upstream terrestrial activities such as mining and agriculture has partially contributed to the decline of the GRT (Pichon 1978; Andréfouët et al. 2013). However, Salary Reef is relatively uncoupled from terrestrial activity and is in better ecological condition than the GRT. This fringing reef system is composed of intertidal flats sloping to a sandy-bottomed lagoon varying 2-5 km in width (~5-15 m depth). The reef crest is periodically emerged during low-tide events, and complex reefs dominate the far reef slope (Randrianarivo et al. 2022, 2023). Further, southwestern Madagascar can be considered a meso- to macrotidal system – tidal ranges can extend up to 5.5 meters (near Besalampy), exposing extensive reef flats and parts of the barrier/fringing reef crest. The Soariake MPA, managed by the Soariake Association with technical assistance from the Wildlife Conservation Society (WCS), was formally protected in 2015 to balance conservation and fishing and encompasses all of Salary Reef and the northern portion of the fringing reef system, but adherence by small-scale artisanal fishers is variable (Zafimahatradraibe et al. 2025). Most boat-based fishing activity occurs on the forereef and in the lagoon, while gleaning can occur both in the shore-adjacent reef flat and on the reef crest itself.

Logger Deployment and Recovery

We deployed ten temperature loggers (ElectricBlue T7.3, Porto, Portugal — all IDs SALEB# “Salary ElectricBlue”) between July 28th and August 10th, 2023, distributed between four reef geomorphic zones (reef flat, lagoon, backreef, and forereef) between 2-4 m depth (Figure 1, Figure 2, Supplemental Table 2). We classified reef flat locations as representative of intertidal gleaning sites, as we were not able to deploy loggers directly on the reef crest. Lagoon, backreef, and forereef sites represented boat fishing locations. Additionally, we selected a site at the Salary Bay dive center to measure the air temperature at 2 m above ground. For all but the air temperature logger, we did not attempt to shade loggers from solar

radiative warming (Bahr et al. 2016; Colin and Johnston 2020; Rich et al. 2024), so as to replicate the microclimate or body temperature that may actually be experienced by exposed sessile marine organisms in each geomorphic reef zone (Kearney 2006; Jimenez et al. 2008; De Frenne et al. 2025). Temperature data should therefore not be interpreted as seawater temperature, but as a record of potential benthic ectotherm body temperature. The simple use of seawater (or air) temperature to understand potentially stressful periods fails to account for the complexities of an organism's heat budget, even in submerged environments.




Fig. 1 A cross section of a typical reef system of southwestern Madagascar with corresponding fishing zones and logger deployment locations is shown. The reef flat and reef crest (not sampled) are the intertidal portions of these reef systems, on which gleaning occurs during low tide events. The reef lagoon, backreef, and forereef are subtidal zones that are mainly fished by boat.




Fig. 2 Logger deployment sites near the village of Salary in southwestern Madagascar are shown. Large colored points show recovered logger locations with their corresponding reef geomorphic zone. Unrecovered loggers are shown outlined in gray. Of the nine deployed loggers on the reef, we recovered four loggers across three of the reef's geomorphic zones. The locations of fishing sites are indicated with labels, while the location of the air temperature logger is not shown. The inset map depicts the location of Salary Reef within Madagascar with accompanying mean yearly sea surface temperatures (2002-2009, Bio-ORACLE, (Assis et al. 2024)).

We affixed loggers to the reef using UV-resistant zip-ties and Splash-Zone two-part marine epoxy (Modern Recreational Technologies, Hickory, NC, USA). We selected crevices for

logger deployment to reduce breakage from wave action and record temperatures of these reef refugia for many targeted fishery species. We recorded GPS coordinates with up to 5 m accuracy (Garmin eTrex 10) and recorded videos of logger placement (GoPro Hero8 Black) to facilitate recovery. Temperature logger accuracy was set to 0.1 °C, recording every 15 minutes. We recovered five loggers after at least one year (Supplemental Table 2). Logger deployment and recovery were accomplished by freediving.

Data Analysis

Temperature Data

Of the ten temperature loggers deployed, we recovered five (four reef loggers, one air temperature logger) after at least 357 days (SALEB8, Flats) and at most after 459 days (SALEB1). Although some loggers collected data for over a year, we focused our comparative site analysis on only the dates where all four recovered water temperature loggers actively collected data (357 days).

To compare our logger-recorded reef temperatures against commonly used remote-sensed products, we further downloaded and extracted seawater temperature data from NOAA OISST (Optimal Interpolated Sea Surface Temperature; (Huang et al. 2021) and Copernicus Global Physics Analysis (European Union-Copernicus Marine Service 2016) at -22.6° S, 43.25° E. OISST records sea surface temperatures were available at 1/4° resolution, while we selected Copernicus temperature records from a depth of 0.5 m at a 1/12° resolution. We calibrated our air temperature data against downscaled ERA5 models of air temperature using the *mcera5* R package (Klinges et al. 2022).

250 Tidal Regime

251 As no tide stations are available in southwestern Madagascar, we used the package
252 *pyTMD* (Sutterley et al. 2024) in Python (v. 3.10.15) to calculate tide heights at the coordinates -
253 22.6° S, 43.25° E between the years 2006-2024. This timescale was chosen to encapsulate the
254 18.6-year lunar cycle to which short-term tidal cycles respond (Denny and Paine 1998). It further
255 illustrates whether the tidal regime recorded in our 2023-2024 intertidal temperature data
256 represents the long-term trend for this location. We used the TPXO9 (v5, 1/30° resolution) atlas
257 model of global tides constituents (Egbert and Erofeeva 2002). For some of our analysis, we
258 grouped tide data by four seasons — early hot-wet (November-January), late hot-wet (February-
259 April), early warm-dry (May-July), and late warm-dry (August-October). Initial exploration of the
260 intertidal logger data indicated that distinct temperature regimes occurred along the four
261 seasonal divisions listed above, and so, we processed some of our tidal data using this
262 seasonality.

263 We used a custom script to determine the relative tide height and tidal exposure regime of the
264 intertidal temperature logger (SALEB8 Flat), based on significant temperature “drops” correlated
265 with rapid cooling or warming of temperature loggers when exposed to air or water (Harley and
266 Helmuth 2003; Gilman et al. 2006). We determined the intertidal temperature logger to have a
267 height of -0.956 m (relative to average height) and +22 minutes behind the modeled tide series
268 obtained via the *pyTMD* package. From this tide height, we reconstructed the immersion status
269 of each time point from the intertidal temperature logger time series. When determining the
270 decadal patterns of low-tide exposure, we used the rounded height of this logger (-1 m, relative
271 average modeled height) as a threshold beyond which we calculated hours of exposure.

Statistical Analysis

All statistical analysis was completed in R (v. 4.2.1). For cross-site comparisons, we filtered available water temperature datasets to only the dates that data had for all sites (July 29th, 2023, to July 20th, 2024). We used a Kruskal-Wallis non-parametric test to determine if median site temperatures were different, and then *post hoc* Bonferroni-adjusted Dunn's test package to determine pairwise site significance.

We used wavelet analysis to determine the temporal patterns in power spectra during the deployment period of our temperature data time series. By decomposing temperature periodicity on a time basis, this method allows for the detection of non-stationary periodicity in environmental time series and provides an advantage over traditional power spectra analysis (Rivest and Gouhier 2015). We used the *WaveletComp* package to decompose and visualize the wavelet spectra using default Morlet wavelets (Roesch and Schmidbauer 2018).

Results

Reef Microclimate Variability

Of the three subtidal data loggers (SALEB1, SALEB3, and SALEB5), the backreef site (SALEB3) had the highest overall mean (27.3 °C) and maximum (33.4 °C) temperatures (Figure 3). It also spent a similar number of hours above the Coral Reef Watch (CRW) bleaching threshold of 30 °C (Southwestern Madagascar Coral Reef Watch threshold, (Skirving et al. 2020) as the shallow lagoon site (SALEB1, Supplemental Table 1). The deep lagoon site (SALEB5), the deepest one overall, had the lowest mean (27.08 °C) and maximum temperatures (33.0 °C), and the fewest number of hours (1089.2 h) above the CRW 30 °C threshold. A Kruskal-Wallis test confirmed that water temperature was significantly different between sites ($H_2 = 160.5$, $P < 0.0001$). *Post-hoc* multiple comparisons indicated significant

differences between all sites, although the significance between the two shallow sites was weak (SALEB1 x SALEB3, Bonferroni-adjusted multiple comparison, $P = 0.0452$), while the deep lagoon site SALEB5 was significantly cooler than both shallower sites (Bonferroni-adjusted multiple comparison, $P < 0.0001$).




Fig. 3 A Temperature time series from the four recovered loggers on Salary Reef, SW Madagascar, are shown. Vertical dashed lines indicate the extent of data used for site comparisons. Data were collected every 15 minutes. **B** The distribution of temperatures at each site is shown. Boxplots indicate mean and interquartile range (25-75%). Asterisks give multiple comparisons (Wilcox's) significance across all loggers – note that in the text, we restrict our multiple comparisons to just subtidal sites, and therefore, there are slight differences in significance level.

We further calculated the daily variance and temperature range of each subtidal logger, and only the temperature range was significantly different between sites (Daily Variance Kruskal-Wallis, $H_2 = 3.69$, $P = 4.72$; Daily Range Kruskal-Wallis, $H_2 = 16.7$, $P = 0.0002$). The daily temperature range was highest at the backreef site SALEB3 and lowest at the shallow lagoon site SALEB1 (Table 1). Post-hoc comparisons demonstrated that the temperature range at the backreef site was only significantly different from the shallow lagoon site (SALEB1 x SALEB3, Bonferroni-adjusted multiple comparisons, $P = 0.000135$). Despite having a mean

temperature of only 0.18/0.21 °C below and a maximum temperature only 0.1/0.4 °C the shallow lagoon and backreef sites respectively, the deep lagoon site had 27% fewer hours above 30 °C compared to these sites.

Table 1 A Summary statistics of temperatures recorded on Salary Reef, calculated between 2023-07-29 and 2024-07-20, and reanalysis products used to compare the accuracy of *in-situ* loggers against commonly used sea surface temperature data sources. **B** To compare our *in-situ* data against the daily reanalysis temperature timeseries, we performed a daily summarization of subtidal logger data. Additional details about *in situ* loggers can be found in Table S2. *Denotes maximum air temperature values that were not calculated due to a short period of logger exposure to direct sunlight

A) Data Source	ID	Zone	Mean (°C) \pm SD	Minimum (°C)	Maximum (°C)	Hours Above 30 °C	Daily Range (°C) \pm SD
In-situ logger	SALEB1	Shallow Lagoon	27.26 \pm 2.43	21.0	33.1	1478	1.49 \pm 0.48
In-situ logger	SALEB3	Backreef	27.29 \pm 2.33	21.2	33.4	1446	1.7 \pm 0.63
In-situ logger	SALEB5	Deep Lagoon	27.08 \pm 2.26	21.0	33.0	1089	1.58 \pm 0.57
In-situ logger	SALEB8	Reef Flat	27.64 \pm 3.57	17.3	48.8	2510	6.42 \pm 6.42
In-situ logger	SALEB10	Air Temperature	26.36 \pm 4.6	12.3	*	1917	11.45 \pm 2.81
B) Data Source	ID	Zone	Mean (°C)	Minimum (°C)	Maximum (°C)	Days Above 30 °C	
Reanalysis	Copernicus Global Ocean Physics Analysis and Forecast		27.34 \pm 2.44	22.91	31.96	15.75	
Reanalysis	NOAA/NCEP I OISST V2		27.22 \pm 2.07	23.05	30.66	9	
Daily summarized in-situ logger	SALEB1	Shallow Lagoon	27.27 \pm 2.39	21.70	31.82	14	

Daily summarized in-situ logger	SALEB3	Backreef	27.3 ± 2.3	21.94	31.59	13.5
Daily summarized in-situ logger	SALEB5	Deep Lagoon	27.09 ± 2.21	21.84	31.32	10.75

The seawater temperatures measured *in situ* summarized as daily means and those measured from reanalysis products (OISST, Copernicus) were not significantly different, indicating that variability driving site-level differences occurs on a sub-daily timescale (Kruskal-Wallis test, $H_2 = 4.722$, $P = 0.0943$, supplemental Fig. 1). Correlation ranged between 0.86 – 0.96 (R^2 , SALEB8 Reef Flat – SALEB3 Shallow Lagoon) for water logger data and Copernicus data, and between 0.76 – 0.92 (R^2 , SALEB8 Reef Flat – SALEB5 Deep Lagoon) for logger data and OISST data (supplemental Fig. 3). However, OISST tended to underpredict the hottest values observed with loggers, while the Copernicus dataset at a 0.5 m depth better matched the distribution of observed data (Table 1, supplemental Fig. 1). Correlations between each 15-minute logger observation and daily reanalysis data were lower than the summarized logger data values, with correlations ranging from 0.62 – 0.93 (R^2 , SALEB8 Reef Flat – SALEB1 Backreef) for water logger data and Copernicus data, and between 0.54 – 0.89 (R^2 , SALEB8 Reef Flat – SALEB5 Deep Lagoon) for water logger data and OISST data (supplemental Fig. 4).

The intertidal temperatures collected at SALEB8 (Reef Flat) were radically different from the other subtidal temperature series due to periods of air exposure during low tide. Both exposure during low tide and the day/night cycles modulated the temperatures experienced at this site, indicated by strong wavelet power levels associated with 8-hour (length of submersion), 12.4-hour (diel tidal cycle), and 24-hour (daily cycle) cycle periods (Figure 5). Vertical bands below the 12.4-hour periods correspond to a combination of tidal harmonics and variable emersion durations under seasonally changing tidal amplification. Average emersion duration was 2.36 hours (median 2.47 hours), although the left-skew of tidal emersion duration indicates longer durations, up to 3.7 hours, are more frequent (supplemental Fig. 7). The three

other subtidal loggers showed power bands over the 12.4- and 24-hour wavelet periods, again indicating the influence of tidal and solar cycles on temperature regimes on the subtidal reef (supplemental Fig. 8). Finally, the air temperature recorded by the temperature logger displayed an obviously strong 24-hour wavelet period power band, corresponding with solar cycles, but also a seasonally moderate 12-hour power band corresponding to atmospheric tides and land-sea breezes that are stronger during the cloudless warm-dry season (supplemental Fig. 8).




Fig. 5 **A** Wavelet analysis of intertidal reef flat (SALEB8) temperature between August 2023 and July 2024. Black outlined areas indicate regions that have strong wavelet power support ($p < 0.01$). **B** Average power spectra illustrate the strongest periods observed over the logger deployment period. Black dots correspond to periods that were strongly statistically significant ($p > 0.01$). Across both plots, four power ridges centered around 8 hours, 12.4 hours, 24 hours, and 2 weeks correspond to tidal exposure, tidal cycle, daily solar cycles, and tidal flushing/upwelling events, respectively. Periods of ~8 hours correspond to the longest tide cycles that exposed the logger for the longest, and the decreasing power of vertical bands indicates the gradually shorter tide cycles that exposed the logger for less and less time. High

power levels between the 128h and 1024h wavelet periods (encompassing a 2-week significant band) may correspond with tidal flushing of the lagoon (Chevalier et al. 2015).

The intertidal temperature data showed a strong seasonal pattern; we recorded extreme temperatures not in the peak wet season/January, but between March to June, and August to October. Loggers recorded a maximum temperature of 48.8 °C on March 24th, 2024, with additional excessive heat days occurring on March 12th, 2024 (48.6 °C) and September 3rd, 2024 (47.5 °C). Between November and February, water temperatures were consistently warmer than the air temperature in the intertidal zone, despite this being the hottest portion of the year in southwest Madagascar. These extreme temperatures can be attributed to low-tide events (Figure 4A). Examining the residuals of the relationship between the intertidal (SALEB8) and lagoon (SALEB5) loggers (linear model, $R^2 = 0.64$, $P < 0.0001$) reveals minimal variation from zero residuals in submerged observations (Figure 4B). However, exposed observation residuals alternate between much warmer (~5-20 °C, March-October) and cooler (~ -5 °C, November-February) than the water temperature in the lagoon. In short, low-tide temperatures were lower than other parts of the year, and indeed lower than water temperature, during the hottest part of the year (wet season, November-March). Water temperature residuals were not significantly different between night and day observations (Figure 4C).




Fig. 4 A Tidal exposure status of the intertidal flat (SALEB8) logger. Air exposure/water submersion status was determined via the application of interpolated logger tide height and offset from tidal model predictions. Logger status predictions in August and September 2023 erroneously identified heat spikes more often associated with low tide as submergence events. Temperature residuals from the lagoon (**B-C**, SALEB5) and air temperature residuals (**D-E**, SALEB10) from intertidal zone temperatures (SALEB8) are also shown. In **B**, nonzero residuals of submerged observations indicate differences in water temperature at the intertidal site. These submerged residuals were not different between day and night observations (**C**). When examining residuals of exposed observations in **D**, nonzero observations indicate times when the exposed intertidal logger was warmer or cooler than the air temperature at Salary. Residuals of exposed observations were significantly different between day and night exposures (**E**).

Residuals of the relationship between observations of the intertidal logger (SALEB8) and the air temperature (SALEB10) (linear model, $R^2 = 0.44$, $P < 0.0001$) show that low-tide logger temperatures were much warmer than air temperature, indicating the importance of additional microclimate factors (e.g., albedo, radiative forcing) on low-tide temperature on intertidal flats (Figure 4D). During the hot, wet season, the intertidal flat logger experienced fewer positive residual spikes with low tide while periodically having negative residuals. However, during the warm, dry season, flat logger temperatures routinely exceed air temperatures. The most extreme negative residuals were recorded during the daytime (Figure 4E). Mean hourly air temperatures recorded using temperature loggers generally matched those from ERA5 microclimate downscaling data ($R^2 = 0.84$, supplemental Fig. 2)

Decadal Tide Patterns

Tides during the field season of 2023-2024 occurred during a weakening in the 18.6-year lunar cycle — low tides were less extreme than at the peak of the cycle in 2015-2016, particularly in the early warm-dry (May-Jul) and early hot-wet seasons (Nov-Jan; Figure 6, supplemental Fig. 5). The maximum tidal range near the Salary reef was 3.4 m. However, through additional data exploration, we found that the tidal range increased to over 5.5 m moving northward to a point off the northwestern coast of the Besalampy district (~16.65 S, 44.42 E)m. However, early warm-dry low tides spend more time below -1 m and have lower overall tides during the day than the night, while the inverse is true during early hot-wet low tides, where nighttime low tides are generally longer and larger. In the late hot-wet and warm-dry months, there is no trend in monthly hours below -1 m or monthly minimum tide across years, indicating that the timing of extreme low tides is consistent along the 18.6-year tidal cycle. Tidal heights that exceed -1 m (and would therefore expose the intertidal flats logger) routinely occur between 09:00 and 15:00 (EAT) across the 18.6-year tidal cycle, encapsulating

solar noon (supplemental Fig. 6). Low tides that expose reef flats below a -1 m elevation are therefore consistent in their diel and annual timings, although interannual extremity varies.




Fig. 6 A Hours spent below -1 m (mean tide, TPXO9 tide model) and **B** minimum tide height (TPXO9 tide model) for each month at Salary Reef between 2006 and 2024 (18 years) across austral seasons (Hot-Wet = November-April, Warm-Dry = May-October) and daily sun cycles. Lines of fit are loess lines. Each point represents the monthly maximum hours below -1 m or

424 minimum tide height (split by day and night), so that there are 24 data points each year (12
425 monthly calculations across day and night cycles).

426 Discussion

427 Through our analysis of *in-situ* temperature data and long-term tidal amplification, we
428 uncovered heterogeneity and extremity across three geomorphic zones within a barrier reef
429 system in southwestern Madagascar. Subtidal sites, while having superficially similar thermal
430 regimes, exhibited up to 27% more heating hours (hours above 30 °C, deep vs. shallow lagoon).
431 Such variance and variation in extremes across temperature series with similar means is
432 common when comparing time series from loggers deployed on reef flats/back reefs and deeper
433 reef sites (Guadayol et al. 2014; Cyronak et al. 2020). However, due to the relatively large tidal
434 range and depth of the lagoon (~15 m max), we see less fine-scale variation in subtidal station
435 temperatures than typical coastal lagoons in microtidal settings (Pichon 1978). Many of the
436 extreme observations across sites (both maxima and minima) were larger than those obtained
437 from two remote-sensed products, indicating that *in-situ* logger observations describe the
438 thermal environment with more fidelity. This is a common observation, as remote-sensed and
439 reanalysis products often mask coastal areas, spatially average, and do not account for other
440 microclimatic processes such as substrate albedo and water residence time (Castillo and Lima
441 2010; Bos and Pinsky 2025; Margaritis et al. 2025; Thompson et al. 2025). Interestingly, the
442 subtidal sites at Salary Reef all had larger temperature maxima (February 2020: 30.7 °C, depth
443 = 25m, February 1964: 30.25 °C, depth = 10m) and minima (April 2020: 21.5 °C, depth = 25m,
444 July 1964 21.45 °C, depth = 10m) than reported from the nearby GRT (Pichon 1978; Godefroid
445 et al. 2024). Variation between January and May at the GRT was much higher than at other
446 times of the year at the same site, including the (on average) cooler months of July and August.
447 This was attributed to the influence of larger tides (observed in our study), increased cyclone
448 activity, and internal waves arising from increased wind activity during this period (Farquhar et

al. 2022; Godefroid et al. 2024; Thompson et al. 2025). Overall, diel heat flux across the reef can be attributed to daytime solar irradiance, which is modulated by the interaction of atmospheric (cloud cover, wind speed, solar azimuth) and oceanic (tidal range, depth, wave height) conditions (Davis et al. 2011; Zhang et al. 2013; Cyronak et al. 2020).

While all subtidal loggers indicated an expected trend pattern of elevated temperature exposure during the hot-wet season (January-February), temperatures recorded from the intertidal flat logger displayed a contrasting pattern of elevated temperatures during the shoulder seasons. Our wavelet analysis of this intertidal logger confirmed a weakening of 24-, 12.4-, and 8-hour bands during the hot-wet season, which is explained by the decreased tide amplification during this period. Further, the lack of warming compared to ambient air temperature in the hot, wet season suggests sufficiently low tides did not occur during the daytime. The relative cooling of logger temperatures throughout the year further indicates convective cooling from wind can reduce thermal exposure below ambient air temperatures, even during the day (Helmuth et al. 2011). However, data from 1961 indicated temperatures of 41 °C in pooled water on the exposed reef crest of the GRT between November and December. Interestingly, this observation timeframe corresponds with the annual minimum tidal amplitude we describe on nearby Salary Reef (Pichon 1978) where we did not observe intertidal temperatures in excess of 38 °C. The maximum decadal tidal amplitude occurred in 1959, and so we can contextualize the 1961 extreme temperatures in this same season as the product of shorter low-tide durations and heights in the present day than in 1961. It is therefore possible that even more extreme temperatures would have been recorded in 1961 if data were recorded during the annual maximum tidal amplitudes of November-January or May-July. Indeed, we quote directly from Pichon (1978):

472 *“The data we have are actually too fragmented to permit an exact appreciation of the*
473 *importance of this warming phenomenon of waters on the reef flat.”*¹

474 Our long-term analysis of the tide series over an 18.6-year cycle indicates that the diel and
475 seasonal timing of longer intertidal exposures are consistent across years, while the length of
476 emersion durations changed year over year. Our data was collected during a relatively low-
477 exposure period of the 18.6-year cycle, suggesting that intertidal logger observations in 2015 or
478 2033 would show hotter, lower, and longer intertidal periods.

479 Bolstered by chronic warming and acute heatwave events, we predict that the next ~16
480 years will see intertidal habitats in southwestern Madagascar experiencing hotter conditions
481 than we observed during our deployment period. This is critical because the two lowest tide
482 windows between February to April and August to October are the most optimal for gleaning,
483 both from a consideration of access to low tidal elevation seafloor and because of the relatively
484 calm seas in this period (Grantham et al. 2021; Farquhar et al. 2022; Maka et al. 2022).
485 Between these tide windows in the warm-dry season, however, wind speeds off southwestern
486 Madagascar increase, which limits the number of days fishermen can access offshore sites and,
487 for gleaners, the exposed offshore reef crest (Pichon 1978; Farquhar et al. 2022). Furthermore,
488 the stationarity of midday low tides throughout the year is a particular risk factor for tropical reefs
489 in southwestern Madagascar. On the Australian Great Barrier Reef, the time forms of the tidal
490 harmonics result in shifting low-tide timing over the year, such that extreme low tides occur only
491 around midday in the winter, potentially offering thermal refuge to intertidal organisms (Buckee
492 et al. 2022).

¹ *“Les données que nous possédons sont actuellement trop fragmentaires pour permettre d’apprécier l’importance exacte de ce phénomène d’échauffement des eaux sur l’ensemble du platier.”*

Inferring Reef Microclimate to Reef Coral and Fishery Resilience

In southwestern Madagascar, gleaning supports a diverse array of species critical to local food security and livelihoods, including sea urchins, bivalves, gastropods, octopus, and fish (Rabesandratana 1985; Herinirina et al. 2023). Among these, sea cucumbers (Bêche-de-Mer, trepang) represent a particularly lucrative commercial resource, harvested alongside more subsistence-oriented species like sea urchin and shellfish. Sea cucumbers, or holothurians, are a relatively abundant target species in protected reef environments and can be found up to the intertidal reef flat zone. Holothurians can be an important part of a gleaner's catch, and additionally, there are major holothurian aquaculture pens in the lagoon of the GRT and numerous other lagoon sites across southwestern Madagascar (McVean et al. 2005; Lavitra et al. 2006; Baker-Médard and Kroger 2024). A majority of holothurians are destined for the export market, and thus can be an important source of currency for gleaners (McVean et al. 2005; Lavitra et al. 2006; Fattebert 2020; Maka et al. 2022). Planktonic larval holothurian densities were the highest between February and April, which correspond with the hottest flat reef temperatures observed (Rasolofonirina and Conand 1998; Lavitra et al. 2006). Further, the highest number of fishers and fishing days occur between April to December, although this seasonality has been attributed to avoiding increased lagoon turbidity due to river flooding in the GRT, which is less of a problem on Salary Reef (Maka et al. 2022).

The underlying coral habitat is also threatened by climate change. Corals and their associated zooxanthellae, Symbiodiniaceae, are able to exhibit moderate tolerance to subtidal heat pulses due to their molecular and cellular plasticity, which can be selected for in highly variable or warm environments (Drury 2020). While some coral colonies exposed to prolonged air exposure during macrotidal regimes have elevated resilience to atmospheric heatwaves (Schoepf et al. 2020; Brown et al. 2024), sufficient heat and air exposure quickly result in intertidal mass mortality across taxa (Glynn 1968, 1976; Schoepf et al. 2015; Ainsworth et al.

2016; Castrillón-Cifuentes et al. 2017; Le Nohaïc et al. 2017; Buckee et al. 2020, 2022; Godefroid et al. 2024). The delineation between resistance/resilience and declines in marine populations is a fine one. Even with local adaptations, *Acropora* coral populations declined dramatically under modeled 3 °C of warming (Lachs et al. 2024). While some might invoke shallow-water/intertidal habitats as a “space-for-time” substitution supporting the potential for organism resilience in a warmer ocean, the fact remains that populations in these reef flat environments are experiencing the warmest temperatures under current conditions, with a hard ceiling of aerial exposure. The hypothetical deadening of shallow-water coastal environments would be devastating for vulnerable gleaning fisheries operating in these marginal environments.

Will intertidal organisms adapt, move, or die in response to an 18.6-year cycle? Previous work investigating intertidal mussels in New Zealand and Washington State, USA, observed a decrease in average mussel intertidal height, a direct product of weakening survival in the upper intertidal and effectively compressing the vertical and horizontal range (Denny and Paine 1998; Sorte et al. 2018). Coupled with rapidly warming air and water temperatures in the WIO region (Ateweberhan and McClanahan 2010; Bruggemann et al. 2012; Lima and Wetthey 2012; Harrington and Otto 2020; Obura et al. 2022), hotter low tides may accelerate the depopulation of tidal flat habitats by gleaned species such as holothurians, whelks, and octopi. Over a longer decadal horizon, sea level rise may alleviate heat exposure of tidal flats by increasing inundation hours, at the expense of lost gleaning opportunities during weak low-tide periods (Harborne 2013). Before 1978, extensive coral cover of the reef flat by branching *Acropora* was reported by Pichon from the GRT, approximately 100 km south of Salary. By 2008, this intertidal coral cover was gone, likely due to a mixture of factors including cyclone activity, fishing gear impacts, poor water quality from Toliara, and river discharges (Harris et al. 2010; Andréfouët et al. 2013; Chevalier et al. 2015). Direct attribution of lost coral cover to warming

atmospheric/water temperatures and decadal tidal cycles is therefore not possible with current data, but our data adds context to coral cover declines in southwestern Madagascar.

Conclusions

Whether modeling or managing, the implication of differential seasonal climatic vulnerability in coastal reef and intertidal flats is that a focus on the putative hottest part of the year (hot-wet season, November-April) risks missing the most sensitive season for intertidal habitats. Microclimatic analysis may best describe conditions at a temporal and spatial scale most relevant to coastal human activities, thereby allowing for more careful considerations of the impacts of climate change on people and the ecosystems they rely on. Previous work assessing the climate sensitivity and resiliency of reef fisheries in Madagascar identified the northwest as the most sensitive region to climate impacts due to elevated SST and solar irradiance, but did not consider the importance of tidal regimes or reef microclimates on gleaning fisheries (Cinner et al. 2012, 2022). It should be highlighted that gleaners in the WIO are already well aware of ongoing seawater warming and its negative impacts on their catch, which, coupled with overexploitation, has resulted in reduced mollusk fishery yields (Cochrane et al. 2019; Alati et al. 2020).

As the climate warms, the reef system of southwestern Madagascar will continue to experience more frequent, long, and intense heatwave events amplified by the semidiurnal, mesotidal system. On this reef system, where gender roles, socioeconomic security, and fishery styles are spatially separated, we also uncovered spatial separation in microclimates and ultimately climate sensitivity. Future work could also assess the extent to which socio-ecological systems will be impacted, especially when considering extremes and spatial structure (White and Wulfinf 2024; Wulfinf and White 2024).

MPAs in Madagascar, including the Soariake MPA within which our study site is located, currently do not account for the impacts of climate change and different fisheries styles. Effective spatial management in coastal reef ecosystems will need to confront different use cases of reef zones that experience different magnitudes and timing of heat stress. In particular, shallow reef areas and the fisheries they support will experience heat stress sooner than deep reef habitats. Therefore, gleaning fisheries will require access to shallow reef zones within MPAs to be managed for their social resiliency. In addition, the promotion of alternative livelihoods to reduce fishing pressure will also be necessary, along with providing a livelihood portfolio in the case of heatwave-induced fishery collapse. By taking a microclimate lens to the thermal landscape of Salary Reef, we advance our spatio-temporal understanding of where extreme heat may be experienced by reef organisms the most, and by extension, the fishers that depend on these reef ecosystems.

Acknowledgements

The authors wish to thank Fred Lucas, Herdinand Haidodiado, and Elizabeth Kroger for the logistical and field support during logger deployment and recovery. Funding to ERW and MBM was provided by NSF grant awards 1923707 and 2409030. Additionally, we would also like to thank Ruscena Wiederholt for their input on earlier versions of this manuscript. Authorization of research was provided by the Direction Générale de la Gouvernance Environnementale of the Republic of Madagascar, n° 357 22 MEDD/SG/DGGE/DAPRNE/SCBE.

Data Availability Statement

Scripts and data will be made available in a public repository upon acceptance for publication.

589 Citations

- 590 Ainsworth TD, Heron SF, Ortiz JC, Mumby PJ, Grech A, Ogawa D, Eakin CM, Leggat W (2016)
591 Climate change disables coral bleaching protection on the Great Barrier Reef. *Science*
592 352:338–342
- 593 Alati VM, Olunga J, Olendo M, Daudi LN, Osuka K, Odoli C, Tuda P, Nordlund LM (2020)
594 Mollusc shell fisheries in coastal Kenya: Local ecological knowledge reveals overfishing.
595 *Ocean & Coastal Management* 195:105285
- 596 Allison EH, Perry AL, Badjeck M-C, Neil Adger W, Brown K, Conway D, Halls AS, Pilling GM,
597 Reynolds JD, Andrew NL, Dulvy NK (2009) Vulnerability of national economies to the
598 impacts of climate change on fisheries. *Fish and Fisheries* 10:173–196
- 599 Andréfouët S, Guillaume MMM, Delval A, Rasoamanendrika FMA, Blanchot J, Bruggemann JH
600 (2013) Fifty years of changes in reef flat habitats of the Grand Récif of Toliara (SW
601 Madagascar) and the impact of gleaning. *Coral Reefs* 32:757–768
- 602 Anthony KRN, Kerswell AP (2007) Coral mortality following extreme low tides and high solar
603 radiation. *Mar Biol* 151:1623–1631
- 604 Assis J, Fernández Bejarano SJ, Salazar VW, Schepers L, Gouvêa L, Fragkopoulou E, Leclercq
605 F, Vanhoorne B, Tyberghein L, Serrão EA, Verbruggen H, De Clerck O (2024) BIO-
606 ORACLE v3.0. Pushing marine data layers to the CMIP6 Earth System Models of
607 climate change research. *Global Ecol Biogeogr* 33:e13813
- 608 Ateweberhan M, McClanahan TR (2010) Relationship between historical sea-surface
609 temperature variability and climate change-induced coral mortality in the western Indian
610 Ocean. *Marine Pollution Bulletin* 60:964–970
- 611 Bahr KD, Jokiel PL, Rodgers KS (2016) Influence of solar irradiance on underwater temperature
612 recorded by temperature loggers on coral reefs. *Limnology and Oceanography: Methods*
613 14:338–342
- 614 Baker-Médard M, Kroger E (2024) Troubling the Waters: Gendered Dispossession, Violence,
615 and Sea Cucumber Aquaculture in Madagascar. *Society & Natural Resources* 37:443–
616 470
- 617 Bos JT, Pinsky ML (2025) Fine resolution satellite sea surface temperatures capture the
618 conditions experienced by corals at monthly but not daily timescales. *Coral Reefs*
619 44:423–434
- 620 Brown KT, Eyal G, Dove SG, Barott KL (2022) Fine-scale heterogeneity reveals
621 disproportionate thermal stress and coral mortality in thermally variable reef habitats
622 during a marine heatwave. *Coral Reefs*
- 623 Brown KT, Martynek MP, Barott KL (2024) Local habitat heterogeneity rivals regional
624 differences in coral thermal tolerance. *Coral Reefs*

625 Bruggemann J, Rodier M, Guillaume M, Andréfouët S, Arfi R, Cinner J, Pichon M, Ramahatratra
626 F, Rasoamanendrika F, Zinke J, McClanahan T (2012) Wicked Social-Ecological
627 Problems Forcing Unprecedented Change on the Latitudinal Margins of Coral Reefs: the
628 Case of Southwest Madagascar. *Ecology and Society* 17:

629 Buckee J, Hetzel Y, Edge W, Verduin J, Pattiaratchi C (2022) Daily timing of low tide drives
630 seasonality in intertidal emersion mortality risk. *Frontiers in Marine Science* 9:

631 Buckee J, Pattiaratchi C, Verduin J (2020) Partial mortality of intertidal corals due to seasonal
632 daytime low water levels at the Houtman Abrolhos Islands. *Coral Reefs* 39:537–543

633 Carter AL, Gilchrist H, Dexter KG, Gardner CJ, Gough C, Roccliffe S, Wilson AMW (2022)
634 Cyclone Impacts on Coral Reef Communities in Southwest Madagascar. *Front Mar Sci*
635 9:

636 Castillo KD, Lima FP (2010) Comparison of in situ and satellite-derived (MODIS-Aqua/Terra)
637 methods for assessing temperatures on coral reefs. *Limnology and Oceanography*:
638 *Methods* 8:107–117

639 Castrillón-Cifuentes AL, Lozano-Cortés DF, Zapata FA (2017) Effect of short-term subaerial
640 exposure on the cauliflower coral, *Pocillopora damicornis*, during a simulated extreme
641 low-tide event. *Coral Reefs* 36:401–414

642 Chevalier C, Devenon J-L, Rougier G, Blanchot J (2015) Hydrodynamics of the Toliara Reef
643 Lagoon (Madagascar): Example of a Lagoon Influenced by Waves and Tides. *coas*
644 31:1403–1416

645 Cinner JE, Bodin Ö (2010) Livelihood Diversification in Tropical Coastal Communities: A
646 Network-Based Approach to Analyzing ‘Livelihood Landscapes.’ *PLoS ONE* 5:e11999

647 Cinner JE, Caldwell IR, Thiault L, Ben J, Blanchard JL, Coll M, Diedrich A, Eddy TD, Everett JD,
648 Folberth C, Gascuel D, Guiet J, Gurney GG, Heneghan RF, Jägermeyr J, Jiddawi N,
649 Lahari R, Kuange J, Liu W, Maury O, Müller C, Novaglio C, Palacios-Abrantes J, Petrik
650 CM, Rabearisoa A, Tittensor DP, Wamukota A, Polnac R (2022) Potential impacts of
651 climate change on agriculture and fisheries production in 72 tropical coastal
652 communities. *Nat Commun* 13:3530

653 Cinner JE, McClanahan TR, Graham NAJ, Daw TM, Maina J, Stead SM, Wamukota A, Brown
654 K, Bodin Ö (2012) Vulnerability of coastal communities to key impacts of climate change
655 on coral reef fisheries. *Global Environmental Change* 22:12–20

656 Cochrane KL, Rakotondrazafy H, Aswani S, Chaigneau T, Downey-Breedy N, Lemahieu A,
657 Paytan A, Pecl G, Plagányi E, Popova E, van Putten EI, Sauer WHH, Byfield V, Gasalla
658 MA, van Gennip SJ, Malherbe W, Rabary A, Rabearisoa A, Ramaroson N,
659 Randrianarimanana V, Scott L, Tsimanaoraty PM (2019) Tools to Enrich Vulnerability
660 Assessment and Adaptation Planning for Coastal Communities in Data-Poor Regions:
661 Application to a Case Study in Madagascar. *Frontiers in Marine Science* 5:

662 Colin PL, Johnston TMS (2020) Measuring Temperature in Coral Reef Environments:
663 Experience, Lessons, and Results from Palau. *Journal of Marine Science and*
664 *Engineering* 8:680

665 Cullen-Unsworth LC, Nordlund LM, Paddock J, Baker S, McKenzie LJ, Unsworth RKF (2014)
666 Seagrass meadows globally as a coupled social–ecological system: Implications for
667 human wellbeing. *Marine Pollution Bulletin* 83:387–397

668 Cyronak T, Takeshita Y, Courtney TA, DeCarlo EH, Eyre BD, Kline DI, Martz T, Page H, Price
669 NN, Smith J, Stoltenberg L, Tresguerres M, Andersson AJ (2020) Diel temperature and
670 pH variability scale with depth across diverse coral reef habitats. *Limnology and*
671 *Oceanography Letters* 5:193–203

672 D’Agata S, Maina JM (2022) Climate change reduces the conservation benefits of tropical
673 coastal ecosystems. *One Earth* 5:1228–1238

674 Davis KA, Lentz SJ, Pineda J, Farrar JT, Starczak VR, Churchill JH (2011) Observations of the
675 thermal environment on Red Sea platform reefs: a heat budget analysis. *Coral Reefs*
676 30:25–36

677 De Frenne P, Beugnon R, Klimes D, Lenoir J, Niittynen P, Pincebourde S, Senior RA, Aalto J,
678 Chytrý K, Gillingham PK, Greiser C, Gril E, Haesen S, Kearney M, Kopecký M, le Roux
679 PC, Luoto M, Maclean I, Man M, Penczykowski R, van den Brink L, Van de Vondel S,
680 De Pauw K, Lembrechts JJ, Kemppinen J, Van Meerbeek K (2025) Ten practical
681 guidelines for microclimate research in terrestrial ecosystems. *Methods in Ecology and*
682 *Evolution* 16:269–294

683 Denny MW, Paine RT (1998) Celestial Mechanics, Sea-Level Changes, and Intertidal Ecology.
684 *The Biological Bulletin* 194:108–115

685 Drury C (2020) Resilience in reef-building corals: The ecological and evolutionary importance of
686 the host response to thermal stress. *Molecular Ecology* 29:448–465

687 Egbert GD, Erofeeva SY (2002) Efficient Inverse Modeling of Barotropic Ocean Tides. *J Atmos*
688 *Oceanic Technol* 19:183–204

689 European Union-Copernicus Marine Service (2016) Global Ocean 1/12° Physics Analysis and
690 Forecast updated Daily.

691 Farquhar S, Nirindrainy AF, Heck N, Saldarriaga MG, Xu Y (2022) The impacts of long-term
692 changes in weather on small-scale fishers’ available fishing hours in Nosy Barren,
693 Madagascar. *Front Mar Sci* 9:

694 Fattebert C (2020) La pêche traditionnelle ou petite pêche maritime à Madagascar. 46

695 Fröcklin S, Torre-Castro M de la, Håkansson E, Carlsson A, Magnusson M, Jiddawi NS (2014)
696 Towards Improved Management of Tropical Invertebrate Fisheries: Including Time
697 Series and Gender. *PLOS ONE* 9:e91161

698 Gilman SE, Harley CDG, Strickland DC, Vanderstraeten O, O’Donnell MJ, Helmuth B (2006)
699 Evaluation of effective shore level as a method of characterizing intertidal wave
700 exposure regimes. *Limnology & Ocean Methods* 4:448–457

701 Glynn PW (1968) Mass mortalities of echinoids and other reef flat organisms coincident with
702 midday, low water exposures in Puerto Rico. *Marine Biol* 1:226–243

703 Glynn PW (1976) Some Physical and Biological Determinants of Coral Community Structure in
704 the Eastern Pacific. *Ecological Monographs* 46:431–456

705 Godefroid M, Vandendriessche M, Todinanahary GGB, Ransquin I, Dubois P (2024) Thermal
706 sensitivity of black corals (Antipatharia: Hexacorallia): Comparisons between sympatric
707 species from a thermally fluctuating site in Madagascar and between allopatric
708 congeners. *Science of The Total Environment* 908:168311

709 Gough CLA, Dewar KM, Godley BJ, Zafindranosy E, Broderick AC (2020) Evidence of
710 Overfishing in Small-Scale Fisheries in Madagascar. *Front Mar Sci* 7:

711 Grantham R, Álvarez-Romero JG, Mills DJ, Rojas C, Cumming GS (2021) Spatiotemporal
712 determinants of seasonal gleaning. *People and Nature* 3:376–390

713 Guadayol Ò, Silbiger NJ, Donahue MJ, Thomas FIM (2014) Patterns in Temporal Variability of
714 Temperature, Oxygen and pH along an Environmental Gradient in a Coral Reef. *PLOS*
715 *ONE* 9:e85213

716 Harborne AR (2013) The ecology, behaviour and physiology of fishes on coral reef flats, and the
717 potential impacts of climate change. *Journal of Fish Biology* 83:417–447

718 Harley CDG, Helmuth BST (2003) Local- and regional-scale effects of wave exposure, thermal
719 stress, and absolute versus effective shore level on patterns of intertidal zonation.
720 *Limnology & Oceanography* 48:1498–1508

721 Harrington LJ, Otto FEL (2020) Reconciling theory with the reality of African heatwaves. *Nat*
722 *Clim Chang* 10:796–798

723 Harris A, Manahira G, Sheppard A, Gough C, Sheppard C (2010) Madagascar's Once Great
724 Barrier Reef – Change In Coral Reef Condition Over 40 Years.

725 Helmuth B, Yamane L, Lalwani S, Matzelle A, Tockstein A, Gao N (2011) Hidden signals of
726 climate change in intertidal ecosystems: What (not) to expect when you are expecting.
727 *Journal of Experimental Marine Biology and Ecology* 400:191–199

728 Herinirina PA, Bigot L, Frouin P, Volandrae HTE, Randriatsara RM, Behivoke F, Ranivoarivelo
729 L, Léopold M (2023) Seagrass macroinvertebrate bycatches support mosquito net trawl
730 fishery in Madagascar. *Aquat Living Resour* 36:18

731 Huang B, Liu C, Banzon V, Freeman E, Graham G, Hankins B, Smith T, Zhang H-M (2021)
732 Improvements of the Daily Optimum Interpolation Sea Surface Temperature (DOISST)
733 Version 2.1. *Journal of Climate* 34:2923–2939

734 Jankowski MW, Gardiner NR, Jones GP (2015) Depth and reef profile: effects on the distribution
735 and abundance of coral reef fishes. *Environ Biol Fish* 98:1373–1386

736 Jimenez IM, Kühl M, Larkum AWD, Ralph PJ (2008) Heat budget and thermal
737 microenvironment of shallow-water corals: Do massive corals get warmer than
738 branching corals? *Limnology and Oceanography* 53:1548–1561

739 Kearney MR (2006) Habitat, environment and niche: what are we modelling? *Oikos* 115:186–
740 191

741 Kekuewa SAH, Courtney TA, Cyronak T, Kindeberg T, Eyre BD, Stoltenberg L, Andersson AJ
742 (2021) Temporal and Spatial Variabilities of Chemical and Physical Parameters on the
743 Heron Island Coral Reef Platform. *Aquat Geochem* 27:241–268

744 Klings DH, Duffy JP, Kearney MR, Maclean IMD (2022) mcera5: Driving microclimate models
745 with ERA5 global gridded climate data. *Methods in Ecology and Evolution* 13:1402–1411

746 Lachs L, Bozec Y-M, Bythell JC, Donner SD, East HK, Edwards AJ, Golbuu Y, Gouezo M,
747 Guest JR, Humanes A, Riginos C, Mumby PJ (2024) Natural selection could determine
748 whether *Acropora* corals persist under expected climate change. *Science* 0:eadi6480

749 Lavitra T, Vaïtilingon D, Rasolofonirina R, Eeckhaut I (2006) Seasonal abundance of sea
750 cucumber larvae at Toliara Great Reef, Madagascar. *SPC Beche-de-mer Information*
751 *Bulletin* 24:35–38

752 Le Manach F, Gough C, Harris A, Humber F, Harper S, Zeller D (2012) Unreported fishing,
753 hungry people and political turmoil: the recipe for a food security crisis in Madagascar?
754 *Marine Policy* 36:218–225

755 Le Nohaïc M, Ross CL, Cornwall CE, Comeau S, Lowe R, McCulloch MT, Schoepf V (2017)
756 Marine heatwave causes unprecedented regional mass bleaching of thermally resistant
757 corals in northwestern Australia. *Sci Rep* 7:14999

758 Leichter JJ, Deane GB, Stokes MD (2005) Spatial and Temporal Variability of Internal Wave
759 Forcing on a Coral Reef. *Journal of Physical Oceanography* 35:1945–1962

760 Lemahieu A, Scott L, Malherbe WS, Mahatante PT, Randrianarimanana JV, Aswani S (2018)
761 Local perceptions of environmental changes in fishing communities of southwest
762 Madagascar. *Ocean & Coastal Management* 163:209–221

763 Lima FP, Wetthey DS (2012) Three decades of high-resolution coastal sea surface temperatures
764 reveal more than warming. *Nat Commun* 3:

765 Maka O, Pascal B, Todinanahary G (2022) Wild sea cucumber fishery in southwestern
766 Madagascar. *SPC Beche-de-mer Information Bulletin* 42:36–45

767 Margaritis G, Kent EC, Foster GL (2025) Intercomparison of satellite-derived SST with logger
768 data in the Caribbean—Implications for coral reef monitoring. *PLOS Climate* 4:e0000480

769 Mawren D, Hermes J, Reason CJC (2022) Marine heatwaves in the Mozambique Channel. *Clim*
770 *Dyn* 58:305–327

771 McClanahan TR, Ateweberhan M, Omukoto J, Pearson L (2009) Recent seawater temperature
772 histories, status, and predictions for Madagascar's coral reefs. *Mar Ecol Prog Ser*
773 380:117–128

774 McVean AR, Hemery G, Walker RCJ, Ralisaona BLR, Fanning E (2005) Traditional sea
775 cucumber fisheries in southwest Madagascar: A case-study of two villages in 2002. SPC
776 Beche-de-mer Information Bulletin 15–19

777 Medina-Valmaseda AE, Rodríguez-Martínez RE, Alvarez-Filip L, Jordan-Dahlgren E, Blanchon
778 P (2020) The role of geomorphic zonation in long-term changes in coral-community
779 structure on a Caribbean fringing reef. *PeerJ* 8:e10103

780 Mejía-Rentería JC, Castellanos-Galindo GA, Osorio-Cano JD, Casella E (2020) Subaerial
781 exposure of coral reefs during spring low tides in the eastern Pacific. *bms* 96:219–220

782 Moustapha S, Chevalier C, Sow B, Pagano M, Devenon J-L (2021) Coupled effects of tide and
783 swell on water renewal in a meso-tidal channel lagoon: Case of the Toliara Lagoon
784 (Madagascar). *Estuarine, Coastal and Shelf Science* 259:107463

785 Obura D, Gudka M, Samoilys M, Osuka K, Mbugua J, Keith DA, Porter S, Roche R, van
786 Hooideonk R, Ahamada S, Araman A, Karisa J, Komakoma J, Madi M, Ravinia I,
787 Razafindrainibe H, Yahya S, Zivane F (2022) Vulnerability to collapse of coral reef
788 ecosystems in the Western Indian Ocean. *Nat Sustain* 5:104–113

789 Pichon M (1978) Recherches sur les peuplements a dominance des anthozoaires dans les
790 recifs coralliens de Tulear. 490

791 Rabesandratana H (1985) Sur quelques utilisations récifales à Madagsacar. Fifth International
792 Reef Congress 6:661–668

793 Ranaivomanana HS, Jaquemet S, Ponton D, Behivoke F, Randriatsara RM, Mahafina J,
794 Léopold M (2023) Intense pressure on small and juvenile coral reef fishes threatens
795 fishery production in Madagascar. *Fisheries Management and Ecology* 30:494–506

796 Randrianarivo M, Botosoamananto RL, Guilhaumon F, Penin L, Todinanahary G, Adjeroud M
797 (2023) Effects of Madagascar marine reserves on juvenile and adult coral abundance,
798 and the implication for population regulation. *Marine Environmental Research*
799 190:106080

800 Randrianarivo M, Guilhaumon F, Tsilavonarivo J, Razakandrainy A, Philippe J,
801 Botosoamananto RL, Penin L, Todinanahary G, Adjeroud M (2022) A contemporary
802 baseline of Madagascar's coral assemblages: Reefs with high coral diversity,
803 abundance, and function associated with marine protected areas. *PLOS ONE*
804 17:e0275017

805 Rasolofonirina R, Conand C (1998) L'exploitation des holothuries dans le sud-ouest de
806 Madagascar, région de Toliarapar. SPC Beche-de-mer Information Bulletin 10:

807 Reid EC, Lentz SJ, DeCarlo TM, Cohen AL, Davis KA (2020) Physical Processes Determine
808 Spatial Structure in Water Temperature and Residence Time on a Wide Reef Flat.
809 *Journal of Geophysical Research: Oceans* 125:e2020JC016543

810 Rich WA, Glanz JS, Mele G, Burt NJ, Carvalho S, Doo SS, Goergen EA, Krieger EC, Li L,
811 Osman EO, Suggett DJ, Johnson MD, Fox MD (2024) Widespread inconsistency in

812 logger deployment methods in coral reef studies may bias perceptions of thermal
813 regimes. PLOS Climate 3:e0000517

814 Rivest EB, Gouhier TC (2015) Complex Environmental Forcing across the Biogeographical
815 Range of Coral Populations. PLOS ONE 10:e0121742

816 Roesch A, Schmidbauer H (2018) WaveletComp: Computational Wavelet Analysis.

817 Schoepf V, Jung MU, McCulloch MT, White NE, Stat M, Thomas L (2020) Thermally Variable,
818 Macrotidal Reef Habitats Promote Rapid Recovery From Mass Coral Bleaching. Front
819 Mar Sci 7:

820 Schoepf V, Stat M, Falter JL, McCulloch MT (2015) Limits to the thermal tolerance of corals
821 adapted to a highly fluctuating, naturally extreme temperature environment. Sci Rep
822 5:17639

823 Skirving W, Marsh B, De La Cour J, Liu G, Harris A, Maturi E, Geiger E, Eakin CM (2020)
824 CoralTemp and the Coral Reef Watch Coral Bleaching Heat Stress Product Suite
825 Version 3.1. Remote Sensing 12:3856

826 Sorte CJB, Bernatchez G, Mislan KAS, Pandori LLM, Silbiger NJ, Wallingford PD (2018)
827 Thermal tolerance limits as indicators of current and future intertidal zonation patterns in
828 a diverse mussel guild. Mar Biol 166:6

829 Stiepani J, Sandig A, Blicharska M (2023) The *Where*, the *How*, and the *Why* of the gleaning
830 fishery: Livelihoods, food security, threats and management on the island of Malalison,
831 Philippines. Ocean & Coastal Management 244:106806

832 Sutterley T, Alley K, Bishop-Taylor R, Brunt K, Howard S, Padman L, Siegfried M (2024)
833 pyTMD.

834 Thompson AM, Stathakopoulos A, Hollister KJ, Lynch AM, Holder JC, Kuffner IB (2025) Acute
835 heat stress and the extirpation of a threatened coral species from a remote, subtropical
836 reef system. Coral Reefs

837 Thorat F, Montie S, Thomsen MS, Tait LW, Pinkerton MH, Schiel DR (2022) Unravelling
838 seasonal trends in coastal marine heatwave metrics across global biogeographical
839 realms. Sci Rep 12:7740

840 Tilley A, Burgos A, Duarte A, dos Reis Lopes J, Eriksson H, Mills D (2021) Contribution of
841 women's fisheries substantial, but overlooked, in Timor-Leste. Ambio 50:113–124

842 de la Torre-Castro M, Fröcklin S, Börjesson S, Okupnik J, Jiddawi NS (2017) Gender analysis
843 for better coastal management – Increasing our understanding of social-ecological
844 seascapes. Marine Policy 83:62–74

845 Wallner-Hahn S, Dahlgren M, de la Torre-Castro M (2022) Linking seagrass ecosystem services
846 to food security: The example of southwestern Madagascar's small-scale fisheries.
847 Ecosystem Services 53:101381

848 Westerman K, Benbow S (2013) The role of women in community-based small-scale fisheries
849 management: the case of the southern Madagascar octopus fishery. *Western Indian*
850 *Ocean Journal of Marine Science* 12:119–132

851 White ER, Baker-Médard M, Vakhitova V, Farquhar S, Ramaharitra TT (2022) Distant water
852 industrial fishing in developing countries: A case study of Madagascar. *Ocean & Coastal*
853 *Management* 216:105925

854 White ER, Wulfinf S (2024) Extreme events and coupled socio-ecological systems. *Ecological*
855 *Modelling* 495:110786

856 Willer DF, Aldridge DC, Gough C, Kincaid K (2023) Small-scale octopus fishery operations
857 enable environmentally and socioeconomically sustainable sourcing of nutrients under
858 climate change. *Nat Food* 1–11

859 Woodson CB, Micheli F, Boch C, Al-Najjar M, Espinoza A, Hernandez A, Vázquez-Vera L,
860 Saenz-Arroyo A, Monismith SG, Torre J (2019) Harnessing marine microclimates for
861 climate change adaptation and marine conservation. *Conservation Letters* 12:e12609

862 Wulfinf S, White ER (2024) Social-ecological models with social hierarchy and spatial structure
863 applied to small-scale fisheries. *Theor Ecol* 17:325–336

864 Yamaguchi M (1975) Sea Level Fluctuations and Mass Mortalities of Reef Animals in Guam,
865 Mariana Islands. 11:

866 Zafimahatradraibe JA, Ranaivomanana LNJ, Rakotomahazo C, Randriamanantsoa B,
867 Hartmann AC, Todinanahary GGB (2025) Authority, capacity, and power to govern:
868 Three marine protected areas co-managed by resource users and non-governmental
869 organizations. *Marine Policy* 177:106647

870 Zhang Z, Falter J, Lowe R, Ivey G, McCulloch M (2013) Atmospheric forcing intensifies the
871 effects of regional ocean warming on reef-scale temperature anomalies during a coral
872 bleaching event. *Journal of Geophysical Research: Oceans* 118:4600–4616

873

874

875


876

877


878

879

Supplemental Material




Supplementary Figure 1 A-B comparison of remote-sensed sea water temperature products (Copernicus SST, NOAA OISST) and *in-situ* reef temperature recorded using loggers, with data summarized as daily means. Logger temperatures were recorded every 15 minutes, so we summarized temperatures as a daily mean to compare them to gridded reanalysis temperature products, which interpolate sea water temperature on a daily time step. OISST records sea surface temperatures, while Copernicus interpolates temperatures at a 0.5 m depth. Pairwise multiple comparisons revealed no statistical significance between any two sites. Dashed vertical lines in **B** indicate the extent of data used for cross-site comparisons.



890

891 **Supplementary Figure 2** Correlation plot between ERA5 2 m air temperatures and logger air
 892 temperatures in Salary village. The red line indicates a 1:1 line, and the blue line indicates the
 893 linear regression of the data. Note that large positive residual observations (>5 °C) are likely
 894 due to the logger being exposed to the direct sun.



895


896

897


898

899

Supplementary Figure 3 Correlation plot between Copernicus and OISST sea surface temperature observations and summarized daily mean logger temperatures at Salary reef. The red line indicates a 1:1 line, and the blue line indicates the linear regression of the data. The color scale of the residuals is pseudo-log to accommodate the range of residuals between sites.




Supplementary Figure 4 Correlation plot between Copernicus and OISST sea surface temperature observations and summarized raw logger temperatures (collected every 15 minutes) at Salary reef. The red line indicates a 1:1 line, and the blue line indicates the linear regression of the data. The color scale of the residuals is pseudo-log to accommodate the range of residuals between sites.




907

908 **Supplementary Figure 5** **A** Hours spent below -1 m (mean tide, TPXO9 tide model) and **B**
 909 minimum tide height (TPOX9 tide model) across 19 years, by month and time of day, on Salary
 910 Reef. Months are color-coded based on the maximum temperature from our intertidal flat
 911 temperature logger (SALEB8) for each month during the observation period (2003-2024), with
 912 red equaling the warmest and blue equaling the coldest.




Supplementary Figure 6 Minimum tide height each month for a given hour of the day and year at Salary reef (TPXO9 tide model). Low tides that exceeded -1 m consistently occur between 09:00 and 15:00.



920

921 **Supplementary Figure 7** Histogram of intertidal logger (SALEB8) inundation duration at the
 922 interpolated tidal height (-0.96m). The dotted vertical line indicates the mean inundation duration
 923 of 2.36 hours.

924



Supplementary Figure 8 Wavelet plots for the subtidal loggers (A-C) and air temperature logger (D). 24-hour horizontal lines indicate the position of power bands associated with daily solar cycles, while the 12.4-cycle corresponds with tidal cycles. The 12-hour cycle identified in the air temperature logger is likely due to semi-diurnal atmospheric tides and land-sea breeze circulation, which are strongest during cloudless seasons (warm-dry season). Black outlines in the wavelet plots indicate regions with pairs of dates/times and cycle periods that have strong power support ($p < 0.01$). Colored dots on the average wavelet plots indicate the strength of support for a significant wavelet period at a given duration.

937 **Table S1** Site data for logger deployments at Salary reef. Of the ten total loggers deployed, we
 938 recovered five (four on the reef and one air temperature logger).

939

Logger ID	Latitude	Longitude	Approximate Depth (m)	Zone	Recovered	Fishing	Deployment Date	Recovery Date	Days of Data
SALE B1	-22.5	43.25	366	Lagoon	Shallow	boat	7/27/2023	10/22/2024	453
SALE B2	-22.5	43.28	1017	Reef Flat		glea	7/28/2023		
SALE B3	-22.5	43.24	945	Backreef	Backreef	boat	7/28/2023	10/31/2024	461
SALE B4	-22.5	43.25	366	Lagoon		boat	7/28/2023		
SALE B5	-22.5	43.25	75	Lagoon	Deep	boat	7/28/2023	7/22/2024	360

SALE	-	43.24	2	Forereef	boat	7/28/2023		
B6	22.5	875						
	863							
SALE	-	43.24	2	Forereef	boat	7/28/2023		
B7	22.5	875						
	863							
SALE	-	43.28	2	Reef Flat	Flat	glea	7/29/2023	7/20/2 357
B8	22.5	1067				ning		024
	53							
SALE	-	43.28	2	Reef Flat		glea	8/10/2023	
B9	22.5	2483				ning		
	493							
SALE	-	43.28		Air	Chez		7/29/2023	8/7/20 375
B10	22.5	5767		Tempera	Fred			24
	542			ture				

940

941

Supplement Text

To determine the immersion regime of the field intertidal temperature logger, we built a function to extract the tidal height of the logger and apply an interpolated time series of immersion/emersion status to the measured intertidal temperatures. We first identified logger immersion points, diagnostic as precipitous drops of at least 3 °C/15 min from temperature peaks that occur during summer low tide events (Harley and Helmuth 2003; Gilman et al. 2006). We manually extracted points that visually fit this drop criterion, as we found other peak-finding algorithms to misidentify smaller drops as immersion points.

To determine the immersion regime of the intertidal flat logger (SALEB8), we developed a custom R function, *tide_findR*, to estimate the local tide height and time offset from the interpolated tide series at -22.6° S, 43.25° E using the *pyTMD* python package (Egbert and Erofeeva 2002; Sutterley et al. 2024); see Methods. Since tidal height and timing can vary spatially, this adjustment was necessary to align logger immersion events with the local tidal cycle.

First, the function shifts the reference tide series forward in time, extracting tide heights corresponding to each recorded immersion timestamp. At each shift, the function calculates the sum of squared differences (least squares) between observed immersion heights and the mean tide height. The time offset that minimizes the least squares is selected as the best estimate of the logger's time shift relative to the reference tide station. While a single time offset may not precisely describe the complexities of water movement within a barrier reef system, this approach approximates the time difference between the open-ocean tide model and experienced tides close to shore, within the Salary reef lagoon.

Using this optimal time shift, the function assigns a tide height to each immersion timestamp and calculates a mean logger height, representing the submergence threshold. This threshold is

966 then used to classify each temperature observation as submerged (immersed) or exposed
967 (emersed), providing a time series of temperature data linked to tidal exposure.

968 Supplement Citations

969 Egbert GD, Erofeeva SY (2002) Efficient Inverse Modeling of Barotropic Ocean Tides. J
970 Atmos Oceanic Technol 19:183–204

971 Gilman SE, Harley CDG, Strickland DC, Vanderstraeten O, O'Donnell MJ, Helmuth B
972 (2006) Evaluation of effective shore level as a method of characterizing intertidal wave
973 exposure regimes. Limnology & Ocean Methods 4:448–457

974 Harley CDG, Helmuth BST (2003) Local- and regional-scale effects of wave exposure,
975 thermal stress, and absolute versus effective shore level on patterns of intertidal
976 zonation. Limnology & Oceanography 48:1498–1508

977 Sutterley T, Alley K, Bishop-Taylor R, Brunt K, Howard S, Padman L, Siegfried M (2024)
978 pyTMD.

979

980

981