

1 Ecosystem dynamics in dry heathlands:  
2 spatial and temporal effects of  
3 environmental drivers on the vegetation

---

4

5 Christian Damgaard, ORCID 0000-0003-3932-4312

6 Department of Ecoscience, Aarhus University, C.F. Møllers Allé 4, 8000 Aarhus C, Denmark

7 Corresponding author: Christian Damgaard, [cf@ecos.au.dk](mailto:cf@ecos.au.dk), +45-30183153

8

9

## 10 **Abstract**

11 To understand and estimate the effects of environmental drivers on temperate dry heathland vegetation,  
12 pin-point cover data from 102 Danish sites sampled during a 16-year period was regressed onto selected  
13 environmental variables. The effects of nitrogen deposition, soil pH, soil C-N ratio, soil type, precipitation  
14 and grazing on the heathland vegetation was modelled in a spatio-temporal structural equation model  
15 using a Bayesian hierarchical model structure. The results suggest that the modelled environmental  
16 variables have important regulating effects on the large-scale spatial variation as well as plant community  
17 dynamics in dry heathlands. The cover of the dwarf shrub *Calluna vulgaris*, which is the characteristic  
18 species of dry heathlands, increased in relatively sandy soils with a relatively high C/N ratio. The cover of  
19 the grass *Avenella flexuosa* increased markedly at relatively high nitrogen deposition and low precipitation.  
20 The cover of other graminoids, of which *Molinia caerulea* is the most abundant, increases with a relatively  
21 low C/N ratio, low grazing pressure, and clayey soils. It was concluded that the modeled environmental  
22 variables are sufficient for predicting the *average* plant community dynamics of dry heathlands.  
23 Consequently, the model may be used to make forecasts for the effect of different management scenarios  
24 at a specific site and thus provide important input to setting up local adaptive management plans.

25 Keywords: Plant community dynamics of dry heathlands; spatial and temporal variation of plant cover;  
26 hierarchical Bayesian models; pin-point cover data; structural equation modelling; ecological predictions,  
27 adaptive management plans.

## 28 **Introduction**

29 To appropriately manage semi-natural dry heathland ecosystems under the influence of climate change  
30 and altered land use practices, it is important to understand and be able to predict the effects of  
31 environmental drivers on the vegetation. Improved quantitative understanding of the effects of the  
32 different environmental drivers will enable us to make quantitative predictions at the local site level to  
33 future ecosystem responses to changes in e.g. climate and grazing regimes, as well as, provide important  
34 input for local adaptive management plans (Damgaard 2022a; 2025b). Consequently, to better understand  
35 the causal relationships underlying the observed large-scale variation and temporal changes in dry  
36 heathlands as well as estimate the effect of the environmental drivers, a structural equation model of the  
37 dry heathland ecosystem was hypothesized and fitted to spatial and temporal ecological data from 102  
38 Danish dry heaths in the period 2007 to 2022.

39 The dynamics of semi-natural heathland ecosystems, and especially *Calluna* dominated heaths, were first  
40 studied by Watt (1947), who gave a detailed account of possible ecological processes that lead to different  
41 spatial vegetation patterns. This line of work has since been extended by several authors (e.g. Gimingham  
42 1978; Gimingham 1988; Gimingham et al. 1981; Løvschal and Damgaard 2022; Usher and Thompson 1993),  
43 who describe the spatial patterns of heathland vegetation at different scales and how they are regulated by  
44 management.

45 The conservation status and resilience of dry heathlands are hypothesized to be highest when a low  
46 nutrient status is maintained, succession to forest or grassland is actively prevented by maintaining  
47 traditional land-use practices, and ensuring spatial heterogeneity in disturbance patterns (Damgaard et al.  
48 2024; Olmeda et al. 2020). An analysis of the Danish dry heathland vegetation from 2004 to 2021 revealed  
49 declines in all three dominant species, *Calluna vulgaris*, *Empetrum nigrum*, and *Avenella flexuosa* (the decline  
50 in *C. vulgaris* was only marginally significant). These species are ecosystem engineers, and their decline

51 threatens the structure and function of heathland ecosystems. In contrast, the increasing cover of purple  
52 moor-grass, a tussock-forming species, may drive substantial ecological shifts (Damgaard 2025c). Overall,  
53 the conservation status of Danish dry heaths is deteriorating and unfavorable in about 40% of the areas  
54 (Nygaard et al. 2020).

55 The *a priori* selection of the studied environmental drivers was based on current ecological knowledge,  
56 where specific environmental drivers, e.g. atmospheric nitrogen deposition, soil type, soil pH and  
57 disturbance, have been shown to have an effect on heathland vegetation (e.g. Aerts et al. 1990; Aerts and  
58 Heil 1993; Britton et al. 2003; Damgaard et al. 2024; Damgaard et al. 2020; De Graaf et al. 2009; Vogels et  
59 al. 2020). However, dry heathland vegetation is expected to be regulated by more environmental drivers  
60 than it was possible to include in this study because of missing data, e.g. species-specific herbivores and  
61 pests and previous natural management actions. Some of these more or less unknown factors can have a  
62 geographical regional structure that may be partly explored using latent geographic factors (Ovaskainen et  
63 al. 2016), and possible significant effects of such latent geographic factors can generate new testable causal  
64 hypotheses. Furthermore, it is expected that soil type and nitrogen deposition may have both direct and  
65 indirect effects by affecting soil pH (Damgaard et al. 2014), and such direct or indirect causal pathways are  
66 often best modelled using structural equation models (SEM) that are fitted to observed ecological data  
67 (Grace et al. 2010). Generally, the effects of changes in environmental variables, e.g. climate, soil physical  
68 properties and disturbance regimes, on plant communities are expected to occur with some time-lags (e.g.  
69 Svenning and Sandel 2013). Furthermore, the regulating environmental variables are expected to vary  
70 considerably among different sites, and it is critical to integrate this large-scale spatial variation into the  
71 analysis of the ecosystem dynamics.

72 The objective of this study is to investigate the effects of selected environmental and ecological drivers  
73 (nitrogen deposition, soil type, soil pH, soil C-N ratio, precipitation, and grazing) on the vegetation at dry  
74 heaths, which is summarized by the multivariate relative cover of six species or aggregated species groups

75 (*Calluna vulgaris*, *Empetrum nigra*, *Avenella flexuosa*, other graminoids, other herbs, and cryptogams).  
76 Previously, similar models have been fitted to wet heathlands (Damgaard 2019; 2025a), as well as, acid  
77 (Damgaard 2022c), and calcareous grassland (Damgaard 2023).

## 78 **Materials and Methods**

### 79 **Dry heathlands**

80 Dry heathlands are semi-natural ecosystems on sandy, nutrient-poor soils, which first became widespread  
81 in prehistoric times under the influence of extensive agricultural practices (Løvschal 2021; Olmeda et al.  
82 2020). In Denmark, the dry heathlands are mainly situated in the western part of Denmark that were not  
83 covered by ice during the last (Weichselian) glacial period. The vegetation is mainly comprised of dwarf  
84 shrubs and graminoids, where especially *C. vulgaris*, *E. nigra*, and *A. flexuosa* are characteristic and often  
85 dominating species (Damgaard 2025c).

86 *Calluna vulgaris* (L.) Hull (heather) is a 10 to 50 cm tall dwarf shrub that is dominant on dry heaths, though  
87 not tolerant to heavy grazing, and prefers at least moderately well-drained soils (Gimingham 1960). The  
88 developmental phases of heather include pioneer, building, mature and degenerate (Usher and Thompson  
89 1993; Watt 1947), and without management actions such as burning, grazing or cutting, *C. vulgaris* plants  
90 degenerate when their age exceeds approximately 30 years, thereby leaving room for heathland  
91 succession.

92 *Empetrum nigrum* L. (black crowberry) is a low-growing, allelopathic, evergreen dwarf shrub typically  
93 forming dense mats on acidic, sandy, and nutrient-poor soils. It is well adapted to wind and salt aerosols. It  
94 is often found in late-successional or climax communities, though it can also act as a pioneer species in  
95 disturbed habitats. The dominant reproductive strategy is through adventitious rooting and sprouting from  
96 the basal parts (Tybirk et al. 2000).

97 *Avenella flexuosa* (L.) Drejer (wavy hair-grass) is a slender, tufted perennial grass typically 20 to 60 cm tall,  
98 which is found on acidic, sandy, and nutrient-poor soils. It is widely distributed across European heathlands  
99 and often increases in abundance following reduced grazing or elevated nitrogen deposition. It is tolerant  
100 of low nutrient availability but can become dominant under disturbance regimes that suppress dwarf  
101 shrubs (Aerts and Heil 1993).

102 The estimated cover and change in cover of the most common species in Danish dry heathland plots are  
103 shown in Table S1 (Damgaard 2025c).

## 104 **Sampling design**

105 Hierarchical time-series data from 102 dry heathland sites (Fig. 1) that had been monitored at least three  
106 times in the period from 2007 to 2022 were used in the analysis. Fifty-nine of the 102 sites are NATURA  
107 2000 habitat sites and are protected under the Habitat Directive (EU 1992). All sites included several plots  
108 classified as dry heathland (EU habitat type: 4030) according to the habitat classification system used for  
109 the European Habitat Directive EU (EU 2013; Olmeda et al. 2020). The area of the sites ranged from 0.14 ha  
110 to 1030 ha, with a median area of 11.7 ha. A total of 883 unique plots were used in the analysis. The  
111 sampling was performed in the summer, and all plots at a site were sampled on the same day. Otherwise,  
112 sampling intensity was irregular among sites and years, but all sites were monitored within a six-year  
113 period. Typically, ten plots were sampled from each site each time. Including resampling over the years, a  
114 total of 3525 plots were used in the analyses. The plots were resampled with GPS-certainty (< 10 meters).

115 The data are a subset of the data collected in the Danish habitat surveillance program NOVANA (Nielsen et  
116 al. 2012; Nygaard et al. 2024).

## 117 **Variables and measurements**

### 118 **Plant cover data**

119 The plant cover, which is the relative projected area covered by a species, was measured for all higher  
120 plants by the pin-point method using a square frame (50 cm X 50 cm) of 16 grid points that were equally  
121 spaced by 10 cm (Nielsen et al. 2012). At each grid point, a thin pin was inserted into the vegetation and  
122 the plant species that were touched by the pin were recorded and used as an estimate of cover (Damgaard  
123 and Irvine 2019; Levy and Madden 1933; Lindquist 1931). Since the pin-point cover data after 2007 were  
124 recorded for each pin separately, the species cover data are readily aggregated into cover data for classes  
125 of species at a higher taxonomic or functional level. At each grid point, the pin may hit different plant  
126 species from the same species class and, in those cases, the hits are only counted as a single hit of the  
127 species class at the grid point.

128 In this study, the species were classified into six groups: *C. vulgaris*, *E. nigra*, and *A. flexuosa*, any other  
129 graminoids, any other herbs or cryptogams. The assumed distribution of pin-point cover data for single  
130 species and the joint distribution for multiple species are outlined in the electronic supplement (Appendix  
131 A).

### 132 **Nitrogen deposition**

133 Nitrogen deposition at each plot was calculated for each year using a spatial atmospheric deposition model  
134 in the period from 2005 to 2014 (Ellermann et al. 2012). The mean site nitrogen deposition ranged from  
135 5.94 kg N ha<sup>-1</sup> year to 21.89 kg N ha<sup>-1</sup> year<sup>-1</sup>, with a mean deposition of 13.80 kg N ha<sup>-1</sup> year<sup>-1</sup>.

136 Anthropogenic nitrogen deposition has reached a maximum in Denmark and is currently decreasing  
137 (Ellermann et al. 2018).

### 138 **Soil pH**

139 Soil pH was measured in randomly selected plots from the uppermost 5 cm of the soil (four samples were  
140 amassed into a single sample). The soils were passed through a 2mm sieve to remove gravel and coarse  
141 plant material, and pH<sub>KCl</sub> was measured on a 1 M KCl-soil paste (1:1). The soil sampling intensity was

142 irregular among sites and years, but typically between one and four plots were sampled from each site at  
143 each time point. When a plot was resampled, the pH at the plot was calculated as the mean of the samples.  
144 In total, 704 independent soil pH values were used in the analysis. The measured soil pH ranged from 2.7 to  
145 6.3, with a mean soil pH of 3.37.

#### 146 **C-N ratio in the soil**

147 Soil C-N ratio was measured in randomly selected plots from the uppermost 5 cm of the soil (four samples  
148 were amassed into a single sample). Total C in each sample was determined by dry combustion and N by  
149 the Kjeldahl method. The soil sampling intensity was irregular among sites and years, but typically between  
150 one and four plots were sampled from each site at each time point. When a plot was resampled, the C-N  
151 ratio at the plot was calculated as the mean of the samples. Measurements outside of the domain [10, 60]  
152 were assessed to be unrealistic, and measurements outside the domain were set to either 10 or 60. The  
153 measured C-N ratio at the site level ranged from 12.2 to 60, with a mean site C-N ratio of 24.8. Moreover,  
154 the average soil C–N ratio on dry heathlands has been observed to decrease in the period from 2004 to  
155 2014 (Strandberg et al. 2018).

156 Since soil C-N ratio is expected to be influenced by the vegetation, e.g. by the amount of slowly  
157 decomposing dwarf shrub litter, soil C-N ratio may be thought of more as an environmental covariable than  
158 an environmental driver.

#### 159 **Soil type**

160 The texture of the topsoil for each site was obtained from a raster based map of Danish soils (Greve et al.  
161 2007). The categorical classification of the soil (JB-nr.) was made on an ordinal scale with decreasing  
162 particle size, 1: coarse sandy soil, 2: fine sand soil, 3: coarse loamy soil, 4: fine loamy soil. There were some  
163 records with other soil types, but because of possible classification errors they were treated as missing  
164 values. The mean soil type was 1.70.

165 **Precipitation**

166 Site-specific precipitation was measured by the average annual precipitation in the period 2001 to 2010,  
167 with a spatial resolution of 10 km (DMI 2014). The annual precipitation ranged from 604 mm to 987 mm,  
168 with a mean precipitation of 824 mm.

169 **Grazing**

170 Land-use was summarized by possible signs of grazing, e.g. the presence of livestock or short vegetation  
171 within fences was recorded by the observer at each plot for each sampling year since 2007 as a binary  
172 variable (sign of grazing = 1, no sign of grazing = 0), i.e. if grazing was 0.5, then this probability may arise by  
173 a number of ways, e.g. if half the plots at the site showed signs of being grazed each year or all plots were  
174 grazed every second year. The mean grazing variable ranged from 0 to 1 among sites, but most sites had no  
175 grazing and the mean grazing intensity at the site level was 0.30. Unfortunately, the grazing variable does  
176 not include information on which animals were used for grazing, stocking densities or grazing duration, and  
177 is therefore a quite imprecise variable that must be interpreted together with general knowledge on the  
178 typically used grazing regime of dry heathlands.

179 **Geographic regions**

180 The 102 dry heathland sites were grouped into six arbitrary geographic regions without using prior  
181 information (Fig. 1). These regions were used to investigate possible latent geographic factors.

182 **Spatio-temporal modelling**

183 To further understand the observed changes in species composition at dry heathlands (Damgaard 2025c),  
184 and the deteriorating conservation status of Danish dry heaths (Nygaard et al. 2020), the observed changes  
185 in pin-point cover data were fitted to site variation in selected abiotic and land-use environmental variables  
186 in a spatio-temporal structural equation model (SEM) (Fig. 2).

187 It was decided to fit the SEM within a Bayesian hierarchical model structure using latent variables to model  
188 the effect of measurement and sampling uncertainties (Fig. 2). This use of a hierarchical model structure is  
189 important, since it has been demonstrated that ignoring measurement and sampling uncertainties may  
190 lead to model and prediction bias (Damgaard 2020; Damgaard and Weiner 2021). Furthermore, it is an  
191 advantage when making ecological predictions to separate measurement and sampling uncertainties from  
192 process uncertainty. The hierarchical SEM approach and the motivation for using it are explained further in  
193 (Damgaard 2025d). The mathematical and statistical details of the spatio-temporal modelling are explained  
194 in the electronic supplement (Appendix B)

195 The procedures for estimating the most important single species cover and change in species cover for all  
196 sampled dry heathland plots since the beginning of the monitoring program in 2004 are explained in the  
197 electronic supplement (Appendix C), and the used code and additional tests of fitting properties etc. may  
198 be found in Appendix D.

## 199 **Results**

200 The selected environmental variables covaried at the 102 dryland heath sites (Fig. S1), which again is  
201 expected to lead to covariance among parameter estimates and affect the fitting properties of the model  
202 negatively. Nevertheless, plots of the mean latent vs. expected logit-transformed cover variables  
203 demonstrated a relatively good fit of the large-scale spatial variation in cover (Fig. 3A; between 48% and  
204 56% of the variation is explained, Table S3), and the model fitted the temporal process of the change in  
205 cover very well (Fig. 3B; > 96% of the variation is explained, Table S3). Furthermore, the Dunn–Smyth  
206 residuals of the marginal observed cover data of the six species classes were approximately normally  
207 distributed (Fig. S2).

208 To prevent possible prediction bias, the different sources of uncertainty, i.e. measurement and sampling  
209 uncertainty when measuring plant cover, nitrogen deposition, soil pH, soil C-N ratio and soil type, as well as

210 the structural uncertainties due to the modelled soil pH, large-scale (among sites) spatial variation and the  
211 temporal processes, were modelled explicitly. The most important source of measurement uncertainty was  
212 the plant cover measurement due to the significant small-scale spatial aggregation of plant species, which  
213 was modelled by the parameter  $\delta$  in the Dirichlet - multinomial mixture distribution. The median estimated  
214 value of  $\delta$  was 0.19 with a relatively narrow credible interval (Table S2). Generally, the estimates of  
215 structural uncertainties of the temporal process were relatively low (Table S2, Fig. 3B), while the estimates  
216 of the structural uncertainties of the large-scale spatial variation were relatively high (Table S2, Fig. 3A).

217 Many of the regression parameters that measure the large-scale spatial and temporal effect of the abiotic  
218 variables on the vegetation were significantly different from zero (Table S2), suggesting that the modelled  
219 environmental and land-use factors have a regulating effect on both the large-scale spatial variation in  
220 cover of the six species classes as well as plant community dynamics in dry heathlands. Soil type had  
221 significant effects on the large-scale spatial variation of all six species classes, whereas the other  
222 explanatory variables had significant effects on at least one species (Fig. 4A, Table S2) and, generally, there  
223 was significant geographic variation among the six assigned Danish regions, where the large-scale cover of  
224 *C. vulgaris* showed a qualitatively different geographic distribution than the other species (Fig. 4A, Table  
225 S2), but without showing a clear pattern (results not shown).

226 The selected environmental drivers, all had significant effects on the temporal variation of all six species  
227 classes, except for grazing and soil type, where three species were significantly affected (Fig. 4B, Table S2).  
228 Notably, relatively high nitrogen deposition was associated with an observed increase in the cover of *A.*  
229 *flexuosa*, and a decrease in the cover of other herbs. Relatively high C/N ratio in the soil was associated  
230 with an observed increase in observed cover of *C. vulgaris*, *E. nigra*, and *A. flexuosa*, and a decrease in the  
231 cover of other graminoids and other herbs. Grazing was associated with a decrease in cover of other  
232 graminoids and other herbs. Finally, relatively high precipitation was associated with a decrease in the  
233 cover of *A. flexuosa* and an increase in cover of cryptogams (Fig. 4B).

234 In this study of dry heathlands, relatively high nitrogen deposition was associated with low soil pH ( $\gamma_N$ , Table  
235 S2), and soil pH was found to be significantly lower on more clayey soils compared to sandy soils ( $\gamma_S$ , Table  
236 S2).

## 237 **Discussion**

### 238 **Environmental drivers and plant community dynamics**

239 Many of the regression parameters that measure the effect of the environmental drivers on the change in  
240 plant species cover were significantly different from zero. These results suggest that the selected modelled  
241 environmental variables have regulating effects on the observed large-scale spatial variation as well as  
242 plant community dynamics in dry heathlands.

243 The cover of the dwarf shrub *C. vulgaris*, which is the characteristic species of dry heathlands, increased in  
244 relatively sandy soils with a relatively high C/N ratio. Notably, the cover of the grass *A. flexuosa* increased  
245 markedly at relatively high nitrogen deposition and low precipitation. These results corroborate earlier  
246 findings, that the ratio between the relative abundances of *C. vulgaris* and *A. flexuosa* at dry heathlands is  
247 associated with a precipitation and soil fertility gradient in Denmark (Damgaard 2015). The cover of other  
248 graminoids, of which *Molinia caerulea* is the most abundant (Table S1), increases with a relatively low C/N  
249 ratio, low grazing pressure, and clayey soils. Overall, these results corroborate the working hypothesis that  
250 the conservation status and resilience of dry heathlands hinge on nutrient-poor sandy soils (Damgaard et  
251 al. 2024). However, note that the current level of nitrogen deposition at dry heathland is within the current  
252 empirical critical load of dry heathlands between 5 and 15 kg N ha<sup>-1</sup> year<sup>-1</sup> (Bobbink et al. 2022), and is  
253 expected to further decrease in the future. The annual precipitation in the future climate in Denmark is  
254 predicted to increase, but with decreasing summer precipitation and longer summer drought periods (DMI  
255 2017). Generally, it is uncertain how this combination of more extreme weather will influence the future  
256 dry heathland vegetation (Olmeda et al. 2020).

257 There were significant effects of grazing on the observed changes in species composition at dry heathlands.  
258 However, except for the negative effects on other graminoids and other herbs, the observed effects on  
259 cover changes were relatively benign. Unfortunately, the Danish ecological surveillance program does not  
260 collect information on stocking rates or seasonality of grazing management, but only on whether grazing  
261 has taken place or not. This lack of resolution prevents a detailed analysis of the effect of the currently  
262 applied grazing on Danish dry heathland vegetation. It has been hypothesized that both grazing  
263 abandonment, which eventually may lead to a succession into a more woody vegetation, and overgrazing,  
264 where grass species are generally favored, may disfavor dwarf shrub species in dry heathlands (Newton et  
265 al. 2009; Olmeda et al. 2020). However, there is still considerable uncertainty as to the effect of grazing on  
266 dry heathland vegetation, and it will be beneficial if future grazing management actions monitor the effects  
267 and regulate stocking rates and seasonality using adaptive management procedures.

268 There were acidifying effects of nitrogen deposition, and soil pH was found to be significantly higher on  
269 sandy soils compared to less sandy soils. The soil acidification effects of nitrogen deposition are due either  
270 to nitrate leaching or removal of base cations from the system by nature management (Williams and  
271 Anderson 1999), and the detrimental effects of this acidification, e.g. reduced biodiversity of higher plants,  
272 have clearly been demonstrated at heathlands in the Netherlands (Bobbink et al. 2022; Vogels et al. 2020),  
273 which generally have received higher nitrogen deposition than Danish heathlands.

274 In conclusion, this study has identified accumulated nitrogen in the soil due to nitrogen deposition and the  
275 associated soil acidification as some of the main threats to the conservation status of dry heathlands. This  
276 conclusion is in concordance with the background information in the EU habitat action plan for dry  
277 heathlands (Olmeda et al. 2020), who further points to the detrimental effects of too little grazing or over-  
278 grazing, and more generally, the abandoning of traditional agricultural practices and the following  
279 encroachments of trees from the border of the heathland areas.

## 280 **Spatial variation**

281 The short-scale spatial aggregation of the three species and the three aggregate classes of other plant  
282 species were modelled by the parameter  $\delta$  in the Dirichlet - multinomial mixture distribution (Appendix A).  
283 The estimated amount of short-scale spatial aggregation significantly increased the measurement  
284 uncertainty of the expected cover data compared to the case of randomly distributed plant species. If this  
285 over-dispersion of the pin-point cover data relative to the random expectation is not taken into account in  
286 the statistical model, then the signal to noise ratio will be severely upward biased and, most likely, will lead  
287 to erroneous conclusions (Damgaard 2013).

288 All species classes had significant regional geographic variation among the six assigned regions (Fig. 1). This  
289 result indicates that some of the large-scale spatial residual variations that could not be explained by the  
290 modelled environmental drivers may be explained by hitherto unexplored factors that differ among the six  
291 regions. In future studies, it will be important to understand the historical causes for the observed large-  
292 scale spatial variation in species abundance, e.g. by collecting and analyzing detailed accounts of site-  
293 specific nature management actions and hydrological data.

## 294 **Uncertainties and application of the model**

295 Generally, a SEM does not allow us to prove hypothesized causal relationships, but it is possible to test  
296 whether specific casual relationships are supported by data. To demonstrate causality, it is necessary to  
297 manipulate the system, e.g. in the form of a manipulated experiment, and observe whether the response  
298 predicted by the SEM actually takes place (Granger 1969; Pearl 2009). Moreover, there is an unknown time  
299 lag between the effect of environmental variables and the change in vegetation cover (e.g. Svenning and  
300 Sandel 2013), so it is uncertain which time period of a changing environmental variables to use in the  
301 model. However, since there typically is a high degree of spatial autocorrelation through time, i.e. sites that  
302 have a relatively high value of an environmental variable at one time period also have a relatively high at a


303 following time period, the model is still able to estimate the relative effects of the environmental variables  
304 among the sites across years.

305 The statistical modelling uncertainty was partitioned into measurement uncertainty and uncertainties due  
306 to the modelled spatial – and temporal processes. The most important source of measurement uncertainty  
307 was the plant cover measurement due to the significant small-scale spatial aggregation of plant species  
308 (see above), but the measurement uncertainty of nitrogen deposition, soil pH, soil C – N ratio, and soil type  
309 was also estimated and, thus, accounted for in the model. Generally, the structural uncertainty of the  
310 temporal processes was relatively small, whereas the large-scale spatial processes entail a higher degree of  
311 structural uncertainty.

312 One of the advantages of partitioning the different types of uncertainties in the SEM is the use of the fitted  
313 SEM for predictive purposes (Damgaard 2022b; Damgaard 2025b), and since the fit of the temporal model  
314 was excellent, it is here suggested that the modeled environmental variables are sufficient for predicting  
315 the *average* plant community dynamics of dry heathlands. This optimistic conclusion is somewhat  
316 surprising in the light of several missing potentially important regulating variables, e.g. previous natural  
317 management actions. Due to the lack of an established causal understanding, caution and humbleness are  
318 required if the fitted model is used for generating local ecological predictions as input to a process of  
319 generating adaptive management plans for specific dry heathlands, since the modelled environmental  
320 variables in this study may be correlated to unknown causal factors of plant community dynamics,  
321 contingent event with large effects, or causal factors where we do not have access to relevant  
322 environmental data. On the other hand, the modelling results provide important information to site  
323 managers on the relative importance of the different environmental factors and management scenarios  
324 (Fig. 5). For example, since nitrogen deposition is invisible it is difficult to assess its effects without a  
325 statistical model and generally the effects of nitrogen deposition tend to be downplayed by site managers.


326 **Figures**

327 Fig. 1. Map of the selected 102 Danish dry heathland sites. The different colors represent a classification of  
328 the different sites into six geographical regions.



329


330 Fig. 2. Outline of hierarchical SEM. The spatial variation in vegetation cover in 2007 is modelled by nitrogen  
 331 deposition (Ndep), soil pH (pH), soil C-N ratio (C/N), soil type and precipitation (Precipit.). The yearly change  
 332 in vegetation cover from 2007 to 2022 (only a single yearly change is shown in the figure) is modelled by all  
 333 the former variables as well as grazing. The black boxes are latent variables and the green ovals are data.  
 334 The black arrows denote large-scale spatial processes, and the red arrows denote temporal processes  
 335 (Appendix B).



336


337 Fig 3. Plots of observed vs. expected logit-transformed cover of the large-scale spatial process (A) and the  
338 temporal (B). The proportions of the variance explained for each species may be found in Table S3. Blue:  
339 *Calluna vulgaris*, yellow: *Empetrum nigra*, green: *Avenella flexuosa*, red: other graminoids, purple: other  
340 herbs, blue: cryptogams.

341 A: large-scale spatial process



342

343 B: temporal process




344

345


346 Fig 4. Standardized regression coefficients of the SEM for the large-scale spatial process (A) and the  
 347 temporal (B).

348 A: large-scale spatial process



349


351 B: temporal process




352

353

354 Fig. 5. Predicted distribution of the cover of *C. vulgaris* and *A. flexuosa* for a specific dry heathland site  
355 (Højsande in West Jutland) after five years under four different scenarios (Damgaard 2022a; 2025b). The  
356 initial cover of *C. vulgaris* and *A. flexuosa* was 0.31 and 0.06, respectively. The scenarios were 1: As now, 2:  
357 reduced N deposition, from 15.5 kg N ha<sup>-1</sup> year<sup>-1</sup> to 8 kg N ha<sup>-1</sup> year<sup>-1</sup> (note that it will require intensive  
358 mangement actions to reduce plant available N in the soil to the level expected at equilibrium under the  
359 reduced N deposition scenario), 3: Higher C/N ratio in soil, from 26 to 40, 4: more grazing 0 to 0.5 (average  
360 of a binary variable where 0 is no grazing) 5: more precipitation, from 987 mm year<sup>-1</sup> to 1100 mm year<sup>-1</sup>.  
361 The other environmental variables were pH in soil: 3.27, soil texture (jb nr.): 1 (coarse sandy soil).



362



363

364

366 **References**

- 367 Aerts R, Berendse F, Caluwe Hd, Schmitz M. 1990. Competition in heathland along an experimental gradient  
368 of nutrient availability. *Oikos*. 57:310-318.
- 369 Aerts R, Heil GW. 1993. *Heathlands. Patterns and processes in a changing environment.*: Kluwer Academic  
370 Publishers.
- 371 Bobbink R, Loran C, Tomassen H. 2022. Review and revision of empirical critical loads of nitrogen for  
372 europe. German Environment Agency.
- 373 Britton A, Marrs R, Pakeman R, Carey P. 2003. The influence of soil-type, drought and nitrogen addition on  
374 interactions between *calluna vulgaris* and *deschampsia flexuosa*: Implications for heathland  
375 regeneration. *Plant Ecol*. 166(1):93-105.
- 376 Damgaard C. 2013. Hierarchical and spatially aggregated plant cover data. *Ecol Inform*. 18:35-39.
- 377 Damgaard C. 2015. Modelling pin-point cover data of complementary vegetation classes. *Ecol Inform*.  
378 30:179-184.
- 379 Damgaard C. 2019. Spatio-temporal structural equation modeling in a hierarchical bayesian framework:  
380 What controls wet heathland vegetation? *Ecosystems*. 22:152-164.
- 381 Damgaard C. 2020. Measurement uncertainty in ecological and environmental models. *Trends Ecol Evol*.  
382 35:871-873.
- 383 Damgaard C. 2022a. Adaptive management plans rooted in quantitative ecological predictions of  
384 ecosystem processes: Putting monitoring data to practical use. *Environmental Conservation*. 49:27-  
385 32.
- 386 Damgaard C. 2022b. Processes and predictions in plant ecological models: Logic and causality. *EcoEvoRxiv*.
- 387 Damgaard C. 2022c. Spatio-temporal modelling of the effect of selected environmental and land-use  
388 factors on acid grassland vegetation. *Journal of Plant Ecology*. 15(2):253-264.
- 389 Damgaard C. 2023. Spatio-temporal modelling of the effect of environmental and land-use factors on  
390 species-rich calcareous grasslands. *Basic and Applied Ecology*. 72:22-29.
- 391 Damgaard C. 2025a. Ecosystem dynamics in wet heathlands: Spatial and temporal effects of environmental  
392 drivers on the vegetation. *Rangeland Ecology & Management*. 100:47-55.
- 393 Damgaard C. 2025b. Local ecological predictions as input to adaptive management of natural plant  
394 communities. *Biol Conserv*. 302:110951.
- 395 Damgaard C. 2025c. Observed vegetation changes in danish dry heathlands since 2004. *Flora*. 327:152728.
- 396 Damgaard C. 2025d. Processes and predictions in ecological models: Logic and causality. *Journal of*  
397 *Forecasting*. 44(5):1658-1665.
- 398 Damgaard C, Bak JL, Strandberg M, Hansen RR. 2024. The resilience of heathland ecosystems: A working  
399 hypothesis. *Acta Oecologica*. 125:104037.
- 400 Damgaard C, Hansen RR, Hui FKC. 2020. Model-based ordination of pin-point cover data: Effect of  
401 management on dry heathland. *Ecol Inform*. 60:101155.
- 402 Damgaard C, Irvine KM. 2019. Using the beta distribution to analyze plant cover data. *J Ecol*. 107:2747-  
403 2759.
- 404 Damgaard C, Strandberg MT, Kristiansen SM, Nielsen KE, Bak JL. 2014. Is *erica tetralix* abundance on wet  
405 heathlands controlled by nitrogen deposition or soil acidification? *Environmental Pollution*. 184:1-  
406 8.
- 407 Damgaard C, Weiner J. 2021. The need for alternative plant species interaction models. *Journal of Plant*  
408 *Ecology*. 14:771-780.

409 De Graaf MCC, Bobbink R, Smits NAC, Van Diggelen R, Roelofs JGM. 2009. Biodiversity, vegetation gradients  
410 and key biogeochemical processes in the heathland landscape. *Biol Conserv.* 142(10):2191-2201.  
411 DMI. 2014. Average annual precipitation in the period 2001 to 2010 with a spatial resolution of 10 km.  
412 Copenhagen: Danmarks Meteorologiske Institut.  
413 Fremtidens klima i danmark. 2017. København: Danmarks Meteorologiske Institut; [accessed].  
414 <https://www.dmi.dk/klima/fremtidens-klima/danmark/>.  
415 Ellermann T, Andersen HV, Bossi B, Christensen J, Løfstrøm P, Monies C, Grundahl L, Geels C. 2012.  
416 Atmosfærisk deposition 2011 – novana. Aarhus: Nationalt Center for Miljø og Energi.  
417 Ellermann T, Nygaard J, Christensen JH, Løfstrøm P, Geels C, Nielsen IE, Poulsen MB, Monies C,  
418 Gyldenkerne S, Brandt J et al. 2018. Nitrogen deposition on danish nature. *Atmosphere.* 9(11).  
419 EU. 1992. Council directive 92/43/eec of 21 may 1992 on the conservation of natural habitats and of wild  
420 fauna and flora. In: Commission E, editor.  
421 EU. 2013. Interpretation manual of european union habitats. Bruxelles: European Commission, DG  
422 Environment, Nature and Biodiversity.  
423 Gimingham C. 1978. Calluna and its associated species: Some aspects of co-existence in communities. *Plant*  
424 *Ecol.* 36(3):179-186.  
425 Gimingham CH. 1960. Biological flora of the british isles. No. 74. Calluna vulgaris (L.) hull. *J Ecol.* 48:455-483.  
426 Gimingham CH. 1988. A reappraisal of cyclical processes in calluna heath. *Vegetatio.* 77(1/3):61-64.  
427 Gimingham CH, Hobbs RJ, Mallik AU. 1981. Community dynamics in relation to management of heathland  
428 vegetation in scotland. *Vegetatio.* 46(1):149-155.  
429 Grace JB, Anderson TM, Olff H, Scheiner SM. 2010. On the specification of structural equation models for  
430 ecological systems. *Ecological Monographs.* 80:67–87.  
431 Granger CWJ. 1969. Investigating causal relations by econometric models and cross-spectral methods.  
432 *Econometrica.* 37(3):424-438.  
433 Greve MH, Greve MB, Bøcher PK, Balstrøm T, Breuning-Madsen H, Krogh L. 2007. Generating a danish  
434 raster-based topsoil property map combining choropleth maps and point information. *Danish*  
435 *Journal of Geography.* 107:1-12.  
436 Levy EB, Madden EA. 1933. The point method of pasture analyses. *New Zealand Journal of Agriculture.*  
437 46:267-279.  
438 Lindquist B. 1931. Den skandinaviska bokskogens biologi. *Svenska Skogsvårdsföreningens Tidskrift.* 3:179-  
439 485.  
440 Løvschal M. 2021. Anthropogenic heathlands: Disturbance ecologies and the social organisation of past  
441 super-resilient landscapes. *Antiquity Project Gallery.* 95:e14.  
442 Løvschal M, Damgaard CF. 2022. Mapping the ecological resilience of atlantic postglacial heathlands. *J Appl*  
443 *Ecol.* n/a(n/a).  
444 Newton AC, Stewart GB, Myers G, Diaz A, Lake S, Bullock JM, Pullin AS. 2009. Impacts of grazing on lowland  
445 heathland in north-west europe. *Biol Conserv.* 142:935-947.  
446 Nielsen KE, Bak JL, Bruus M, Damgaard C, Ejrnæs R, Fredshavn JR, Nygaard B, Skov F, Strandberg B,  
447 Strandberg M. 2012. Naturdata.Dk - danish monitoring program of vegetation and chemical plant  
448 and soil data from non-forested terrestrial habitat types. *Biodiversity & Ecology.* 4:375.  
449 Kontrolovervågning af terrestriske habitatnaturtyper 2004 – 2022. Novana. 2024. Aarhus Universitet: DCE –  
450 Nationalt Center for Miljø og Energi; [accessed]. [www.novana.au.dk](http://www.novana.au.dk).  
451 Nygaard B, Damgaard C, Bladt J, Ejrnæs R. 2020. Fagligt grundlag for vurdering af bevaringsstatus for  
452 terrestriske naturtyper. Artikel 17-rapporteringen 2019. Aarhus Universitet, DCE – Nationalt Center  
453 for Miljø og Energi.  
454 Olmeda C, ŠeffEROVÁ V, Underwood E, Millan L, Gil T, Naumann S. 2020. Eu action plan to maintain and  
455 restore to favourable conservation status the habitat type 4030 european dry heaths. European  
456 Commission.

457 Ovaskainen O, Roy DB, Fox R, Anderson BJ. 2016. Uncovering hidden spatial structure in species  
458 communities with spatially explicit joint species distribution models. *Methods in Ecology and*  
459 *Evolution*. 7(4):428-436.

460 Pearl J. 2009. *Causality. Models reasoning, and inferences*. 2, editor. Cambridge: Cambridge University  
461 Press.

462 Strandberg M, Nielsen KE, Damgaard C. 2018. Habitat monitoring reveals decreasing morlayer c:N ratios in  
463 danish heathlands. *Ecological Indicators*. 89:538-542.

464 Svenning J-C, Sandel B. 2013. Disequilibrium vegetation dynamics under future climate change. *Am J Bot*.  
465 100:1–21.

466 Tybirk K, Nilsson M-C, Michelsen A, Kristensen HL, Shevtsova A, Tune Strandberg M, Johansson M, Nielsen  
467 KE, Riis-Nielsen T, Strandberg B et al. 2000. Nordic *empetrum* dominated ecosystems:  
468 Function and susceptibility to environmental changes. *AMBIO: A Journal of the Human*  
469 *Environment*. 29(2):90-97, 98.

470 Usher MB, Thompson DBA. 1993. Variation in the upland heathlands of great britain: Conservation  
471 importance. *Biol Conserv*. 66(1):69-81.

472 Vogels JJ, Weijters MJ, Bobbink R, Bijlsma R-J, Lamers LPM, Verberk WCEP, Siepel H. 2020. Barriers to  
473 restoration: Soil acidity and phosphorus limitation constrain recovery of heathland plant  
474 communities after sod cutting. *Applied Vegetation Science*. 23(1):94-106.

475 Watt AS. 1947. Pattern and process in the plant community. *J Ecol*. 35:1-22.

476 Williams BL, Anderson HA. 1999. The role of plant and soil processes in determining the fate of atmospheric  
477 nitrogen. In: Langan SJ, editor. *The impact of nitrogen deposition on natural and semi-natural*  
478 *ecosystems*. Dordrecht: Kluwer.

479

480

481 **Electronic supplements**

482 *The following electronic supplements may be downloaded from <https://osf.io/4ezan/>*

483 Table S1 – Cover and change in cover of single species at Danish dry heathlands

484 Table S2 - Marginal distribution of parameters

485 Table S3 – Proportion of variance explained

486 Fig. S1 – Pairwise scatter plot of variables

487 Fig. S2 - Dunn–Smyth residuals

488 Appendix A - Distribution of pin-point plant cover data

489 Appendix B - Spatio-temporal modelling

490 Appendix C - Estimating species cover and change in species cover

491 Appendix D – Mathematica notebook (a free reader may be downloaded from

492 <https://www.wolfram.com/player/>)

493