

Born this way: individuality is seeded before birth and robust to environmental stress

Short title: Individuality is seeded before birth, robust to environmental stress

James H. Gallagher¹, Ammon D. Perkes¹, Chia-Chen Chang^{1,2}, Emma S. Chirila¹, Karen Kacevas¹, Kate L. Laskowski¹

¹ Department of Evolution and Ecology, University of California Davis, Davis CA, 95616

² Department of Biological Sciences, National University of Singapore, Singapore, 119077

Corresponding author: James H. Gallagher (jhgallagher@ucdavis.edu)

Keywords: Development, animal personality, variance partitioning, developmental plasticity, individual variation

Abstract

Behavioral individuality, or consistent individual differences in behavior, are pervasive across the animal world and have major ecological and evolutionary consequences. Nevertheless, we still have a limited understanding of what drives individuality and how it emerges during ontogeny. Here, we subjected clonal individuals to a ubiquitous yet critical environmental challenge—the threat of predation—to disentangle the developmental mechanisms of individuality. Under such a salient environmental stressor, among-individual differences may collapse or expand depending on whether there is a single or multiple optimal strategies, demonstrating that individuality itself is a developmentally plastic trait. If, however, individuality is determined before birth, we may expect it to be resistant to environmental influences. We continuously tracked the behavior of genetically identical fish (Amazon mollies, *Poecilia formosa*), reared with or without predation stress, all day, every day, from birth through their first month of life, providing unprecedented insight into the trajectories of behavioral development in response to this key environmental cue. Predation shifted mean-level behaviors, with predator-exposed individuals swimming more slowly and spending more time near their refuges. However, the magnitude of individuality (as evidenced by repeatability) increased similarly over development in both treatments, indicating that individuality crystallizes robustly over time, even under stress and in a vacuum of genetic variation. Predator-reared fish also exhibited greater within-individual variation in refuge use, suggesting decreased behavioral predictability or disrupted developmental canalization in response to stress. Surprisingly, maternal identity, but not maternal behavior, was the strongest contributor to variance in swimming velocity (accounting for two thirds of variation), pointing to maternal effects as a key pre-birth source of behavioral variation. Variance in refuge use however was only negligibly explained by maternal identity, highlighting that fundamental behaviors may have very different developmental mechanisms. Collectively, our results show that individuality persists despite environmental stress and is likely seeded before birth through non-genetic factors. Even in the face of a shared environmental challenge, the behavioral trajectories of individuals are unique.

Introduction

Understanding what shapes individuality has been a long-standing, fundamental question in the field of biology that has permeated many aspects of culture, from philosophy to science fiction. Individuality—consistent differences in behavior among individuals—is ubiquitous across the animal kingdom^{1–5}, can have important fitness implications for individuals^{6–11}, and can shape the evolutionary trajectories of populations^{12–14}, even driving speciation^{15,16}. Where do individual differences in behavior come from? While this question has been

historically posed as the “nature vs nurture” debate, this framing is now widely regarded as a false dichotomy, as both genetic and environmental variation will almost always interact to influence an individual's behavior. Considerable effort has been spent controlling for these two factors in order to pinpoint the drivers of behavioral individuality. Nevertheless, there is a growing body of work indicating that even when genetic and environmental variation is minimized, individuals still exhibit consistent differences in their behaviors^{17–20}, even at birth^{21,22}. Some of this variation may be due to variation in pre-birth influences, such as maternal effects or developmental stochasticity, which may play an underappreciated role in generating variation among individuals^{5,20,23–28}. This type of non-genetic variation may help set the seed around which further behavioral individuality crystallizes. Regardless of where such initial differences arise, the continued persistence of individuality despite our best efforts to minimize it suggests that we still lack a full understanding of the key mechanisms underlying its emergence. To identify these mechanisms, we must consider not only *what* shapes behavior, but also *how* behavioral differences emerge over the course of life^{12,29}.

Development is an iterative, path-dependent process, where early experiences may lead to lasting behavioral differences^{30,31}. Thus, tracking how individuals respond to environmental challenges during development can reveal the processes that shape their behaviors. One predominant environmental feature that individuals must contend with is that of risk, most notably in the form of predation. Because most animals are subject to predation risk, their behavioral responses to this risk can heavily impact their fitness. Across taxa, predation stress during development has been shown to have profound effects on individual behavior, brain function, and personality^{32–43}, making predation a salient environmental candidate for shaping behavioral individuality. However, we still do not understand how predation shapes the *emergence* of individuality from birth throughout ontogeny. This is an important knowledge gap because exposure to predation stress may impact patterns of individual behavioral variation in different ways. On one hand, predation stress may collapse variation attributed to among-

individual differences, making individuals behave more similar to each other, if there is an optimal behavioral strategy for survival in a predator-rich environment ⁴⁴. Alternatively, predation stress could instead expand among-individual differences ³³, as there may be many strategies to mitigate risk in the face of predation. A final possibility is that individual differences may not be impacted by predation stress during development, implying a predominant role for pre-birth factors in driving behavioral individuality. Disentangling these possibilities requires closely tracking individual behavior from birth throughout development, across different developmental environments.

Studying the emergence of individuality over development can be very challenging due to the presence of confounding genetic variation in animals, as well as an inability to reliably measure the behavior of individuals from birth in standardized conditions throughout ontogeny, either due to features of the study species (e.g., some species require parental care introducing uncontrolled variation in experience among offspring) or technological limitations in following individuals at such early points in their lives. To fully understand the processes that shape behavioral variation, we must be able to solve these challenges and intimately track the details of the timing and rate at which behaviors emerge in individuals ⁴⁵.

Here, we used an ideal animal study system in conjunction with high-resolution tracking to bypass historical experimental limitations and deeply study the emergence of behavioral individuality over ontogeny. The Amazon molly (*Poecilia formosa*) is an all-female freshwater fish that reproduces via gynogenesis, resulting in individuals that are genetically identical to each other and their mothers ^{46–49}, allowing us to minimize the effects of confounding (among-individual) genetic variation, analogous to twin studies in humans ⁵⁰. They are also live-bearing without any parental care, so individuals can be isolated from birth with minimal consequence to standardize life experience from day one. We combined this optimal study species with custom tracking technology (using cameras connected to individual Raspberry Pi computers ^{21,51}) to record and extract behaviors from individuals (N=107) at high temporal resolution (1s resolution

for 12 hours every day) over the first 28 days of their lives. These fish were placed in one of two environmental conditions—with or without exposure to predator cues—to determine how stress shapes behavior and the magnitude of individuality across development. We also standardized the environments that these individuals' mothers experienced in the week prior to breeding (with none of them facing predation stress at any time) and used the same tracking approach to measure the behavior of all mothers before they gave birth (N=13, for seven days) to test whether maternal behavior and/or identity explain offspring behavior (Fig 1). This resulted in over 30,000 hours of data-rich recordings (>100 million data points) that we then quantified using a convolutional neural network-based pipeline to track the position of each individual, all day, every day. Using these data we extracted ecologically relevant behavioral metrics, such as swimming velocity and time spent near a refuge in the tank, that are likely to be affected by predation threat^{52–56}.




Fig 1. Experimental design. A) Adult Amazon mollies were each placed in identical individual tanks in our custom tracking system and recorded using cameras attached to Raspberry Pi computers for 12 hours each day over one week. B) Fish were then placed in individual tanks with a male Atlantic molly to initiate reproduction. C) Offspring born from the breeding tanks were split evenly by brood and placed into individual tanks in one of two closed-water tracking systems—with and without water containing predator cues—and continuously recorded for four weeks.

In our study, we asked a series of questions to disentangle the developmental drivers of behavioral variation. First, we asked how average behavior shifts over ontogeny in the face of a significant environmental challenge (threat of predation). Predator-exposed fish may adjust fitness-relevant behaviors, such as swimming velocity and time spent near their shelter, and/or show different behavioral developmental trajectories from control fish. Second, we investigated which factors (developmental environment, maternal effects, and remaining non-genetic among-individual variation) best explain behavioral variance. By determining what shapes behavioral variation, we will also learn whether different behaviors share the same developmental mechanisms. Finally, and of prime interest here, we tested whether and how predation alters the magnitude of individuality—quantified as the relative proportion of among-individual variance—across developmental time compared to animals that were reared under benign conditions. This will reveal whether individuality itself is developmentally plastic or determined before birth through non-genetic factors. Our powerful study system, large individual sample sizes, and high temporal resolution tracking provides a uniquely comprehensive understanding of the fundamental yet elusive processes that generate individuality.

Results

Predation stress during development impacts mean-level behavior

We used generalized linear mixed models to test for behavioral differences between treatments by including treatment, time, and their interaction as our fixed effects of interest. We did not find evidence for significant treatment-by-time interactions in either swimming velocity (treatment x time: post.mean = 0.002 [95% CI: -0.01, 0.01]; Fig 2A) or refuge use (treatment x time: post.mean = 0.01 [95% CI: -0.004, 0.02]; Fig 2B), indicating that behaviors did not change differently over ontogeny based on environment. However, we did find significant main effects of treatment for both swimming velocity (post.mean = -0.18 scaled pixels per second [95% CI:

-0.34, -0.02]) and refuge use (post.mean = 0.24 scaled proportion [95% CI: 0.03, 0.43]), with predator-reared fish moving more slowly and spending more time near their refuge (Fig 2). For velocity, these differences between the two treatments appeared small, but were present on the first day of life (post.mean = 1.21 scaled pps [95% CI: 0.74, 1.75]). Unlike velocity, there were no significant differences in refuge use on day one (post.mean = 0.22 scaled proportion [95% CI: -0.01, 0.48]). Neither velocity nor refuge use were predicted by body size (Table S1), and body size was not predicted by rearing environment (length: post.mean = 0.06 [95% CI: -1.71, 1.67]; width: post.mean = -0.16 [95% CI: -0.45, 0.10]).




Fig 2. Average-level behaviors (velocity (A), in scaled pixels per second, and refuge use (B), in proportion scaled) of individuals over developmental time, reared with (orange) and without (blue) predation threat. Thin lines represent the behaviors of individuals over time, while thick lines represent mean behaviors of individuals in a given developmental treatment. Note that for median velocity (A), a single individual's day one data is not shown (but are included in analysis) due to its unusually high value (10.12 scaled pps) that would hinder visual clarity of the plot if included.

Different behaviors have different sources of variance

We used a combination of forward model selection and variance partitioning to determine how developmental environment (treatment), among-individual differences, and/or maternal identity explained the behavioral variance. Using model selection, we found that shared variance components across treatments were best supported for both velocity and refuge use, indicating that there were not significant differences in the magnitude of among-individual and among-mother variance between the treatments (see supplemental information). For velocity, maternal identity explained the majority of the total variance across all of development ($ICC = 0.659 [0.368, 0.893]$), while differences among-individuals explained a much lower amount ($ICC = 0.059 [0.020, 0.113]$; Fig 3A). However, for refuge use, patterns were reversed, with among-individual differences ($ICC = 0.267 [0.214, 0.323]$) explaining much more of the variance than maternal identity ($ICC = 0.021 [0.001, 0.081]$), which was minimal (Fig 3B).

Because maternal identity strongly contributed to variation in velocity (Fig 3C; but not refuge use, Fig 3D), we tested whether these effects could be explained by maternal behavior. For both velocity and refuge use, maternal behavior did not predict offspring behavior (velocity: $\text{post.mean} = 0.005 [95\% \text{ CI: } -0.02, 0.03]$; refuge use: $\text{post.mean} = 0.02, [95\% \text{ CI: } -0.128, 1.41]$), nor did maternal identity predict body size (length: $\text{post.mean} = 0.546 [95\% \text{ CI: } <0.001, 3.926]$; width: $\text{post.mean} = 0.004 [95\% \text{ CI: } <0.001, 0.021]$; Figs 3E-F).




Fig 3. Sources of variation across behaviors (left = velocity, right = refuge use). For velocity (A), most of the variance was explained by maternal identity, providing strong evidence of maternal effects. For refuge use (B), maternal identity contributed minimally to variance, with variation among individuals explaining a much larger share, and variation within individuals (i.e., residual) explaining the largest amount. C-D show the relationship between control and predator siblings from each mother (units = scaled behavior). Grey dots indicate the mean behavioral values of all offspring from a given mother. Error bars show the standard error of means for control (blue) and predator (orange) offspring of that mother. Maternal identity is a much stronger predictor of velocity than refuge use, as indicated by the lack of overlap between offspring behavior from different mothers (error bars). E-F show the lack of association between maternal behavior and offspring behavior, for both velocity (E) and refuge use (F). Boxplots represent individual mothers, while blue and orange dots show mean behaviors over developmental time for each individual offspring of that mother, reared with (orange) or without (blue) exposure to predator cues.

Individuality persists despite predation stress during development

We used generalized linear mixed models to explore patterns of individual behavioral variation, estimating variance components at each day (among- and within-individual) to then calculate repeatability of behavior over developmental time, our proxy for individuality, for each treatment separately, while controlling for among-mother variance. We found that for both velocity and refuge use, individuality persisted, regardless of environmental treatment (Fig 4A-B). In both treatments, individuals began life with low repeatability, which gradually increased over the next four weeks, resulting in individuals with fairly repeatable behaviors by the time they were one month old. Refuge use was consistently more repeatable than velocity. Notably, repeatability and among-individual variance did not differ by developmental treatment, as indicated by overlapping confidence intervals. However, for refuge use, within-individual variance was greater in predator-reared individuals (Fig 4D), indicating that predator-reared fish are less predictable in this behavior, within any given day compared to control-reared fish.




Fig 4. Extracted variance components over developmental time, colored by treatment. Repeatability—the proportion of variance attributed to variation between individuals (the metric of individuality)—increased over the first month of life for velocity (A) and refuge use (B), and this pattern did not differ between treatments, indicating the persistence of individuality in the face of predation stress. C-D show variance components as posterior medians, separated into among- and within-individual (i.e., residual) components. For velocity (C), both among- and within-individual variance did not change over time or between treatments, aside from day one of life, where within-individual variance was much higher, particularly in the control group. For

refuge use (D), among-individual variance also did not differ by treatment. Within-individual variance was highest during the first week of life, followed by a sharp decrease and plateau for the rest of development. However, within-individual variance was consistently greater in predator-reared fish over developmental time. Error bars on all plots indicate 95% confidence intervals.

Discussion

We found that both developmental environment (Fig 2) and maternal identity (Fig 3) are sufficient to generate behavioral variation, even in the absence of genetic differences. However, the relative contribution of each source differed dramatically between behaviors, indicating that distinct developmental mechanisms underlie each trait. For example, most of the variance in velocity was explained by maternal identity, while most of the variance in refuge use was explained by variation within individuals, with among-individual variation playing a notable role as well. Perhaps surprisingly, environmental stress did not disrupt the emergence of behavioral individuality (repeatable differences among individuals) across development, suggesting that individuality is seeded before birth rather than developmentally plastic in response to an individual's environment (Fig 4A-B). However, individuals reared under stress did show greater variability (within-individual variation) in their refuge use behavior, indicating that stress does increase intraindividual variability or disrupt developmental canalization (Fig 4D). Here, we discuss the biological implications of how developmental environment shapes behavior and individuality, the role of other factors—notably maternal effects—in explaining this variation, and the contrasting mechanisms underlying different behaviors. Finally, we discuss possible limitations, place our findings in the context of other recent studies, and propose directions for future research.

Exposure to predation during development impacted mean-level behavior, with predator-reared individuals moving more slowly and spending more time near their refuges, showing that our animals responded to the cues of predation stress as we expected (Fig 2). However, the

timing of when these differences first emerged varied by behavior. For velocity, behavioral differences between treatments were immediately apparent on day one (Fig 2A); for refuge use, these differences were not present at the start of life, instead arising over the first week of development (Fig 2B). Considering the clonal nature of Amazon mollies, this reflects pure behavioral plasticity (that may be adaptive), suggesting that stress alone can alter the way in which behaviors emerge over ontogeny. Our comprehensive approach to investigating the effects of predation on the emergence of behavior supports previous work in various non-clonal taxa showing that exposure to predation during development shapes behavior^{32–34,36,37,41}.

In both environmental conditions, repeatability (the magnitude of individuality) increased over ontogeny, suggesting that individuality crystallizes over time potentially through self-reinforcing behavioral patterns or internal feedback loops^{21,57}. Perhaps surprisingly, this strengthening of individuality over time was not impacted by predation, with repeatability increasing similarly over development in both treatments (Fig 4A-B). This indicates that the seeds for individuality may be set before birth, and suggests that there may be many optimal behavioral strategies to overcome environmental challenges. While our results are contrary to other studies where exposure to predation either suppressed or expanded among-individual variance (e.g.,^{37–39}), our work measured behavior continuously over development, rather than at set intervals which can only provide snapshots of behavior. This temporally rich approach allowed us to estimate individual variance components daily, improving the accuracy of our repeatability calculations and revealing additional insights into how behavioral variation emerges throughout ontogeny⁴⁵. One example of this can be seen in our estimates of behavioral predictability (i.e., within-individual variance or intraindividual variability^{58,59}). Although among-individual variance did not differ between experimental treatments, we did uncover temporal and treatment-level changes in within-individual variation by allowing residual variance to fluctuate over time in our model (i.e., heterogeneous residual variance; see methods for more information on our statistical approach). This revealed that while individuals in both treatments behaved

more predictably as they aged, predator-reared fish showed lower predictability (i.e., greater intraindividual variability) in refuge use within a given day throughout development (Fig 4D). Lower predictability in predator-reared individuals may reflect either an adaptive increase in flexible use of the refuge throughout the day as individuals assess immediate risk, or a disruption of developmental canalization, in response to risk (see more on this below). We encourage others to account for heterogeneous residual variance over time, as this not only provides useful insights into the often underappreciated yet important within-individual variation^{58,60–63}, but also directly impacts the calculation of repeatability (which is dependent on correctly estimating residual variance at a given time point).

If individuality remains robust throughout environmental stress during development, what then drives it? We found that despite attempting to minimize potential maternal effects, for activity-related behavior (i.e., velocity), maternal identity predicted offspring behavior more so than any other component (including variance attributed to individual identity; Fig 3A). This study was not designed to fully test for the effects of differential maternal experience on offspring behavior because the mothers used in this experiment were all reared in similar conditions. Nevertheless, our results suggest that maternal effects do play a notable role in shaping behavior, and may be a key candidate for setting the seeds of individuality before birth. Despite this strong evidence of maternal effects (i.e., the similarity of clonal siblings to each other), we did not see any association between maternal and offspring behavior (Fig 3E-F), indicating that the influence of mothers was not mediated by what they did, but rather by who they were (although maternal behavior was collected when fish were adults rather than during early development). This points to maternal effects with a non-behavioral basis, such as maternal provisioning, physiological state, and/or inherited epigenetic variation through changes in DNA methylation or maternal hormone transfer^{21,23,24,64,65}, as mechanisms through which behavioral variation emerges, even in the absence of genetic or environmental variation.

Individuality may begin with cryptic differences among mothers that bias developmental trajectories from the very start of life.

Interestingly, our two representative behaviors, velocity and refuge use, emerged differently over ontogeny and appeared driven by different factors, suggesting contrasting underlying mechanisms. Variation in velocity was overwhelmingly explained by maternal identity (Fig 3A), while variation in refuge use was not explained by maternal identity at all (Fig 3B). As for developmental environment, differences between treatments were more apparent from the beginning of life for velocity, but not for refuge use, which emerged over the first week of life. Additionally, predation stress increased behavioral variability (within-individual variance) throughout development for refuge use, but not velocity. These differences in how behavioral traits arise during ontogeny and what drives their variation suggests that major suites of ecologically relevant behaviors can have different mechanisms of emergence. Behaviors relating to space or refuge use may be far more context-dependent than swimming speed. For example, while individual humans have highly consistent walking speeds⁶⁶, decisions about where and when to move, such as crossing a street, vary much more within individuals based on context (e.g., how much traffic is present). Even fully understanding the developmental processes underlying one suite of behaviors may not inform others, and we should be careful to not generalize across traits when studying the mechanisms of animal behaviors.

Although we attempted to control for as many variables as possible, there remain potential limitations. Our study focused on isolating the effects of the environment on behavioral development, so we minimized genetic variation and carefully controlled experimental environments. However, it is impossible to eliminate all factors that could contribute to variation in behavior. Small genetic differences between individuals could still arise even among individuals within the same clonal lineage (e.g., either germ-line or somatic mutations⁶⁷), and individual experiences, particularly of the mothers, are impossible to fully standardize from the first second of life.

We end this discussion by placing our results in the context of other studies, proposing future work that could address new questions that arose from our findings. First, our repeatability estimates differed somewhat from those in other recent work in Amazon mollies. Repeatability of both behaviors increased over ontogeny, with refuge use reaching higher levels by the end of the first month of life (>0.5). However, for velocity, repeatability remained fairly low (under 0.2) until week four. Such values differ from previous work in this species, where repeatability of velocity was greater than 0.4 for the entire first month of development (measured only in control conditions;²¹, and instead more closely matched levels found in other taxa (e.g., crickets) when using highly inbred lines⁶⁸. One likely reason for the discrepancy between studies on Amazons is that here we had a larger number of both individuals and mothers, allowing us to more accurately account for variance that was attributed to differences between mothers—of which there was a tremendous amount for velocity (Fig 3A). While repeatability remained relatively low for velocity over the first three weeks, there was a greater rate of increase over the final week of the experiment, a pattern that mirrors results from the aforementioned work²¹. If that trajectory were to continue, repeatability of velocity in our fish may have reached much higher levels over time (as was the case in Laskowski et al. 2022).

The persistence of variation among individual Amazon mollies, despite their lack of genetic differences, raises additional questions about the role that genetic variation actually plays in generating individuality. One hallmark of sexual reproduction is its ability to generate variation among individuals. Yet here (and in other recent work^{21,22}) we found a striking degree of among-individual variation even when genetic differences were minimized. Future work could reintroduce genetic variation in a fully standardized environment to better understand the contribution of allelic variation (or lack thereof) to the onset of individuality (e.g., comparing clonal Amazon mollies to their parental species, which naturally contain more genetic variation via sexual reproduction). For example, a recent study compared repeatability of behavior between inbred and outbred lines of field crickets, and found reduced repeatability in inbred

lines⁶⁸. Whether this pattern holds true in other taxa—and in truly clonal (i.e., parthenogenetic, gynogenetic) species, including the Amazon molly—would make for interesting and important future work. Other ecologically important environmental factors, such as temperature^{69–71} or resource availability⁷², could also be systematically manipulated to further assess their impact on individuality. Given our evidence for maternal effects without behavioral transmission, a key next step is to explore how maternal environment shapes offspring behavior using a multigenerational design that investigates the impacts of maternal experience on mechanisms that canalize individual behavioral trajectories.

In this study, we disentangled the developmental drivers of individuality to understand the mechanisms that make individuals unique. Despite facing a strong environmental stressor that affected mean-level behaviors, individuality persisted, suggesting that development does not result in a single dominant phenotype, but rather an acceptable range. Even though mean values may be shifted up or down, the magnitude of variation among individuals does not change—behavioral individuality appears to be a robust and intrinsic feature of animal life.

Methods

Maternal behavior and breeding

To track the behavior of potential mothers of experimental fish, we placed adult Amazon mollies in individual observation tanks and monitored them for one week (Fig 1A). All adult fish were of similar age, from the same clonal lineage, and were reared in the same conditions (social housing with similar densities, food and environmental regimes, etc.). Observation tanks were made of white acrylic (Perspex) and had identical environments (dimensions = 15x19x15 cm), each containing white gravel along two of the four sides of the tank, as well as a standpipe covered in a sponge filter that fish also use as a refuge. All fish experienced the same feeding schedule (fed a standard amount of Tetra-Min flake fish food and brine shrimp twice per day),

daily light cycle (12:12 light:dark), water temperature (26 ± 2 °C), and water depth (10 cm). The observation tank area was also surrounded by white curtains to reduce potential disturbance from external stimuli. Above each set of four tanks was a mounted camera attached to a Raspberry Pi 4B computer that recorded individual behaviors (see “Tracking behavior” below).

After one week, we transferred each adult fish to an individual 5-gallon glass breeding tank containing an artificial plant for shelter and a male Atlantic molly (*Poecilia mexicana*) to initiate breeding (Fig 1B). Amazon mollies are gynogenetic, meaning that while all individuals are genetically identical to their mother, they still require mating with a male from a closely related species to trigger embryogenesis (his genetic material is unused in the offspring^{46,48}). One week later, we removed males from the tanks. Throughout the breeding period, all fish experienced the same conditions (including temperature, light cycle, and food access). This resulted in 13 mothers that gave birth to offspring used in the rest of the study.

Offspring experimental setup

When fish were born in the breeding tanks, we transferred them that day (the first day of their lives) to their observation I tanks in the tracking system (Fig 1C). We transferred all newborn fish in a standardized way, by first herding them by net to a small container of water, then moving them to their respective individual tanks, thus limiting air exposure. We took individuals from each brood and distributed them evenly between observation tanks that were part of two closed circulating water systems: one containing a live electric yellow lab cichlid (*Labidochromis caeruleus*), and one without (control). Each closed water system was connected to a sump tank below the observation tanks; in the predator system, this sump tank contained the cichlid, which was fed whole Tetra-Min flake food, blood worms, and brine shrimp daily. All tanks in a given filtration system shared water, so fish in the predator system were consistently receiving chemical (but not visual) cues from the cichlid (cues from this cichlid elicit anti-

predator behavior in adult Amazon mollies⁷³). To deliver alarm cues to experimental fish, which are released from the skin following mechanical damage^{74–78}, we fed the cichlid an anesthetized small black molly (*Poecilia sphenops*) twice per week—once between Monday and Wednesday, and once between Thursday and Friday—on random days during those two windows and at random times of day to reduce habituation. Experimental conditions for offspring were identical to those used for tracking the behavior of the adults (mothers), except fish were fed powdered instead of whole Tetra-Min flake fish food. Tanks contained aquarium gravel along the walls to help promote natural foraging behaviors, and each group of tanks was surrounded by curtains to reduce outside disturbance. There were always multiple individuals in a given shared water system, so fish did not perceive complete social isolation^{21,22}, which can lead to altered behaviors in some fish⁷⁹. We allowed fish to develop in their observation tanks undisturbed for four weeks, resulting in data from 107 individuals.

Tracking behavior

For efficiency and reproducibility, all videos were processed within one of two automated pipelines (see supplemental code;⁸⁰), specialized either for adults or for juveniles. All videos were first cropped to exclude adjacent tanks using ffmpeg (version 6.1.1, FFmpeg Developers), based on manually labeled points. We also used labeled center-points to assign the (up to) four tracks in each video to specific individuals. For tracking adults, we used a neural network (efficientNet b3⁸¹), trained and implemented using SLEAP⁸². For tracking babies, we similarly began with a neural network (LEAP⁸³) using SLEAP⁸², which was generally accurate, but struggled when small fish were above gravel. To address this, we also used a custom tracker based on a mixture of gaussians background subtractor⁸⁴, implemented using openCV⁸⁵ in python. Because this takes advantage of frame-to-frame differences, it was successful tracking fish over gravel. Both sets of tracks were filtered to remove extremely high-density pixels (which

likely represent false detections of stationary objects). To combine these two streams, we used a custom implementation of the Viterbi algorithm⁸⁶, which seeks to estimate the most likely path of some unknown and hidden value (in this case, the location of fish) based on probabilistic observed values (the two parallel detections of fish). From these post-processed tracks we calculated the median velocity, proportion of frames “active” (defined as moving > 5 pixels/second), median velocity when “active,” and proportion of frames detected in the corner with the standpipe (which acts as a refuge). Videos were manually inspected by an expert observer (JHG) for major issues that would result in inaccurate tracking (i.e., missing or moved fish, camera movement). We manually scored 1-minute segments from a randomly selected day within the first and last week for 6 groups (a total of 18,000 frames). The mean false detection rate was only 6.2% (+/- .03% SEM), suggesting for >90% of frames, the fish was either accurately detected or not visible (e.g., under the refuge).

Measuring body size

To estimate individual body size of offspring at the end of the experiment, we used a custom python script to extract crops of each individual (based on their detected position) on day 27 and manually labeled the head, tail-peduncle, and body width (at its widest point) using Supervisely, a web-based computer vision platform⁸⁷. The identity and treatment of fish in these crops were blinded during labeling.

Statistical analysis

All statistical analysis was conducted using R accessed through R Studio (R Studio version 2023.12.1, R version 4.3.3). We selected which behaviors we would investigate by eliminating highly correlated behaviors. This resulted in two behaviors, median velocity (swimming speed) and proportion of time spent near the standpipe (refuge use), that we used

for analysis. In order to understand which factors shape mean-level behaviors and behavioral variance, we first ran Bayesian bivariate mixed models using the MCMCglmm package⁸³, separately for each behavior (velocity and refuge use) where our bivariate response was the same behavior in each treatment. We used a bivariate model first to determine whether our random effects should be estimated independently by treatment. All models included treatment (with or without predator exposure), day of life, hour, and the interaction between treatment and day as fixed effects, and individual and maternal ID as random effects because these were both of primary interest to our research question. We first conducted model selection to determine any additional random effects structure of our models (i.e. inclusion of random intercepts and slopes for individual and/or mother), choosing the least complex model within 2 of the lowest DIC value. To account for differences in behavior due to time of year (given the unpredictability of births, fish were placed into the tanks as they were born which staggered throughout the year), we next compared our models with and without a time of year variable as a fixed effect. We performed hierarchical clustering using the stats package to translate the time of year variable into batches; this revealed four primary clusters, which we used for our time of year batch variable. The best supported models for both behaviors included random effects (individual and maternal IDs) pooled between treatments, as opposed to independent random effects that were estimated uniquely for each treatment. This suggested that variance components did not differ by treatment, and therefore a univariate model was most appropriate to answer our questions moving forward.

We used the same model selection process above to determine the best structure for the univariate model. We also compared our models to one that allowed for heterogeneous residual variance over time, as this could impact repeatability calculations and reveal cryptic yet relevant information about the effects of the developmental environment on behavioral variability. For both behaviors, heterogeneous residual variance improved the models' fit. Thus, the final model for velocity included treatment, day, their interaction, hour, and time of year as

fixed effects, random intercepts for individual and maternal ID (family = gaussian), and daily heterogeneous residual variation. For refuge use, the final model included treatment, day, their interaction, and hour as fixed effects (time of year did not improve the model fit), random intercepts for individual and maternal ID (family = gaussian), and daily heterogeneous residual variation. All models were validated by ensuring variables of interest had low autocorrelation values (<0.1) and high effective sample sizes (>1500), and by visually inspecting trace and density plots. To aid with statistical interpretation of differences between treatment groups (shown in Supplemental Information), we calculated probability of direction (pd) using the package bayestestR (a pd value > 0.975 is equivalent to a frequentist p -value of < 0.05 ^{88,89}).

To test for the effects of body size on behavior, we ran the full models described above with and without the inclusion of individual length and width as fixed effects. Because body size data was collected at the end of the experiment (day 27), any individuals that were not recorded through the end of the experiment could not have their body size measured, so we ran the models on a subset of the data that only included individuals with body size measurements ($N = 90$ individuals).

To test for differences in mean behaviors between treatments on day one of life, we ran the same models on a subset of the data (day one) but excluded day as a fixed effect. We again conducted model selection to determine which other fixed effects should be included for the day one comparison (time of year and hour). Our final model for velocity included treatment and hour as fixed effects, and random intercepts for individual and maternal ID (family = gaussian), and for refuge use included treatment as a fixed effect, and random intercepts for individual and maternal ID (family = gaussian).

To estimate the proportion of variance explained overall (i.e. over the entire observation period and across both treatments) by different sources, we calculated marginalized repeatability (i.e., intraclass correlation) by dividing the variance attributed to each variable (individual ID, maternal ID, and residual) by the total variance^{90,91}. For example, to calculate

maternal ICC, we used the following equation: maternal ID variance / (maternal ID variance + individual ID variance + residual variance; see supplementary information for further details). To determine whether maternal behavior predicted offspring behavior, we ran our full mixed model (as above), this time with the inclusion of maternal behavior (separately for velocity and refuge use) on a subset of individuals that had data for their own and their mother's behavior (N = 86 individuals).

To more deeply investigate how exposure to predation may alter patterns of individual behavior specifically, we ran separate univariate mixed models for each treatment (control, predator) and each behavior, which allowed us to partition variance components (among- versus within-individual variance) and estimate behavioral repeatability, our proxy for individuality, within each treatment. We ran models separately by treatment because even though variance averaged over development did not differ by treatment, our aim here was to understand how multiple variance components, including within-individual variation, changed throughout development, which our pooled treatment model above cannot estimate. As for the previous models, we also conducted model selection to determine the random and fixed effects structure. Because our goal was to compare control and predator models directly, we chose the most complex model structure that was the best fit for either treatment group, then used this structure for both models to ensure that important sources of variation were not excluded. The best supported model structure for velocity included developmental day, hour, and time of year as fixed effects, random intercepts and slopes for individual ID (across days), random intercepts and slopes for maternal ID (across days), and daily heterogeneous residual variances. For refuge use, the best supported models included developmental day and hour as fixed effects, random intercepts and slopes for individual ID, random intercepts for maternal ID, and daily heterogeneous residual variances. Models were again validated by ensuring variables of interest had low autocorrelation and high effective sample sizes.

Estimating variance components and repeatability across time series data is challenging. When random slopes are present, meaning individuals are changing rank-order over time, conditional repeatability can be used to estimate the magnitude of among-individual variance at any given time point^{90,91}; however, in these models, residual (i.e., within-individual) variance is typically described by a single component regardless of time. However, with growing evidence of the importance of within-individual behavioral variance^{60,63,92}, we chose to use heterogeneous residual models that can estimate within-individual variance at a given time point. Indeed, in our study, the best supported models (as described above) included both random intercepts, slopes and heterogeneous residual variance. Therefore, we expanded the calculations of conditional repeatability proposed by Schielzeth and Nakagawa⁹⁰ and Nakagawa et al.⁹¹ to also account for heterogeneous residual variance, estimated separately for each day. Our final repeatability equation included the variance attributable to individual intercepts ($\sigma_{\alpha 0}^2$), slopes ($\sigma_{\alpha 1}^2(x_1^*)^2$), their covariance ($2\rho\sigma_{\alpha 0}\sigma_{\alpha 1}x_1^*$), and time-specific heterogeneous residual variance ($\sigma_{\epsilon_{x^*}}^2$):

$$R_{c_{x^*}} = \frac{\sigma_{\alpha 0}^2 + \sigma_{\alpha 1}^2(x_1^*)^2 + 2\rho\sigma_{\alpha 0}\sigma_{\alpha 1}x_1^*}{\sigma_{\alpha 0}^2 + \sigma_{\alpha 1}^2(x_1^*)^2 + 2\rho\sigma_{\alpha 0}\sigma_{\alpha 1}x_1^* + \sigma_{\epsilon_{x^*}}^2}$$

For all the models above, we used parameter-expanded priors from similar studies^{21,61}; we also verified that these expanded priors were not biasing our interpretation (using MCMCglmm default priors did not change results). To investigate the effects of treatment and maternal identity on offspring body size, we used mixed models (separately for length and width) with body size as our dependent variable, treatment and time of year as fixed effects, and maternal ID as a random effect (family = gaussian).

Acknowledgements

We thank all members of the Laskowski lab, especially Abigail Trocinski, for assistance with fish husbandry throughout the study. We would also like to thank John Liu for constructing the tracking system. Lastly, we acknowledge Bubbles, our electric yellow lab cichlid, for continuously providing predator cues during the experiment. This work was funded by a Sloan Research Fellowship and an NSF grant (IOS2100625) to KLL.

Declaration of interests

The authors declare no competing interests.

References

1. Sih, A., Bell, A., and Johnson, J.C. (2004). Behavioral syndromes: an ecological and evolutionary overview. *Trends Ecol. Evol.* 19, 372–378.
2. Sih, A., Bell, A.M., Johnson, J.C., and Ziemba, R.E. (2004). Behavioral syndromes: an integrative overview. *Q. Rev. Biol.* 79, 241–277.
3. Bell, A.M., Hankison, S.J., and Laskowski, K.L. (2009). The repeatability of behaviour: a meta-analysis. *Anim. Behav.* 77, 771–783.
4. Dall, S.R.X., Bell, A.M., Bolnick, D.I., and Ratnieks, F.L.W. (2012). An evolutionary ecology of individual differences. *Ecol. Lett.* 15, 1189–1198.
5. de Bivort, B.L. (2025). The developmental origins of behavioral individuality. *Annu. Rev. Cell Dev. Biol.* <https://doi.org/10.1146/annurev-cellbio-101323-025423>.
6. Scherer, U., Ehlman, S.M., Bierbach, D., Krause, J., and Wolf, M. (2023). Reproductive individuality of clonal fish raised in near-identical environments and its link to early-life behavioral individuality. *Nat. Commun.* 14, 7652.
7. Dall, S.R.X., Houston, A.I., and McNamara, J.M. (2004). The behavioural ecology of personality: consistent individual differences from an adaptive perspective. *Ecol. Lett.* 7, 734–739.
8. Moiron, M., Laskowski, K.L., and Niemelä, P.T. (2020). Individual differences in behaviour explain variation in survival: a meta-analysis: Meta-analysis of behaviour and survival. *Ecol. Lett.* 23, 399–408.
9. Smith, B.R., and Blumstein, D.T. (2008). Fitness consequences of personality: a meta-analysis. *Behav. Ecol.* 19, 448–455.
10. Dingemanse, N., and Réale, D. (2005). Natural selection and animal personality. *Behaviour* 142, 1159–1184.
11. Li, R., Yang, C., Lou, Y., Lü, J., and Lyu, N. (2024). Animal personalities influence male mating behavior and reproductive fitness. *Behav. Ecol.* 36, arae094.
12. Laskowski, K.L., Chang, C.-C., Sheehy, K., and Aguiñaga, J. (2022). Consistent Individual Behavioral Variation: What Do We Know and Where Are We Going? *Annu. Rev. Ecol. Evol. Syst.* 53, 161–182.
13. Wagner, G.P., and Altenberg, L. (1996). Perspective: Complex Adaptations and the Evolution of Evolvability. *Evolution* 50, 967–976.
14. Duckworth, R.A. (2009). The role of behavior in evolution: a search for mechanism. *Evol. Ecol.* 23, 513–531.
15. Ingley, S.J., and Johnson, J.B. (2014). Animal personality as a driver of reproductive isolation. *Trends Ecol. Evol.* 29, 369–371.
16. Wolf, M., and Weissing, F.J. (2012). Animal personalities: consequences for ecology and evolution. *Trends Ecol. Evol.* 27, 452–461.
17. Schuett, W., Dall, S.R.X., Baeumer, J., Kloesener, M.H., Nakagawa, S., Beinlich, F., and Eggers, T. (2011). Personality variation in a clonal insect: the pea aphid, *Acyrtosiphon pisum*. *Dev. Psychobiol.* 53, 631–640.
18. Kain, J.S., Zhang, S., Akhund-Zade, J., Samuel, A.D.T., Klein, M., and de Bivort, B.L. (2015). Variability in thermal and phototactic preferences in *Drosophila* may reflect an adaptive bet-hedging strategy. *Evolution* 69, 3171–3185.
19. Freund, J., Brandmaier, A.M., Lewejohann, L., Kirste, I., Kritzler, M., Krüger, A., Sachser, N., Lindenberger, U., and Kempermann, G. (2013). Emergence of individuality in genetically identical mice. *Science* 340, 756–759.
20. Vogt, G., Huber, M., Thiemann, M., van den Boogaart, G., Schmitz, O.J., and Schubart, C.D. (2008). Production of different phenotypes from the same genotype in the same

- environment by developmental variation. *J. Exp. Biol.* *211*, 510–523.
21. Laskowski, K.L., Bierbach, D., Jolles, J.W., Doran, C., and Wolf, M. (2022). The emergence and development of behavioral individuality in clonal fish. *Nat. Commun.* *13*, 6419.
 22. Bierbach, D., Laskowski, K.L., and Wolf, M. (2017). Behavioural individuality in clonal fish arises despite near-identical rearing conditions. *Nat. Commun.* *8*, 15361.
 23. Mousseau, T.A., and Fox, C.W. (1998). The adaptive significance of maternal effects. *Trends Ecol. Evol.* *13*, 403–407.
 24. Richards, E.J. (2006). Inherited epigenetic variation--revisiting soft inheritance. *Nat. Rev. Genet.* *7*, 395–401.
 25. Mulder, E.J.H., Robles de Medina, P.G., Huizink, A.C., Van den Bergh, B.R.H., Buitelaar, J.K., and Visser, G.H.A. (2002). Prenatal maternal stress: effects on pregnancy and the (unborn) child. *Early Hum. Dev.* *70*, 3–14.
 26. Vogt, G. (2015). Stochastic developmental variation, an epigenetic source of phenotypic diversity with far-reaching biological consequences. *J. Biosci.* *40*, 159–204.
 27. Oates, A.C. (2011). What's all the noise about developmental stochasticity? *Development* *138*, 601–607.
 28. Johnson, J., Sheehy, K., and Laskowski, K.L. (2025). Playing dice with behavior: drivers of stochastic individuality. *Trends Ecol. Evol.* <https://doi.org/10.1016/j.tree.2025.10.003>.
 29. Stamps, J.A., and Krishnan, V.V. (2014). Individual differences in the potential and realized developmental plasticity of personality traits. *Frontiers in Ecology and Evolution* *2*. <https://doi.org/10.3389/fevo.2014.00069>.
 30. Monaghan, P. (2007). Early growth conditions, phenotypic development and environmental change. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* *363*, 1635–1645.
 31. Zippel, M.N., Chang Kuo, D., Meng, X., Reichard, T.M., Guess, K., Vogt, C.C., Moeller, A.H., and Sheehan, M.J. (2025). Competitive social feedback amplifies the role of early life contingency in male mice. *Science* *387*, 81–85.
 32. Bucklaew, A., and Dochtermann, N.A. (2021). The effects of exposure to predators on personality and plasticity. *Ethology* *127*, 158–165.
 33. Urszán, T.J., Garamszegi, L.Z., Nagy, G., Hettyey, A., Török, J., and Herczeg, G. (2018). Experience during development triggers between-individual variation in behavioural plasticity. *J. Anim. Ecol.* *87*, 1264–1273.
 34. Bell, A.M., and Sih, A. (2007). Exposure to predation generates personality in threespined sticklebacks (*Gasterosteus aculeatus*). *Ecol. Lett.* *10*, 828–834.
 35. Edenbrow, M., and Croft, D.P. (2013). Environmental and genetic effects shape the development of personality traits in the mangrove killifish *Kryptolebias marmoratus*. *Oikos* *122*, 667–681.
 36. Tariel, J., Plénet, S., and Luquet, E. (2020). How do developmental and parental exposures to predation affect personality and immediate behavioural plasticity in the snail *Physa acuta*? *Proc. Biol. Sci.* *287*, 20201761.
 37. Scherer, U., Laskowski, K.L., Kressler, M.M., Ehlman, S.M., Wolf, M., and Bierbach, D. (2024). Predator exposure early in life shapes behavioral development and individual variation in a clonal fish. *Sci. Rep.* *14*, 1–10.
 38. Ehlman, S.M., Halpin, R., Jones, C., Munson, A., Pollack, L., and Sih, A. (2019). Intermediate turbidity elicits the greatest antipredator response and generates repeatable behaviour in mosquitofish. *Anim. Behav.* *158*, 101–108.
 39. Toscano, B.J., Gatto, J.V., and Griffen, B. (2014). Effect of predation threat on repeatability of individual crab behavior revealed by mark-recapture. *Behav. Ecol. Sociobiol.* *68*, 519–527.
 40. Sommer-Trembo, C., Zimmer, C., Jourdan, J., Bierbach, D., and Plath, M. (2016). Predator experience homogenizes consistent individual differences in predator avoidance. *J. Ethol.* *34*, 155–165.

41. Płaskonka, B., Zaborowska, A., Mikulski, A., and Pietrzak, B. (2024). Predation risk experienced by tadpoles shapes personalities before but not after metamorphosis. *Ecol. Evol.* 14, e70532.
42. Broder, E.D., and Angeloni, L.M. (2014). Predator-induced phenotypic plasticity of laterality. *Anim. Behav.* 98, 125–130.
43. Krama, T., Munkevics, M., Krams, R., Grigorjeva, T., Trakimas, G., Jöers, P., Popovs, S., Zants, K., Elferts, D., Rantala, M.J., et al. (2023). Development under predation risk increases serotonin-signaling, variability of turning behavior and survival in adult fruit flies *Drosophila melanogaster*. *Front. Behav. Neurosci.* 17, 1189301.
44. Dingemanse, N.J., Van der Plas, F., Wright, J., Réale, D., Schrama, M., Roff, D.A., Van der Zee, E., and Barber, I. (2009). Individual experience and evolutionary history of predation affect expression of heritable variation in fish personality and morphology. *Proc. Biol. Sci.* 276, 1285–1293.
45. Dupont, L., Thierry, M., Zinger, L., Legrand, D., and Jacob, S. (2024). Beyond reaction norms: the temporal dynamics of phenotypic plasticity. *Trends Ecol. Evol.* 39, 41–51.
46. Scharl, M., Wilde, B., Schlupp, I., and Parzefall, J. (1995). Evolutionary Origin of a Parthenoform, The Amazon Molly *Poecilia formosa*, on the Basis of a Molecular Genealogy. *Evolution* 49, 827–835.
47. Schlupp, I., and Plath, M. (2005). Male mate choice and sperm allocation in a sexual/asexual mating complex of *Poecilia* (Poeciliidae, Teleostei). *Biol. Lett.* 1, 169–171.
48. Schlupp, I., Riesch, R., and Tobler, M. (2007). Amazon mollies. *Curr. Biol.* 17, R536–R537.
49. Lampert, K.P., and Scharl, M. (2008). The origin and evolution of a unisexual hybrid: *Poecilia formosa*. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 363, 2901–2909.
50. Polderman, T.J.C., Benyamin, B., de Leeuw, C.A., Sullivan, P.F., van Bochoven, A., Visscher, P.M., and Posthuma, D. (2015). Meta-analysis of the heritability of human traits based on fifty years of twin studies. *Nat. Genet.* 47, 702–709.
51. Jolles, J.W. (2021). Broad-scale applications of the Raspberry Pi: A review and guide for biologists. *Methods Ecol. Evol.* 12, 1562–1579.
52. Lehtiniemi, M. (2005). Swim or hide: predator cues cause species specific reactions in young fish larvae. *J. Fish Biol.* 66, 1285–1299.
53. Brown, G.E. (2003). Learning about danger: chemical alarm cues and local risk assessment in prey fishes. *Fish Fish (Oxf.)* 4, 227–234.
54. Heczko, E.J., and Seghers, B.H. (1981). Effects of alarm substance on schooling in the common shiner (*Notropis cornutus*, Cyprinidae). *Environ. Biol. Fishes* 6, 25–29.
55. Lawrence, B.J., and Smith, R.J. (1989). Behavioral response of solitary fathead minnows, *Pimephales promelas*, to alarm substance. *J. Chem. Ecol.* 15, 209–219.
56. Mathis, A., Chivers, D.P., and Smith, R.J.F. (1993). Population differences in responses of fathead minnows (*Pimephales promelas*) to visual and chemical stimuli from predators. *Ethology* 93, 31–40.
57. Sih, A., Mathot, K.J., Moirón, M., Montiglio, P.-O., Wolf, M., and Dingemanse, N.J. (2015). Animal personality and state-behaviour feedbacks: a review and guide for empiricists. *Trends Ecol. Evol.* 30, 50–60.
58. Biro, P.A., and Adriaenssens, B. (2013). Predictability as a personality trait: consistent differences in intraindividual behavioral variation. *Am. Nat.* 182, 621–629.
59. Stamps, J.A., Briffa, M., and Biro, P.A. (2012). Unpredictable animals: individual differences in intraindividual variability (IIV). *Anim. Behav.* 83, 1325–1334.
60. Westneat, D.F., Wright, J., and Dingemanse, N.J. (2015). The biology hidden inside residual within-individual phenotypic variation: The biology of residual phenotypic variance. *Biol. Rev. Camb. Philos. Soc.* 90, 729–743.
61. Dingemanse, N.J., and Dochtermann, N.A. (2013). Quantifying individual variation in behaviour: mixed-effect modelling approaches. *J. Anim. Ecol.* 82, 39–54.

62. Mitchell, D.J., Beckmann, C., and Biro, P.A. (2021). Understanding the unexplained: The magnitude and correlates of individual differences in residual variance. *Ecol. Evol.* *11*, 7201–7210.
63. Alonzo, S.H. (2015). Integrating the how and why of within-individual and among-individual variation and plasticity in behavior. *Curr. Opin. Behav. Sci.* *6*, 69–75.
64. Mousseau, T.A., Uller, T., Wapstra, E., and Badyaev, A.V. (2009). Evolution of maternal effects: past and present. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* *364*, 1035–1038.
65. Ensminger, D.C., Langkilde, T., Owen, D.A.S., MacLeod, K.J., and Sheriff, M.J. (2018). Maternal stress alters the phenotype of the mother, her eggs and her offspring in a wild-caught lizard. *J. Anim. Ecol.* *87*, 1685–1697.
66. Veilleux, L.N., Robert, M., Ballaz, L., Lemay, M., and Rauch, F. (2011). Gait analysis using a force-measuring gangway: intrasession repeatability in healthy adults. *J. Musculoskelet. Neuronal Interact.* *11*, 27–33.
67. Lushai, G., De Barro, P.J., David, O., Sherratt, T.N., and Maclean, N. (1998). Genetic variation within a parthenogenetic lineage. *Insect Mol. Biol.* *7*, 337–344.
68. Sweep, Z.J., Royaute, R., and Dochtermann, N. (2025). Genetic and environmental sources of behavioral individuality: a test of the standard model. *bioRxiv*, 2025.08.27.672710. <https://doi.org/10.1101/2025.08.27.672710>.
69. Hageter, J., Waalkes, M., Starkey, J., Copeland, H., Price, H., Bays, L., Showman, C., Lavery, S., Bergeron, S.A., and Horstick, E.J. (2021). Environmental and molecular modulation of motor individuality in larval zebrafish. *Front. Behav. Neurosci.* *15*, 777778.
70. Nakayama, S., Laskowski, K.L., Klefoth, T., and Arlinghaus, R. (2016). Between- and within-individual variation in activity increases with water temperature in wild perch. *Behav. Ecol.* *27*, 1676–1683.
71. Briffa, M., Bridger, D., and Biro, P.A. (2013). How does temperature affect behaviour? Multilevel analysis of plasticity, personality and predictability in hermit crabs. *Anim. Behav.* *86*, 47–54.
72. Honegger, K.S., Smith, M.A.-Y., Churgin, M.A., Turner, G.C., and de Bivort, B.L. (2020). Idiosyncratic neural coding and neuromodulation of olfactory individuality in *Drosophila*. *Proc. Natl. Acad. Sci. U. S. A.* *117*, 23292–23297.
73. Aguiñaga, J., Gomulkiewicz, R., and Watts, H.E. (2021). Effect of social information on an individual's assessment of its environment. *Anim. Behav.* *178*, 267–277.
74. Wisenden, B., Rugg, M., Korpi, N., and Fuselier, L. (2009). Lab and field estimates of active time of chemical alarm cues of a cyprinid fish and an amphipod crustacean. *Behaviour* *146*, 1423–1442.
75. Wisenden, B.D. (2014). Chemical cues that indicate risk of predation. In *Fish Pheromones and Related Cues* (John Wiley & Sons, Inc), pp. 131–148.
76. Wisenden, B.D. (2019). Evidence for incipient alarm signalling in fish. *J. Anim. Ecol.* *88*, 1278–1280.
77. Chivers, D.P., and Smith, R.J.F. (1998). Chemical alarm signalling in aquatic predator-prey systems: A review and prospectus. *Écoscience* *5*, 338–352.
78. Aguiñaga, J., Jin, S., Pesati, I., and Laskowski, K.L. (2024). Behavioral responses of a clonal fish to perceived predation risk. *PeerJ* *12*, e17547.
79. Franck, D., Hannes, R.P., Lanfermann, H., and Ribowski, A. (1985). Effects of social isolation on aggressiveness in fish with special reference to the swordtail (*Xiphophorus helleri*). *Behav. Processes* *10*, 415–427.
80. Mölder, F., Jablonski, K.P., Letcher, B., Hall, M.B., Tomkins-Tinch, C.H., Sochat, V., Forster, J., Lee, S., Twardziok, S.O., Kanitz, A., et al. (2021). Sustainable data analysis with Snakemake. *F1000Res.* *10*, 33.
81. Tan, M., and Le, Q. (2019). EfficientNet: Rethinking Model Scaling for Convolutional Neural Networks. In *International Conference on Machine Learning (PMLR)*, pp. 6105–6114.

82. Pereira, T.D., Tabris, N., Matsliah, A., Turner, D.M., Li, J., Ravindranath, S., Papadoyannis, E.S., Normand, E., Deutsch, D.S., Wang, Z.Y., et al. (2022). SLEAP: A deep learning system for multi-animal pose tracking. *Nat. Methods* 19, 486–495.
83. Pereira, T.D., Aldarondo, D.E., Willmore, L., Kislin, M., Wang, S.S.-H., Murthy, M., and Shaevitz, J.W. (2019). Fast animal pose estimation using deep neural networks. *Nat. Methods* 16, 117–125.
84. Zivkovic, Z. (2004). Improved adaptive Gaussian mixture model for background subtraction. In *Proceedings of the 17th International Conference on Pattern Recognition, 2004. ICPR 2004 (IEEE)*, pp. 28–31 Vol.2.
85. Bradski, G.R., and Kaehler, A. (2015). *Learning OpenCV 2nd ed.* (O'Reilly Media).
86. Viterbi, A. (1967). Error bounds for convolutional codes and an asymptotically optimum decoding algorithm. *IEEE Trans. Inf. Theory* 13, 260–269.
87. Supervisely (2023). Supervisely Computer Vision platform.
88. Makowski, D., Ben-Shachar, M., and Lüdtke, D. (2019). BayestestR: Describing effects and their uncertainty, existence and significance within the Bayesian framework. *J. Open Source Softw.* 4, 1541.
89. Makowski, D., Ben-Shachar, M.S., Chen, S.H.A., and Lüdtke, D. (2019). Indices of effect existence and significance in the Bayesian framework. *Front. Psychol.* 10, 2767.
90. Schielzeth, H., and Nakagawa, S. (2022). Conditional repeatability and the variance explained by reaction norm variation in random slope models. *Methods Ecol. Evol.* 13, 1214–1223.
91. Nakagawa, S., Westneat, D.F., Mizuno, A., Ajoy, Y.G.A., Dochtermann, N.A., Laskowski, K., Pick, J.L., Réale, D., Williams, C., Wright, J., et al. (2025). Understanding different types of repeatability and intra-class correlation for an analysis of biological variation.
92. Westneat, D.F., Schofield, M., and Wright, J. (2013). Parental behavior exhibits among-individual variance, plasticity, and heterogeneous residual variance. *Behav. Ecol.* 24, 598–604.