

Extending the open-source R package {spatsoc} with measures of intragroup social dynamics identified by a scoping review

Alec L. Robitaille (corresponding author)
Department of Biology, Memorial University of Newfoundland and Labrador, St. John's, Newfoundland and Labrador, Canada
robit.alec@gmail.com

Quinn M. R. Webber
University of Guelph

Eric Vander Wal
Department of Biology, Memorial University of Newfoundland and Labrador, St. John's, Newfoundland and Labrador, Canada

2026-10-23

1 Abstract

1.1 Background

Beyond proximity-based social networks and home range overlap, animal telemetry data can also be used to measure intragroup social dynamics including individual position within groups, individual and group-level movement directions, leadership patterns and lagged follower behaviours. We used a scoping review of literature across domains, including behavioural ecology, collective movement, and GISciences, to identify widely used metrics for measuring intragroup social dynamics not openly available in the R programming language.

1.2 Results

The scoping review identified a series of metrics for measuring intragroup social dynamics commonly used in the literature and not available in the R programming language at the time of the review. We developed a suite of new functions for the R package {spatsoc} to make these metrics readily available in the R programming language for broader application across study species and systems. We present a case study to illustrate these new functions for measuring intragroup social dynamics with animal telemetry data.

1.3 Conclusions

The open availability of these new and flexible functions in {spatsoc} will allow researchers to more comprehensively measure the multifaceted social behaviours in their study systems. Users can now start by detecting spatiotemporal groups and analysing social networks, then move to investigating the behaviours of individuals, dyads and groups within animal social populations.

2 Keywords

Animal social dynamics, spatiotemporal groups, collective movement, leader-follower behaviour, fission-fusion, GPS, telemetry data, movement data

3 Background

Many facets of group living need to be considered to characterize animal social systems [1]. Beyond associations, additional types of dyadic relationships including genetic, affiliative, agonistic, and cooperative [2] and the interactions between spatial phenotypes, social phenotypes, and environments [3]. Using direct observations, researchers have studied intragroup social dynamics including social foraging [4–6], spatial position within groups [7,8] and the influence of individual traits such as age, sex, and dominance [9–11]. Unfortunately, direct observations can be labour-intensive to collect and susceptible to observer bias [12], with low spatial and temporal resolution making them imperfect for studies of collective movement [1].

Pairing advanced remote tracking technologies with emerging methods expands opportunities to measure intragroup social dynamics. The availability and resolution of remote tracking data continues to improve [13–15]. There are over two decades of research on measuring interactions between moving individuals from the study of “moving point objects” in GISciences [16–19] to group coordination [20,21], leadership, and decision making [22–24]. Despite this, we suggest the lack of availability of open source tools [25–27] is the missing piece between the call to integrate technology enhanced research with multidisciplinary approaches to better study animal social systems [1,28].

The R package {spatsoc} was developed to derive social networks from animal telemetry data using aggregated association rates, interindividual distances, and home range overlap [29]. The methods in {spatsoc} have been used to measure social cohesion [30], human-wildlife conflict [31], socioecology and resource availability [32], community structure [33], the influence of stress on social proximity [34], and social patterns in non-social species [35]. Despite the substantive and varied applications of {spatsoc}, users lacked sufficient and accessible functionality for measuring the behaviour of individuals within social groups.

We first conducted a scoping review to identify metrics applicable to spatiotemporal data focusing on dyadic interactions, leadership patterns, and dominance hierarchies. Of the resulting metrics, we selected those with a high evidence of use, and lack of availability in the R programming language, either through R packages or supplemental materials in the literature. Here we present 19 new functions in the R package {spatsoc} [29] for measuring intragroup dynamics. We will demonstrate these new functions through a case study and discuss biological interpretations.

4 Implementations

4.1 Scoping review

We conducted a scoping review to identify metrics used in ecology studies for measuring dyadic interactions, dominance hierarchies and leadership patterns. We focused on metrics applicable to spatiotemporal data such as GPS telemetry. The scoping review expands on the review conducted

by [36] to also include metrics that measure group-level dynamics, leadership patterns, and lagged-follower behaviours, and to explicitly establish their open source availability in the R programming language. We established a list of nine benchmark papers (Supplement A Table S 1) that were relevant to our objectives and verified whether each paper was indexed in at least one of our selected databases (Web of Science and Scopus). We used an iterative search method to fine-tune the search terms measuring the total number of papers returned and sensitivity of the search. Sensitivity is defined as the proportion of benchmark papers returned from each database [4.2.7 37]. Search terms were developed in four categories: topic, population, method and exclusion. Topic included terms associated with dyadic interactions, dominance hierarchies and leadership patterns. Population included terms associated with animals, wildlife and ecology. Method included terms associated with data collection and data types. Finally, a list of exclusion terms were used to remove irrelevant papers. Each category of terms was combined with OR operators and all four categories of terms were combined with AND operators. The resulting set of papers used in further steps was the combination of the last iteration of search terms applied to both of the Web of Science (using the “TS” field) and Scopus (using the “TITLE-ABS-KEY” field) databases (Supplement A Table S 2). The temporal range of articles was limited to 1980-2023 using the PY (Web of Science) or PUBYEAR (Scopus) fields. The resulting set of papers were imported into Covidence [38] for deduplication, title and abstract screening, full text screening, and data extraction.

Articles were screened based on the following criteria for exclusion:

- Paper does not describe dyadic or group metric related to leadership or dominance (e.g. [39])
- Paper describes metric that is not applicable to spatiotemporal data (e.g. [40] used focal observations and video recordings)
- Paper does not describe metric beyond static interaction (e.g. [41] used home range or spatial overlap)
- Paper does not describe metric beyond group size (e.g. [42])
- Paper full text is not available in English (e.g. [43])
- Paper is a thesis
- Paper is a book chapter (e.g. [44])

Despite the intended focus on animal ecology, some papers discussed movement data more generically or, alternatively, specifically focused on human or robotic movement data (e.g. [45]). These papers were retained when the metrics described were deemed applicable to animal spatiotemporal movement data. Papers using mathematical models or simulations were largely beyond the scope of this review but papers were retained when the parameters used as inputs could be applied to spatiotemporal data (e.g. [46]). Review papers were read for building the scope of the introduction and discussion but not included in data extraction unless they explicitly discussed dyadic measures (e.g. [1]).

The goals of data extraction were to record the breadth of metrics for measuring dyadic interactions, dominance hierarchies and leadership patterns applicable to spatiotemporal data, and the degree of code availability. General information about each paper was recorded including the type of study (empirical, simulation, algorithm, model, or review), study species and study region.

Metrics discussed in each paper were listed including data input required, citations and interpretation. Code availability was described including location of code (e.g. Zenodo, or GitHub), programming languages and software packages used. The data extraction form is in Supplement A Table Table S 3.

Metrics identified were aggregated using a manual process to identify synonymous terms. This aggregation process was required due to variability in defining metrics during data extraction and differences between authors in describing metrics (e.g. distance to group “center”, “centroid”, or “center of mass”). Each paper’s study location was geocoded using the {tidygeocoder} package [47] and the study species were parsed using the {rgnparser} package [48] which provides the GParser [49] taxonomic parser in R. We calculated frequencies for each categorical field including programming language, software package, analysis code availability, geographic region, study species and metric used. For many fields, authors did not provide sufficient information to determine how metrics were calculated, e.g. not including the programming language or software package used, and as a result, counts are presented as an overall frequency including papers with missing fields.

Metrics were selected for inclusion in {spatsoc} based on their relative importance (as measured by frequency of use in the scoping review), and their lack of availability in R when the review was conducted (Supplement Section 11). Some metrics were specifically not chosen for inclusion such as metrics relating to posture, metrics relating to static interaction or home range overlap and metrics based on statistical models such as Hidden Markov model and Granger causality. A full list of metrics not selected is in Supplement Section 12.

4.2 Software

The R package {spatsoc} was originally developed to detect spatiotemporal groups in animal telemetry data for building social networks [29]. In this new version, {spatsoc} provides an additional suite of functions to measure individual, dyadic and group level behaviours within spatiotemporal groups. These measures of intragroup dynamics are derived from three geospatial operations (distance, direction, and centroid) calculated in internal functions built on top of the R spatial packages {sf} [50], {lwgeom} [51] and {s2} [52]. Improving on past versions, {spatsoc} now accommodates data in both projected and geographic coordinate reference systems.

5 Results

The final set of search terms (Supplement A Table S 2) applied to both Web of Science and Scopus returned a database of 3878 articles between 1980 and 2023. Manual deduplication removed 128 articles and Covidence [38] identified an additional 799 duplicated articles. 2527 articles were excluded during title and abstract screening and 275 articles were excluded by full text screening. The resulting database included 141 studies (see Supplement B for detailed PRISMA).

Data extraction revealed a total of 119 unique metrics including metrics used in articles that may not explicitly measure intragroup social dynamics. For example, if an article measured an individual’s position within the group and also their step length, both metrics were recorded. The review indicated variability in the methodological definitions relating to intragroup social

dynamics. Differences in defining spatiotemporal groups were largely centered on spatial and temporal thresholds (Supplement B Table S 12). Similarly, fission-fusion dynamics were defined by spatial and temporal thresholds [e.g. 53,54], but also how to handle missed observations and temporary separations [e.g. 55,56] (Supplement B Table S 14). Definitions of group-level leadership patterns were based on measures of leadership including tendency to be at the front of the group [e.g. 57,58,59], directional correlation delay [e.g. 60,61], and movement initiations [30,e.g. 62,63] (Supplement B Table S 13).

Details in the methods and supplemental materials were used to assess the availability of the extracted metrics in open source programming languages commonly used in ecology. Accompanying code was not available in a supplement or linked repository for 82% of articles and 67% of articles did not indicate any software package (e.g. R package, Python library) used. R was the most common programming language followed by MATLAB and Python (Supplement B Table S 5) in the database of articles. Metrics already provided by existing open source software packages (e.g. DI provided by {wildlifeDI} [45]), or code accompanying articles (e.g. FLICA code provided alongside [64]) were not selected for inclusion in {spatsoc}. Finally, metrics were prioritized for inclusion in {spatsoc} according to measured frequency of use in the database of articles and external searches were conducted to ensure no availability in R packages were missed by the scoping review (see Supplement B Table S 10 Supplement C). The resulting set of metrics provided by new functions in the R package {spatsoc} for measuring intragroup social dynamics is shown in Table Table 1.

Table 1: Description of new functions in the R package {spatsoc} for measuring intragroup dynamics categorized by family. All available functions in the R package {spatsoc} including new and existing functions are listed in Table 15.

Function	Description
Spatial interface	
get_geometry()	Setup data for geometry interface
Edge-list generation	
edge_delay()	Directional correlation delay based edge-lists
edge_alignment()	Directional alignment based edge-lists
edge_direction()	Direction based edge-lists
edge_zones()	Behavioural zones based edge-lists
Dyad	
fusion_id()	Identify fusion events
Centroid	
centroid_group()	Group centroid
centroid_dyad()	Dyad centroid
centroid_fusion()	Fusion event centroid
Direction	
direction_step()	Direction step
direction_to_centroid()	Direction to group centroid
direction_to_leader()	Direction to group leader
direction_group()	Group mean direction
direction_polarization()	Polarization
Distance	
distance_to_centroid()	Distance to group centroid
distance_to_leader()	Distance to group leader
Leadership	
leader_direction_group()	Leadership along group direction
leader_edge_delay()	Leadership in directional correlation delay

6 Discussion: case study

The expected input data structure for functions in {spatsoc} is a `data.table` with columns representing the individual identifiers (e.g. “ID”), the date and time (e.g. “datetime”) and the

coordinates (e.g. “X”, “Y”) of each location. Other tabular data formats in R can be converted to `data.table` with the function `as.data.table`. Many of the functions in `{spatsoc}` process input data using `{data.table}`’s modify-by-reference to avoid making copies of input data and, instead, directly adding or updating columns in place. Functions that return the input `data.table` with an added column do not need to be reassigned (e.g. `group_pts` and `group_times`). In contrast, functions that reshape the input data, return aggregated results or require an internal merge need to be reassigned (e.g. `edge_dist` and `centroid_dyad`). Each function’s documentation describes if it uses modify-by-reference or if its output needs to be reassigned. The FAQ vignette lists all functions in these two categories and users are encouraged to read more about `{data.table}`’s modify-by-reference in their vignette about Reference Semantics.

Table 2: Expected input data structure for `{spatsoc}` functions: a `data.table` with columns representing the individual identifier (“ID”), date and time (“datetime”) and coordinates (“X” and “Y”) of each location.

ID	X	Y	datetime
A	715851	5505340	2016-11-01 00:00:54
B	699640	5509638	2016-11-01 00:01:02
C	710205	5505888	2016-11-01 00:00:44
D	700875	5490954	2016-11-01 00:00:47
E	701672	5504286	2016-11-01 00:00:48
F	705583	5513813	2016-11-01 00:00:48

Functions previously available in `{spatsoc}` can be used to identify spatiotemporal grouping, generate distance based edge-lists and conduct data-stream permutations (see existing functionality in [29]; Table S 15). These functions have allowed users to detect interactions between individuals, measure co-occurrence within and across species, and generate social networks from animal telemetry data [30,35,e.g. 65,66].

Spatiotemporal grouping can be performed given animal movement data, temporal and spatial thresholds, and one of three grouping functions: `group_pts()`, `group_lines()` or `group_polys()`. These return a unique identifier for each spatiotemporal group defined by point-based distances, linear trajectory overlap and home range overlap, respectively.

`get_geometry()` is a new helper function added to `{spatsoc}` to provide an alternative interface for providing coordinates and, optionally, coordinate reference systems. Instead of providing the names to columns for the X and Y coordinates using the `coords` argument, the `get_geometry()` function adds a simple feature geometry list column (`sfc`) using the `{sf}` package [50]. The geometry method is used by default for all `{spatsoc}` functions that require coordinates when the `coords` argument is left null. This interface also extends `{spatsoc}` to allow users to provide coordinates in any coordinate reference systems, including both projected and longlat degrees, supported by the `{sf}` package. The following examples and case study will illustrate the geometry interface throughout.

For example, the suggested `group_pts()` workflow with the example data from the package and the `get_geometry()` helper function to identify spatiotemporal groups (Table 3):

```
# Load packages
library(data.table)
library(spatSOC)

# Read example data
DT <- fread(system.file("extdata", "DT.csv", package = "spatSOC"))

# Cast the character column to POSIXct
DT[, datetime := as.POSIXct(datetime, tz = 'UTC')]

# Set variables
temporal_threshold <- '20 minutes'
spatial_threshold <- 50
id <- 'ID'
coords <- c('X', 'Y')
utm <- 32736

# Setup DT for geometry interface
get_geometry(DT, coords = coords, crs = utm)

# Temporal grouping
group_times(DT, datetime = 'datetime', threshold = temporal_threshold)

# Spatial grouping with timegroup
group_pts(
  DT,
  threshold = spatial_threshold,
  id = id,
  coords = coords,
  timegroup = 'timegroup'
)
```

Table 3: Spatiotemporal groups measured using `group_times()` and `group_pts()`. `group_times()` returns a 'timegroup' column indicating temporal groups within the specified temporal threshold. `group_pts()` returns a 'group' column indicating spatiotemporal groups within the provided timegroup and specific distance threshold. Example output showing a subset of three spatiotemporal groups.

ID	datetime	geometry	timegroup	group
B	2016-11-08 12:00:43	POINT (697250 5510781)	91	1487
G	2016-11-08 12:00:54	POINT (697243 5510778)	91	1487
B	2016-12-02 20:00:53	POINT (699459 5510354)	383	1779
G	2016-12-02 20:00:48	POINT (699445 5510330)	383	1779
E	2016-12-12 02:00:27	POINT (697934 5507852)	494	5542
H	2016-12-12 02:00:48	POINT (697928 5507853)	494	5542

Users can also measure interindividual distance using the `edge_dist()` function (Table 4) and identify nearest neighbours using `edge_nn()` function (Table 5). For example:

```
# Measure interindividual distance
interindividual_dist <- edge_dist(
  DT,
  threshold = spatial_threshold,
  id = id,
  timegroup = 'timegroup',
  returnDist = TRUE,
  fillNA = TRUE
)
```

Table 4: Distance based edge-lists measured using `edge_dist()`. Using the ‘timegroup’ column returned by `group_times()`, `edge_dist()` returns a distance based edge-list where individuals in the column ‘ID1’ are within the provided distance threshold from ‘ID2’. Optionally, measured distances can be returned in the column ‘distance’ if ‘returnDist’ is TRUE and NAs can be returned where an individual is not within the provided distance threshold if ‘fillNA’ is TRUE. Example output showing six observations of one timegroup with distances returned between focal individual ‘ID1’ and neighbour ‘ID2’.

timegroup	ID1	ID2	distance
1436	B	H	48.12
1436	C	NA	NA
1436	E	G	13.19
1436	E	J	12.36
1436	F	NA	NA
1436	G	E	13.19

```
# Identify nearest neighbours
nearest_neighbours <- edge_nn(
  DT,
  id = id,
  timegroup = 'timegroup',
  returnDist = TRUE
)
```

Table 5: Distance based edge-lists measured using `edge_nn()`. Using the ‘timegroup’ column returned by `group_times()`, `edge_nn()` returns a distance based edge-list where individuals in the column ‘NN’ are the nearest neighbour to focal individuals in the column ‘ID’. Optionally, measured distances can be returned in the column ‘distance’ if ‘returnDist’ is TRUE and a distance threshold can be provided using ‘threshold’. Example output showing six observations of one timegroup with distances returned between focal individual ‘ID’ and nearest neighbour ‘NN’.

timegroup	ID	NN	distance
1424	J	B	64.19
1424	G	E	20.53
1424	H	E	3389.49
1424	I	F	43.92
1424	B	G	27.13
1424	E	G	20.53

Building on these, {spatsoc}'s new functions provide more detailed measures of an individual's behaviour with respect to their conspecifics. We aim to provide users with accessible, flexible functions that will help to identify leader-follower patterns, fission-fusion dynamics and potentially dominance behaviours from animal telemetry data in their ecological systems. Sufficient data quality and resolution are a continued consideration when determining if a particular data set is suited to the new functions in {spatsoc}. For example, the directional correlation delay functions `leader_direction_delay` and `edge_delay` depend on a sufficiently high temporal resolution to capture correlations in lagged movement directions between individuals that are biologically relevant. While the example data used in the case study may not be sufficient for biological interpretation for all functions, it is used for illustrative purposes due to its accessibility in the package. We refer users to the references accompanying each function for more details on ensuring their data is appropriate.

6.1 Position within groups

Extending {spatsoc}'s abilities to identify spatiotemporal groups, a new set of functions are now available to measure individual's relative position within spatiotemporal groups. After identifying spatiotemporal groups with e.g. `group_pts()`, we can measure the group centroid, as defined by the mean of individual locations in a group, using `centroid_group()` (Table 6).

```
# Measure group centroids
centroid_group(DT)
```

Table 6: Group centroids measured using `centroid_group()`. Using the 'group' column returned by `group_pts()`, `centroid_group()` returns the column 'centroid' representing the centroid of all individual's locations in each spatiotemporal group. Example output showing a subset of the centroids for three spatiotemporal groups.

ID	timegroup	group	geometry	centroid
B	463	1859	POINT (698424 5511571)	POINT (698410 5511576)
G	463	1859	POINT (698396 5511582)	POINT (698410 5511576)
I	471	8728	POINT (698754 5506683)	POINT (698752 5506679)
J	471	8728	POINT (698749 5506674)	POINT (698752 5506679)
B	643	2039	POINT (699011 5511657)	POINT (699003 5511660)
G	643	2039	POINT (698995 5511664)	POINT (699003 5511660)

Then we can measure each individual's distance and direction to the group centroid using `distance_to_centroid()` and `direction_to_centroid()` (Figure 1). The distance to group centroid is the geographic distance from the focal individual to the group centroid. The direction to group centroid is the direction from the focal individual to the group centroid. The rank of individuals' distances to the group centroid can be optionally returned using the argument `return_rank`. Spatial position within groups with respect to the group centroid has been studied in relation to predation risk [67,68], leadership [61,69] and group cohesion [30,70].

```
# Measure distance to group centroid
distance_to_centroid(DT, return_rank = TRUE)

# Measure direction to group centroid
direction_to_centroid(DT)
```




Figure 1: A) Distance and B) direction to group centroids measured using `distance_to_centroid()` and `direction_to_centroid()`. Using the ‘centroid’ column returned by `centroid_group`, `distance_to_centroid()` and `direction_to_centroid()` return the columns ‘distance_centroid’ and ‘direction_centroid’ representing the focal individual’s distance and direction to the group centroid. Optionally, the rank distance to the group centroid can be returned by `distance_to_centroid()` if the argument ‘return_rank’ is true.

6.2 Directions

Next, we can use movement directions to measure how individuals and groups move together. We can measure an individual’s movement direction using `direction_step()` and the mean direction of each spatiotemporal group identified by `group_pts()` using `direction_group()` (Table 7).

```
# Measure movement direction
direction_step(DT, id = id)

# Measure mean movement direction of each spatiotemporal group
direction_group(DT)
```

Table 7: Individual and group mean directions measured using `direction_step()` and `direction_group()`. `direction_step()` returns the column ‘direction’ representing the movement direction of each individual to their next relocation. Using the spatiotemporal groups returned by `group_pts`, `direction_group()` returns the column ‘group_direction’ representing the mean direction of all individuals in each spatiotemporal group. Example output showing the movement directions of individuals and mean direction of two spatiotemporal groups.

ID	timegroup	group	direction	group_direction
E	623	5629	2.4033 [rad]	2.687 [rad]
H	623	5629	3.0019 [rad]	2.687 [rad]
I	623	5629	2.6582 [rad]	2.687 [rad]
C	1426	3640	0.1258 [rad]	0.168 [rad]
F	1426	3640	0.1819 [rad]	0.168 [rad]
I	1426	3640	0.1963 [rad]	0.168 [rad]

We can measure polarization, the uniformity of movement directions in a group of individuals, using `direction_polarization()` (Figure 2). Polarization is on a scale of 0-1 where values near 0 show that individuals in a group are moving in different directions (or “non-aligned”) and values near 1 show that individuals are moving in similar directions or (“aligned”). Polarization in the movement direction of groups of animals has been used to study cognition [71], learning [72], the influence of habitat [73] and individual differences [69] in collective behaviour.

```
# Measure direction polarization
direction_polarization(DT)
```




Figure 2: Direction polarization measured using `direction_polarization()`. Using the ‘direction’ column returned by `direction_step()` and ‘group’ column returned by `group_pts()`, `direction_polarization()` returns a column named ‘polarization’ indicating the polarization in directions in each spatiotemporal group. Ranges of directions shown for a sample of groups in four equal interval categories of direction polarization. Values of polarization range from 0 (non-alignment) to 1 (alignment).


The interindividual direction, or direction from a focal individual to a neighbour, can be measured using `edge_direction()` (Figure 3). Like `edge_alignment()`, the interindividual directions are returned in an edge-list format. Interindividual direction has been used to study risk exposure in schooling fish [74], coordinated movement [75] and mass migration [76].

```
# Measure interindividual direction
dyad_directions <- edge_direction(
  edges = interindividual_dist,
  DT = DT,
  id = id
)
```

The directional alignment, or relative difference between two individuals’ directions, can be measured using `edge_alignment()` (Figure 3). The differences in movement directions are returned in an edge-list format similar to `edge_dist()` and other edge-list generating functions. Directional alignment has been used to study leadership [77], decision making [24], and information transfer [78].

```
# Measure directional alignment
directional_align <- edge_alignment(
  DT = DT,
  id = id,
  signed = FALSE
)
```

A)



B)

timegroup	ID1	ID2	direction_dyad	direction_diff
1	F	E	-1.06 [rad]	0.08 [rad]
1	E	F	2.08 [rad]	0.08 [rad]
2	F	E	1.99 [rad]	0.34 [rad]
2	E	F	-1.15 [rad]	0.34 [rad]
3	F	E	2.86 [rad]	0.57 [rad]
3	E	F	-0.28 [rad]	0.57 [rad]

Figure 3: A) Example locations for individuals ‘E’ (purple) and ‘F’ (blue) showing concurrent movement paths in timegroups 1, 2 and 3. B) Interindividual direction and `directional_align()` using `edge_direction()` and `edge_alignment()`. Using distance based edge-lists returned by `edge_dist()` or `edge_nn()` and the ‘timegroup’ column returned by `group_times()`, `edge_direction()` returns a column named ‘direction_dyad’ representing the direction between the focal individual (‘ID1’) and the neighbouring individual (‘ID2’). Using the ‘direction’ column returned by `direction_step()`, the `edge_alignment()` function returns a column named ‘direction_diff’ representing the absolute difference in movement directions between two individuals (‘ID1’ and ‘ID2’). Optionally, the signed difference can be returned by `edge_alignment()` if the argument ‘signed’ is true.

6.3 Behavioural zones

The behavioural zones [20], or non-overlapping behavioural regions around a focal individual, can be measured using `edge_zones()` (Figure 4). The distance from the focal individual to each

neighbour is used to assign the neighbours to behavioural zones. In addition, the direction from the focal individual to each neighbour can be used to determine if the neighbour is within a “blind volume” outside of the individual’s visual perception. [20] defined three non-overlapping zones. The “zone of repulsion” is at a minimum distance around the focal individual within which neighbours are expected to move away to avoid collisions. The “zone of orientation” is around the focal individual beyond the “zone of repulsion” within which neighbours are expected to orient themselves to the movement of their neighbours. The “zone of attraction” is beyond the “zone of orientation” where neighbours are expected to be attracted to the position of the focal individual. The `edge_zones()` function is designed to allow for flexible definitions of zones including distance thresholds, zone labels and blind volume thresholds. The behavioural zones metric has been used to study the impact of individual movement speed and predator evasion [79] and identifying rules of interaction in large animal groups [80].

```
# Define zone thresholds, labels and blind volume
zone_thresholds <- c(10, 25, 50)
zone_labels <- c('repulsion', 'orientation', 'attraction')
blind_volume <- 3

# Measure behavioural zones
zones <- edge_zones(
  dyad_directions,
  zone_thresholds = zone_thresholds,
  zone_labels = zone_labels,
  blind_volume = blind_volume
)
```




Figure 4: Behavioural zones measured using `edge_zones()`. Using the ‘distance’ column returned by `edge_dist()`, the `edge_zones()` function returns the column ‘zone’ representing the behavioural zone of the neighbour (ID2) with respect to the focal individual (ID1). Optionally, the ‘direction_dyad’ column returned by `edge_direction()` can be used to define the blind volume, or the range of interindividual directions that are beyond the visual perception of the focal individual. Example shows positions of neighbour ‘E’ in behavioural zones with respect to focal individual ‘G’ across the sample period.

6.4 Position along group direction

An individual’s position relative to the mean group direction can be calculated with `leader_direction_group()`. We first calculate each individual’s movement direction with `direction_step()`, the group’s centroid with `centroid_group()` and the group’s mean direction with `direction_group()`. Then `leader_direction_group()` rotates the coordinate system around the group centroid by the group’s mean direction for each spatiotemporal group. This new axis represents individual’s front-back position within the group along the group’s mean direction [57,62,70] (Figure 5). The rank of individual positions within spatiotemporal groups [81] can be optionally returned when the argument ‘return_rank’ is true.

```
# Measure movement direction
direction_step(DT, id = id)

# Measure mean movement direction of each spatiotemporal group
direction_group(DT)

# Measure position along group direction
leader_direction_group(DT, return_rank = TRUE)
```




Figure 5: A) Example spatiotemporal group showing relative positions and movement directions of individuals (small points and grey arrows) with respect to the group centroid and mean group direction (large point and black arrow). B) Position along the group direction measured using `leader_direction_group()`. Using the 'group_direction' column returned by `direction_group()` and 'centroid' column returned by `centroid_group()`, `leader_direction_group()` returns a column named 'position_group_direction' representing the front-back position along the mean group direction. Optionally, the rank position along the mean group direction if the argument 'return_rank' is true.

Taking this simple, dynamic definition of leadership, we can identify the distance and direction of individuals to the leader of each spatiotemporal group using `distance_to_leader()` and `direction_to_leader()` (Figure 6). The distance to leader is the geographic distance from the focal individual to the group's leader and the direction to leader is the direction from the focal individual to the group's leader. The front-back position within the group, and distance and direction to the group leader has been used to study interaction rules [82], leadership [62,77],

risk avoidance [83], and how reproduction influences social structure [58]. In addition, a user can calculate an individual's time spent leading by calculating the rate each individual was in the first ranked position out of the total observations. Other definitions of leadership in the scoping review also include the directional correlation delay, described below, and movement initiations (Supplement B Table S 13).

```
# Measure distance to leader
distance_to_leader(DT)

# Measure direction to leader
direction_to_leader(DT)
```




Figure 6: A) Distance and B) direction to group leaders measured using `distance_to_leader` and `direction_to_leader`. Using the ‘rank_position_group_direction’ column returned by `leader_direction_group()`, `distance_to_leader()` and `direction_to_leader()` return columns named ‘distance_leader’ and ‘direction_leader’ indicating the distance and direction of the focal individual to the leader of the spatiotemporal group. Note: for individuals identified as leaders using ‘position_group_direction’, the distance is 0 and the direction is NA.

6.5 Dyadic fusion events

Dyadic fusion events, or temporal periods where two individuals are within a given threshold distance from each other, can be identified using `fusion_id()` (Figure 7). Given the variability in defining dyadic fission-fusion dynamics in the literature (Table S 14), the `fusion_id()` function flexibly allows users to set specific thresholds and definitions that match their study species and research questions. This will also allow users to easily compare their results to previously published work by using the same thresholds and definitions. The `threshold` argument defines


the spatial distance threshold to establish a fusion event. The `n_min_length` argument defines the minimum number of successive fixes that are required to establish a fusion event. The `n_max_missing` defines the maximum number of allowable missed observations for either individual in a dyad within a fusion event. The `allow_split` argument defines if fusion events allow a temporary spatial splitting for one observation without resulting in a fission event. Identification of fusion events has been used to study resource seasonality [84], information transfer [85], decision making [86], and habitat selection [87].

```
# Identify dyads
dyad_id(interindividual_dist, id1 = 'ID1', id2 = 'ID2')

# Identify dyadic fusion events
fusion_events <- fusion_id(
  interindividual_dist,
  threshold = spatial_threshold,
  n_min_length = 3,
  n_max_missing = 1,
  allow_split = TRUE
)

# Measure centroids of dyadic fusion events
centroid_fusion(
  interindividual_dist,
  DT,
  id = id
)
```

A)



B)

timegroup	ID1	ID2	distance
1	J	NA	NA
2	J	C	11.26 [m]
3	J	C	9.98 [m]
4	J	C	35.89 [m]
5	J	NA	NA

Figure 7: A) Example dyadic fusion event for dyad of individuals 'J' and 'C' with black points representing the centroid of fusion observations returned by `centroid_fusion()`. B) A dyadic fusion event identified using `fusion_id`. Using the distance based edge-lists measured using `edge_dist()` and the dyad id returned by `dyad_id()`, `fusion_id()` returns a column named 'fusionID' representing the dyadic fusion events defined with user specified constraints: the distance threshold ('threshold'), the minimum number of observations defining a fusion event ('n_min_length'), the maximum number of missing observations from either individual in a fusion event ('n_max_missing') and if a single observation where individuals are beyond the distance threshold should result in a fission event ('allow_split'). In B), a dyadic fusion event is identified for individuals 'J' and 'C' given they are within the distance threshold at timegroups 2, 3, and 4. Individual 'J' is shown at timegroups 1 and 5 with no neighbours within the threshold spatial distance (ID2 is NA).

6.6 Directional correlation delay

The directional correlation delay can be calculated with `leader_direction_delay()` to measure the correlation between movement directions of a focal individual and a neighbour within a specified temporal window [23]. The directional correlation delay time represents the delay that maximizes the mean directional correlation delay across the sampling period. The delay time can be used to indicate if an individual is leading their neighbour (where the delay time is positive)

or following their neighbour (where the delay time is negative). The window argument controls the number of time steps around the focal time (e.g. window = 2 defines the window to include $t_{-2}, t_{-1}, t_0, t_1, t_2$). The min_nobs argument sets the minimum number of valid observations for the focal individual in the temporal window required for each focal observation. The splitBy argument is used to define sampling periods within which the directional correlation delay and delay time are calculated. leader_direction_delay() uses the fusion events defined with fusion_id() to ensure that dyads are within a minimum distance for a specified minimum number of timesteps. The mean directional correlation delay is calculated across all fusion events in the sampling period for each dyad and delay time defined by the window argument. The directional correlation delay time can be used to measure leadership patterns and generate hierarchical leadership networks depending on the study system [23,88–90].

The momentary directional correlation delay is adapted from the global directional correlation delay [62] and can be measured with edge_delay(). In contrast to the global directional correlation delay [23], the momentary directional correlation delay [62] measures the correlation between movement directions within dyads at each time step. The edge_delay() function uses the fusion events defined with fusion_id() like leader_direction_delay() and returns the momentary directional correlation delay time that maximizes the directional correlation delay at each time step. The window and min_nobs arguments are shared with leader_direction_delay(). The momentary leadership index can be calculated from the output of edge_delay by taking the mean of the directional correlation delay time for each focal individual (ID1) at each time step [62].

```
# Measure global directional correlation delay
leader_delay <- leader_direction_delay(
  edges = fusion_events,
  DT = DT,
  window = 3,
  id = id,
  min_nobs = 2
)
```

```
# Measure momentary directional correlation delay
delay <- edge_delay(
  edges = fusion_events,
  DT = DT,
  window = 3,
  id = id,
  min_nobs = 2
)

# Calculate momentary leadership index
delay[, momentary_leadership_index := mean(direction_delay),
  by = .(ID1, timegroup)]
```

A)

ID1	ID2	direction_delay	direction_cor	max_direction_cor
E	H	-3	0.1416	FALSE
E	H	-2	0.0802	FALSE
E	H	-1	0.1542	FALSE
E	H	0	0.7253	TRUE
E	H	1	0.1712	FALSE
E	H	2	0.0443	FALSE
E	H	3	0.0309	FALSE

B)

timegroup	ID1	ID2	direction_cor	direction_delay
1	E	H	0.7080	-1
1	H	E	0.9854	-2
2	E	H	0.9827	0
2	H	E	0.9827	0
3	E	H	0.9982	0
3	H	E	0.9982	0
4	E	H	0.8876	2
4	H	E	0.7291	-3

Figure 8: A) Example global directional correlation delay [23] for focal individual ‘E’ (ID1) and neighbour ‘H’ (ID2) indicating individual ‘E’ had a movement direction most similar to individual ‘H’ at the temporal delay of 0. Using the fusion events identified by `fusion_id()` and the ‘direction’ column returned by `direction_step()`, `leader_direction_delay()` returns a column named ‘direction_cor’ indicating the correlation of directions between ID1 at time t and ID2 at time $t + \tau$ where τ represents the temporal delay returned in the column named ‘direction_delay’, and a column named ‘max_direction_cor’ indicating the temporal delay that maximizes the mean directional correlation delay. B) Example momentary directional correlation delay [62] for individuals ‘E’ and ‘H’ during four timesteps of a fusion event. Using the fusion events identified by `fusion_id()` and the ‘direction’ column returned by `direction_step()`, `edge_delay()` returns a column named ‘direction_cor’ representing the maximum directional correlation delay at each time step for each dyad and a column named ‘direction_delay’ representing the corresponding temporal delay that maximizes the directional correlation delay at the focal timestep within the temporal window specified by the user. Notably, the momentary directional correlation delay for ID2-ID1 is not the inverse of ID1-ID2, and therefore both individual’s perspectives are returned. The momentary directional correlation delay is calculated in the code example but is not shown here for simplicity.

7 Conclusion

An ever expanding collection of methods have been described for studying animal social systems. The scoping review identified hundreds of metrics applied to species across the globe. Despite this, a lack of their accessibility in open source software risks limiting the application of these methods broadly. The open source package {spatsoc} [29] for the R programming language [91] has improved accessibility to methods for spatiotemporal grouping and social network analysis with animal telemetry data [30,35,e.g. 65,66]. Despite this, proximity-based association is only one dimension and studying intragroup social dynamics will allow researchers to further measure animal social systems. The new functions in {spatsoc} and the accompanying case study presented here make openly available a series of metrics that, while commonly used in the literature, are largely absent from R packages. Paired with the continued development of animal tracking technology [15,92] and novel data collection methods [93,94], the open availability of these metrics in R will broaden their application across research questions, including, among others, long-standing ecological concepts such as:

- Social foraging theory and producer-scrounger dynamics where “producers” are those who discover patches and “scroungers” are those who join patches discovered by others [4,95,96]. This concept can be applied to animal telemetry data by evaluating the position of individuals within groups, the individual’s arrival timing to forage patches and the characteristics of forage patches. Individuals balance risk of predation and foraging success by adjusting their spatial position within groups [6,11], and the foraging strategy selected by an individual can be influenced by the size of forage patches [97].
- Leader-follower behaviour in groups of moving animals is studied both to determine factors that causally influence leadership, and for the influence of leader-follower strategies on foraging success and downstream fitness. Applied to animal telemetry data, leadership can be defined simply by identifying the individual at the foremost position along the front-to-back axis of the group’s direction [62,70] or by calculating the directional correlation delay between individuals and building leadership hierarchies [23]. Leadership in a group of moving animals can inform animal decision making, group coordination, and migratory patterns [61,62,82,98].
- Individual traits can influence behaviours related to intragroup dynamics. Dominance is related to foraging success [11,99] and choice of foraging strategy [10,100,101]. Dominant, aggressive, grooming and other interindividual behaviours are typically measured using focal observations. However, some metrics have been developed for remotely estimating dominance behaviours. The directional correlation delay can be used to derive hierarchies of leadership which can be a proxy for dominance relationships in certain species [88,90]. Dominance relationships can also be inferred from animal telemetry data using movement initiations [64], approach-avoidance behaviours [24] and displacement events [102].

The lack of open source code and transparency of methods alongside research (Table S 4, Table S 5, Table S 6) continues to slow the progress of science by limiting the broader application of new methods. We hope that the availability of these metrics in {spatsoc} will allow researchers to apply them to diverse study species, systems or research questions, including novel applications that could have management and conservation implications for vulnerable species [103].

8 Availability and requirements

Project name: spatsoc

Project home page: <https://github.com/ropensci/spatsoc>

Operating system(s): Platform independent

Programming language: R (≥ 3.4)

Other requirements: R packages adehabitatHR ($\geq 0.4.21$), data.table ($\geq 1.15.0$), igraph, sf, lwgeom ($\geq 0.2.15$), CircStats, stats, units ($\geq 0.8-6$), rlang. System dependencies GDAL ($\geq 2.0.1$), GEOS ($\geq 3.4.0$), PROJ ($\geq 4.8.0$), sqlite3

License: GPL-3

Any restrictions to use by non-academics: None

9 Declarations

9.1 Ethics approval and consent to participate

Not applicable

9.2 Consent for publication

Not applicable

9.3 Availability of data and materials

All code used to produce this manuscript including figures and tables are available on GitHub at <https://github.com/robital/ec/extendspatsoc-application-paper> and on Zenodo at <https://doi.org/10.5281/zenodo.19712779>. The data used in this case study are included with the package and can be imported with:

```
library(spatsoc)

library(data.table)

DT <- fread(system.file("extdata", "DT.csv", package = "spatsoc"))
```

The code for producing the scoping review results, figures, tables and manuscript was developed as a reproducible pipeline with the R package {targets} [104]. Figures and tables were constructed using {ggplot2} [105], {ggdist} [106], {patchwork} [107] and {tintytable} [108]. The manuscript was produced using {quarto} [109]. The {spatsoc} package gratefully depends on the R packages {adehabitatHR} [110], {data.table} [111], {igraph} [112], {sf} [50], {lwgeom} [51], {CircStats} [113], {units} [114], and {rlang} [115].

9.4 Competing interests

The authors declare that they have no competing interests.

9.5 Funding

ALR received a NSERC Canada Graduate Research Scholarship, QMRW received a NSERC Discovery Grant, EVW received a NSERC Discovery Grant.

9.6 Authors' contributions

ALR, QMRW and EVW contributed to conceptualization and writing. EVW contributed to methodology, funding acquisition, project administration and supervision. ALR contributed to writing (original draft), software, investigation, formal analysis, data curation and visualization. All authors read and approved the final manuscript.

9.7 Acknowledgements

We thank Edzer Pebesma and Dewey Dunnington for support with bug reports and feature requests for the `{sf}`, `{s2}` and `{lwgeom}` packages related to the development of this new functionality in `{spatoc}`. We thank Martin Leclerc, Rick Heeres and Shane Frank for testing the package's new functionality ahead of its release. We thank members of the Wildlife Evolutionary Ecology Lab for their comments on previous versions of this manuscript. We thank the rOpenSci organisation including staff, editors, reviewers and contributors for their continued efforts to foster a community around open data, software and reproducibility.

References

1. King AJ, Fehlmann G, Biro D, Ward AJ, Fürtbauer I. Re-wilding Collective Behaviour: An Ecological Perspective. *Trends in Ecology & Evolution*. Elsevier; 2018;33:347–57. <https://doi.org/10.1016/j.tree.2018.03.004>
2. Farine DR, Whitehead H. Constructing, conducting and interpreting animal social network analysis. *Journal of Animal Ecology*. 2015;84:1144–63. <https://doi.org/10.1111/1365-2656.12418>
3. Webber QMR, Albery GFF, Farine DRR, Pinter-Wollman N, Sharma N, Spiegel O, et al. Behavioural ecology at the spatial-social interface. *BIOLOGICAL REVIEWS*. 111 RIVER ST, HOBOKEN 07030-5774, NJ USA: WILEY; 2023;98:868–86. <https://doi.org/10.1111/brev.12934>
4. Barnard CJ, Sibly RM. Producers and scroungers: A general model and its application to captive flocks of house sparrows. *Animal Behaviour*. 1981;29:543–50. [https://doi.org/10.1016/S0003-3472\(81\)80117-0](https://doi.org/10.1016/S0003-3472(81)80117-0)
5. Giraldeau L-A, Lefebvre L. Exchangeable producer and scrounger roles in a captive flock of feral pigeons: A case for the skill pool effect. *Animal Behaviour*. 1986;34:797–803. [https://doi.org/10.1016/S0003-3472\(86\)80064-1](https://doi.org/10.1016/S0003-3472(86)80064-1)
6. Hirsch BT, Malpass E, Di Blanco YE. Interindividual spacing affects the finder's share in ring-tailed coatis (*Nasua nasua*). *Behavioral Ecology*. 2020;31:232–8. <https://doi.org/10.1093/beheco/arz181>
7. Krause J. Differential fitness returns in relation to spatial position in groups. *Biological reviews of the Cambridge Philosophical Society*. 1994;69:187–206. <https://doi.org/10.1111/j.1469-185x.1994.tb01505.x>

8. Mónus F, Barta Z. The Effect of Within-Flock Spatial Position on the Use of Social Foraging Tactics in Free-Living Tree Sparrows. *Ethology*. 2008;114:215–22. <https://doi.org/10.1111/j.1439-0310.2007.01472.x>
9. Furuichi T. Interindividual distance and influence of dominance on feeding in a natural Japanese macaque troop. *Primates*. 1983;24:445–55. <https://doi.org/10.1007/BF02381678>
10. King AJ, Cowlishaw G. Leaders, followers and group decision-making. *Communicative & Integrative Biology*. 2009;2:147–50. <https://doi.org/10.4161/cib.7562>
11. Teichroeb JA, White MMJ, Chapman CA. Vervet (*chlorocebus pygerythrus*) intragroup spatial positioning: Dominants trade-off predation risk for increased food acquisition. *International Journal of Primatology*. 2015;36:154–76. <https://doi.org/10.1007/s10764-015-9818-4>
12. Smith JE, Pinter-Wollman N. Observing the unwatchable: Integrating automated sensing, naturalistic observations and animal social network analysis in the age of big data. *Journal of Animal Ecology*. 2021;90:62–75. <https://doi.org/10.1111/1365-2656.13362>
13. Kays R, Crofoot MC, Jetz W, Wikelski M. Terrestrial animal tracking as an eye on life and planet. *Science*. 2015;348:aaa2478. <https://doi.org/10.1126/science.aaa2478>
14. Kays R, Davidson SC, Berger M, Bohrer G, Fiedler W, Flack A, et al. The Movebank system for studying global animal movement and demography. *Methods in Ecology and Evolution*. 2022;13:419–31. <https://doi.org/10.1111/2041-210X.13767>
15. Nathan R, Monk CT, Arlinghaus R, Adam T, Alós J, Assaf M, et al. Big-data approaches lead to an increased understanding of the ecology of animal movement. *Science*. American Association for the Advancement of Science; 2022;375:eabg1780.
16. Andersson M, Gudmundsson J, Laube P, Wolle T. Reporting Leaders and Followers among Trajectories of Moving Point Objects. *GeoInformatica*. 2008;12:497–528. <https://doi.org/10.1007/s10707-007-0037-9>
17. Dodge S, Weibel R, Lautenschütz A-K. Towards a taxonomy of movement patterns. *Information Visualization*. 2008;7:240–52. <https://doi.org/10.1057/PALGRAVE.IVS.9500182>
18. Laube P. Analysing point motion: Spatio-temporal data mining of geospatial lifelines [PhD thesis]. University of Zurich; 2005.
19. Miller JA. Exploring the utility of movement parameters to make inferences about dynamic interactions in moving objects. *Transactions in GIS*. 2021;25:2701–20. <https://doi.org/10.1111/tgis.12789>
20. Couzin ID, Krause J, James R, Ruxton GD, Franks NR. Collective Memory and Spatial Sorting in Animal Groups. *Journal of Theoretical Biology*. 2002;218:1–11. <https://doi.org/10.1006/jtbi.2002.3065>
21. Herbert-Read JE, Perna A, Mann RP, Schaerf TM, Sumpter DJT, Ward AJW. Inferring the rules of interaction of shoaling fish. *Proceedings of the National Academy of Sciences*.

- Proceedings of the National Academy of Sciences; 2011;108:18726–31. <https://doi.org/10.1073/pnas.1109355108>
22. King AJ, Sueur C. Where Next? Group Coordination and Collective Decision Making by Primates. *International Journal of Primatology*. 2011;32:1245–67. <https://doi.org/10.1007/s10764-011-9526-7>
 23. Nagy M, Ákos Z, Biro D, Vicsek T. Hierarchical group dynamics in pigeon flocks. *Nature*. Nature Publishing Group; 2010;464:890–3. <https://doi.org/10.1038/nature08891>
 24. Strandburg-Peshkin A, Farine DR, Couzin ID, Crofoot MC. Shared decision-making drives collective movement in wild baboons. *Science*. American Association for the Advancement of Science; 2015;348:1358–61. <https://doi.org/10.1126/science.aaa5099>
 25. Hampton SE, Anderson SS, Bagby SC, Gries C, Han X, Hart EM, et al. The Tao of open science for ecology. *Ecosphere*. 2015;6:art120. <https://doi.org/10.1890/ES14-00402.1>
 26. Powers SM, Hampton SE. Open science, reproducibility, and transparency in ecology. *Ecological Applications*. 2019;29:e01822. <https://doi.org/10.1002/eap.1822>
 27. Wilson G, Bryan J, Cranston K, Kitzes J, Nederbragt L, Teal TK. Good enough practices in scientific computing. *PLOS Computational Biology*. Public Library of Science; 2017;13:e1005510. <https://doi.org/10.1371/journal.pcbi.1005510>
 28. Couzin ID, Heins C. Emerging technologies for behavioral research in changing environments. *Trends in Ecology & Evolution*. Elsevier; 2023;38:346–54. <https://doi.org/10.1016/j.tree.2022.11.008>
 29. Robitaille AL, Webber QMR, Vander Wal E. Conducting social network analysis with animal telemetry data: Applications and methods using spatsoc. *Methods in Ecology and Evolution*. 111 RIVER ST, HOBOKEN 07030-5774, NJ USA: WILEY; 2019;10:1203–11. <https://doi.org/10.1111/2041-210X.13215>
 30. Bracken AM, Christensen C, O’Riain MJ, Fürtbauer I, King AJ. Flexible group cohesion and coordination, but robust leader–follower roles, in a wild social primate using urban space. *Proceedings of the Royal Society B: Biological Sciences*. Royal Society; 2022;289:20212141. <https://doi.org/10.1098/rspb.2021.2141>
 31. Boudreau MR, Gantchoff MG, Ramirez-Reyes C, Conlee L, Belant JL, Iglay RB. Using habitat suitability and landscape connectivity in the spatial prioritization of public outreach and management during carnivore recolonization. *Journal of Applied Ecology*. 2022;59:757–67. <https://doi.org/10.1111/1365-2664.14090>
 32. Peignier M, Webber QMR, Koen EL, Laforge MP, Robitaille AL, Vander Wal E. Space use and social association in a gregarious ungulate: Testing the conspecific attraction and resource dispersion hypotheses. *ECOLOGY AND EVOLUTION*. 111 RIVER ST, HOBOKEN 07030-5774, NJ USA: WILEY; 2019;9:5133–45. <https://doi.org/10.1002/ece3.5071>

33. Sunga J, Webber QMR, Broders HG. Influence of number of individuals and observations per individual on a model of community structure. *PLOS ONE*. Public Library of Science; 2021;16:e0252471. <https://doi.org/10.1371/journal.pone.0252471>
34. Keshavarzi H, Lee C, Dyll TR, Johnson M, Campbell DLM. Shared stressful experiences affect social proximity in Merino sheep. *Biology Letters*. 2023;19:20220396. <https://doi.org/10.1098/rsbl.2022.0396>
35. Heeres RW, Leclerc M, Frank S, Kopatz A, Pelletier F, Zedrosser A. Are nonsocial species more social than we think? Seasonal patterns in sociality in a solitary terrestrial carnivore. *Animal Behaviour*. Elsevier; 2024;216:107–30.
36. Joo R, Etienne M-P, Bez N, Mahévas S. Metrics for describing dyadic movement: A review. *Movement Ecology*. 2018;6:26. <https://doi.org/10.1186/s40462-018-0144-2>
37. Collaboration for Environmental Evidence. Guidelines and Standards for Evidence synthesis in Environmental Management. AS Pullin, GK Frampton, B Livoreil, G Petrokofsky, editors. 2022.
38. Veritas Health Innovation. Covidence systematic review software. Melbourne, Australia; 2025.
39. Papageorgiou D, Farine DR. Shared decision-making allows subordinates to lead when dominants monopolize resources. *Science Advances*. 2020;6:eaba5881. <https://doi.org/10.1126/sciadv.aba5881>
40. Allanic M, Hayashi M, Furuichi T, Matsuzawa T. Social influences on grooming site preferences in wild bonobos (*Pan paniscus*) at Wamba, DRC. *Primates*. 2020;61:213–23. <https://doi.org/10.1007/s10329-019-00788-z>
41. Carter KD, Seddon JM, Frère CH, Carter JK, Goldizen AW. Fission–fusion dynamics in wild giraffes may be driven by kinship, spatial overlap and individual social preferences. *Animal Behaviour*. 2013;85:385–94. <https://doi.org/10.1016/j.anbehav.2012.11.011>
42. del Mar Delgado M, Arlettaz R, Bettiga C, Brambilla M, De Gabriel Hernando M, España A, et al. Spatio-temporal variation in the wintering associations of an alpine bird. *Proceedings of the Royal Society B: Biological Sciences*. 2021;288:20210690. <https://doi.org/10.1098/rspb.2021.0690>
43. Bernard A, Krafft B. L’attraction pour la soie : base de la cohésion du groupe et des comportements collectifs chez les araignées sociales. *Comptes Rendus Biologies*. 2002;325:1153–7. [https://doi.org/10.1016/S1631-0691\(02\)01521-4](https://doi.org/10.1016/S1631-0691(02)01521-4)
44. Bernstein IS, Blue SK. Dominance relationships, dominance hierarchies and rankings. *Encyclopedia of Animal Behavior*. Elsevier; 2019. p. 455–60. <https://doi.org/10.1016/B978-0-12-809633-8.20774-7>
45. Long JA, Webb SL, Harju SM, Gee KL. Analyzing Contacts and Behavior from High Frequency Tracking Data Using the wildlifeDI R Package. *Geographical Analysis*. 2022;54:648–63. <https://doi.org/10.1111/gean.12303>

46. Long JA. Quantifying Spatial-temporal Interactions from Wildlife Tracking Data: Issues of Space, Time, and Statistical Significance. *Procedia Environmental Sciences*. 2015;26:3–10. <https://doi.org/10.1016/j.proenv.2015.05.004>
47. Cambon J, Hernangómez D, Belanger C, Possenriede D. Tidygeocoder: An R package for geocoding. *Journal of Open Source Software*. 2021;6:3544. <https://doi.org/10.21105/joss.03544>
48. Chamberlain S, Nitta JH, Sagouis A. Rgnparser: Parse scientific names. 2023. <https://doi.org/10.32614/CRAN.package.rgnparser>
49. Mozzherin D, Marsden T, Ower G, Juillerat P, Pereira HL. GNparser – a powerful scientific names parser. 2026. <https://doi.org/10.5281/zenodo.18221581>
50. Pebesma E. Simple features for R: Standardized support for spatial vector data. *The R Journal*. 2018;10:439–46. <https://doi.org/10.32614/RJ-2018-009>
51. Pebesma E. Lwgeom: Bindings to selected ‘liblwgeom’ functions for simple features. 2025.
52. Dunnington D, Pebesma E, Rubak E. S2: Spherical geometry operators using the S2 geometry library. 2025. <https://doi.org/10.32614/CRAN.package.s2>
53. Nishikawa M, Suzuki M, Sprague DS. Activity and social factors affect cohesion among individuals in female Japanese macaques: A simultaneous focal-follow study. *American Journal of Primatology*. 2014;76:694–703. <https://doi.org/10.1002/ajp.22263>
54. Haydon DT, Morales JM, Yott A, Jenkins DA, Rosatte R, Fryxell JM. Socially informed random walks: Incorporating group dynamics into models of population spread and growth. *Proceedings of the Royal Society Biological Sciences Series B*. 2008;275:1101–9. <https://doi.org/10.1098/rspb.2007.1688>
55. Lesmerises F, Johnson CJ, St-Laurent M-H. Landscape knowledge is an important driver of the fission dynamics of an alpine ungulate. *Animal Behaviour*. 2018;140:39–47. <https://doi.org/10.1016/j.anbehav.2018.03.014>
56. Della Libera K, Strandburg-Peshkin A, Griffith SC, Leu ST. Fission–fusion dynamics in sheep: The influence of resource distribution and temporal activity patterns. *Royal Society Open Science*. 2023;10:230402.
57. Andrienko N, Andrienko G, Barrett L, Dostie M, Henzi P. Space Transformation for Understanding Group Movement. *IEEE Transactions on Visualization and Computer Graphics*. 2013;19:2169–78. <https://doi.org/10.1109/TVCG.2013.193>
58. Pérez-Barbería FJ, Walker DM. Dynamics of social behaviour at parturition in a gregarious ungulate. *Behavioural Processes*. 2018;150:75–84. <https://doi.org/10.1016/j.beproc.2018.01.013>
59. Jolles JW, Laskowski KL, Boogert NJ, Manica A. Repeatable group differences in the collective behaviour of stickleback shoals across ecological contexts. *Proceedings of the Royal Society B: Biological Sciences*. 2018;285:20172629. <https://doi.org/10.1098/rspb.2017.2629>

60. Pettit B, Perna A, Biro D, Sumpter DJT. Interaction rules underlying group decisions in homing pigeons. *Journal of The Royal Society Interface*. Royal Society; 2013;10:20130529. <https://doi.org/10.1098/rsif.2013.0529>
61. Kano F, Sasaki T, Biro D. Collective attention in navigating homing pigeons: Group size effect and individual differences. *Animal Behaviour*. 2021;180:63–80. <https://doi.org/10.1016/j.anbehav.2021.08.004>
62. Quera V, Beltran FS, Gimeno E, Dolado R. Motion leadership and local interaction in two species of freshwater fish (*Danio rerio* and *Hyphessobrycon herbertaxelrodi*). *J Fish Biol*. John Wiley and Sons Inc; 2023;102:856–69. <https://doi.org/10.1111/jfb.15315>
63. Strandburg-Peshkin A, Papageorgiou D, Crofoot MC, Farine DR. Inferring influence and leadership in moving animal groups. *Philosophical Transactions of the Royal Society B: Biological Sciences*. Royal Society; 2018;373:20170006. <https://doi.org/10.1098/rstb.2017.0006>
64. Amornbunchornvej C, Brugere I, Strandburg-Peshkin A, Farine DR, Crofoot MC, Berger-Wolf TY. Coordination Event Detection and Initiator Identification in Time Series Data. *ACM Trans Knowl Discov Data*. 2018;12:53:1–33. <https://doi.org/10.1145/3201406>
65. Albery GF, Becker DJ, Firth JA, De Moor D, Ravindran S, Silk M, et al. Density-dependent network structuring within and across wild animal systems. *Nature Ecology & Evolution*. Nature Publishing Group; 2025;9:2002–13. <https://doi.org/10.1038/s41559-025-02843-z>
66. Merkle JA, Poulin M-P, Caldwell MR, Laforge MP, Scholle AE, Verzuh TL, et al. Spatial-social familiarity complements the spatial-social interface: Evidence from Yellowstone bison. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2024;379:20220530. <https://doi.org/10.1098/rstb.2022.0530>
67. Heesen M, Macdonald S, Ostner J, Schülke O. Ecological and Social Determinants of Group Cohesiveness and Within-Group Spatial Position in Wild Assamese Macaques. *Ethology*. 2015;121:270–83. <https://doi.org/10.1111/eth.12336>
68. Herbert-Read JE, Rosén E, Szorkovszky A, Ioannou CC, Rogell B, Perna A, et al. How predation shapes the social interaction rules of shoaling fish. *Proceedings of the Royal Society B: Biological Sciences*. 2017;284:20171126. <https://doi.org/10.1098/rspb.2017.1126>
69. Jolles JW, Boogert NJ, Sridhar VH, Couzin ID, Manica A. Consistent Individual Differences Drive Collective Behavior and Group Functioning of Schooling Fish. *Current Biology*. 2017;27:2862–2868.e7. <https://doi.org/10.1016/j.cub.2017.08.004>
70. Harel R, Loftus JC, Crofoot MC. Locomotor compromises maintain group cohesion in baboon troops on the move. *Proceedings of the Royal Society B: Biological Sciences*. 2021;288:20210839. <https://doi.org/10.1098/rspb.2021.0839>
71. Wang W, Escobedo R, Sanchez S, Sire C, Han Z, Theraulaz G. The impact of individual perceptual and cognitive factors on collective states in a data-driven fish school model. Gov N, editor. *PLOS Computational Biology*. 2022;18:e1009437. <https://doi.org/10.1371/journal.pcbi.1009437>

72. Vega-Trejo R, Boussard A, Wallander L, Estival E, Buechel SD, Kotrschal A, et al. Artificial selection for schooling behaviour and its effects on associative learning abilities. *Journal of Experimental Biology*. 2020;223:jeb235093. <https://doi.org/10.1242/jeb.235093>
73. Strandburg-Peshkin A, Farine DR, Crofoot MC, Couzin ID. Habitat and social factors shape individual decisions and emergent group structure during baboon collective movement. Carr C, editor. *eLife*. eLife Sciences Publications, Ltd; 2017;6:e19505. <https://doi.org/10.7554/eLife.19505>
74. Lemasson BH, Haefner JW, Bowen MD. Schooling Increases Risk Exposure for Fish Navigating Past Artificial Barriers. *PLOS ONE*. Public Library of Science; 2014;9:e108220. <https://doi.org/10.1371/journal.pone.0108220>
75. Herbert-Read JE. Understanding how animal groups achieve coordinated movement. *Journal of Experimental Biology*. 2016;219:2971–83. <https://doi.org/10.1242/jeb.129411>
76. Torney CJ, Lamont M, Debell L, Angohiatok RJ, Leclerc L-M, Berdahl AM. Inferring the rules of social interaction in migrating caribou. *Philosophical Transactions of the Royal Society B: Biological Sciences*. The Royal Society; 2018;373:20170385.
77. Šárová R, Špinka M, Panamá JLA, Šimeček P. Graded leadership by dominant animals in a herd of female beef cattle on pasture. *Animal Behaviour*. 2010;79:1037–45. <https://doi.org/10.1016/j.anbehav.2010.01.019>
78. Sumpter DJT, Szorkovszky A, Kotrschal A, Kolm N, Herbert-Read JE. Using activity and sociability to characterize collective motion. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 2018;373:20170015. <https://doi.org/10.1098/rstb.2017.0015>
79. Klamser PP, Romanczuk P. Collective predator evasion: Putting the criticality hypothesis to the test. *PLOS Computational Biology*. Public Library of Science; 2021;17:e1008832. <https://doi.org/10.1371/journal.pcbi.1008832>
80. Lukeman R, Li Y-X, Edelstein-Keshet L. Inferring individual rules from collective behavior. *Proceedings of the National Academy of Sciences*. Proceedings of the National Academy of Sciences; 2010;107:12576–80. <https://doi.org/10.1073/pnas.1001763107>
81. Burns ALJ, Herbert-Read JE, Morrell LJ, Ward AJW. Consistency of Leadership in Shoals of Mosquitofish (*Gambusia holbrooki*) in Novel and in Familiar Environments. Chaline N, editor. *PLoS ONE*. 2012;7:e36567. <https://doi.org/10.1371/journal.pone.0036567>
82. Weesner A, Bentley I, Fullerton J, Kloepper L. Interaction rules guiding collective behaviour in echolocating bats. *Anim Behav*. Academic Press; 2023; <https://doi.org/10.1016/j.anbehav.2023.09.009>
83. Josephs N, Bonnell T, Dostie M, Barrett L, Henzi SP. Working the crowd: Sociable vervets benefit by reducing exposure to risk. *Behavioral Ecology*. 2016;27:988–94. <https://doi.org/10.1093/beheco/arw003>

84. Baden AL, Webster TH, Kamilar JM. Resource seasonality and reproduction predict fission-fusion dynamics in black-and-white ruffed lemurs (*Varecia variegata*). *Am J Primatol*. John Wiley and Sons Inc.; 2016;78:256–79. <https://doi.org/10.1002/ajp.22507>
85. Barocas A, Golden HN, Harrington MW, McDonald DB, Ben-David M. Coastal latrine sites as social information hubs and drivers of river otter fission–fusion dynamics. *Animal Behaviour*. 2016;120:103–14. <https://doi.org/10.1016/j.anbehav.2016.07.016>
86. Merkle JA, Sigaud M, Fortin D. To follow or not? How animals in fusion–fission societies handle conflicting information during group decision-making. Grether G, editor. *Ecology Letters*. 2015;18:799–806. <https://doi.org/10.1111/ele.12457>
87. Fortin D, Fortin M-E, Beyer HL, Duchesne T, Courant S, Dancose K. Group-size-mediated habitat selection and group fusion–fission dynamics of bison under predation risk. *Ecology*. 2009;90:2480–90. <https://doi.org/10.1890/08-0345.1>
88. Flack A, Ákos Z, Nagy M, Vicsek T, Biro D. Robustness of flight leadership relations in pigeons. *Animal Behaviour*. 2013;86:723–32. <https://doi.org/10.1016/j.anbehav.2013.07.005>
89. Nagy M, Vásárhelyi G, Pettit B, Roberts-Mariani I, Vicsek T, Biro D. Context-dependent hierarchies in pigeons. *Proceedings of the National Academy of Sciences*. *Proceedings of the National Academy of Sciences*; 2013;110:13049–54. <https://doi.org/10.1073/pnas.1305552110>
90. Ákos Z, Beck R, Nagy M, Vicsek T, Kubinyi E. Leadership and Path Characteristics during Walks Are Linked to Dominance Order and Individual Traits in Dogs. *PLOS Computational Biology*. Public Library of Science; 2014;10:e1003446. <https://doi.org/10.1371/journal.pcbi.1003446>
91. R Core Team. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing; 2025.
92. Beardsworth CE, Gobbens E, van Maarseveen F, Denissen B, Dekinga A, Nathan R, et al. Validating ATLAS: A regional-scale high-throughput tracking system. *Methods in Ecology and Evolution*. 2022;13:1990–2004. <https://doi.org/10.1111/2041-210X.13913>
93. Koger B, Deshpande A, Kerby JT, Graving JM, Costelloe BR, Couzin ID. Quantifying the movement, behaviour and environmental context of group-living animals using drones and computer vision. *Journal of Animal Ecology*. 2023;92:1357–71. <https://doi.org/10.1111/1365-2656.13904>
94. Pedrazzi L, Naik H, Sandbrook C, Lurgi M, Fürtbauer I, King AJ. Advancing animal behaviour research using drone technology. *Animal Behaviour*. 2025;222:123147. <https://doi.org/10.1016/j.anbehav.2025.123147>
95. Vickery WL, Giraldeau L-A, Templeton JJ, Kramer DL, Chapman CA. Producers, scroungers, and group foraging. *The American Naturalist*. 1991;137:847–63. <https://doi.org/10.1086/285197>
96. Giraldeau L-A, Caraco T. *Social foraging theory*. Princeton University Press; 2000.

97. Hansen M, Ward A, Fürtbauer I, King A. Environmental quality determines finder-joiner dynamics in socially foraging three-spined sticklebacks (*Gasterosteus aculeatus*). *BEHAVIORAL ECOLOGY AND SOCIOBIOLOGY*. 2016;70:889–99. <https://doi.org/10.1007/s00265-016-2111-5>
98. Flack A, Nagy M, Fiedler W, Couzin ID, Wikelski M. From local collective behavior to global migratory patterns in white storks. *Science*. American Association for the Advancement of Science; 2018;360:911–4.
99. Caraco T, Barkan C, Beacham JL, Brisbin L, Lima S, Mohan A, et al. Dominance and social foraging: A laboratory study. *Animal Behaviour*. 1989;38:41–58. [https://doi.org/10.1016/S0003-3472\(89\)80064-8](https://doi.org/10.1016/S0003-3472(89)80064-8)
100. Aplin LM, Morand-Ferron J. Stable producer–scrounger dynamics in wild birds: Sociability and learning speed covary with scrounging behaviour. *Proceedings of the Royal Society B: Biological Sciences*. Royal Society; 2017;284:20162872. <https://doi.org/10.1098/rspb.2016.2872>
101. Lee AGE, Ounsley JP, Coulson T, Bullock JM, Cowlshaw G. Information use and resource competition: An integrative framework. *Proceedings of the Royal Society B: Biological Sciences*. 2016;283:20152550. <https://doi.org/10.1098/rspb.2015.2550>
102. Evans JC, Devost I, Jones TB, Morand-Ferron J. Inferring dominance interactions from automatically recorded temporal data. Hebets E, editor. *Ethology*. 2018;124:188–95. <https://doi.org/10.1111/eth.12720>
103. Ramos A, Bousquet CAH, Sueur C. How leadership could be used to manage domestic and wild ungulate herds. *Applied Animal Behaviour Science*. 2021;239:105326. <https://doi.org/10.1016/j.applanim.2021.105326>
104. Landau WM. The targets R package: A dynamic Make-like function-oriented pipeline toolkit for reproducibility and high-performance computing. *Journal of Open Source Software*. 2021;6:2959.
105. Wickham H. *Ggplot2: Elegant graphics for data analysis*. Springer-Verlag New York; 2016.
106. Kay M. ggdist: Visualizations of distributions and uncertainty. 2025. <https://doi.org/10.5281/zenodo.3879620>
107. Pedersen TL. Patchwork: The composer of plots. 2025. <https://doi.org/10.32614/CRAN.package.patchwork>
108. Arel-Bundock V. Tinytable: Simple and configurable tables in ‘HTML’, ‘LaTeX’, ‘markdown’, ‘word’, ‘PNG’, ‘PDF’, and ‘typst’ formats. 2025. <https://doi.org/10.32614/CRAN.package.tinytable>
109. Allaire J, Dervieux C. Quarto: R interface to ‘quarto’ markdown publishing system. 2024. <https://doi.org/10.32614/CRAN.package.quarto>
110. Calenge C. adehabitatHR: Home range estimation. 2024. <https://doi.org/10.32614/CRAN.package.adehabitatHR>

111. Barrett T, Dowle M, Srinivasan A, Gorecki J, Chirico M, Hocking T, et al. Data.table: Extension of 'data.frame'. 2025. <https://doi.org/10.32614/CRAN.package.data.table>
112. Csárdi G, Nepusz T, Traag V, Horvát S, Zanini F, Noom D, et al. igraph: Network analysis and visualization in R. 2026. <https://doi.org/10.5281/zenodo.7682609>
113. Lund U, Agostinelli C. CircStats: Circular statistics, from "topics in circular statistics" (2001). 2025. <https://doi.org/10.32614/CRAN.package.CircStats>
114. Pebesma E, Mailund T, Hiebert J. Measurement units in R. R Journal. 2016;8:486–94. <https://doi.org/10.32614/RJ-2016-061>
115. Henry L, Wickham H. Rlang: Functions for base types and core R and 'tidyverse' features. 2026. <https://doi.org/10.32614/CRAN.package.rlang>
116. Amornbunchornvej C. mFLICA: An R package for inferring leadership of coordination from time series. SoftwareX. 2021;15:100781. <https://doi.org/10.1016/j.softx.2021.100781>
117. Papadopoulou M, Garnier S, King AJ. swaRmverse: An R package for the comparative analysis of collective motion. Methods in Ecology and Evolution. 2025;16:29–39. <https://doi.org/10.1111/2041-210X.14460>

10 Supplement: Scoping review methods

Table S 1: Benchmark papers used to evaluate iterative search results for scoping review

Citation
Nagy, Máté, Zsuzsa Ákos, Dora Biro, and Tamás Vicsek. 2010. “Hierarchical Group Dynamics in Pigeon Flocks.” <i>Nature</i> 464 (7290): 890–93. https://doi.org/10.1038/nature08891.
Strandburg-Peshkin, Ariana, Damien R. Farine, Iain D. Couzin, and Margaret C. Crofoot. 2015. “Shared Decision-Making Drives Collective Movement in Wild Baboons.” <i>Science</i> 348 (6241): 1358–61.
Merkle, J.A., M. Sigaud, and D. Fortin. 2015. “To Follow or Not? How Animals in Fusion–Fission Societies Handle Conflicting Information during Group Decision-Making.” <i>Ecology Letters</i> 18 (8): 799–806.
Andersson, Mattias, Joachim Gudmundsson, Patrick Laube, and Thomas Wolle. 2007. “Reporting Leadership Patterns among Trajectories.” In <i>Proceedings of the 2007 ACM Symposium on Applied Computing</i> , 3–7. SAC ’07. New York, NY, USA: Association for Computing Machinery. https://doi.org/10.1145/1244002.1244004.
Flack, Andrea, Máté Nagy, Wolfgang Fiedler, Iain D. Couzin, and Martin Wikelski. 2018. “From Local Collective Behavior to Global Migratory Patterns in White Storks.” <i>Science</i> 360 (6391): 911–14.
Bracken, A.M., C. Christensen, M.J. O’Riain, I. Fürtbauer, and A.J. King. 2022. “Flexible Group Cohesion and Coordination, but Robust Leader-Follower Roles, in a Wild Social Primate Using Urban Space.” <i>Proc. R. Soc. B Biol. Sci.</i> 289 (1967). https://doi.org/10.1098/rspb.2021.2141.
Long, Jed A., Trisalyn A. Nelson, Stephen L. Webb, and Kenneth L. Gee. 2014. “A Critical Examination of Indices of Dynamic Interaction for Wildlife Telemetry Studies.” <i>Journal of Animal Ecology</i> 83 (5): 1216–33. https://doi.org/10.1111/1365-2656.12198.
Dodge, Somayeh, Robert Weibel, and Anna-Katharina Lautenschütz. 2008. “Towards a Taxonomy of Movement Patterns.” <i>Information Visualization</i> 7 (3–4): 240–52. https://doi.org/10.1057/PALGRAVE.IVS.9500182.
Laube, Patrick, Stephan Imfeld, and Robert Weibel. 2005. “Discovering Relative Motion Patterns in Groups of Moving Point Objects.” <i>International Journal of Geographical Information Science</i> 19 (6): 639–68. https://doi.org/10.1080/13658810500105572.

Table S 2: Search strings for scoping review using Web of Science and Scopus

Source	String
Web of Science	TS=(“dominance hierarch*” OR “leadership” OR “dominance” OR “collective move*” OR “fusion-fission” OR “fission-fusion” OR (“fission*” AND “fusion*”) OR (“producer*” AND “scrounger”) OR “producer-scrounger” OR “leader-follower” OR “leader follower” OR (“leader*” AND “follower*”) OR (“finder*” AND “joiner*”) OR “group cohesion” OR “dynamic interaction” OR “leadership pattern*” OR “collective behav*” OR “spatiotemporal group*” OR “relative motion” OR “group dynamics”) AND TS=(“ecolog*” OR “animal*” OR “wildlife”) AND TS=((“spatial*” AND “network*”) OR “track*” OR “gps” OR “telemetry” OR “PIT” OR “video” OR “RFID” OR “biotelem*” OR “spatiotemporal” OR “spatio-temporal” OR “GPS” OR “movement pattern*”) NOT TS=(“human group*” OR “robot*” OR “genetic algorithm” OR “cell movement” OR “cellular” OR “sport*” OR “team sport*” OR “task force” OR “food web*” OR “host plant*” OR “fossil” OR “urbanism” OR “psychiatry” OR “bile”) AND PY=(1980-2023)
Scopus	TITLE-ABS-KEY (“dominance hierarch*” OR “leadership” OR “dominance” OR “collective move*” OR “fusion-fission” OR “fission-fusion” OR (“fission*” AND “fusion*”) OR (“producer*” AND “scrounger”) OR “producer-scrounger” OR “leader-follower” OR “leader follower” OR (“leader*” AND “follower*”) OR (“finder*” AND “joiner*”) OR “group cohesion” OR “dynamic interaction” OR “leadership pattern*” OR “collective behav*” OR “spatiotemporal group*” OR “relative motion” OR “group dynamics”) AND TITLE-ABS-KEY (“ecolog*” OR “animal*” OR “wildlife”) AND TITLE-ABS-KEY ((“spatial*” AND “network*”) OR “track*” OR “gps” OR “telemetry” OR “PIT” OR “video” OR “RFID” OR “biotelem*” OR “spatiotemporal” OR “spatio-temporal” OR “GPS” OR “movement pattern*”) AND NOT TITLE-ABS-KEY (“human group*” OR “robot*” OR “genetic algorithm” OR “cell movement” OR “cellular” OR “sport*” OR “team sport*” OR “task force” OR “food web*” OR “host plant*” OR “fossil” OR “urbanism” OR “psychiatry” OR “bile”) AND PUBYEAR > 1980

Table S 3: Data extraction form for scoping review to identify metrics for studying intragroup dynamics

Category	Field
General	Covidence #
	Study ID
	Title
Study information	Type of study (Empirical, simulation, algorithm, model, review, other)
	Species
	Region
Dyadic social metrics	Metric used or described
	Software package(s) used
	Citation for metric if not introduced in this paper
	Interpretation
	Data input required
	Definitions of “leader”, “initiator”, etc
Dominance	Dominance metric
	Interpretation
	Citation for dominance metric if not introduced in this paper
Code availability	Analysis code availability
	Programming language
	Analysis code link
	Analysis code availability location (e.g. GitHub, Zenodo, FigShare)
	Other comments about code availability
Misc	Discussion points
	Etc

11 Supplement: Scoping review results

Results

PRISMA

- 3870 references imported for screening as 3870 studies
 - 128 duplicates identified manually
 - 799 duplicates identified by Covidence
- 2943 studies screened against title and abstract

- 2527 studies excluded
- 416 studies assessed for full-text eligibility
 - 275 studies excluded
 - 184 Does not discuss dyadic measure
 - 86 Metric not applicable to spatiotemporal data
 - 2 Book chapter
 - 2 Full text not in English
 - 1 Thesis
 - 0 studies ongoing
 - 0 studies awaiting classification
- 141 studies included

Table S 4: Count of code availability in references screened for full text extraction of scoping review

Code availability	N
No	115 (82%)
Yes	26 (18%)

Out of 141 articles, 52 did not list a programming language or GUI program. Of the 89 that did, 21 articles listed more than one programming language or GUI program, therefore the total count below will be greater than the number of articles.

Table S 5: Count of programming language or GUI program used in references screened for full text extraction of scoping review

Programming language	N
R	62 (36.26%)
NA	52 (30.41%)
MATLAB	15 (8.77%)
Python	6 (3.51%)
CUDA	4 (2.34%)
Ucinet	3 (1.75%)
Perl	3 (1.75%)
C++	3 (1.75%)
NetLogo	2 (1.17%)
SPSS	2 (1.17%)
SchoolTracker	1 (0.58%)
Bonsai	1 (0.58%)
Keras	1 (0.58%)
TensorFlow	1 (0.58%)
SYSTAT	1 (0.58%)
Fortran	1 (0.58%)
Mathematica	1 (0.58%)
Brodgar	1 (0.58%)
Stata	1 (0.58%)
ArcGIS	1 (0.58%)
Pascal	1 (0.58%)
Matlab	1 (0.58%)
Hawth Tools for ArcGIS	1 (0.58%)
Wolfram Mathematica	1 (0.58%)
Ctrax	1 (0.58%)
StreamPix 5	1 (0.58%)
MoveMine	1 (0.58%)
Java	1 (0.58%)
Netlogo	1 (0.58%)

Out of 141 articles, 113 did not list a software package. Of the 28 that did, 14 articles listed more than one software package , therefore the total count below will be greater than the number of articles.

Table S 6: Count of software packages used in references screened for full text extraction of scoping review

(a)

Software package used	N
NA	113 (66.86%)
wildlifeDI	9 (5.33%)
geosphere	3 (1.78%)
adehabitat	2 (1.18%)
moveVis	2 (1.18%)
ctmm	2 (1.18%)
CircStats	2 (1.18%)
corrMove	2 (1.18%)
adehabitatHR	2 (1.18%)
numpy	2 (1.18%)
h5py	2 (1.18%)
matplotlib	2 (1.18%)
pathlib	2 (1.18%)
swaRm	1 (0.59%)
forecast (cross correlation)	1 (0.59%)
move	1 (0.59%)
moveUD	1 (0.59%)
boot	1 (0.59%)
MASS	1 (0.59%)
BBMM	1 (0.59%)
momentuHMM	1 (0.59%)
adehabitatLT	1 (0.59%)
DescTools	1 (0.59%)
splancs	1 (0.59%)
vegan	1 (0.59%)
EloRating	1 (0.59%)
compete	1 (0.59%)
changepoint	1 (0.59%)
igraph	1 (0.59%)
rethinking	1 (0.59%)
pdist	1 (0.59%)
EDM4WR	1 (0.59%)

Table S 7: Count of references screened for full text extraction across temporal period (1980-2023) of scoping review




Table S 8: Map of study regions georeferenced using {tidygeocoder} from references screened for full text extraction of scoping review




Table S 9: Count of species parsed using {rgnparser} in references screened for full text extraction of scoping review

Species	N
<i>Columba livia</i>	9 (7.09%)
<i>Danio rerio</i>	6 (4.72%)
<i>Rangifer tarandus</i>	6 (4.72%)
<i>Gasterosteus aculeatus</i>	5 (3.94%)
<i>Papio anubis</i>	5 (3.94%)
<i>Poecilia reticulata</i>	5 (3.94%)
<i>Macaca fuscata</i>	4 (3.15%)
<i>Ovis aries</i>	4 (3.15%)
<i>Bos taurus</i>	3 (2.36%)
<i>Ateles geoffroy</i>	3 (2.36%)
<i>Equus ferus</i>	3 (2.36%)
<i>Hemigrammus rhodostomus</i>	3 (2.36%)
<i>Loxodonta africana</i>	3 (2.36%)
<i>Odocoileus virginianus</i>	3 (2.36%)
<i>Bison bison</i>	2 (1.57%)
<i>Capra aegagrus</i>	2 (1.57%)
<i>Cervus elaphus</i>	2 (1.57%)
<i>Ciconia ciconia</i>	2 (1.57%)
<i>Equus hemionus</i>	2 (1.57%)
<i>Homo sapiens</i>	2 (1.57%)
<i>Panthera pardus</i>	2 (1.57%)
<i>Varecia variegata</i>	2 (1.57%)
<i>Acinonyx jubatus</i>	1 (0.79%)
<i>Acryllium vulturinum</i>	1 (0.79%)
<i>Ailurus fulgens</i>	1 (0.79%)
<i>Anser anser</i>	1 (0.79%)
<i>Astyanax mexicanus</i>	1 (0.79%)
<i>Aves NA</i>	1 (0.79%)
<i>Callisaurus draconoides</i>	1 (0.79%)
<i>Canis familiaris</i>	1 (0.79%)
<i>Canis lupus</i>	1 (0.79%)
<i>Canis lupus familiaris</i>	1 (0.79%)

Table S 10: Count of metrics used to measure intragroup dynamics in references screened for full text extraction of scoping review

Metric	N
interindividual distance	61 (14.66%)
nearest neighbour distance	19 (4.57%)
directional correlation delay	18 (4.33%)
speed	17 (4.09%)
position within group	16 (3.85%)
direction	15 (3.61%)
fission fusion	15 (3.61%)
zones	15 (3.61%)
distance to group centroid	14 (3.37%)
directional alignment	9 (2.16%)
DI	7 (1.68%)
group centroid	7 (1.68%)
polarization	7 (1.68%)
Don	6 (1.44%)
Prox	6 (1.44%)
velocity	5 (1.20%)
voronoi	5 (1.20%)
group velocity	5 (1.20%)
rank position within group	5 (1.20%)
direction to group centroid	5 (1.20%)
group size	5 (1.20%)
distance	5 (1.20%)
group direction	5 (1.20%)
acceleration	4 (0.96%)
time spent leading group	4 (0.96%)
Lixn	4 (0.96%)
interindividual direction	3 (0.72%)
HAI	3 (0.72%)
flock	3 (0.72%)
trend setting	3 (0.72%)
group speed	3 (0.72%)
total	45 (10.22%)

Table S 11: Count of metrics used to measure dominance in references screened for full text extraction of scoping review

Dominance metric	N
NA	112 (77.24%)
focal observations	14 (9.66%)
direction correlation delay	5 (3.45%)
fleeing events	2 (1.38%)
survey	2 (1.38%)
approach-avoidance events	2 (1.38%)
egalitarian leadership index	1 (0.69%)
movement initiations	1 (0.69%)
proportion of paired flights led	1 (0.69%)
spatially constrained trend-setter	1 (0.69%)
video observations	1 (0.69%)
displacement events	1 (0.69%)
directional consistency index	1 (0.69%)
feeding-queuing events	1 (0.69%)

Table S 12: Review of definitions of spatiotemporal groups in references screened for full text extraction of scoping review

study_id	definition
Quaglietta 2014	interaction if within 100 m and 1 hour
Mielke 2020	group defined as all individuals within visual contact of focal individual
Leighty 2008	proximity defined as 8 m; approach/avoidance after event defined as changes in interindividual distance after an event
Hansen 2015	fish considered isolated if not within three body lengths of any other fish; group defined by within three body lengths
Bista 2022	temporal threshold 2 hour, distance threshold 100 m
Pinacho-Guendulain 2017	group using 30 m chain rule
Lemasson 2014	group when within 5 body lengths
Inoue 2019	proximity defined as two body lengths away

Table S 13: Review of definitions of leadership in references screened for full text extraction of scoping review

study_id	definition
Rahman 2020	leadership as tendency to be at the front of the group
Quera 2023	following Krause et al. (2000) and Collignon et al. (2019), the authors define a motion leader as an individual who initiates movement towards a certain direction and is followed by other group members
Andrienko 2013	leader is an object that moves in front of others; trend setter is an object whose movements are copied by others
Amornbunchornvej 2018	following relation is where two individuals perform the same sequence of actions with some fixed delay; initiator is individual who first performs a sequence of actions and all other individuals follow
Kano 2021	leader from directional correlation delay
Pettit 2013	leadership based on directional correlation delay; leadership also based on tendency to use route taken on solo flight
Pérez-Barbería 2018	leadership is ranked position along front-back axis
Laube 2005	leadership extends flock with constance over previous time steps
Milner 2021	leader is animal that is dominant to at least one and subordinate to none
Strandburg-Peshkin 2018	an individual leads if it has repeated influence either directly or hierarchically on the behaviour of others
Andersson 2008	leader if and only if individual doesnt follow anyone and is followed by N others
Jolles 2017	leader in terms of propagation of movement changes in the group
King 2015	leader-follower dynamics from velocity-cross correlation
Bracken 2022	leader initiates movement away from another individual (follower) who joins or who doesn't and leader returns
Flack 2018	leaders are birds who are ahead of the flock on average and followers those who are behind
ACM 2007	leader IFF x does not follow anyone and is followed by N others; leadership pattern if x is followed by minimum M others for K time unit intervals
Jolles 2018	measure of leadership as proportion of time spent in front of group centroid

Table S 14: Review of definitions of fission-fusion in references screened for full text extraction of scoping review

study_id	definition
Baden 2016	NA
Aguilar-Melo 2018	fission when greater than or equal to 1 individual were not observed with a group for 2 consecutive scans; fusion when greater than or equal to 1 who were previously not observed with group are observed with group for 2 consecutive scans
Barocas 2016	fission and fusion defined as joining and leaving group in subsequent observations
Haydon 2008	fusion where solitary to grouped and fission where grouped to solitary
Fortin 2009	group/fusion when within 100 m; fission when two or more consecutive locations separated by > 100 m
Body 2015	fusion greater than or equal to 2 groups merging into one; fission one group splitting into greater than or equal to 2
DellaLibera 2023	define a fission and fusion event if one or more individuals left or joined the group, respectively. in the rare cases of missing data, missing individuals are not considered as having changed group membership, and individual 'disappearances' and 'reappearances' in the dataset are not considered as fission-fusion events
Wielgus 2020	group where simultaneously within 1 km or ≥ 1 km for ≤ 2 hr; fusion where together then different by one time step; fission reverse
Nishikawa 2014	converged when IID less than 20 m; separated when IID greater than 20 m; converged and separated synonymous with fission fusion
Krueger 2014	NA
Lesmerises 2018	grouped/fission fusion if within 50 m for two relocations; to minimize the false fission events, considered dyad valid if together at t0, t2, but apart at t1; if one missing GPS location, dyad valid if before and after missing loc distance was < 50 m
Merkle 2015	group defined as within 100 m within 24 hours

12 Supplement: Selected metrics

12.1 Selected metrics

The following metrics were not selected due to their availability in existing R packages:

- {wildlifeDI}: DI, Don, Prox, Lixn, Cs, DI in direction, DI in displacement, HAI, IAB, Cr, Ca

- {corrMove}: diffusive movement correlation, drift movement correlation, overall component movement correlation
- {changePoint}: change point analysis

The following metrics were not selected due to their availability in code accompanying the publication:

- mFLICA provided in code accompanying [64] and R package {mFLICA} [116]
- StickyDBSCAN provided in code accompanying [56]

The follow metrics were not selected due to the expected input data differing from existing {spatsoc} functions (XYT):

- all metrics relating to posture

The selected metrics are listed in Table 1. All existing functions in {spatsoc} are listed below in Table S 15.

Table S 15: Description of all functions (new and previously existing) available in the R package {spatsoc} categorized by family.

Function	Description	Category
Spatial interface		
get_geometry()	Setup data for geometry interface	New
Edge-list generation		
edge_dist()	Distance based edge-lists	Existing
edge_nn()	Nearest neighbour based edge-lists	Existing
edge_delay()	Directional correlation delay based edge-lists	New
edge_alignment()	Directional alignment based edge-lists	New
edge_direction()	Direction based edge-lists	New
edge_zones()	Behavioural zones based edge-lists	New
Dyad		
dyad_id()	Identify dyads	Existing
fusion_id()	Identify fusion events	New
Centroid		
centroid_group()	Group centroid	New
centroid_dyad()	Dyad centroid	New
centroid_fusion()	Fusion event centroid	New
Direction		
direction_step()	Direction step	New
direction_to_centroid()	Direction to group centroid	New
direction_to_leader()	Direction to group leader	New
direction_group()	Group mean direction	New
direction_polarization()	Polarization	New
Distance		
distance_to_centroid()	Distance to group centroid	New
distance_to_leader()	Distance to group leader	New
Leadership		
leader_direction_group()	Leadership along group direction	New
leader_edge_delay()	Leadership in directional correlation delay	New
Temporal grouping		
group_times()	Group times	Existing
Spatial grouping		
group_by_h1n1_h3n2()	Group by influenza A/H1N1 and A/H3N2	Existing

Distance, direction and rank distance to group centroid

The scoping review found 21 studies that used the group centroid, distance to group centroid, direction to group centroid, or rank distance to group centroid metrics, across 14 study species. 4 studies had code available. The {swaRm} package provides functionality to calculate the group centroid and the distance to the group centroid. At the time when the scoping review was conducted, to the best of our knowledge, there were no R packages that provide the rank to distance to group centroid metric.

Polarization, directional alignment and interindividual direction

The scoping review found 20 studies that used the polarization, directional alignment or interindividual direction metrics, across 11 study species. 4 studies had code available. The {CircStats} and the {swaRm} packages provide functionality to calculate the polarization order parameter. At the time when the scoping review was conducted, to the best of our knowledge, there were no R packages that provide the directional alignment or interindividual direction metrics. After the scoping review was conducted, the {swaRmverse} package was released which provides the direction to nearest neighbour metric [117].

Behavioural zones

The scoping review found 13 studies that used the behavioural zones metric, across 9 study species. 4 studies had code available. At the time when the scoping review was conducted, to the best of our knowledge, there were no R packages that provide the behavioural zones metric.

Position within group, rank position within group, distance and direction to leader

The scoping review found 18 studies that used the position within group, rank position within group, distance to leader, or direction to leader metrics across 14 study species. Only 1 study had code available. At the time when the scoping review was conducted, to the best of our knowledge, there were no R packages that provide these metrics.

Directional correlation delay

The scoping review found 17 studies that used the directional correlation delay metric, across 12 study species. Only 1 study had code available using a combination of MATLAB (closed source), CUDA (closed source), and Python (open source). At the time when the scoping review was conducted, to the best of our knowledge, there were no R packages that provide this metric.