

A Protocol for Standardizing Measurements and Enabling Global Harmonization of Herbarium
Leaf Reflectance Spectra

**Dawson M. White,^{1,2,21} Flavia M. Durgante,³ Matthew W. Austin,⁴ J. Antonio Guzmán Q.,²
Jonathan A. Kennedy,¹ Cornelius Onyedikachi Nichodemus,⁵ Natalie I. Ahlstrand,⁶ Chris
Baraloto,⁷ Billur Bektaş,⁸ Khalil Boughalmi,⁹ Warren Cardinal-McTeague,¹⁰ Thomas L.P.
Couvreur,⁹ Riley P. Fortier,⁴ Ryan S. Fuller,⁴ Olwen M. Grace,¹¹ Rykkar Jackson,¹⁰ Kevin
Karbstein,^{12,13} Shan Kothari,¹⁴ Aaron K. Lee,¹⁵ Jéssica Lira Viana,¹⁶ Elizabeth Martinez-
Victoria,² José Eduardo Meireles,⁵ Thomas H. Murphy,⁷ Carlos Rodrigues-Vaz^{9,17}, Daniel
Spalink,¹⁸ Hanna Tuomisto,^{16,19} Caroline C. Vasconcelos,^{3,20} Jeannine Cavender-Bares^{1,2,21}**

¹Harvard University Herbaria, 22 Divinity Ave., Cambridge, MA 02138 USA

²Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA
02138, USA.

³Herbarium, National Institute of Amazonian Research, Manaus, AM 69067-375, Brazil

⁴Herbarium, Missouri Botanical Garden, 4344 Shaw Blvd., St. Louis, MO 63110 USA

⁵School of Biology and Ecology, University of Maine, Orono, ME 04469, USA

⁶Natural History Museum Denmark, University of Copenhagen, Copenhagen, 1354, Denmark

⁷International Center for Tropical Botany, Florida International University, Miami, FL 33133,
USA

⁸Department of Environmental Systems Science, Institute of Integrative Biology, ETH Zurich,
8092, Zurich, Switzerland

⁹DIADE, Univ Montpellier, CIRAD, IRD, Montpellier, France

¹⁰Department of Forest and Conservation Sciences, University of British Columbia, 2424 Main
Mall, Vancouver BC, V6T 1Z4, Canada

¹¹Royal Botanic Garden Edinburgh, 20A Inverleith Row, Edinburgh EH3 5LR, United Kingdom

¹²Professorship of Integrative Plant Taxonomy, Friedrich Schiller University of Jena, Jena,
Germany

¹³Department of Natural History, Florida Museum of Natural History, Gainesville, FL 32611,
USA

¹⁴Department of Renewable Resources, University of Alberta, Edmonton, AB T6G 2E3, Canada

¹⁵Department of Plant and Microbial Biology, University of Minnesota, St. Paul, MN 55108,
USA

¹⁶Department of Biology, Aarhus University, Aarhus, 8000, Denmark

¹⁷Muséum National d'Histoire Naturelle, Institut de Systématique, Évolution, et Biodiversité
(ISYEB), Centre National de la Recherche Scientifique, Sorbonne Université, École Pratique des
Hautes Études-Paris, Sciences et Lettres, Université des Antilles, C.P. 39, 57 rue Cuvier, 75005
Paris, France.

¹⁸S.M. Tracy Herbarium, Department of Ecology and Conservation Biology, Texas A&M
University, College Station, TX 77843, USA

¹⁹Department of Biology, University of Turku, 20014 Turku, Finland

²⁰Department of Forest Sciences, Federal University of Amazonas, Manaus, AM 69080-900
Brazil

²¹Authors for correspondence (dawson.white@gmail.com; jcavender@fas.harvard.edu)

Abstract—Reflectance spectroscopy offers a powerful approach to integrate high-throughput phenotypic data from herbarium specimens into the digital landscapes of ecology, evolution, and systematics. Spectral digitization complements specimen metadata capture and photographic imaging by generating high-dimensional optical data from herbarium tissues that encode information about plant chemistry and structure. Because inconsistencies in instrumentation and measurement practices can increase noise and limit dataset compatibility, the International Herbarium Spectral Digitization Working Group (IHerbSpec) has published the *Protocol for Spectral Digitization of Herbarium Specimens*, currently in version 1.3, as an open resource that will continue to evolve through community use and collaboration (<https://iherbspec.github.io/protocol>). The protocol defines a stepwise measurement workflow, standardized filename conventions, structured metadata tables with controlled vocabularies, and guidance on tissue selection, materials, and instrumentation quality control to support spectral datasets that are interoperable across projects and reusable through well-documented metadata. By embedding standardized practices at the point of data collection, it provides a scalable foundation for data synthesis and new quantitative insights into plant diversity across taxonomic, geographic, and temporal scales.

Keywords—Data harmonization, data standards, digital extended specimen, FAIR principles, spectral fingerprint, phenomics, near-infrared spectroscopy, NIRS

The more than 400 million botanical collections stored in global herbaria represent a vast and irreplaceable repository of plant diversity, together with rich environmental and ecological metadata (Davis, 2023; Thiers, 2025). These specimens underpin our understanding of organismal morphology, taxonomy, and biogeography, and are increasingly used to generate derived genetic and phenotypic datasets that expand their scientific impact and potential to address societal challenges (Lendemer et al., 2020; Kunkel et al., 2025).

Digitization of herbarium collections in recent decades has centered on creating digital specimen records through whole-specimen imaging, label transcription, georeferencing, and databasing. These efforts have laid the foundation for a global metaherbarium that makes specimens broadly discoverable and accessible for research, expanding both the scale of traditional herbarium-based studies and the range of stakeholders and scientific domains in which specimens are used (Davis, 2023; Antonelli et al., 2026). Spectral digitization, by contrast, complements these efforts by capturing reflectance spectra from specimens as a distinct product: a spectral fingerprint of plant tissues. Rather than documenting specimen appearance, reflectance spectra record high-dimensional signals associated with tissue morphology, creating quantitative phenomic datasets that can be analyzed computationally as part of the digital extended specimen (Hardisty et al., 2022; Cavender-Bares et al., 2026) and integrated with other specimen-derived datasets (Neto-Bradley et al., 2026). Spectral digitization can be integrated into other digitization workflows or carried out separately. Indeed, several collaborating institutions have found it preferable to perform spectral measurements after specimen databasing has been completed.

Reflectance spectroscopy—here referring to point measurements of plant organs made using a contact probe (Fig. 1b)—was developed largely through studies of fresh leaf tissues and is now well established as a high-throughput means of linking phenotype, environment, and

evolutionary history at different scales (Serbin et al., 2014; Cavender-Bares et al., 2016, 2025; Vance et al., 2016; Kothari and Schweiger, 2022; Kothari et al., 2023; Stefanski et al., 2025). Spectral measurements across the visible, near-, and shortwave infrared regions (350–2500 nm) capture integrated optical signals related to tissue structure and chemistry (Curran, 1989; Jacquemoud et al., 1996). As a result, reflectance spectra provide a high-dimensional data source for modeling taxonomy, plant traits, and biological processes that could otherwise require substantially more time-intensive, costly, and often destructive wet-lab measurements. Because a single standardized spectral measurement can be reused across multiple analytical frameworks, it can support both current applications and future analyses as predictive models continue to improve.

Leaf reflectance spectra have demonstrated utility in discriminating taxa and cytotypes (Abasolo et al., 2013; Durgante et al., 2013; Lang et al., 2017; Stasinski et al., 2021; Buono and Albach, 2023; Vasconcelos et al., 2025), estimating leaf traits (Costa et al., 2018; Kothari et al., 2023), and revealing adaptive differentiation and evolutionary processes (Cavender-Bares et al., 2016; Meireles et al., 2020; Fine et al., 2021; Hernández-Leal et al., 2025). Importantly, multiple studies have now demonstrated that these same applications extend to herbarium specimens that are decades to centuries old (Kühn et al., 2024; Boughalmi et al., 2025; Lee et al., 2026; Neto-Bradley et al., 2026; White et al., 2026). These advances have signaled herbarium spectral digitization as a robust approach for phenotyping plant biodiversity at unprecedented spatial and temporal scales; thus expanding the relevance of collections to broader sectors of biology, and science generally, and the available funding sources for biological collections. For a recent review and perspective on use cases justifying the value of herbarium spectral digitization, see Cavender-Bares et al. (2026).

A core strength of herbarium spectral digitization lies in the opportunity to represent plant phenotypic diversity across regions, taxa, and time in harmonized datasets generated across herbaria (Cavender-Bares et al., 2026). Yet, coordinated standards for data collection, metadata, and databasing are needed to ensure that spectral datasets remain findable, accessible, interoperable, and reusable, rather than becoming fragmented, poorly documented, difficult to interpret, or excessively noisy due to undocumented differences among measurement setups (Wilkinson et al., 2016). Comparable challenges in plant trait research have been addressed through standardized measurement and databasing protocols (Kattge et al., 2011; Enquist et al., 2016; Pérez-Harguindeguy et al., 2016).

To document botanical variation present in herbaria through spectral measurements—and to ensure that these data are interpretable and reusable—standards must explicitly capture both the measurement environment and the characteristics of the specimens themselves to enable assessments of data quality. Spectral reflectance measurements do not represent a single, invariant biological signal comparable to a DNA sequence (Cruickshank and Munck, 2011). Instead, they integrate the molecular and structural properties of all the materials that interact with incident light, and this includes not only the target plant tissue but also the background material and potential contaminants, such as mounting materials or epiphylls. This complexity is compounded by variation among herbarium tissues in age, preservation history, mounting technique, and degree of degradation (Kühn et al., 2024; Cavender-Bares et al., 2026; White et al., 2026).

To address these challenges at an early stage of data generation, we established the International Herbarium Spectral Digitization Working Group (IHerbSpec), a global consortium dedicated to advancing best practices for herbarium-based reflectance spectroscopy and fostering coordinated, cross-institutional collaboration.

Here, we present the *Protocol for the Spectral Digitization of Herbarium Specimens* (the IHerbSpec Protocol for short), a community-developed standard that defines minimum requirements, recommended practices, a step-by-step workflow, metadata schema, and guidelines for quality control and data management. The IHerbSpec Protocol is a living document deployed at iherbspec.github.io with versioned releases archived on Zenodo (IHerbSpec, 2026). It is designed to foster collaboration and provide a foundation for interoperable spectral datasets that can be confidently aggregated, reused, and integrated into the global biodiversity data ecosystem for ecological and evolutionary research.

METHODS AND RESULTS

Knowledge exchange and consensus building

The IHerbSpec Protocol was developed collaboratively by the International Herbarium Spectral Digitization Working Group (IHerbSpec), established in fall 2024. The group was assembled to represent a cross-section of the herbarium and spectral biology communities, including herbarium-based curators and researchers interested in spectral digitization and members of the ASCEND Biology Integration Institute (NSF-DBI-2021898; spectralbiology.org) interested in advancing the use of preserved plant specimens.

The development of a measurement and metadata capture protocol was among the first objectives for the working group. Activities began with a systematic comparison and consolidation of existing herbarium spectroscopy workflows conducted through a series of virtual webinars between December 2024 and April 2025. Ten distinct workflows were presented, several of which are published (Durgante et al., 2013; Laliberté and Soffer, 2018; Kothari et al., 2023; Quinteros Casaverde et al., 2024; Mersni et al., 2025; White et al., 2026). Members also shared approaches to specimen digitization and data management at their

institutions. These exchanges provided a common foundation for identifying shared practices, sources of variability, and priorities for coordination. The process culminated in an in-person workshop at the Harvard University Herbaria in May 2025, where remaining points of divergence were discussed and resolved collectively.

Decisions during protocol development were made by consensus rather than formal voting. Agreement was reached through iterative discussion, refinement of draft text, and collective review, with the explicit goal of protocol compliance (see *Embedding quality control and interoperability into the protocol*). The resulting protocol is a community-developed standard authored by 31 members from 23 institutions across three continents.

Publication and overview of the protocol

The IHerbSpec Protocol is written using Quarto (Allaire et al., 2025) and developed through a public GitHub repository (github.com/IHerbSpec/iherbspec.github.io) where community feedback is organized and incorporated via Discussions and Issues tools. The source files are compiled to HTML and published as a living document at iherbspec.github.io, ensuring that the online version reflects the most current updates. The GitHub repository is linked to Zenodo (<https://doi.org/10.5281/zenodo.18451589>), where versioned releases, including a compiled PDF, are permanently archived and assigned persistent DOIs. This structure maintains a single, version-controlled source of the protocol that supports transparent revision and community input, while generating both web and citable PDF outputs to ensure long-term preservation and stable scholarly reference. The protocol is currently distributed under a Creative Commons Attribution 4.0 International (CC BY 4.0) license.

The website, GitHub repository, and Zenodo archive provide a suggested citation for the protocol, including the protocol title, version number, and DOI. We recommend that users cite

the specific protocol version they followed, together with this paper, to ensure that methodological references remain both reproducible and connected to the broader rationale for the standard.

The broad goal of the IHerbSpec Protocol is for its standards to be implemented across global contexts and scales—from small, hypothesis-driven projects to large institutional digitization initiatives. To support this range of applications, the protocol distinguishes between required and recommended elements within its measurement workflow and metadata schema, providing a common structural framework while accommodating differences in project goals and institutional capacity (see *Embedding quality control and interoperability into the protocol*).

The protocol is organized into modular sections that separate procedural guidance, documentation standards, and supporting rationale. It begins with a preface describing its development and consensus process, followed by six core sections and two appendices detailing workflow, metadata, file conventions, instrumentation, tissue selection, and sampling thresholds (Table 1).




Fig. 1. Specimen and tissue selection workflow and bench measurement setup. (A)

Specimen and tissue selection decision tree. The first step (top) prioritizes specimens that are generally suitable for measurement (mature, intact, flat, and with minimally contaminated areas), unless the specimen is of unique importance. The second step prioritizes tissues that permit insertion of a thin non-reflective black background behind the target tissue. As stated in Part I of the IHerbSpec Protocol, insertion of a black background is required whenever possible to minimize spectral contamination from mounting materials. However, this is not always feasible for certain mounted or otherwise valuable specimens; in such cases, measurements may still be

201 collected provided that background conditions and potential sources of contamination are
202 explicitly documented in the metadata. The final step emphasizes selecting tissue areas that are
203 as representative and uncontaminated as possible (e.g., avoiding glue, epiphylls, or herbivory
204 where feasible). (B) Bench setup for spectral measurement. Figure A reproduced from
205 IHerbSpec Protocol (CC BY 4.0).

Accepted Manuscript




Fig. 2. Schematic of the IHerbSpec Protocol v1.3 measurement workflow. Steps 1–4

illustrate instrument and materials setup, measurement of white and black references and calibrated standards, and recording of session-level metadata associated with these activities. Steps 5–9 illustrate the tissue measurement sequence, filename and data quality validation, scoring of specimen- and tissue-level metadata, and annotation of project information on herbarium specimens. The protocol requires a minimum of three and a recommendation of five or more measurements (see Appendix I of the protocol) for both adaxial and abaxial leaf surfaces across the specimen, providing sufficient tissue is available and suitable for measurement. Plants with small, terete, or otherwise challenging leaves for measurement might not be able to meet this requirement (see *Vision for Future Versions* section). Figure reproduced from IHerbSpec Protocol (CC BY 4.0).

Informatics standards and metadata

The IHerbSpec metadata schema (Parts 3 and 4) was developed to support discovery, harmonization, and reuse of spectral data across projects, consistent with FAIR principles (Wilkinson et al., 2016). During the in-person and virtual meetings, the working group discussed informatics needs for spectral digitization, including digital storage, integration with collection management systems, data entry interfaces, and suitable formats for data creation, preservation, and dissemination. An informatics subgroup reviewed existing biodiversity data standards, particularly those maintained by Biodiversity Information Standards (TDWG; <https://www.tdwg.org>), including Darwin Core (Wieczorek et al., 2012) and its MeasurementOrFact extension.

However, these frameworks did not adequately capture the session–specimen–tissue hierarchy desired for herbarium spectral measurements. The IHerbSpec metadata schema was therefore developed as a relational model encompassing session-, specimen-, and tissue-level information, with explicit attention to measurement configuration and tissue characteristics (<https://iherbspec.github.io/protocol/part4-metadata.html#section4-1>). This approach aligns with the goals of the BIOFAIR data network, which emphasizes community-driven, domain-aware metadata standards that support interoperability, machine readability, and reuse across biological research infrastructures (Kunkel et al., 2025).

To support different project needs, the metadata schema is designed as a simple spreadsheet (“flat file”) for data capture and dissemination, and as a structured model suitable for integration into relational databases. The spreadsheet with all metadata fields is provided on the IHerbSpec website for immediate use or adaptation <https://iherbspec.github.io/protocol/tables>.

Although customized to capture the contextual detail needed for quality control, interpretation, and downstream aggregation of herbarium spectral data, the metadata schema was designed with future integration as a Darwin Core extension in mind. Where possible, existing Darwin Core terms are reused (e.g., `scientificName`, `identifiedBy`), and newly defined fields for spectral-specific information (e.g., `tissueDevelopmentalStage`, `hasGlue`, `measurementFlags`) follow Darwin Core naming conventions and semantics. This design supports immediate usability while laying the groundwork for potential future development of a Darwin Core extension for spectral data.

Data sharing and project-level data integration

Alongside measurement and metadata standards, the protocol provides data management guidance to support long-term preservation, interoperability, and reuse. Projects are strongly encouraged to archive and share unprocessed spectral data files together with a metadata spreadsheet because these files usually record measurement and instrument configuration settings in their headers along with the unprocessed data. Unprocessed spectra are typically output as text files containing reflectance values in columnar format, with instrument- and software-generated metadata recorded in file headers. The protocol also defines recommended filename conventions in Part 3 that ensure minimum traceability to the project, specimen, tissue class, and background conditions (<https://iherbspec.github.io/protocol/part3-filenames.html>).

Projects may also share tabular, analysis-ready files (e.g., .csv) containing specimens in rows and spectral bands and metadata fields in columns. Although these files are more user-friendly for data consumers, they may reflect prior processing steps, such as radiance-to-reflectance conversion, data extrapolation, or resampling, and may omit metadata such as exact wavelength centers, spectral resolution, detector overlap regions, and measurement conditions.

All processing steps applied to shared tabular files should therefore be clearly documented in the data archive to maintain transparency and reproducibility.

If projects are collecting trait data from herbarium specimens for model calibration or validation, those measurements should also be archived with stable specimen identifiers so they can be linked to the corresponding spectra and voucher records (see *Embedding spectra within institutional digitization workflows* section). Because general trait databases are not always designed to preserve specimen-level linkages, we recommend sharing validation trait datasets alongside spectral data, or in repositories that maintain specimen identifiers, as part of broader digital extended specimen data architectures (see *Scaling spectral digitization into the global metaherbarium* section).

To support discovery, citation, and long-term accessibility, the working group established the IHerbSpec Dataverse (<https://dataverse.harvard.edu/dataverse/iherbspec/>) as a primary project-level repository for herbarium spectral data. The Dataverse is freely available, widely used in the research community, and supports persistent identifiers (DOIs) and versioning, and user-defined licensing to communicate how data can be used. Submitted datasets may be minimally curated to help ensure conformance with the protocol's required components and metadata fields.

The IHerbSpec Dataverse is designed to accommodate datasets generated both within and outside formal institutional digitization pipelines, including projects based on specimens that lack barcodes, images, or fully curated database records. Its flexible deposit structure allows users to archive spectra, a metadata spreadsheet, validation trait measurements, and related metadata together while preserving specimen-level identifiers and links to institutional occurrence records (see the *Embedding spectra within institutional digitization workflows*

section). In this way, the Dataverse serves as one mechanism to support the aggregation, preservation, and reuse of herbarium spectral data and to link it to the corresponding herbarium specimens across projects and institutions. In addition, IHerbSpec and the IHerbSpec Dataverse act as signposts that enhance the discoverability of herbarium spectral datasets that are stored in other repositories.

Embedding quality control and interoperability into the protocol

The IHerbSpec Protocol embeds quality control throughout its structure, beginning with the definition and overview of the protocol's *required* versus *recommended* elements in Part 1, extending through the measurement workflow, filename conventions, and metadata schema (Parts 2–4), and reinforced by guidance on instrumentation, materials, and tissue selection (Parts 5 and 6; Table 1; Fig. 1).

Compliance with the IHerbSpec Protocol requires implementation of all required procedural elements and metadata fields. The required elements define the minimum standard for producing interoperable datasets with sufficient documentation to support evaluation, reuse, and integration across projects. Deviations from required elements may nevertheless be appropriate for particular specimens, instruments, or research objectives. Such deviations do not preclude valuable scientific use, but they should be explicitly documented so that downstream users can assess their implications for interpretation, comparison, and data integration. Recommended elements provide additional measurement context that can further strengthen interoperability and downstream analyses but are not required for protocol compliance. Collectively, these fields improve transparency and facilitate future harmonization across instruments, optical configurations, and measurement workflows.

To support consistency across institutions, the working group identified several components requiring explicit standardization. These include the measurement and archiving of white and black reference standards, the use of non-reflective black backgrounds to minimize contamination from mounting materials (Fig. 3), a minimum number of measurements per specimen, and standardized metadata and file naming conventions.

A key quality-control decision concerned sampling intensity. While a single carefully collected spectrum per leaf surface can adequately represent a specimen, it does not capture variance. The protocol therefore requires a minimum of three measurements per abaxial and adaxial leaf surface per specimen, when feasible, with on-screen verification that each measurement is free of anomalous or error-related spectral patterns (Fig. 2; Fig. 4). Five measurements per surface are recommended (or more if feasible given the taxon and project scope) to better capture within-specimen spectral variation. Given the importance of inspecting spectral data quality and consistency, the protocol emphasizes that replicate measurements can be taken from any combination of multiple leaves or a single spot on a leaf depending on the availability of suitable material. Appendix I of the protocol further discusses these decisions (<https://iherbspec.github.io/protocol/appendix1.html>).

Beyond measurement procedures, the hierarchical metadata schema (session-, specimen-, and tissue-level metadata) is designed to communicate the essential contextual information required to assess data quality and interoperability. Instrument-related fields—such as `instrumentModel`, `opticalSetupDescription`, and `measurementSettings`—are explicitly required because different instruments and optical configurations can introduce batch effects if not documented and harmonized (Meireles et al., 2020). Additional required fields such as `tissueDevelopmentalStage`, `hasGlue`, and

hasNonGlueContamination (e.g. chemical preservatives, insecticides), and recommended fields such as controlled-vocabulary measurementFlags (e.g. AlcoholPresent) and free-text tissueNotes, are tied specifically to the measurement area to enable precise downstream filtering. Finally, to support training and reduce subjectivity, Appendix II of the protocol provides additional guidelines for scoring tissue characteristics and scoring examples linked to images for 51 specimens from 11 herbaria (See IHerbSpec Protocol Appendix II: <https://iherbspec.github.io/protocol/appendix2.html>). Metadata should also be subject to quality checks in line with the quality assurance protocols applied to other data in the digitization pipeline.

By documenting methodological choices, measurement conditions, and specimen characteristics through its structured metadata schema, the IHerbSpec Protocol supports data interoperability and reuse. Compliance requires completion of all required elements, whereas recommended elements provide additional context that can strengthen quality control and facilitate cross-project integration. This framework allows measurements from scientifically important but challenging specimens to be retained and transparently evaluated, supporting inclusive data capture without sacrificing downstream filtering and integration across projects and herbaria globally.




Fig. 3. The effect of background on tissue spectral profiles. The IHerbSpec Protocol

requires placement of a low-reflectance black background beneath tissues to minimize background contamination in reflectance data. (A) Reflectance spectra of reference and background materials measured during a session, including the white reference panel (White, colored blue), the black background (Black), gray backgrounds, and representative herbarium mounting papers. These background materials exhibit distinct spectral signatures, particularly outside the visible range (400–700 nm). Plots show full-range (350–2500 nm), unprocessed reflectance data measured with the Spectral Evolution NaturaSpec and Surface Probe. (B) Difference spectra showing adaxial leaf reflectance measured on a black background minus

reflectance measured at the same leaf surface location on alternative backgrounds, illustrating wavelength-dependent background effects on tissue measurements. (C–D) Reflectance profiles of the same leaf measured on different backgrounds for two target classes: leaf abaxial (TCAB; panel C), and leaf adaxial (TCAD; panel D) surfaces. Measurements taken over black and gray backgrounds consistently show lower reflectance than those taken over white reference material and white to yellowed herbarium papers. Across panels, background effects are most pronounced in the 800–1800 nm region, corresponding mostly to the near-infrared (1100–2000 nm), with weaker but detectable effects in the visible (400–700 nm) and longer-wavelength SWIR (2000–2500 nm).




Fig. 4. Instrument and spatial heterogeneity in herbarium leaf reflectance

measurements. Reflectance spectra collected from the same dried leaf (*Betula papyrifera*; [NEBC00636882](#); tissue condition scoring in IHerbSpec Protocol [Appendix II](#)) using two spectroradiometers with different optical configurations: a Spectral Evolution NaturaSpec with a 2 mm field of view and an SVC HR-1024i with a 22 mm field of view. Spectra were trimmed to 400–2400 nm and plotted with the *ggplot2* R package (Wickham, 2016). Colored circles on the leaf image indicate measurement locations corresponding to the plotted spectra (circles not drawn to scale). NaturaSpec measurements include normal lamina tissue, midvein, pathogen-damaged tissue, off-target measurement including some black background, and reference measurements of the black background and white standard. The SVC measurement represents a standard lamina measurement integrating reflectance over a substantially larger sampling area. Spectral regions are shaded to indicate the visible range (VIS, 400–700 nm; rainbow band), near-infrared (NIR, ~700–1400 nm), and shortwave infrared (SWIR1, ~1400–1900 nm; SWIR2, ~2000–2500 nm). Differences among spectra illustrate how optical configuration and fine-scale

384 tissue heterogeneity influence reflectance profiles across the visible to shortwave infrared
385 spectrum.
386

Accepted Manuscript

Embedding spectra within institutional digitization workflows

A central goal in developing the protocol was ensuring its suitability for adoption under real-world institutional digitization workflows. Although expanded metadata capture will slow specimen throughput, the seventeen required metadata fields are essential for quality control filtering and harmonizing spectral data across institutions in accordance with FAIR principles (Wilkinson et al., 2016). Where possible, institutional software for digitization (“front ends”) can streamline metadata capture by automatically propagating repetitive elements (e.g., session-level fields like `projectId` and `instrumentModel`).

The protocol’s filename conventions and hierarchical metadata schema (Parts 3 and 4; Table 1) help prevent spectral data files from becoming detached from their project, specimen, or measurement conditions, safeguarding their long-term discoverability, interpretability, and reuse alongside the enduring stewardship of the specimens themselves. Thorough metadata capture also honors the substantial institutional effort invested in specimen digitization.

Critically, each specimen and corresponding extended dataset must be associated with a stable, traceable identifier. In the protocol, this linkage is supported by required specimen-level metadata fields that are incorporated into the filename structure: `herbariumCode`, which identifies the herbarium or collection housing the specimen, and `specimenId`, which identifies the specimen itself. The `specimenId` field is intentionally flexible because institutions differ in the identifiers they use for internal specimen tracking; it may contain, for example, a GUID, GBIF occurrence ID, barcode, catalog number, accession number, or collector-based identifier such as “Thorne24070a.” To reduce ambiguity, protocol v1.3 adds a required

`specimenIdClass` field specifying the type or source of the identifier provided in `specimenId`.

Stable specimen identifiers also provide the foundation for linking spectra, trait measurements, and other specimen-derived datasets to institutional specimen records. At the institutional level, these associations may be implemented through herbarium database annotations or outward links indicating that spectral data are available, who generated them, and where they are archived.

In practice, taking spectral measurements is most straightforward on unmounted specimens or free tissues stored in herbarium packets, where all tissue surfaces are accessible, black backgrounds can be easily used, and tissues can be thoroughly inspected for quality. IHerbSpec therefore recommends that specimen preparation workflows include storing detached leaf material in specimen packets, a practice that may already be in place to support destructive sampling and dissection. Institutions may also consider integrating spectral capture with barcode assignment prior to mounting, ensuring stable linkage between specimen and data and enabling access to the whole specimen for spectral measurement.

Importantly, incorporating spectral readings into digitization workflows—whether alongside imaging or specimen label transcription—will require a dedicated workstation, trained personnel, and time and monetary investment. While high-throughput photographic imaging can achieve rates less than 10 seconds per specimen (Davis et al., 2021), IHerbSpec-compliant spectral measurements typically require approximately 5–10 minutes per specimen, even though each individual spectral scan takes only about 2–10 seconds. This is due to the extra time needed for tissue selection, adaxial and abaxial measurements, metadata entry, and quality-control review. Consequently, organizing spectral digitization as a separate workflow is often ideal.

Most current herbarium spectroscopy projects use full-range (350–2,500 nm), high-resolution field spectroradiometers, which typically cost approximately \$70,000–\$90,000 USD. More compact, lower-resolution instruments with reduced spectral ranges can cost less than \$2,000 USD, but their analytical capacity for herbarium applications remains under active study and may differ from full-range systems (Durgante et al., 2026). Additional costs include calibration standards, background materials, software, and routine instrument maintenance. Institutions interested in incorporating spectral measurements into digitization workflows should be mindful of these added resource investments.

The guidelines embedded in the protocol (Parts 5, 6, and Appendix II) were designed to be readable and to support training and consistent implementation. For specimens already databased, spectral digitization can proceed in coordinated batches, with suitability of each specimen and tissue assessed prior to measurement (Fig. 1) and any deviations from recommended conditions explicitly documented in the tissue-level metadata fields (e.g., `measurementFlags`, `tissueNotes`).

Scaling spectral digitization into the global metaherbarium

Scaling herbarium spectral digitization beyond individual institutions requires interoperable mechanisms for publishing and discovering associations between specimen records and externally archived datasets. Within the IHerbSpec metadata schema, the required `specimenId` field provides the primary identifier for establishing these linkages, with `specimenIdClass` documenting the type or source of that identifier. Together, these fields enable resolvable connections among institutional repositories, GBIF, iDigBio, and other aggregators.

Extended specimen integration can currently be implemented using outward linking models, such as Darwin Core Resource Relationship records, within existing biodiversity platforms or collection management systems. Resource Relationship metadata is already supported within GBIF's Integrated Publishing Toolkit (IPT) for representing associations to external resources and distributing these associations alongside specimen occurrence records. Symbiota-based portals aggregate these metadata and can allow users to manage associations via the 'Linked Resources' data entry panel where the `isReferencedBy` relationship type may be used to express linkages to external spectral datasets (e.g., New England Botanical Society, 2024).

This distributed approach allows spectral datasets to remain hosted in domain-specific repositories while remaining discoverable through biodiversity portals. IHerbSpec is actively working toward the development of a Darwin Core extension to provide a more standardized mechanism for associating spectral datasets and metadata with specimen records. Connecting spectral data through established biodiversity infrastructures and extended specimen networks is essential for realizing their broader scientific value and supporting their integration and reuse as herbarium spectroscopy scales toward a global metaherbarium (Davis, 2023; Kunkel et al., 2025).

DISCUSSION

Taken together, herbaria represent the largest scientific database of plant diversity across taxonomic, geographic, and temporal scales (Thiers, 2025). Spectral digitization transfers high-dimensional phenotypic information from these collections directly into the digital landscape of modern biodiversity science. Here, we present a community-developed protocol that standardizes

measurement, metadata capture, and data sharing to ensure that these inherently expansive collections can be mobilized as interoperable spectral datasets across institutions and projects.

The development of shared spectral datasets and analytical frameworks strengthens connections among scientific communities. For example, spectral data can be used to model structural and chemical traits that would be difficult or impossible to measure directly, but that can provide valuable information for systematists, ecologists and ecophysiologicalists (Jetz et al., 2016; Cavender-Bares et al., 2026). Spectral data may also help guide molecular sampling by screening specimens for tissue condition or chemical signatures associated with DNA preservation, potentially helping prioritize material for genomic work while reducing unnecessary destructive sampling. Spectral species libraries derived from dried specimens can also advance conservation by supporting taxonomic identification and trait estimations of preserved material collected during field surveys—enabling more robust evaluation of species composition and functional diversity. This is especially important for reducing taxonomic uncertainty and geographic biases in biodiversity knowledge by expanding access to analytical tools, digital infrastructure, and training in megadiverse tropical countries, where many scientifically valuable herbaria remain underrepresented in global databases and research networks (Delves et al., 2024; Thiers, 2024; Zhigila et al., 2026). Wider deployment of affordable spectrometers could enable these collections to generate and mobilize new phenotypic data locally.

Another goal for herbarium spectral digitization is to create a bridge to remote sensing. As modeling approaches advance, translating dry-leaf spectra into estimates of fresh functional traits may enable more direct integration with airborne and satellite observations (e.g., Asner and Martin, 2008; Singh et al., 2015; Wang et al., 2019, 2020; Sapes et al., 2024). Standardized

spectral libraries from herbaria have the potential to contribute geographically distributed reference datasets to calibrate canopy-level trait models. In turn, the advancement of remotely sensed biodiversity patterns can guide targeted botanical exploration, herbarium collection, and floristic and systematic research.

Finally, expanding spectral digitization enhances both the accessibility and enduring relevance of herbarium collections. Rather than rendering physical collections obsolete, digital extensions underscore their continued necessity: advances in technology and digital data architectures will create new opportunities to generate, link, and reuse biodiversity data, while drawing new scientific communities into collections-based research and opening additional avenues for investment and funding. These datasets will be most verifiable, repeatable, and extensible themselves when they are anchored to vouchered specimens that can be revisited and remeasured as technologies evolve. By embracing new analytical applications—while maintaining rigorous standards for specimen stewardship—the herbarium community can demonstrate that collections are not static archives of past utility, but dynamic research infrastructures that support emerging disciplines and address ongoing ecological, evolutionary, and conservation challenges. In doing so, spectral digitization provides new pathways to safeguard and steward collections in perpetuity, diversify their user and funding bases, and contribute to their digital repatriation (McAlvay et al., 2021; Pinheiro et al., 2024) under CARE principles (Carroll et al., 2020; Hart et al. 2026). Together, these advances extend the scientific and societal reach of natural history collections and reinforce their central role in addressing global challenges.

Limitations

Some aspects of metadata capture, including assessment of tissue condition and developmental stage, involve subjective judgment and may vary among operators. The protocol provides guidelines and examples in Appendix II of the IHerbSpec Protocol (<https://iherbspec.github.io/protocol/appendix2.html>) aimed at minimizing subjectivity. Differences in instrumentation and optical configuration can also introduce batch effects, even when protocols are followed. Continued development of validation, harmonization, and anomaly-detection methods will therefore be essential for cross-institutional analyses. In addition, although the IHerbSpec metadata schema was designed for immediate community use, it has not yet undergone formal governance within established biodiversity standards frameworks; coordination with Biodiversity Information Standards (TDWG) will be necessary to ensure long-term alignment.

Although spectral measurements are meant to be non-destructive, specimen handling and use of a contact probe do entail some risk of damaging the specimens themselves, either through excess illumination or mechanical damage, summarized in Table 2. These risks can be mitigated through training of spectral digitizers and review of protocol guidelines.

Projects aimed at developing predictive models for traits such as leaf nitrogen and phosphorus concentrations, carbon fractions (e.g., lignin and cellulose), or micronutrients may require targeted destructive sampling of selected specimens to generate ground-truth data for model building and validation. This is because the taxonomic and ecological breadth over which current models can be confidently applied remains uncertain, and more trait data are needed to improve model generalizability (Cavender-Bares et al., 2026). Recent work shows that models can remain robust across specimen age and species but may perform less reliably when phenological phases or functional groups represented in the target data are absent from the

training set (Lee et al., 2026; Nichodemus and Meireles, 2026). Although herbarium specimens can provide reliable sources of traits such as LMA, leaf thickness, and stable carbon or nitrogen isotopes (Körner et al., 2016; Perez et al., 2020), other observed or spectrally estimated traits may not closely approximate trait values of living leaf tissue because of drying, preservation, and degradation effects, limiting their direct biological interpretation. We therefore, envision model building and validation drawing on complementary sources of ground-truth data, including recently-dried and pressed leaves (Serbin et al., 2014; Kothari et al., 2023), silica-dried leaf material (Kothari et al., 2024), and controlled herborization experiments that explicitly quantify changes associated with specimen processing (Quinteros Casaverde et al., 2024; Jackson, 2025).

Finally, measuring very small, thin, or structurally complex photosynthetic tissues (e.g., graminoids, cushion plants, succulents, terete or needle-like leaves) may require justified deviations from the protocol's minimum sampling requirements, such as collecting fewer replicate measurements when tissue size and/or optical geometry limits the feasibility of making multiple high-quality measurements.

Vision for Future Versions

The IHerbSpec Protocol provides a standardized measurement approach designed to evolve alongside advances in instrumentation, digital infrastructure, and analytical methodologies. Continued community use and empirical evaluation across taxa, specimen conditions, instruments, and project scales will be essential for refining metadata standards, improving data harmonization, and expanding guidance for measurements.

The current protocol focuses primarily on vascular plant leaves, reflecting the extensive body of work on leaf spectra in plant physiology, functional ecology, and remote sensing (Wessman et al., 1988; Gamon and Surfus, 1999; Cavender-Bares et al., 2025). Although the

protocol does define tissue class codes for additional plant structures (see IHerbSpec Protocol Table 4.6: *Tissue Descriptor Codes*; e.g., `bract`, `OuterBark`, `PhotosyntheticSucculentStem`), measurement frameworks for these other tissues, as well as other herbarium collections like lichens and bryophytes, remain to be developed (but see (Hadlich et al., 2018, 2025; Guzmán Q. et al., 2020; Stasinski, 2021). Establishing specific protocols for these materials represents a key direction for future community coordination.

More broadly, the session–specimen–tissue metadata hierarchy developed here may provide a useful template for other specimen-derived datasets beyond leaf reflectance spectra, consistent with calls for community-driven, domain-aware standards for extended specimen data (Kunkel et al., 2025). Similar standardization efforts will be important for other herbarium-based measurements, including image-derived traits, molecular sampling, chemical assays, and handheld X-ray fluorescence spectroscopy for estimating elemental concentrations in dried leaves (van der Ent et al., 2019). Fields such as `tissueDevelopmentalStage`, `hasGlue`, `hasNonGlueContamination`, and `measurementFlags` may be broadly useful for documenting specimen suitability and measurement context across these applications.

Ongoing development is facilitated through the public GitHub repository (<https://github.com/IHerbSpec/iherbspec.github.io>), which hosts the protocol code and deploys the IHerbSpec website (<https://iherbspec.github.io>), providing a transparent platform for revision and community engagement. As participation expands, structured governance—such as subcommittees and leadership teams—will be necessary to balance inclusivity with efficient coordination. Researchers, students, and collections professionals interested in contributing are encouraged to visit the IHerbSpec Contact page (<https://iherbspec.github.io/contact.html>) to learn about joining the working group and participating in community discussions.

ACKNOWLEDGEMENTS

We gratefully acknowledge botanical collectors as well as the herbaria, curators, collection managers, and staff who steward biodiversity collections. We thank Michaela Schmul for helpful comments on the manuscript. This project was supported by the U.S. National Science Foundation Biology Integration Institute ASCEND (DBI-2021898), the Harvard University Herbaria, iDigBio (DBI-2027654), and the Missouri Botanical Garden's *Revolutionizing Species Identification* project. FD is funded by the Amazonas State Research Support Foundation (FAPEAM) *Women in Science Program* (SpectraPop project, process number 01.02.016301.04984/2024-17), the National Council for Scientific and Technological Development (CNPq) *Pró-Amazônia* (HerbSpectra-Amazon project, process number 385465/2025-4), and the German Federal Ministry of Research, Technology, and Space (BMBF) grant 01LK2101D. JEM is funded by NSF DEB-2442433. BB is funded by ETH Zurich Career Seed Award (25-1 Seed-006). CCV acknowledges support from the International Association for Plant Taxonomy (IAPT), 2025 research grant round. JLV is supported by Danish National Research Foundation grant DNRF179 to HT.

AUTHOR CONTRIBUTIONS

JCB initiated IHerbSpec and the creation of a protocol. IHerbSpec conceptualized the protocol and DMW led its development. DMW wrote this manuscript with contributions from MWA, FMD, JAG, JAK, and CON. KB drafted Table 2 with input from EMV. All authors contributed to protocol development, reviewed the manuscript, and approved the final version.

LITERATURE CITED

- Abasolo, M., D. J. Lee, C. Raymond, R. Meder, and M. Shepherd. 2013. Deviant near-infrared spectra identifies *Corymbia* hybrids. *Forest Ecology and Management* 304: 121–131.
- Allaire, J. J., C. Teague, C. Scheidegger, Y. Xie, C. Dervieux, and G. Woodhull. 2025. Quarto.
- Antonelli, A., C. Davis, I. Larridon, D. P. Little, R. J. Smith, and S. A. Smith. 2026. The digital biodiversity revolution. *New Phytologist* 251: 595–601.
- Asner, G. P., and R. E. Martin. 2008. Spectral and chemical analysis of tropical forests: Scaling from leaf to canopy levels. *Remote Sensing of Environment* 112: 3958–3970.
- Boughalmi, K., P. G. Santacruz Endara, L. A. Bennett, M. Ecartot, S. Bazan, D. Bastianelli, L. Bonnal, and T. L. P. Couvreur. 2025. Discrimination of Annonaceae using herbarium leaf reflectance spectra under limited sample size conditions. 2025.09.02.673631.
- Buono, D., and D. C. Albach. 2023. Infrared spectroscopy for ploidy estimation: An example in two species of *Veronica* using fresh and herbarium specimens. *Applications in Plant Sciences* 11: e11516.
- Carroll, S. R., I. Garba, O. L. Figueroa-Rodríguez, J. Holbrook, R. Lovett, S. Materechera, M. Parsons, et al. 2020. The CARE Principles for Indigenous Data Governance. *Data Science Journal* 19.
- Cavender-Bares, J., J. E. Meireles, J. J. Couture, M. A. Kaproth, C. C. Kingdon, A. Singh, S. P. Serbin, et al. 2016. Associations of leaf spectra with genetic and phylogenetic variation in oaks: Prospects for remote detection of biodiversity. *Remote Sensing* 8: 17–17.
- Cavender-Bares, J., J. E. Meireles, J. Pinto-Ledezma, Peter B. Reich, M. Schuman, P. A. Townsend, and A. Trowbridge. 2025. Spectral biology across scales in changing environments. *Ecology* 105.
- Cavender-Bares, J., D. M. White, N. I. Ahlstrand, M. W. Austin, D. Bastianelli, S. Bazan, K. Boughalmi, et al. 2026. Next-generation specimen digitization: capturing reflectance spectra from the world's herbaria for modeling plant biology across time, space, and taxa. *New Phytologist* 251: 645–665.
- Costa, F. R., C. Lang, D. R. Almeida, C. V. Castilho, and L. Poorter. 2018. Near-infrared spectrometry allows fast and extensive predictions of functional traits from dry leaves and branches. *Ecological Applications* 28: 1157–1167.
- Cruickshank, R. H., and L. Munck. 2011. It's barcoding Jim, but not as we know it. *Zootaxa* 2933: 55–56.
- Curran, P. J. 1989. Remote sensing of foliar chemistry. *Remote Sensing of Environment* 30: 271–278.

- 646 Davis, C. C. 2023. The herbarium of the future. *Trends in Ecology & Evolution* 38: 412–423.
- 647 Davis, C. C., J. A. Kennedy, and C. J. Grassa. 2021. Back to the future: A refined single-user
648 photostation for massively scaling herbarium digitization. *TAXON* 70: 635–643.
- 649 Delves, J., J. Albán-Castillo, A. Cano, C. Fernández Aviles, E. Gagnon, P. Gonzáles, S. Knapp,
650 et al. 2024. Small and in-country herbaria are vital for accurate plant threat assessments:
651 A case study from Peru. *Plants, People, Planet* 6: 174–185.
- 652 Durgante, F. M., N. Higuchi, A. Almeida, and A. Vicentini. 2013. Species Spectral Signature:
653 Discriminating closely related plant species in the Amazon with Near-Infrared Leaf-
654 Spectroscopy. *Forest Ecology and Management* 291: 240–248.
- 655 Durgante, F. M., C. da C. Vasconcelos, H. L. Hadlich, C. L. Mallmann, R. de O. Perdiz, K. C.
656 M. da Cunha, S. R. Chaves, et al. 2026. Species recognition in herbaria and standing trees
657 using an accessible portable spectrometer.
- 658 Enquist, B. J., R. Condit, R. K. Peet, M. Schildhauer, and B. M. Thiers. 2016.
659 Cyberinfrastructure for an integrated botanical information network to investigate the
660 ecological impacts of global climate change on plant biodiversity. *PeerJ Preprint*.
- 661 van der Ent, A., G. Echevarria, A. J. Pollard, and P. D. Erskine. 2019. X-Ray Fluorescence
662 Ionomics of Herbarium Collections. *Scientific Reports* 9: 4746.
- 663 Fine, P. V. A., D. Salazar, R. E. Martin, M. R. Metz, T. M. Misiewicz, and G. P. Asner. 2021.
664 Exploring the links between secondary metabolites and leaf spectral reflectance in a
665 diverse genus of Amazonian trees. *Ecosphere* 12: e03362.
- 666 Gamon, J. A., and J. S. Surfus. 1999. Assessing leaf pigment content and activity with a
667 reflectometer. *New Phytologist* 143: 105–117.
- 668 Guzmán Q., J. A., K. Laakso, J. C. López-Rodríguez, B. Rivard, and G. A. Sánchez-Azofeifa.
669 2020. Using visible-near-infrared spectroscopy to classify lichens at a Neotropical Dry
670 Forest. *Ecological Indicators* 111: 105999.
- 671 Hadlich, H. L., F. M. Durgante, J. dos Santos, N. Higuchi, J. Q. Chambers, and A. Vicentini.
672 2018. Recognizing Amazonian tree species in the field using bark tissues spectra. *Forest*
673 *Ecology and Management* 427: 296–304.
- 674 Hadlich, H. L., J. Schöngart, F. Wittmann, C. C. Vasconcelos, C. L. Mallmann, M. L. G. Conde,
675 P. Amaral de Sá, et al. 2025. Exploring the potential of field spectroscopy for tree species
676 identification in different Amazonian forest ecosystems. *Global Ecology and*
677 *Conservation* 64: e03970.
- 678 Hardisty, A. R., E. R. Ellwood, G. Nelson, B. Zimkus, J. Buschbom, W. Addink, R. K. Rabeler,
679 et al. 2022. Digital Extended Specimens: Enabling an Extensible Network of Biodiversity
680 Data Records as Integrated Digital Objects on the Internet. *BioScience* 72: 978–987.

- 681 Hart, R., V. Fonseca-Kruel, E. Dalcin, E. da Silva Luís Alexandre, M. Pace, M. Schnull,
682 L. Beltran-Rodríguez, et al. 2026. Repositories of biocultural diversity: Toward best
683 practices for empowering ethnobotany in digital herbaria. *Plants, People, Planet* 8: 1095–
684 1103.
- 685 Hernández-Leal, M. S., J. A. Guzmán Q., A. González-Rodríguez, and J. Cavender-Bares. 2025.
686 Foliar spectral signatures reveal adaptive divergence in live oaks (*Quercus* section
687 *Virentes*) across species and environmental niches. *New Phytologist* 248: 370–388.
- 688 IHerbSpec. 2026. Protocol for Spectral Digitization of Herbarium Specimens v1.2.1.
- 689 Jackson, R. 2025. Effects of specimen degradation on herbarium leaf spectroscopy. University of
690 British Columbia.
- 691 Jacquemoud, S., S. L. Ustin, J. Verdebout, G. Schmuck, G. Andreoli, and B. Hosgood. 1996.
692 Estimating leaf biochemistry using the PROSPECT leaf optical properties model. *Remote*
693 *Sensing of Environment* 56: 194–202.
- 694 Jetz, W., J. Cavender-Bares, R. Pavlick, D. Schimel, F. W. Davis, G. P. Asner, R. Guralnick, et
695 al. 2016. Monitoring plant functional diversity from space. *Nature Plants* 2: 1–5.
- 696 Kattge, J., K. Ogle, G. Bönsch, S. Díaz, S. Lavorel, J. Madin, K. Nadrowski, et al. 2011. A
697 generic structure for plant trait databases. *Methods in Ecology and Evolution* 2: 202–213.
- 698 Körner, C., S. Leuzinger, S. Riedl, R. T. Siegwolf, and L. Streule. 2016. Carbon and nitrogen
699 stable isotope signals for an entire alpine flora, based on herbarium samples. *Alpine*
700 *Botany* 126: 153–166.
- 701 Kothari, S., R. Beauchamp-Rioux, E. Laliberté, and J. Cavender-Bares. 2023. Reflectance
702 spectroscopy allows rapid, accurate and non-destructive estimates of functional traits
703 from pressed leaves. *Methods in Ecology and Evolution* 14: 385–401.
- 704 Kothari, S., S. E. Hobbie, and J. Cavender-Bares. 2024. Rapid estimates of leaf litter chemistry
705 using reflectance spectroscopy. *Canadian Journal of Forest Research* 54: 978–991.
- 706 Kothari, S., and A. K. Schweiger. 2022. Plant spectra as integrative measures of plant
707 phenotypes. *Journal of Ecology* 110: 2536–2554.
- 708 Kühn, P., T. Proß, C. Römermann, K. Wesche, and H. Bruelheide. 2024. Using near-infrared
709 spectroscopy to predict nitrogen and phosphorus concentrations of herbarium specimens
710 under different storage conditions. *Plant Methods* 20: 19.
- 711 Kunkel, D. M., B. L. Long-Fox, C. Pittman, J. Portmann, M. Sheik, J. M. Bates, A. Bentley, et
712 al. 2025. Integrating biological and environmental data to solve key scientific and societal
713 challenges. *BioScience*: biaf150.
- 714 Laliberté, E., and R. Soffer. 2018. Measuring spectral reflectance and transmittance (350-2500

- 715 nm) of small and/or narrow leaves using the Spectra Vista Corporation (SVC) DC-R/T
716 Integrating Sphere V.3. *protocols.io*.
- 717 Lang, C., D. R. A. Almeida, and F. R. C. Costa. 2017. Discrimination of taxonomic identity at
718 species, genus and family levels using Fourier Transformed Near-Infrared Spectroscopy
719 (FT-NIR). *Forest Ecology and Management* 406: 219–227.
- 720 Lee, A. K., L. Vander Esch, J. Oong, B. Fredericksen, J. Ramirez-Lerma, J. A. Guzmán, D.
721 White, et al. 2026. Robust estimation of key leaf traits from reflectance spectroscopy of
722 herbarium specimens. *Methods in Ecology and Evolution*.
- 723 Lendemer, J., B. Thiers, A. K. Monfils, J. Zaspel, E. R. Ellwood, A. Bentley, K. LeVan, et al.
724 2020. The Extended Specimen Network: A Strategy to Enhance US Biodiversity
725 Collections, Promote Research and Education. *BioScience* 70: 23–30.
- 726 McAlvay, A. C., C. G. Armstrong, J. Baker, L. B. Elk, S. Bosco, N. Hanazaki, L. Joseph, et al.
727 2021. Ethnobiology Phase VI: Decolonizing Institutions, Projects, and Scholarship.
728 *Journal of Ethnobiology* 41: 170–191.
- 729 Meireles, J. E., J. Cavender-Bares, P. A. Townsend, S. Ustin, J. A. Gamon, A. K. Schweiger, M.
730 E. Schaepman, et al. 2020. Leaf reflectance spectra capture the evolutionary history of
731 seed plants. *New Phytologist* 228: 485–493.
- 732 Mersni, J., S. Bazan, E. Baby, V. Bonnal, and D. Bastianelli. 2025. Protocol for NIR spectra
733 acquisition on herbarium specimen.
- 734 Neto-Bradley, B. M., P. Bonnet, H. Goëau, A. Joly, J. Cavender-Bares, and D. A. Coomes. 2026.
735 Using reflectance spectra and Pl@ntNet to identify herbarium specimens: a case study
736 with *Lithocarpus*. *New Phytologist* 251: 800–810.
- 737 New England Botanical Society. 2024. Harvard University Herbaria: Vascular Plants of North
738 America (Harvard:NEBC). *Occurrence ID: 55f53526-cc56-4b3a-8bb7-670db12c035b*.
739 Website
740 [https://midwestherbaria.org/portal/collections/individual/index.php?occid=33010605&cli](https://midwestherbaria.org/portal/collections/individual/index.php?occid=33010605&clid=0)
741 [d=0](https://midwestherbaria.org/portal/collections/individual/index.php?occid=33010605&clid=0) [accessed 13 July 2026].
- 742 Nichodemus, C. O., and J. E. Meireles. 2026. Reflectance spectra capture temporal variation in
743 functional traits and leaf phenology. 2026.04.21.719921.
- 744 Perez, T. M., J. Rodriguez, and J. Mason Heberling. 2020. Herbarium-based measurements
745 reliably estimate three functional traits. *American Journal of Botany* 107: 1457–1464.
- 746 Pérez-Harguindeguy, N., S. Díaz, E. Garnier, S. Lavorel, H. Poorter, P. Jaureguiberry, M. S.
747 Bret-Harte, et al. 2016. Corrigendum to: New handbook for standardised measurement of
748 plant functional traits worldwide. *Australian Journal of Botany* 64: 715–716.
- 749 Pinheiro, F. da C., R. C. Forzza, P. M. Leitman, and J. Prado. 2024. The REFLORA Program:

- 750 implementation, repatriation, and creation of the REFLORA Virtual Herbarium as a tool
751 for biodiversity studies. *Biota Neotropica* 24: e20241701.
- 752 Quinteros Casaverde, N. L., S. P. Serbin, and D. C. Daly. 2024. Changes in spectral signature of
753 leaves after desiccation: Implications for the prediction of leaf traits and plant-soil
754 interaction in herbarium samples. 2024.01.21.576284.
- 755 Sapes, G., L. Schroeder, A. Scott, I. Clark, J. Juzwik, R. A. Montgomery, J. A. Guzmán Q, and J.
756 Cavender-Bares. 2024. Mechanistic links between physiology and spectral reflectance
757 enable previsual detection of oak wilt and drought stress. *Proceedings of the National*
758 *Academy of Sciences* 121: e2316164121.
- 759 Serbin, S. P., A. Singh, B. E. McNeil, C. C. Kingdon, and P. A. Townsend. 2014. Spectroscopic
760 determination of leaf morphological and biochemical traits for northern temperate and
761 boreal tree species. *Ecological Applications* 24: 1651–1669.
- 762 Singh, A., S. P. Serbin, B. E. McNeil, C. C. Kingdon, and P. A. Townsend. 2015. Imaging
763 spectroscopy algorithms for mapping canopy foliar chemical and morphological traits
764 and their uncertainties. *Ecological Applications* 25: 2180–2197.
- 765 Stasinski, L. 2021. Assessing Plant and Lichen Diversity Using Reflectance Spectra. University
766 of Maine, Orono, Maine.
- 767 Stasinski, L., D. M. White, P. R. Nelson, R. H. Ree, and J. E. Meireles. 2021. Reading light: leaf
768 spectra capture fine-scale diversity of closely related, hybridizing arctic shrubs. *New*
769 *Phytologist* 232: 2283–2294.
- 770 Stefanski, A., E. E. Butler, L. J. Williams, R. Bermudez, J. A. Guzmán Q., A. Larson, P. A.
771 Townsend, et al. 2025. All the light we cannot see: Climate manipulations leave short and
772 long-term imprints in spectral reflectance of trees. *Ecology* 106: e70048.
- 773 Thiers, B. M. 2024. Strengthening Partnerships to Safeguard the Future of Herbaria.
774 *Diversity* 16: 36.
- 775 Thiers, B. M. 2025. The World's Herbaria 2024: A Summary Report Based on Data from
776 Index Herbariorum. New York Botanical Garden, New York.
- 777 Vance, C. K., D. R. Tolleson, K. Kinoshita, J. Rodriguez, and W. J. Foley. 2016. Near Infrared
778 Spectroscopy in Wildlife and Biodiversity. *Journal of Near Infrared Spectroscopy* 24: 1–
779 25.
- 780 Vasconcelos, C. C., I. D. K. Ferraz, F. M. Durgante, F. Wittmann, J. Schöngart, M. T. F.
781 Piedade, J. L. C. Camargo, and M. H. Terra-Araujo. 2025. *Ecclinusa nervosa*
782 (Sapotaceae, Chrysophylloideae), a New Species Discovered in Central Amazonia.
783 *Systematic Botany* 50: 55–66.
- 784 Wang, Z., A. Chlus, R. Geygan, Z. Ye, T. Zheng, A. Singh, J. J. Couture, et al. 2020. Foliar

- 785 functional traits from imaging spectroscopy across biomes in eastern North America.
786 *New Phytologist* 228: 494–511.
- 787 Wang, Z., P. A. Townsend, A. K. Schweiger, J. J. Couture, A. Singh, S. E. Hobbie, and J.
788 Cavender-Bares. 2019. Mapping foliar functional traits and their uncertainties across
789 three years in a grassland experiment. *Remote Sensing of Environment* 221: 405–416.
- 790 Wessman, C. A., J. D. Aber, D. L. Peterson, and J. M. Melillo. 1988. Remote sensing of canopy
791 chemistry and nitrogen cycling in temperate forest ecosystems. *Nature* 335: 154–156.
- 792 White, D. M., J. Cavender-Bares, C. C. Davis, J. A. Guzmán Q., S. Kothari, J. M. Robles, and J.
793 E. Meireles. 2026. Seeing herbaria in a new light: leaf reflectance spectroscopy unlocks
794 trait and classification modeling in plant biodiversity collections. *New Phytologist* 251:
795 811–829.
- 796 Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York.
- 797 Wieczorek, J., D. Bloom, R. Guralnick, S. Blum, M. Döring, R. Giovanni, T. Robertson, and D.
798 Vieglais. 2012. Darwin Core: An Evolving Community-Developed Biodiversity Data
799 Standard. *PLOS ONE* 7: e29715.
- 800 Wilkinson, M. D., M. Dumontier, Ij. J. Aalbersberg, G. Appleton, M. Axton, A. Baak, N.
801 Blomberg, et al. 2016. The FAIR Guiding Principles for scientific data management and
802 stewardship. *Scientific Data* 3: 160018.
- 803 Zhigila, D. A., R. J. Schmidt-Knapik, B. M. Thiers, S. D. Abdul, S. Abdullahi, A. A.
804 AbdulRahaman, E. I. Aigbokhan, et al. 2026. Biodiversity science is improved when
805 silent herbaria speak. *Plants, People, Planet* 8: 1378–1386.

806 TABLES

807 Table 1: Structure and content of the IHerbSpec Protocol

Protocol Part	Content Description
Preface	Abstract, frontmatter, and details about IHerbSpec.
Part 1: Overview	Overview of the core measurement procedure and associated metadata fields, distinguishing <i>required</i> and <i>recommended</i> elements of the protocol.
Part 2: Workflow	Stepwise workflow for spectral measurement and metadata collection (Fig. 2).
Part 3: Filename Conventions and Formats	Standardized ‘Full’ and ‘Simple’ file-naming conventions for spectral data files (tissue, background, and calibration spectra), designed to link measurements and ensure traceability. The simple filename convention is streamlined for use during measurement sessions and should be converted into full filenames at the point of data archival and sharing to enable interoperability.
Part 4: Metadata and Databasing	Metadata schema for herbarium spectral reflectance data. Three metadata tables are organized into logical groups of session-, specimen-, and tissue metadata. Additional tables describe controlled vocabularies for tissue classes (e.g. <i>adaxial</i> leaf, <i>abaxial</i> leaf, bract, etc), tissue developmental stage (e.g. <i>young</i> , <i>mature</i> , <i>unknown</i> , etc.), tissue descriptors (e.g. <i>AlcoholPresent</i> , <i>LichenPresent</i> , etc.), and background and white reference class codes (e.g. <i>PaperBackground</i> , <i>BlackBackground</i>). The part concludes with guidance for data management, archiving, and sharing, including recommendations to use the IHerbSpec Dataverse community repository.
Part 5: Instrumentations and Materials	Guidance on instrumentation, materials, and measurement configuration.
Part 6: Tissue Selection	Decision tree and considerations for selecting suitable herbarium tissues for spectral measurement.
Appendix I	Data-driven guidance on the number of measurements per specimen.
Appendix II	Examples and guidance for recording biological and preparation-related sources of variation affecting spectral data in the IHerbSpec metadata.

Accepted Manuscript

Table 2: Estimated incidence of accidental specimen damage during contact-based spectroscopy of herbarium collections.

Type of destructive effect	Estimated relative frequency (500 scanned specimens*)	Mitigation measures
Leaf detachment (<i>detachment at the petiole–stem junction</i>)	2-3 / 500	Prefer already detached leaves (e.g., mounted or stored in protective sleeves or packets) when available; insert background materials carefully to avoid mechanical stress at attachment points (leaf base, petiole base), particularly in fragile tissues.
Major cracks (<i>visible structural fracture across lamina</i>)	5 / 500	Select flat, structurally intact leaf areas; assess tissue resistance before probe contact; maintain proper probe alignment with minimal pressure (especially when using small ~10 mm probes); avoid lateral movement once contact is made; and use extra care near fragile (e.g., deteriorated margins, damaged areas) or unsupported regions (e.g., areas near prominent veins).
Superficial cracks (<i>minor surface fissures without full rupture</i>)	10 / 500	
Probe perforation	2 / 500	
Partial border breakage (<i>fragment loss along leaf margin</i>)	5 / 500	Test instrumentation on thin or delicate leaves prior to measurement; use low light intensity with longer integration times to minimize heat exposure; and regularly clean the probe surface to prevent transfer of particles, dust, or residues between specimens.
Burning or discoloration of leaf from heat of contact probe	1 / 500	

*Data compiled from Boughalmi et al. (2025) and additional spectroscopy campaigns conducted by the same research group. Frequencies represent the number of damaged specimens relative to a cumulative total of 500 herbarium specimens scanned across two spectroscopy campaigns

815 conducted between 2024 and 2026 at Muséum National d'Histoire Naturelle (P), Naturalis
816 Biodiversity Center (WAG, L), and Université de Montpellier (MPU).
817

Accepted Manuscript

FIGURE LEGENDS

Fig. 1. Specimen and tissue selection workflow and bench measurement setup. (A)

Specimen and tissue selection decision tree. The first step (top) prioritizes specimens that are generally suitable for measurement (mature, intact, flat, and with minimally contaminated areas), unless the specimen is of unique importance. The second step prioritizes tissues that permit insertion of a thin non-reflective black background behind the target tissue. As stated in Part I of the IHerbSpec Protocol, insertion of a black background is required whenever possible to minimize spectral contamination from mounting materials. However, this is not always feasible for certain mounted or otherwise valuable specimens; in such cases, measurements may still be collected provided that background conditions and potential sources of contamination are explicitly documented in the metadata. The final step emphasizes selecting tissue areas that are as representative and uncontaminated as possible (e.g., avoiding glue, epiphylls, or herbivory where feasible). (B) Bench setup for spectral measurement. Figure A reproduced from IHerbSpec Protocol (CC BY 4.0).

Fig. 2. Schematic of the IHerbSpec Protocol v1.3 measurement workflow. Steps 1–4

illustrate instrument and materials setup, measurement of white and black references and calibrated standards, and recording of session-level metadata associated with these activities. Steps 5–9 illustrate the tissue measurement sequence, filename and data quality validation, scoring of specimen- and tissue-level metadata, and annotation of project information on herbarium specimens. The protocol requires a minimum of three and a recommendation of five or more measurements (see Appendix I of the protocol) for both adaxial and abaxial leaf surfaces across the specimen, providing sufficient tissue is available and suitable for measurement. Plants

with small, terete, or otherwise challenging leaves for measurement might not be able to meet this requirement (see *Vision for Future Versions* section). Figure reproduced from IHerbSpec Protocol (CC BY 4.0).

Fig. 3. The effect of background on tissue spectral profiles. The IHerbSpec Protocol requires placement of a low-reflectance black background beneath tissues to minimize background contamination in reflectance data. (A) Reflectance spectra of reference and background materials measured during a session, including the white reference panel (White, colored blue), the black background (Black), gray backgrounds, and representative herbarium mounting papers. These background materials exhibit distinct spectral signatures, particularly outside the visible range (400–700 nm). Plots show full-range (350–2500 nm), unprocessed reflectance data measured with the Spectral Evolution NaturaSpec and Surface Probe. (B) Difference spectra showing adaxial leaf reflectance measured on a black background minus reflectance measured at the same leaf surface location on alternative backgrounds, illustrating wavelength-dependent background effects on tissue measurements. (C–D) Reflectance profiles of the same leaf measured on different backgrounds for two target classes: leaf abaxial (TCAB; panel C), and leaf adaxial (TCAD; panel D) surfaces. Measurements taken over black and gray backgrounds consistently show lower reflectance than those taken over white reference material and white to yellowed herbarium papers. Across panels, background effects are most pronounced in the 800–1800 nm region, corresponding mostly to the near-infrared (1100–2000 nm), with weaker but detectable effects in the visible (400–700 nm) and longer-wavelength SWIR (2000–2500 nm).

Fig. 4. Instrument and spatial heterogeneity in herbarium leaf reflectance measurements.

Reflectance spectra collected from the same dried leaf (*Betula papyrifera*; [NEBC00636882](#); tissue condition scoring in IHerbSpec Protocol [Appendix II](#)) using two spectroradiometers with different optical configurations: a Spectral Evolution NaturaSpec with a 2 mm field of view and an SVC HR-1024i with a 22 mm field of view. Spectra were trimmed to 400–2400 nm and plotted with the *ggplot2* R package (Wickham, 2016). Colored circles on the leaf image indicate measurement locations corresponding to the plotted spectra (circles not drawn to scale). NaturaSpec measurements include normal lamina tissue, midvein, pathogen-damaged tissue, off-target measurement including some black background, and reference measurements of the black background and white standard. The SVC measurement represents a standard lamina measurement integrating reflectance over a substantially larger sampling area. Spectral regions are shaded to indicate the visible range (VIS, 400–700 nm; rainbow band), near-infrared (NIR, ~700–1400 nm), and shortwave infrared (SWIR1, ~1400–1900 nm; SWIR2, ~2000–2500 nm). Differences among spectra illustrate how optical configuration and fine-scale tissue heterogeneity influence reflectance profiles across the visible to shortwave infrared spectrum.