Symbiont interactions bias measures of arthropod biodiversity

Emily Dovydaitis[1,2,*], Chloé Schmidt[3], Michael Gerth[1,2,*]

[1]Department of Zoology, Institute for Biology, Faculty of Natural Sciences I, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany
[2]German Centre for Integrative Biodiversity Research (iDiv), Leipzig, Germany
[3]Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada
[*]Corresponding author

Corresponding authors

Emily Dovydaitis (primary) - emily.dovydaitis@zoologie.uni-halle.de
Michael Gerth - michael.gerth@zoologie.uni-halle.de

Abstract

Although arthropods comprise the majority of all named species, their biodiversity is relatively understudied. In this Perspective, we highlight how interactions between arthropods and their bacterial symbionts can pose challenges for biodiversity research, especially for estimating taxonomic, functional, and genetic diversity. We also argue that recent technological developments and biomonitoring schemes provide excellent opportunities to learn more about arthropod-symbiont interactions, consequently leading to a better understanding of arthropod biodiversity.

Introduction

With one million described species[1] and about 5-10 million species in total[2,3], arthropods account for the majority of animal species. Many arthropods are hosts to specialised, bacterial symbionts. Although the overall symbiont incidence is difficult to ascertain, at least half of all terrestrial arthropod species are estimated to carry inherited endosymbionts[4]. Bacterial symbionts play a key role in shaping their arthropod hosts' diversity[5]. For example, many arthropods feed on nutritionally unbalanced food sources (e.g. xylem, phloem, blood, wood) that lack essential nutrients. Such species typically depend on obligate symbionts to supply amino acids or vitamins, thereby shaping their hosts' feeding niches[6]. Obligate symbionts are found in 100% of their host conspecifics. Other symbionts are facultative for their hosts, changing their incidence depending on the season[7], host population[8], and presence of natural enemies[9]. Facultative symbionts can manipulate host reproduction[10,11], confer adaptations to abiotic stressors[12], enhance detoxification[12], provide defense against natural enemies[139], and even supplement host nutrition[14,15]. Symbionts impact many aspects of host biology, as well as population and community dynamics[16], influencing arthropod ecology and evolution. Despite increasing efforts to catalogue host-symbiont interactions[17], we know very little about the functional role of most arthropod associates beyond model systems.

Arthropod communities are crucial for the functioning of most ecosystems[18,19], but many species are declining[20,21]. Despite their importance, arthropods are severely under-represented in biodiversity research and conservation monitoring[22]. Many taxa have yet to be described by science and countless described species are poorly understood[23]. Limited data on arthropods makes it difficult to establish historical baselines, understand phenological shifts, or estimate temporal shifts in abundance[24]. However, recent technological developments may revert this trend[22]. Molecular methods can inform the monitoring and conservation of community composition, occupancy, species richness, and species interactions[22]. Innovations such as megabarcoding of bulk samples[25] or incorporating environmental DNA and remote sensing[26] can make large scale biodiversity studies more accessible and standardised long-term monitoring of arthropods more feasible. These technological advances, alongside novel network analysis approaches[27], also offer exciting new opportunities to detect, monitor, and study arthropod symbionts, contributing to a better understanding of arthropod biodiversity, and how host-symbiont interactions contribute to ecosystem functioning and responses to global change.

Although symbiosis is pervasive in arthropods, its significance has so far received little attention in biodiversity research. However, with the potential for wider implementation of molecular methods for monitoring arthropod biodiversity, it is increasingly important to consider symbionts and how they may interfere with biodiversity measurements. In this perspective, we will discuss how symbionts have the potential to affect biodiversity metrics in ways that obscure patterns of conservation interest (Figure 1). We argue that considering the effects of microbial symbionts is crucial for interpreting arthropod biodiversity data. First, we illustrate how

symbionts influence estimates of *taxonomic diversity* by affecting the amplification and inheritance patterns of molecular markers. Next, we show that many traits used to estimate *functional diversity* in arthropod communities are directly and dynamically altered by symbionts. We argue that symbionts can bias estimates of *genetic diversity* in arthropod populations, while simultaneously providing insight into their host's functional genetic diversity. Finally, we highlight how studying symbiotic relationships can improve predictions of biodiversity change in response to global environmental change.

Taxonomic diversity

Taxonomic diversity is used to inventory the species in an area, underpinning conservation lists[28], the basis of international agreements, and nature legislation[29]. The first step of estimating taxonomic diversity is species identification—a non-trivial task for arthropods. Although arthropod species are best delimited through integrative taxonomy methods[30], DNA barcoding is increasingly replacing other approaches[31]. In principle, DNA barcoding is possible with many loci; however, in practice, a partial sequence of the mitochondrial locus *cytochrome c oxidase subunit 1* (COI) is usually the only marker employed for barcoding arthropods[32]. In metabarcoding approaches, bulk samples of arthropods (e.g., from a malaise trap) are homogenized, DNA is extracted from the homogenate, and COI is amplified via the polymerase chain reaction (PCR). High-throughput sequencing allows cost-effective recovery of COI haplotypes which can then be assigned to a species by comparing it to a reference database (e.g., BOLD[33]). Alternatively, when references are lacking or incomplete, haplotypes can be clustered into operational taxonomic units (OTUs) based on sequence identity thresholds (often 97% similarity[34]).

Inherited symbionts can interfere with DNA barcoding protocols, directly impacting taxonomic diversity estimates. This happens for two main reasons. First, most bacterial symbionts carry a homologue of the mitochondrial COI locus which can be targeted by universal COI barcoding primers, meaning that symbiont DNA competes with host DNA for barcoding primers[35]. Fortunately, symbiont sequences can be detected and removed from the data *in silico*[36,37]. Given that PCR biases are more pronounced for some taxa than for others[38], they may systematically change the relative abundance of reads in the sample, with higher potential for rare species to go undetected. Such effects could be difficult to detect and avoid in large-scale arthropod metabarcoding studies, particularly those involving dark taxa[39] where the existence or prevalence of symbionts is unknown.

Second, like mitochondria, most common bacterial symbionts of arthropods are strictly maternally inherited. This introduces linkage disequilibrium between symbionts and mitochondrial haplotypes (i.e., symbionts and specific mitochondrial variants are inherited together), and can cause indirect selection on mitochondrial DNA (mtDNA)[40]. Because of this, symbionts can artificially inflate or reduce species diversity estimates that are based exclusively

on mtDNA[41]. The spread of symbionts in arthropod populations, for example through reproductive manipulation[42], therefore leads to the spread of specific mitochondrial haplotypes[43–45]. This phenomenon is very widespread in arthropods and generally leads to reduced mitochondrial diversity across individuals carrying a symbiont[40]. A uniform mitochondrial haplotype associated with symbiont presence can be mistaken for a taxonomic lineage distinct from individuals without the symbiont, and inflate species richness estimates. Hosts with facultative symbionts are especially prone to skewed estimates, because their incidence is more likely to vary between and within populations. Conspecifics may be infected with distinct bacterial species or strains. This can result in multiple mitochondrial haplotypes across one species, where each symbiont species or strain is associated with a different mitochondrial haplotype[46,47], once again biasing species richness estimates upwards. Conversely, symbionts can also lead to decreased species richness estimates based on mtDNA. When symbionts spread into novel species through hybrid introgression, the associated mitochondria may also spread, resulting in multiple species sharing the same symbiont and mtDNA haplotypes[48,49].

These effects of linkage disequilibrium between symbionts and mitochondria have been discussed since the advent of DNA barcoding for species identification > 20 years ago[40]. Examples have been reported from many arthropod lineages, including flies (Diptera)[47,50], butterflies and moths (Lepidoptera)[44,51–53], bees and wasps (Hymenoptera)[54], and beetles (Coleoptera)[55], which together comprise the bulk of arthropod species. Although there are currently no metastudies quantifying the extent to which symbionts impact species richness estimates, taxon-specific studies suggest that taxonomic diversity estimates based on mtDNA will be widely affected by symbionts[48]. Notably, symbionts may be gained, lost, or replaced throughout arthropod evolutionary history[56–58]. Although current symbiont presence can be detected through molecular methods and microscopy, the existence of previous or even ancient symbiont infections in a host lineage are not always known or easily detected. Thus, it is impossible to conclusively assess if, and to what extent, symbionts have had an effect on mitochondrial inheritance patterns in their hosts.

The inclusion of complementary nuclear markers for taxonomic identification would mitigate these issues. Adding nuclear markers to existing barcoding pipelines requires little additional effort and new protocols can be informed by available primer sets amplifying suitable nuclear barcoding loci. While these additional PCR protocols, just like COI amplification protocols, come with their own biases and challenges[59], including nuclear loci will typically be more informative than COI alone[60,61]. Using multiple loci for molecular species identification is already practiced in other taxa. For example, the internal transcribed spacer (ITS) is used in combination with secondary markers in fungal metabarcoding[62]. For metabarcoding arthropod communities, using additional nuclear markers would allow independent verification of diversity measurements obtained from COI barcodes, and thus alleviate biases that stem from interactions with symbionts or other sources.

Functional diversity

Functional diversity focuses on the diversity of organismal traits in a community that shapes ecosystem processes[63]. Although arthropod functional traits[64] are primarily determined by host genetics, symbionts can secondarily modify many traits related to life history, physiology, and behavior[65] (Supplemental Material Table S1). For example, symbionts heavily influence arthropod feeding niches, corresponding to the feeding guild functional trait (Table S1). Symbionts can enable hosts to exploit otherwise unavailable food sources[66,67] by aiding digestion of plant biomass[68,69], supplementing nutrients[57,70,71], and preserving nutritional resources[72], contributing to productivity and nutrient cycling at the ecosystem level. Many nutritional symbionts are obligate and therefore confer the same functional traits to all individuals of a host species. While important contributors to overall arthropod functional diversity, given their ubiquity, obligate symbionts are unlikely to bias functional diversity estimates.

On the other hand, the presence of facultative symbionts, their frequencies within and across populations[8], and the degree of their influence on functional traits[73] can change over time. This means that sampling a single population or a single time point does not capture the ways in which facultative symbionts modify host functional traits. As an example, *Bemisia tabaci,* an important agricultural pest and plant-disease vector, carries the inherited intracellular symbiont *Rickettsia.* Individuals carrying *Rickettsia* develop faster (functional trait[64]: relative growth rate) and produce more eggs (clutch size) than symbiont-free conspecifics[74]. As with other facultative endosymbionts[8], the incidence rate of *Rickettsia* has changed over time. In a North American population of *B. tabaci*, the incidence of *Rickettsia* changed in less than 20 years from nearly absent to near fixation, and subsequently declined to about one third[75]. At the same time, the impact of *Rickettsia* shifted from enhancing whitefly fitness to becoming costly and resulting in the production of non-hatching eggs[75]. As one of many bacterial symbiont lineages[76], *Rickettsia* is found in about one quarter of known arthropod species[4], so similar spatiotemporal variation in functional traits is likely to be observed across host species (Supplementary Table S1).

Facultative symbionts also complicate the compilation of arthropod trait databases. Unlike other taxa with established functional trait databases (e.g., plants: TRY[77]; mammals: PanTHERIA[78], birds: AVONET[79], fish: FishBase[80] (https://www.fishbase.se/), arthropod functional trait databases are relatively recent developments and are undergoing standardization[81]. Growing data availability necessitates and is facilitating the development of scalable biodiversity databases with individual-level traits recorded at specific locations and times. For example, the Botanical Information and Ecology Network which contains nearly 26 million unique land plant trait observations[82]. In general, however, most trait databases record static, species-level functional traits[83]. Without an abundance of samples and replicates, it is difficult for trait databases to capture intraspecific variation across space and time[84], such as those induced by facultative symbionts. As arthropod and symbiont biodiversity data accumulate, database

curators can build on scalable infrastructure developed in other taxonomic groups with longer histories of biodiversity data curation to capture intraspecific variation at higher spatiotemporal resolutions[82].

Therefore, if a functional trait is known to be influenced by symbionts, we encourage curators to, at a minimum, add this information to arthropod trait databases, so future researchers are aware and know to consider the role of symbionts. For example, the recently curated iSymBase database integrates insect functional traits that are attributed to symbionts alongside (meta)genomes[85]. If arthropod functional trait measurements shift over time, researchers can then investigate symbionts as a possible driver and to examine their incidence over time. Possible methods for detecting symbionts include PCR, multi-locus sequence typing (MLST), or fluorescence in situ hybridization (FISH) microscopy[86]. Although there are countless symbiont species, many of which are host-specific, we recommend looking for common facultative symbionts such as *Wolbachia*, *Rickettsia*, *Cardinium*, *Spiroplasma*, and *Arsenophonus* as a starting point. Ultimately, without knowledge of symbiont presence and how symbionts shape arthropod phenotypes, estimates of trait diversity may not reflect functional diversity related to ecosystem processes of interest.

Genetic diversity

The potential for a large influx of arthropod genetic data available from eDNA and metabarcoding efforts has generated interest in using these data to estimate genetic diversity for research and monitoring[32,87,88]. Genetic diversity is targeted for conservation due to its importance for adaptive evolution and population persistence in the short- and long-term[89] (https://www.cbd.int/gbf). Although multiple types of genetic diversity can be evaluated for conservation purposes (e.g., gene-specific diversity, functional genetic diversity, or neutral genome-wide genetic diversity), neutral genome-wide diversity is one of the most widely used[90]. Neutral genome-wide diversity is useful for broad-scale, multispecies assessments because it can be linked to population fitness, capacity to respond to environmental change, and common environmental perturbations such as habitat loss and fragmentation that affect population demography and species extinction risk[91].

Despite its widespread usage, the mitochondrial DNA barcoding gene COI is not useful for measuring conservation-relevant genetic diversity outside of specific purposes[92]. Although any single gene will fail to capture genome-wide diversity or the evolutionary history of species[93], the unique biology of mitochondria (non-neutral, maternally inherited, no recombination), can decouple its diversity from that of the nuclear genome[94]. In arthropods, maternally inherited symbionts shape mitochondrial variation in ways that can further decouple it from nuclear genome-wide variation, generating mito-nuclear discordance[52,95,96]. As noted in the *Taxonomic diversity* section, many common symbionts are maternally inherited, generating strong linkage disequilibrium between symbiont and mitochondrial genomes. Inherited symbionts, especially

facultative symbionts, can cause indirect selection on the mitochondria[40] that may reduce or increase mitochondrial diversity and mitochondrial effective population sizes[97]. Selective sweeps, where symbionts (and thus linked mitochondrial haplotypes) increase their frequency or become fixed in the host population, reduce mitochondrial genetic diversity[41]. This means that estimating genetic diversity from the COI gene or other mitochondrial barcoding loci will not reflect population demography or be associated with adaptive potential in general, and this disconnect from conservation goals is likely stronger in arthropod species with maternally inherited symbionts.

However, symbiont genetic diversity itself may be of interest[98]. Because symbionts moderate and generate arthropod phenotypes, the genetic diversity of symbiont genomes may reflect functional genetic diversity of their hosts, that is, genetic diversity with known relationships to adaptive traits. For example, in aphids, defense phenotypes (e.g., production of toxins against parasitoids) vary depending on symbiont genotype, and cannot be predicted based on symbiont presence alone[99,100]. Understanding genetic variation across symbiont genomes may therefore be important for predicting host response to enemy exposure. Mobile genetic elements (MGEs), such as plasmids, phages, and transposons, contribute to high genetic diversity across bacterial strains[101]. MGEs enable bacteria to store a variety of genes at the population level, thus accumulating variation and potentially leading to trait innovation[101]. The new bacterial traits that emerge from MGEs can lead to variation in the services arthropod hosts receive from their symbionts[102]. Genetic variation in symbionts can drive phenotypic differences in arthropod hosts[103], meaning it forms a component of host functional genetic diversity.

Predicting biodiversity change in response to global change

Identifying factors that determine the spatial distribution of species and how they may change in response to human-driven climate change is of central interest in biodiversity research. Abiotic environmental variables such as temperature and precipitation are often used in species distribution models to predict species range boundaries under future climate change scenarios. Biotic interactions also play an important role in shaping species' ranges and range shift dynamics[104]. For example, symbionts can facilitate range expansion[5] or enhance persistence by enabling their hosts to adapt to environmental change[56,105] and natural enemies[106–108]. On the contrary, symbionts may also limit their hosts' range by constraining the host to a particular plant or temperature range[109,110]. When predicting arthropod biodiversity change, it is important to consider their dynamic relationships with symbionts and how these relationships can be altered by environmental change.

Environmental factors mediate host-symbiont interactions[12,111–113]. Here, we focus on temperature to outline how environmental effects on symbionts may impede predictions of biodiversity change. Temperature directly affects symbiont titres[110,114], impacts host fitness[111], and often alters symbiont-mediated host phenotypes such as thermal preference[115], protection

against parasitoids[112,116] and reproductive manipulation[117]. Temperature can also influence the efficiency of vertical transmission[118] and the overall incidence of symbionts in host populations[7]. These effects of temperature on symbionts can add considerable uncertainty to modeling and predicting arthropod species distributions, as we suggest in the examples below.

Wolbachia, the most common heritable endosymbiont (present in 50% of terrestrial arthropod species[4]), is a common facultative associate of *Drosophila melanogaster* flies. Laboratory experiments suggest that flies carrying *Wolbachia* prefer cooler temperatures compared to symbiont-free flies, and differences in thermal preference between flies carrying different *Wolbachia* strains can be as high as 3°C[119]. Similar trends are found across different *Wolbachia* strains and various host species[120], indicating a general pattern of *Wolbachia*-governed thermal preference in arthropods. In natural populations of *Drosophila melanogaster*, the predominant *Wolbachia* strain was very recently replaced (< 100 years ago) across the globe[121,122]. Given the effects of strain[119] and titer[123] on host temperature preference found in laboratory experiments, it is possible that the temperature preference of natural populations also changed because of this replacement. This illustrates an inherent difficulty of estimating species distributions under climate change: upon loss or exchange of symbionts, the temperature optima for hosts may also shift. Furthermore, the occurrence of recent, rapid symbiont turnover in other arthropods suggests that this is likely a common dynamic[124–127]. Although the role of symbiosis in climate change response is well documented and regularly considered in organisms such as corals[128,129] and plant-associated fungi[130,131], it is still a nascent topic in terrestrial arthropod studies[17,115].

Host-symbiont interactions are also mutable. Host species can escape the constraints of their symbionts through replacements, in which they acquire a new symbiont[56]. In some cases, ecological shifts occur with the help of symbionts, such as a phytophagous (plant-eating) arthropod switching from one plant resource to another[56]. Likewise, symbionts can undergo host-shifting, by switching to a different, novel host species[132]. If a symbiont is adept at host-shifting, like *Wolbachia*, it will be able to increase its overall incidence and spread globally[132]. The acquisition and loss of symbionts helped generate current global biodiversity patterns[6] and continues to influence our ability to predict future ones. Through host shifting, facultative symbionts may swiftly and unpredictably change the temperature preference and therefore the geographic distribution of their hosts.

Even in seemingly static obligate symbioses with long term co-evolved partners, symbiont replacements and interactions with secondary, facultative symbionts may alter the thermal limits of their hosts in an unpredictable fashion. For nearly 200 million years aphids have codiversified with the nutritional symbiont *Buchnera*, relying on the bacteria to complement their plant sap diet with essential amino acids[58]. Short periods of heat strongly affect *Buchnera*, which in turn strongly reduces aphid fitness[133]. This is in part due to *Buchnera*'s reduced genome. Like many obligate endosymbionts, *Buchnera* experiences weak purifying selection, leading to an accumulation of mutations[134]. These mutations can contribute to thermal

instability of proteins, triggering heat sensitivity in symbionts[135]. The heat sensitivity of *Buchnera* thus limits the thermal flexibility of their aphid hosts, which has been cited as a major reason why so few aphid species colonize the tropics[136]. In some aphid species, the acquisition of facultative symbionts may help overcome these constraints by buffering the negative effects of heat on *Buchnera*[137,138]. In other aphid species, the obligate symbiont *Buchnera* was replaced[58]. Understanding the loss and gain of symbionts can help explain a host's capacity to survive in a changing environment.

Moving forward: incorporating symbionts into biodiversity research

For many commonly applied approaches to assess arthropod biodiversity, symbionts may represent an obstacle or confounding factor not usually accounted for in biodiversity studies. At the same time, our limited knowledge on widespread arthropod-symbiont interactions means it is likely that additional symbiont-mediated effects on biodiversity metrics are yet to be discovered. Furthermore, symbionts naturally fluctuate in frequency, regularly shift host species, and impact various host traits, making their impact on biodiversity estimates dynamic and difficult to predict. For these reasons, we believe it is necessary to better integrate symbiotic interactions into arthropod biodiversity research.

This may be achieved without major changes to study design. For example, the taxonomy of arthropods in biodiversity studies is often ascertained through DNA barcoding. The importance of molecular identification tools will likely increase with the advent of large-scale arthropod biomonitoring schemes[139–142] (https://pureportal.inbo.be/en/projects/metabug-monitoring-insects-with-malaise-traps-and-dna-metabarcodi/). Such projects present huge opportunities to collect barcoding data not only from arthropods, but also their symbionts. Technically, this would involve just one additional PCR per sample (targeting a bacterial barcoding locus), which may be multiplexed for convenience[143]. Adopting such new laboratory protocols would naturally result in costs and time for optimization in any lab. We anticipate that the tools developed by the microbiome community over decades of microbial DNA barcoding would strongly facilitate this adoption[144–146].

To better understand the role of symbionts in arthropod communities and predict responses to environmental change, symbionts can be directly integrated into modeling approaches such as ecological network analyses and species distribution models. Ecological network analyses can contribute to provide a mechanistic understanding of the roles symbionts play in their communities[27]. For example, symbionts can be treated as independent nodes, where they directly or indirectly interact with other nodes (species) in the community[27]. Nodes representing hosts can also be labeled as uninfected or infected, to capture differences between those with and without symbionts[27]. Likewise, the use of k-partite or multilayer networks can help predict responses to disturbances such as environmental stress or symbiont loss[27]. Species distribution models are widely used to map and predict changes in species geographic ranges. For host

345 species where symbiont incidence varies across populations, symbiont presence or absence can
346 be used as a predictor variable in addition to environmental variables. Composite species
347 distribution models conducted on data subdivided at the level of subspecies, geographic
348 regions, or genetic clusters can better model intraspecific variation[147,148]. This approach can also
349 be used to account for varied host-environment relationships due to symbiont presence or
350 genotype. Expanding focus from single host-symbiont interactions, joint species distribution
351 models can be applied to study how multiple symbiont species may simultaneously affect host
352 distributions[149].
353
354 Given symbionts' broad importance for arthropod phenotypic diversity, distributions, and
355 capacity to adapt to environmental change, symbiont communities can and should be included
356 in large scale biodiversity monitoring efforts, and ultimately integrated into conservation
357 planning and policy[150]. Directly targeting symbiont communities for biodiversity monitoring is in
358 line with the increased recognition of the importance of microbial conservation, as evidenced
359 by the newly established Microbial Conservation Specialist Group (MCSG) in the International
360 Union for Conservation of Nature[151]. Moreover, complementing existing arthropod monitoring
361 schemes with symbiont metabarcoding would give unprecedented insight into the diversity of
362 arthropod-symbiont interactions[16] . When considering the existing labor, travel, and equipment
363 costs of arthropod monitoring programs, the additional sequencing costs for symbiont
364 barcoding are negligible. In light of the extensive and invasive nature of arthropod sampling,
365 such additional data generation would also be a sensible and frugal use of biological resources.
366 Examining the temporal, spatial, and environmental variation in symbiont interactions would
367 provide a better, more integrative understanding of arthropod biodiversity and community
368 functioning[27].

Acknowledgements

We thank the Evolution and Biodiversity research group for their valuable input and support.

Author Contributions

ED, CS, and MG conceptualized the project. ED wrote the first draft of the manuscript and created the visual. ED, CS, and MG contributed to reviewing and editing the subsequent manuscript drafts. All authors have all read and approved the manuscript.

Competing Interests

The authors declare no financial or non-financial competing interests.

Funding

MG is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – 497854142. The work was supported by the German Centre for Integrative Biodiversity Research (iDiv), funded by the DFG (FZT 118, 202548816).

References

1. Li, X. & Wiens, J. J. Estimating Global Biodiversity: The Role of Cryptic Insect Species. *Syst. Biol.* **72**, 391–403 (2023).

2. Ødegaard, F. How many species of arthropods? Erwin's estimate revised. *Biological Journal of the Linnean Society* **2**, 583–597 (1999).

3. Stork, N. E. How Many Species of Insects and Other Terrestrial Arthropods Are There on Earth? *Annu. Rev. Entomol.* 31–45 (2018).

4. Weinert, L. A., Araujo-Jnr, E. V., Ahmed, M. Z. & Welch, J. J. The incidence of bacterial endosymbionts in terrestrial arthropods. *Proceedings of the Royal Society B: Biological Sciences* **282**, (2015).

5. Holt, J. R., Cavichiolli de Oliveira, N., Medina, R. F., Malacrinò, A. & Lindsey, A. R. I. Insect–microbe interactions and their influence on organisms and ecosystems. *Ecol. Evol.* **14**, 1–19 (2024).

6. Cornwallis, C. K. *et al.* Symbioses shape feeding niches and diversification across insects. *Nat. Ecol. Evol.* **7**, 1022–1044 (2023).

7. Martins, M., César, C. S. & Cogni, R. The effects of temperature on prevalence of facultative insect heritable symbionts across spatial and seasonal scales. *Front. Microbiol.* **14**, (2023).

8. Hoffmann, A. A. & Cooper, B. S. Changes in the frequency of facultative endosymbionts in insect populations: overview and applications. *Entomologia Generalis* **45**, 351–368 (2025).

9. Oliver, K. M., Smith, A. H. & Russell, J. A. Defensive symbiosis in the real world - advancing ecological studies of heritable, protective bacteria in aphids and beyond. *Funct. Ecol.* **28**, 341–355 (2014).

10. Engelstädter, J. & Hurst, G. D. D. The ecology and evolution of microbes that manipulate host reproduction. *Annu. Rev. Ecol. Evol. Syst.* **40**, 127–149 (2009).

11. Hurst, G. D. D. & Frost, C. L. Reproductive parasitism: Maternally inherited symbionts in a biparental world. *Cold Spring Harb. Perspect. Biol.* **7**, 1–21 (2015).

12. Lemoine, M. M., Engl, T. & Kaltenpoth, M. Microbial symbionts expanding or constraining abiotic niche space in insects. *Curr. Opin. Insect Sci.* **39**, 14–20 (2020).

13. Łukasik, P., van Asch, M., Guo, H., Ferrari, J. & Charles, H. Unrelated facultative endosymbionts protect aphids against a fungal pathogen. *Ecol. Lett.* **16**, 214–218 (2013).

14. Lindsey, A. R. I. *et al.* The intracellular symbiont Wolbachia alters Drosophila development and metabolism to buffer against nutritional stress. *PLoS Genet.* **21**, 01–20 (2025).

15. Preuß, L. *et al.* Rapid spread of nutritional Wolbachia symbionts across and within solitary bee species. *bioRxiv* (2026).

16. Łukasik, P. & Kolasa, M. R. With a little help from my friends: The roles of microbial symbionts in insect populations and communities. *Philosophical Transactions of the Royal Society B: Biological Sciences* **379**, (2024).

17. Graber, L. C. & Moreau, C. S. Insect–microbiome interactions in a changing world. *Curr. Opin. Insect Sci.* **75**, (2026).

18. Wilson, E. O. The little things that run the world* (the importance and conservation of invertebrates. *Conserv. Biol.* **1**, 344–346 (1987).

19. Schowalter, T. D. *Insect Ecology*. (Academic Press, San Diego, CA, 2016).

20. Eisenhauer, N., Bonn, A. & A. Guerra, C. Recognizing the quiet extinction of invertebrates. *Nature Communications* **10**, (2019).

21. Scherber, C., Kirse, A., Kempel, A., Ott, D. & Hartke, T. Insect declines: challenges and solutions. *Curr. Opin. Insect Sci.* **75**, (2026).

22. Van Klink, R. *et al.* Towards a toolkit for global insect biodiversity monitoring. *Philosophical Transactions of the Royal Society B: Biological Sciences* **379**, (2024).

23. The science of insect taxonomy: Prospects and needs. in *Insect Biodiversity* 499–526 (John Wiley & Sons, Ltd, Chichester, UK, 2017).

24. Didham, R. K. *et al.* Interpreting insect declines: seven challenges and a way forward. *Insect Conserv. Divers.* **13**, 103–114 (2020).

25. Meier, R., Hartop, E., Pylatiuk, C. & Srivathsan, A. Towards holistic insect monitoring: Species discovery, description, identification and traits for all insects. *Philosophical Transactions of the Royal Society B: Biological Sciences* **379**, (2024).

26. Li, Y. *et al.* Combining environmental DNA and remote sensing for efficient, fine-scale mapping of arthropod biodiversity. *Philosophical Transactions of the Royal Society B: Biological Sciences* **379**, (2024).

27. Morente, M., Hackett, T. D. & McLean, A. H. C. Mapping microbial symbiont impacts using ecological networks. *Trends Ecol. Evol.* **41**, 384–394 (2026).

28. Salem, H. *et al.* Vitamin supplementation by gut symbionts ensures metabolic homeostasis in an insect host. *Proceedings of the Royal Society B: Biological Sciences* **281**, (2014).

29. Garnett, S. T. & Christidis, L. Taxonomy anarchy hampers conservation. *Nature* **546**, 25–27 (2017).

30. Schlick-Steiner, B. C. *et al.* Integrative taxonomy: A multisource approach to exploring biodiversity. *Annu. Rev. Entomol.* **55**, 421–438 (2010).

31. Jinbo, U., Kato, T. & Ito, M. Current progress in DNA barcoding and future implications for entomology. *Entomol. Sci.* **14**, 107–124 (2011).

32. Yu, D. W. *et al*. Biodiversity soup: Metabarcoding of arthropods for rapid biodiversity assessment and biomonitoring. *Methods Ecol. Evol.* **3**, 613–623 (2012).

33. Ratnasingham, S. & Hebert, P. D. N. BOLD: The Barcode of Life Data System. *Mol. Ecol. Notes* 355–364 (2007).

34. Meier, R. *et al*. "Dark taxonomy": A new protocol for overcoming the taxonomic impediments for dark taxa and broadening the taxon base for biodiversity assessment. *Cladistics* **41**, 223–238 (2025).

35. Park, E. & Poulin, R. Widespread Torix Rickettsia in New Zealand amphipods and the use of blocking primers to rescue host COI sequences. *Sci. Rep.* **10**, (2020).

36. Smith, M. A. *et al*. Wolbachia and DNA barcoding insects: Patterns, potential, and problems. *PLoS One* **7**, (2012).

37. Pilgrim, J. *et al*. Torix Rickettsia are widespread in arthropods and reflect a neglected symbiosis. *Gigascience* **10**, (2021).

38. Bleidorn, C. & Henze, K. A new primer pair for barcoding of bees (Hymenoptera: Anthophila) without amplifying the orthologous coxA gene of Wolbachia bacteria. *BMC Res. Notes* **14**, (2021).

39. Page, R. D. M. DNA barcoding and taxonomy: Dark taxa and dark texts. *Philosophical Transactions of the Royal Society B: Biological Sciences* **371**, (2016).

40. Hurst, G. D. D. & Jiggins, F. M. Problems with mitochondrial DNA as a marker in population, phylogeographic and phylogenetic studies: The effects of inherited symbionts. *Proceedings of the Royal Society B: Biological Sciences* **272**, 1525–1534 (2005).

41. Deng, J. *et al*. Wolbachia-driven selective sweep in a range expanding insect species. *BMC Ecol. Evol.* **21**, (2021).

42. Perlman, S. J., Hodson, C. N., Hamilton, P. T., Opit, G. P. & Gowen, B. E. Maternal transmission, sex ratio distortion, and mitochondria. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 10162–10168 (2015).

43. Duplouy, A., Hurst, G. D. D., O'Neill, S. L. & Charlat, S. Rapid spread of male-killing wolbachia in the butterfly hypolimnas bolina. *J. Evol. Biol.* **23**, 231–235 (2010).

44. Miyata, M., Nomura, M. & Kageyama, D. Elusive origin of mitochondria shared by two sister species of Eurema butterflies. *BMC Ecol. Evol.* **25**, (2025).

45. Kodandaramaiah, U., Simonsen, T. J., Bromilow, S., Wahlberg, N. & Sperling, F. Deceptive single-locus taxonomy and phylogeography: Wolbachia-associated divergence in mitochondrial DNA is not reflected in morphology and nuclear markers in a butterfly species. *Ecol. Evol.* **3**, 5167–5176 (2013).

46. Sun, X. J., Xiao, J. H., Cook, J. M., Feng, G. & Huang, D. W. Comparisons of host mitochondrial, nuclear and endosymbiont bacterial genes reveal cryptic fig wasp species

and the effects of Wolbachia on host mtDNA evolution and diversity. *BMC Evol. Biol.* **11**, (2011).

47. Symula, R. E. *et al.* Influence of host phylogeographic patterns and incomplete lineage sorting on within-species genetic variability in Wigglesworthia species, obligate symbionts of tsetse flies. *Appl. Environ. Microbiol.* **77**, 8400–8408 (2011).

48. Whitworth, T. L., Dawson, R. D., Magalon, H. & Baudry, E. DNA barcoding cannot reliably identify species of the blowfly genus Protocalliphora (Diptera: Calliphoridae). *Proceedings of the Royal Society B: Biological Sciences* **274**, 1731–1739 (2007).

49. Klopfstein, S., Kropf, C. & Baur, H. Wolbachia endosymbionts distort DNA barcoding in the parasitoid wasp genus Diplazon (Hymenoptera: Ichneumonidae). *Zool. J. Linn. Soc.* **177**, 541–557 (2016).

50. Dyer, K. A., Burke, C. & Jaenike, J. Wolbachia-mediated persistence of mtDNA from a potentially extinct species. *Mol. Ecol.* **20**, 2805–2817 (2011).

51. Narita, S., Nomura, M., Kato, Y. & Fukatsu, T. Genetic structure of sibling butterfly species affected by Wolbachia infection sweep: Evolutionary and biogeographical implications. *Mol. Ecol.* **15**, 1095–1108 (2006).

52. Gaunet, A. *et al.* Two consecutive Wolbachia-mediated mitochondrial introgressions obscure taxonomy in Palearctic swallowtail butterflies (Lepidoptera, Papilionidae). *Zool. Scr.* **48**, 507–519 (2019).

53. Charlat, S. *et al.* The joint evolutionary histories of Wolbachia and mitochondria in Hypolimnas bolina. *BMC Evol. Biol.* **9**, (2009).

54. Gerth, M., Geißler, A. & Bleidorn, C. Wolbachia infections in bees (Anthophila) and possible implications for DNA barcoding. *Syst. Biodivers.* **9**, 319–327 (2011).

55. Jäckel, R., Mora, D. & Dobler, S. Evidence for selective sweeps by Wolbachia infections: Phylogeny of Altica leaf beetles and their reproductive parasites. *Mol. Ecol.* **22**, 4241–4255 (2013).

56. Sudakaran, S., Kost, C. & Kaltenpoth, M. Symbiont Acquisition and Replacement as a Source of Ecological Innovation. *Trends Microbiol.* **25**, 375–390 (2017).

57. Koga, R. & Moran, N. A. Swapping symbionts in spittlebugs: Evolutionary replacement of a reduced genome symbiont. *ISME Journal* **8**, 1237–1246 (2014).

58. Chong, R. A. & Moran, N. A. Evolutionary loss and replacement of Buchnera, the obligate endosymbiont of aphids. *ISME Journal* **12**, 898–908 (2018).

59. Krehenwinkel, H., Kennedy, S. R., Rueda, A., Lam, A. & Gillespie, R. G. Scaling up DNA barcoding – Primer sets for simple and cost efficient arthropod systematics by multiplex PCR and Illumina amplicon sequencing. *Methods Ecol. Evol.* **9**, 2181–2193 (2018).

60. Hartig, G. *et al.* Oligonucleotide primers for targeted amplification of single-copy nuclear genes in apocritan Hymenoptera. *PLoS One* **7**, (2012).

61. Wahlberg, N., Peña, C., Ahola, M., Wheat, C. W. & Rota, J. PCR primers for 30 novel gene regions in the nuclear genomes of lepidoptera. *Zookeys* **2016**, 129–141 (2016).

62. Lücking, R. *et al*. Unambiguous identification of fungi: Where do we stand and how accurate and precise is fungal DNA barcoding? *IMA Fungus* **11**, (2020).

63. Petchey, O. L. & Gaston, K. J. Functional diversity: Back to basics and looking forward. *Ecol. Lett.* **9**, 741–758 (2006).

64. Moretti, M. *et al*. Handbook of protocols for standardized measurement of terrestrial invertebrate functional traits. *Funct. Ecol.* **31**, 558–567 (2017).

65. Feldhaar, H. Bacterial symbionts as mediators of ecologically important traits of insect hosts. *Ecol. Entomol.* **36**, 533–543 (2011).

66. Salem, H. & Kaltenpoth, M. Beetle-Bacterial Symbioses: Endless Forms Most Functional. *Annu. Rev. Entomol. 2022* **67**, 2021 (2022).

67. Kaltenpoth, M., Flórez, L. V., Vigneron, A., Dirksen, P. & Engl, T. Origin and function of beneficial bacterial symbioses in insects. *Nat. Rev. Microbiol.* (2025).

68. Salem, H. *et al*. Drastic Genome Reduction in an Herbivore's Pectinolytic Symbiont. *Cell* **171**, 1520-1531.e13 (2017).

69. Reis, F. *et al*. Bacterial symbionts support larval sap feeding and adult folivory in (semi-)aquatic reed beetles. *Nat. Commun.* **11**, (2020).

70. Nikoh, N. *et al*. Evolutionary origin of insect-Wolbachia nutritional mutualism. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 10257–10262 (2014).

71. Hansen, A. K. & Moran, N. A. The impact of microbial symbionts on host plant utilization by herbivorous insects. *Mol. Ecol.* **23**, 1473–1496 (2014).

72. Shukla, S. P. *et al*. Microbiome-assisted carrion preservation aids larval development in a burying beetle. *Proc. Natl. Acad. Sci. U. S. A.* **115**, 11274–11279 (2018).

73. Nakayama, S. *et al*. Can maternally inherited endosymbionts adapt to a novel host? Direct costs of Spiroplasma infection, but not vertical transmission efficiency, evolve rapidly after horizontal transfer into D. melanogaster. *Heredity (Edinb).* **114**, 539–543 (2015).

74. Himler, A. G. *et al*. Rapid spread of a bacterial symbiont in an invasive whitefly is driven by fitness benefits and female bias. *Science (1979).* **332**, 254–256 (2011).

75. Bockoven, A. A. *et al*. What Goes Up Might Come Down: the Spectacular Spread of an Endosymbiont Is Followed by Its Decline a Decade Later. *Microb. Ecol.* **79**, 482–494 (2020).

76. Duron, O. *et al*. The diversity of reproductive parasites among arthropods: Wolbachia do not walk alone. *BMC Biol.* **6**, (2008).

77. Kattge, J. *et al*. TRY - a global database of plant traits. *Glob. Chang. Biol.* **17**, 2905–2935 (2011).

78. Jones, K. E. *et al.* PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecological Archives* **90**, 2648 (2009).

79. Tobias, J. A. *et al.* AVONET: morphological, ecological and geographical data for all birds. *Ecol. Lett.* **25**, 581–597 (2022).

80. Froese, R. , D. Pauly. Fishbase.

81. Brousseau, P. M., Gravel, D. & Handa, I. T. On the development of a predictive functional trait approach for studying terrestrial arthropods. *Journal of Animal Ecology* **87**, 1209–1220 (2018).

82. Enquist, B. J. *et al.* BIEN: A biodiversity informatics ecosystem advancing open and reproducible workflows for plant observation, plot and trait data. *Methods Ecol. Evol.* **17**, 1556–1584 (2026).

83. Middleton-Welling, J. *et al.* A new comprehensive trait database of European and Maghreb butterflies, Papilionoidea. *Sci. Data* **7**, (2020).

84. Kattge, J. *et al.* TRY plant trait database – enhanced coverage and open access. *Glob. Chang. Biol.* **26**, 119–188 (2020).

85. Wang, Z. *et al.* iSymBase: an integrative functional-genomic platform for ecological exploration of insect symbionts. *ISME Communications* **6**, (2026).

86. Nevalainen, L. B. M. & Newton, I. L. G. Detection and Assessment of Wolbachia pipientis Infection. *Methods Mol. Biol.* **2626**, 291–307 (2023).

87. Takenaka, M., Uchida, N., Hasebe, Y., Yano, K. & Ueki, G. Environmental DNA metabarcoding for insect biodiversity surveys: Current status and future prospects for potential applications in highly diverse insect taxa. *Entomol. Sci.* **29**, (2026).

88. Emerson, B. C. *et al.* Collective and harmonized high throughput barcoding of insular arthropod biodiversity: Toward a Genomic Observatories Network for islands. *Mol. Ecol.* **32**, 6161–6176 (2023).

89. Secretariat of the Convention on Biological Diversity. Kunming-Montreal Global Biodiversity Framework. (2026).

90. Ellegren, H. & Galtier, N. Determinants of genetic diversity. *Nat. Rev. Genet.* **17**, 422–433 (2016).

91. Schmidt, C., Hoban, S. & Jetz, W. Conservation macrogenetics: harnessing genetic data to meet conservation commitments. *Trends in Genetics* **39**, 816–829 (2023).

92. Rubinoff, D. Utility of mitochondrial DNA barcodes in species conservation. *Conservation Biology* **20**, 1026–1033 (2006).

93. Nichols, R. Gene trees and species trees are not the same. *Trends Ecol. Evol.* **16**, 358–364 (2001).

94. Galtier, N., Nabholz, B., Glémin, S. & Hurst, G. D. D. Mitochondrial DNA as a marker of molecular diversity: A reappraisal. *Mol. Ecol.* **18**, 4541–4550 (2009).

95. Toews, D. P. L. & Brelsford, A. The biogeography of mitochondrial and nuclear discordance in animals. *Mol. Ecol.* **21**, 3907–3930 (2012).

96. Park, E. & Poulin, R. Extremely divergent COI sequences within an amphipod species complex: A possible role for endosymbionts? *Ecol. Evol.* **12**, 1–15 (2022).

97. Dyer, K. A. & Jaenike, J. Evolutionary dynamics of a spatially structured host-parasite association: Drosophila innubila and male-killing Wolbachia. *Evolution (N. Y).* **59**, 1518-1528. (2005).

98. Russell, S. L. & Cavanaugh, C. M. Intrahost genetic diversity of bacterial symbionts exhibits evidence of mixed infections and recombinant haplotypes. *Mol. Biol. Evol.* **34**, 2747–2761 (2017).

99. McLean, A. H. C. *et al.* Multiple phenotypes conferred by a single insect symbiont are independent. *Proceedings of the Royal Society B: Biological Sciences* **287**, 1–9 (2020).

100. Panossian, B. *et al.* Phage toxin variants are linked to protection specificity in a defensive symbiont. *Mol. Biol. Evol.* **43**, (2026).

101. Weisberg, A. J. & Chang, J. H. Mobile Genetic Element Flexibility as an Underlying Principle to Bacterial Evolution. *Annu. Rev. Microbiol.* **56**, 33 (2026).

102. Patel, V. *et al.* Mobile elements create strain-level variation in the services conferred by an aphid symbiont. *Environ. Microbiol.* **25**, 3333–3348 (2023).

103. Panossian, B. *et al.* Phenotypic divergence is driven by mobile genetic elements in a heritable insect symbiont. *Proceedings of the National Academy of Sciences* **123**, (2026).

104. Osmolovsky, I., Xirocostas, Z. A., Chiarenza, G. M. & Moles, A. T. Counterintuitive Range Shifts May Be Explained by Climate Induced Changes in Biotic Interactions. *Glob. Chang. Biol.* **31**, (2025).

105. Engl, T. *et al.* Ancient symbiosis confers desiccation resistance to stored grain pest beetles. *Mol. Ecol.* **27**, 2095–2108 (2017).

106. Xie, J., Butler, S., Sanchez, G. & Mateos, M. Male killing Spiroplasma protects Drosophila melanogaster against two parasitoid wasps. *Heredity (Edinb).* **112**, 399–408 (2014).

107. Jaenike, J., Unckless, R., Cockburn, S. N., Boelio, L. M. & Perlman, S. J. Adaptation via Symbiosis: Recent Spread of a Drosophila Defensive Symbiont. *Science (1979).* **329**, 212–215 (2010).

108. Xie, J., Winter, C., Winter, L. & Mateos, M. Rapid spread of the defensive endosymbiont Spiroplasma in Drosophila hydei under high parasitoid wasp pressure. *FEMS Microbiol. Ecol.* **91**, 1–11 (2015).

109. Bennett, G. M. & Moran, N. A. Heritable symbiosis: The advantages and perils of an evolutionary rabbit hole. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 10169–10176 (2015).

110. Fan, Y. & Wernegreen, J. J. Can't Take the Heat: High Temperature Depletes Bacterial Endosymbionts of Ants. *Microb. Ecol.* **66**, 727–733 (2013).

111. Kikuchi, Y. *et al.* Collapse of insect gut symbiosis under simulated climate change. *mBio* **7**, (2016).

112. Patel, V., Kucuk, R. A., Haines-Eitzen, B. R., Russell, J. A. & Oliver, K. M. Emergent symbiont strains provide thermally robust protection against co-evolved and novel parasitoids of introduced pea aphids. *ISME Journal* **20**, (2026).

113. Beraldo, C. S., Van Nouhuys, S. & Duplouy, A. How environmental conditions affect the acquisition, establishment, and persistence of microbial endosymbiosis in insects. *EcoEvoRxiv* (2026).

114. Corbin, C., Jones, J. E., Chrostek, E., Fenton, A. & Hurst, G. D. D. Thermal sensitivity of the Spiroplasma–Drosophila hydei protective symbiosis: The best of climes, the worst of climes. *Mol. Ecol.* **30**, 1336–1344 (2021).

115. Corbin, C., Heyworth, E. R., Ferrari, J. & Hurst, G. D. D. Heritable symbionts in a world of varying temperature. *Heredity (Edinb).* **118**, 10–20 (2017).

116. Doremus, M. R. *et al.* Breakdown of a defensive symbiosis, but not endogenous defences, at elevated temperatures. *Mol. Ecol.* **27**, 2138–2151 (2018).

117. Montenegro, H. & Klaczko, L. B. Low temperature cure of a male killing agent in Drosophila melanogaster. *J. Invertebr. Pathol.* **86**, 50–51 (2004).

118. Hague, M. T. J. *et al.* Temperature effects on cellular host-microbe interactions explain continent-wide endosymbiont prevalence. *Current Biology* **32**, 878-888.e8 (2022).

119. Truitt, A. M., Kapun, M., Kaur, R. & Miller, W. J. Wolbachia modifies thermal preference in Drosophila melanogaster. *Environ. Microbiol.* **21**, 3259–3268 (2019).

120. Hague, M. T. J., Caldwell, C. N. & Cooper, B. S. Pervasive effects of Wolbachia on host temperature preference. *mBio* **11**, 1–15 (2020).

121. Riegler, M., Sidhu, M., Miller, W. J. & O'Neill, S. L. Evidence for a global Wolbachia replacement in Drosophila melanogaster. *Current Biology* **15**, 1428–1433 (2005).

122. Strunov, A. *et al.* Historic Museum Samples Provide Evidence for a Recent Replacement of Wolbachia Types in European Drosophila melanogaster. *Mol. Biol. Evol.* **40**, (2023).

123. Strunov, A., Schönherr, C. & Kapun, M. Wolbachia effects on thermal preference of natural Drosophila melanogaster are influenced by host genetic background, Wolbachia type, and bacterial titer. *Environ. Microbiol.* **26**, (2024).

124. Schuler, H. *et al.* Evidence for a recent horizontal transmission and spatial spread of Wolbachia from endemic Rhagoletis cerasi (Diptera: Tephritidae) to invasive Rhagoletis cingulata in Europe. *Mol. Ecol.* **22**, 4101–4111 (2013).

125. Kriesner, P., Hoffmann, A. A., Lee, S. F., Turelli, M. & Weeks, A. R. Rapid Sequential Spread of Two Wolbachia Variants in Drosophila simulans. *PLoS Pathog.* **9**, (2013).

126. Turelli, M. *et al.* Rapid Global Spread of wRi-like Wolbachia across Multiple Drosophila. *Current Biology* **28**, 963-971.e8 (2018).

127. Bakovic, V., Schebeck, M., Telschow, A., Stauffer, C. & Schuler, H. Spatial spread of wolbachia in rhagoletis cerasi populations. *Biol. Lett.* **14**, (2018).

128. Goulet, T. L. & Goulet, D. Climate Change Leads to a Reduction in Symbiotic Derived Cnidarian Biodiversity on Coral Reefs. *Front. Ecol. Evol.* **9**, (2021).

129. Parker, K. E. & Quigley, K. M. The coral microbiome under climate change: shifting symbiotic interactions. *Microbiol. Aust.* (2026).

130. Kivlin, S. N., Emery, S. M. & Rudgers, J. A. Fungal symbionts alter plant responses to global change. *Am. J. Bot.* **100**, 1445–1457 (2013).

131. Bennett, A. E., Ee, A. & Classen, T. Climate change influences mycorrhizal fungal-plant interactions, but conclusions are limited by geographical study bias. (2020).

132. Sanaei, E., Charlat, S. & Engelstädter, J. Wolbachia host shifts: routes, mechanisms, constraints and evolutionary consequences. *Biological Reviews* **96**, 433–453 (2021).

133. Zhang, B., Leonard, S. P., Li, Y. & Moran, N. A. Obligate bacterial endosymbionts limit thermal tolerance of insect host species. *PNAS* **116**, 24712–24718 (2019).

134. Moran, N. A. & Bennett, G. M. The tiniest tiny genomes. *Annu. Rev. Microbiol.* **68**, 195–215 (2014).

135. Perreau, J. & Moran, N. A. Genetic innovations in animal–microbe symbioses. *Nat. Rev. Genet.* **23**, 23–39 (2022).

136. Perkovsky, E. & Wegierek, P. Aphid-Buchnera-Ant symbiosis; Or why are aphids rare in the tropics and very rare further south? *Earth Environ. Sci. Trans. R. Soc. Edinb.* **107**, 297–310 (2018).

137. Heyworth, E. R., Smee, M. R. & Ferrari, J. Aphid Facultative Symbionts Aid Recovery of Their Obligate Symbiont and Their Host After Heat Stress. *Front. Ecol. Evol.* **8**, (2020).

138. Tougeron, K. *et al.* Still standing: The heat protection delivered by a facultative symbiont to its aphid host is resilient to repeated thermal stress. *Current Research in Insect Science* **3**, (2023).

139. Karlsson, D., Hartop, E., Forshage, M., Jaschhof, M. & Ronquist, F. The Swedish malaise trap project: A 15 year retrospective on a countrywide insect inventory. *Biodivers. Data J.* **8**, (2018).

140. Hobern, D. Bioscan: Dna barcoding to accelerate taxonomy and biogeography for conservation and sustainability. *Genome* **64**, 161–164 (2021).

141. Streitberger, M. *et al.* The German insect monitoring scheme: Establishment of a nationwide long-term recording of arthropods. *Basic Appl. Ecol.* **80**, 81–91 (2024).

142. Research Institute for Nature and Forest. METABUG: Monitoring insects with Malaise traps and DNA metabarcoding. . (2026).

143. Nowak, K. H. *et al.* What lurks in the dark? An innovative framework for studying diverse wild insect microbiota. *Microbiome* **13**, (2025).

144. Gibson, J. *et al.* Simultaneous assessment of the macrobiome and microbiome in a bulk sample of tropical arthropods through DNA metasystematics. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 8007–8012 (2014).

145. Combrink, L. *et al.* Best practice for wildlife gut microbiome research: A comprehensive review of methodology for 16S rRNA gene investigations. *Front. Microbiol.* **14**, (2023).

146. Whelan, F. J. & Hall, L. J. An ode to the 16S rRNA gene: its history, importance, caveats and future in microbiome research. *Microb. Genom.* **12**, (2026).

147. Gonzalez, S. C., Soto-Centeno, J. A. & Reed, D. L. Population distribution models: Species distributions are better modeled using biologically relevant data partitions. *BMC Ecol.* **11**, (2011).

148. Gotelli, N. J. & Stanton-Geddes, J. Climate change, genetic markers and species distribution modelling. *J. Biogeogr.* **42**, 1577–1585 (2015).

149. Ovaskainen, O., Łukasik, P. & Roslin, T. Joint species distribution modelling of insect microbiota: Time to jump onto a new opportunity. *EcoEvoRxiv* (2026).

150. Peixoto, R. S. *et al.* Harnessing the microbiome to prevent global biodiversity loss. *Nat. Microbiol.* **7**, 1726–1735 (2022).

151. Gilbert, J. A. *et al.* Launching the IUCN Microbial Conservation Specialist Group as a global safeguard for microbial biodiversity. *Nat. Microbiol.* **10**, 2359–2360 (2025).

Figures

Figure 1.




Figure title: Bacterial symbionts bias how the taxonomic, functional, and genetic diversity of arthropods are estimated.

Legend: *Taxonomic diversity:* Due to linkage disequilibrium between mitochondria and bacterial symbionts, species richness estimates based on mitochondrial DNA (mtDNA) may be higher or lower than those made using nuclear DNA (nuDNA). *Functional diversity:* Although many traits overlap between species with and without symbionts, symbionts can confer or alter many arthropod functional traits (e.g., nutrition, defense, reproduction). *Genetic diversity:* Selective sweeps of mitochondria alongside vertically transmitted, bacterial symbionts can reduce genetic diversity estimates based on mtDNA.

Supplementary information: Symbiont interactions bias measures of arthropod biodiversity

Emily Dovydaitis[1,2,*], Chloé Schmidt[3], Michael Gerth[1,2,*]

[1]Department of Zoology, Institute for Biology, Faculty of Natural Sciences I, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany

[2]German Centre for Integrative Biodiversity Research (iDiv), Leipzig, Germany

[3]Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada

*To whom correspondence should be addressed

Table S1: Examples of how symbionts shape arthropod functional traits.

The functional traits are grouped by category (morphology, feeding, life history, physiology, behaviour). Each trait includes an example, along with the relevant host and symbiont species. The categories and traits are based on a handbook of invertebrate functional traits[1]. Functional traits not included in the handbook are indicated with an asterisk (*). The table is comprehensive, but not exhaustive or systematic (i.e., it covers a broad range of examples, but does not include all relevant studies.)

Category	Functional Trait	Example	Host	Symbiont	Ref.
Morphology	Body size	Symbiont presence associated with larger progeny.	Red spider mite *Tetranychus urticae*	*Wolbachia*	[2]
		Higher host body size and water content compared to aposymbiotic conspecifics.	Rice weevil *Sitophilus oryzae*	*Sodalis pierantonius*	[3]
		Loss of symbiont results in stunted growth and reduced body size.	Southern green stink bug *Nezara viridula*	Gut bacterial symbiont (unnamed)	[4]
	Respiration system	Presence of symbiont triggers development of an extensive tracheal network that envelops the host's gut.	Bean bug *Riptortus pedestris*	*Burkholderia insecticola*	[5]
	Colour	Aposymbiotic offspring are paler compared to those with symbiont.	West Indian sweet potato weevil *Euscepes postfasciatus*	*Nardonella*	[6]
		After losing its symbiont, the host turns a yellowish colour.	Southern green stink bug *Nezara viridula*	Gut bacterial symbiont (unnamed)	[4]
	Cuticle hardness*	Symbiont provisions tyrosine, which is used for host cuticle synthesis.	Superfamily Curculionoidea	*Nardonella*	[7]
	Sex traits*	Presence of symbiont induces feminization in its host, including appearance of female genital apertures.	Various isopod species (e.g., *Armadillidium vulgare*, *A. nasatum*)	*Wolbachia*	[8,9]
Feeding	Feeding guild	Symbiont provisions amino acids and vitamins for sap-feeding hosts, whose diet is high in sugar but low in other nutrients.	Various Auchenorrhyncha and Sternorrhyncha species	Various bacterial symbionts *(Sulcia muelleri, Sodalis glossinidius, Buchnera aphidicola* etc.)	[10–14]
		Symbiont helps its host procure B vitamins, which are lacking from the host's blood-based diet.	Bed bug *Cimex lectularius*	*Wolbachia*	[15]
		The host's ability to digest and live on a diet of lignocellulose depends on its symbionts.	Eastern subterranean termite *Reticulitermes flavipes*	Various protist and bacterial gut microbes	[16]
	Ingestion rate	Symbiont infection increases feeding rate and weight gain in the host.	Silkworm *Bombyx mori*	*Cordyceps militari*	[17]
		After mating with symbiont infected males, host females have a lower feeding frequency.	Fruit fly *Drosophila melanogaster*	*Wolbachia*	[18]

Life history	Ontogeny	Symbiont proliferates dramatically after metamorphosis and synthesizes amino acids needed for building a protective exoskeleton.	*Cereal weevil* *Sitophilus spp.*	*Sodalis pierantonius*	[19]
		Individuals with symbionts developed faster than those without.	Silverleaf whitefly *Bemisia tabaci*	*Rickettsia*	[20]
	Clutch size	Individuals with symbionts produce more offspring compared to uninfected individuals.	Silverleaf whitefly *Bemisia tabaci*	*Rickettsia*	[20]
		Individuals with symbionts have more eggs in their ovaries compared to aposymbiotic individuals.	Fruit fly *Drosophila neotestacea*	*Spiroplasma*	[21]
	Egg size	Maternal fitness and egg size increase when infected with one symbiont, but decrease when infected by a different symbiont.	*Kelly's citrus thrips* *Pezothrips kellyanus*	*Cardinium* (increase), *Wolbachia* (decrease)	[22]
		Egg size increases with symbiont infection, which in turn promotes fertilization.	Red spider mite *Tetranychus urticae*	*Wolbachia*	[2]
	Egg production*	Symbiont plays a role in egg production. Following antibiotics treatment, the host has lower fecundity and eggs are no longer viable.	*Rice water weevil* *Lissorhoptrus oryzophilus*	*Wolbachia*	[23]
	Life span	Individuals with the symbiont infection were more likely to survive into adulthood.	Silverleaf whitefly *Bemisia tabaci*	*Rickettsia*	[20]
		Symbiont infection decreases larval survival and adult longevity.	Mill moth *Ephestia kuehniella*	*Wolbachia*	[24]
		Symbiont infection decreases longevity.	Fruit fly *Drosophila melanogaster*	*Spiroplasma*	[25,26]
	Age at maturity	Individuals with symbiont reach adulthood faster than those without.	Silverleaf whitefly *Bemisia tabaci*	*Rickettsia*	[27]
	Reproductive mode	Parthenogenesis: Symbiont infection is linked to parthenogenesis	Broad-nosed weevils Tribe *Naupactini*	*Wolbachia*	[28,29]

		(Asexual reproduction, where embryos develop from unfertilized eggs.)	Parasitoid wasp *Pnigalio soemius*	*Rickettsia*	30
		Cytoplasmic incompatibility (CI): Symbiont infection triggers CI (mating between infected males and uninfected females results in embryo death.)	Spider mite *Eotetranychus suginamensis*	*Cardinium*	31
			Parasitoid wasp *Encarsia inaron*	*Cardinium, Wolbachia*	32
		Male-killing (MK): Symbiont infection results in MK (males die during embryogenesis).	Ladybird beetle *Adalia bipunctata*	*Rickettsia*	33,34
			Pierre's acraea butterfly *Acraea encedana*	*Wolbachia*	35
			Fruit fly *Drosophila melanogaster*	*Spiroplasma*	36
	Voltinism	Univoltine and multivoltine populations are associated with different symbiont strains.	Plum curculios *Conotrachelus nenuphar*	*Wolbachia*	37
Physiology	Resting metabolic rate	Facultative symbionts are energetically costly and increase the standard metabolic rate of the host.	Pea aphid *Acyrthosiphon pisum*	*Hamiltonella defensa, Regiella insecticola*	38
	Relative growth rate	Killing the symbiont reduces the host's growth rate.	Cotton stainers *Dysdercus fasciatus* Red firebugs *Pyrrhocoris apterus*	*Actinobacteria*	39
		Hosts without symbionts have lower growth rates.	Saw-toothed grain beetle *Oryzaephilus surinamensis*	*Sulcia*-like bacteria	40
	Desiccation resistance	Symbiont contributes to host's cuticle production and increases cuticle thickness, thus preventing transpiration and enhancing desiccation resistance.	Saw-toothed grain beetle *Oryzaephilus surinamensis*	*Sulcia*-like bacteria; *Sodalis pierantonius*	3,40

			Rice weevil *Sitophilus oryzae*		
		Artificially introduced symbiont confers desiccation resistance to the host.	Yellow fever mosquito *Aedes aegypti*	*Wolbachia*	[41]
	Salinity resistance	Symbionts of hosts living in arid, saline-alkali areas are more resistant to saline-alkali stress.	Beetles Assorted Coleoptera	Various bacterial species	[42]
	Temperature tolerance	Shifts in symbiont heat sensitivity contribute to variation in heat tolerance of the host.	Aphids *Aphis gossypii, Ap. fabae, Acyrthosiphon kondoi, Ac. pisum*	*Buchnera*	[43]
		Hosts infected with different symbiont strains have different temperature preferences. Individuals infected with symbionts prefer lower temperatures compared to aposymbiotic individuals.	Fruit fly *Drosophila melanogaster*	*Wolbachia*	[44]
	Pesticide tolerance*	Hosts infected with symbiont have increased susceptibility to pesticides.	Silverleaf white fly *Bemisia tabaci* Brown plant hopper *Nilaparvata lugens*	*Arsenophonus, Rickettsia, Wolbachia*	[45–47]
	Nitrogen recycling*	Symbionts located in bacterial pouches are used for nitrogen recycling.	Slender ants *Tetraponera* spp.	Various bacterial species	[48]
	Pathogen resistance*	Symbiont protects the host from fungal infections.	Beewolf *Philanthus triangulum*	*Streptomyces*	[49]
		Symbiont protects the host from RNA viral infections.	Fruit fly *Drosophila melanogaster*	*Wolbachia*	[50]

	Protection from natural enemies*	Symbiont protects its host from parasitic nematodes.	Fruit fly *Drosophila neotestacea*	*Spiroplasma*	[21]
		Symbiont protects its host from parasitoid wasps.	Fruit fly *Drosophila melanogaster*	*Spiroplasma*	[51]
Behaviour	Activity time	Depending on the symbiont infection, the time the host sleeps increases or decreases.	Fruit fly *Drosophila melanogaster*	*Wolbachia* (increases), *Spiroplasma* (decreases)	[52,53]
		Nymphs with a higher symbiont titer spend more time resting whereas those without symbionts spend more time wandering.	Japanese common stinkbug *Megacopta punctatissima*	*Candidatus* Ishikawaella capsulata	[54]
	Aggregation	Symbionts produce the components of host aggregation pheromones.	Source bark beetle *Ips typographus* Desert locust *Schistocerca gregaria*	*Grosmannia spp.* *Pantoea agglomerans*	[55,56]
		Presence of microbes, not host pheromones, causes hosts to aggregate.	Firebrats *Thermobia domestica*	*Mycotypha microspora, Enterobacter cloacae*	[57]
	Dispersal mode	Flight activity of the host is contingent on the amino acids and vitamins procured by symbiont.	Grain weevil *Sitophilus oryzae*, *S. zeamais*	Enterobacteriaceae gut endosymbiont	[58]
		Long-distance dispersal via ballooning is influenced by symbionts. Certain symbionts decrease long-distance ballooning.	Money spider *Erigone atra*	*Wolbachia, Rickettsia Spiroplasma, Cardinium*	[59]
	Locomotion speed	Gut symbiont increases intrinsic movement, dispersal, and velocity.	Black bean aphid *A. fabae*	*Serratia symbiotica*	[60]

	Sociality	By enabling hosts to subsist on nutrient-poor wood, symbionts partially enabled cooperative societies in wood-dwelling insects.	Cockroaches, termites, beetles Blattodea, Isoptera, Coleoptera: *Passalidae*, Coleoptera: *Curculionidae,* Coleoptera: *Tenebrionidae*	Various bacterial, fungal, protistan symbionts	[61]
	Defensive behavior*	Hosts infected with symbiont show less aggression and escape behavior when in the presence of parasitoid wasps.	Pea aphid *Acyrthosiphon pisum*	*Hamiltonella defensa*	[62]
	Kin recognition*	Conspecifics with a more similar gut bacteria avoided mating with each other.	Fruit fly *D. melanogaster*	Gut bacteria	[63]

References

1. Moretti, M. *et al.* Handbook of protocols for standardized measurement of terrestrial invertebrate functional traits. *Functional Ecology* **31**, 558–567 (2017).
2. Wybouw, N., Van Reempts, E., Zarka, J., Zélé, F. & Bonte, D. Egg provisioning explains the penetrance of symbiont-mediated sex allocation distortion in haplodiploids. *Heredity* **131**, 221–229 (2023).
3. Kanyile, S. N., Engl, T., Heddi, A. & Kaltenpoth, M. Endosymbiosis allows *Sitophilus oryzae* to persist in dry conditions. *Front. Microbiol.* **14**, 1199370 (2023).
4. Kikuchi, Y. *et al.* Collapse of Insect Gut Symbiosis under Simulated Climate Change. *mBio* **7**, e01578-16 (2016).
5. Jang, S. *et al.* Dual oxidase enables insect gut symbiosis by mediating respiratory network formation. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2020922118 (2021).
6. Kuriwada, T. *et al.* Biological Role of *Nardonella* Endosymbiont in Its Weevil Host. *PLoS ONE* **5**, e13101 (2010).
7. Anbutsu, H. *et al.* Small genome symbiont underlies cuticle hardness in beetles. *Proc. Natl. Acad. Sci. U.S.A.* **114**, (2017).
8. Bouchon, D., Rigaud, T. & Juchault, P. Evidence for widespread *Wolbachia* infection in isopod crustaceans: molecular identification and host feminization. (1998).
9. Cordaux, R., Michel-Salzat, A., Frelon-Raimond, M., Rigaud, T. & Bouchon, D. Evidence for a new feminizing *Wolbachia* strain in the isopod *Armadillidium vulgare*: evolutionary implications. *Heredity* **93**, 78–84 (2004).

10. Moran, N. A., Tran, P. & Gerardo, N. M. Symbiosis and Insect Diversification: an Ancient Symbiont of Sap-Feeding Insects from the Bacterial Phylum *Bacteroidetes*. *Appl Environ Microbiol* **71**, 8802–8810 (2005).
11. Hansen, A. K. & Moran, N. A. Aphid genome expression reveals host–symbiont cooperation in the production of amino acids. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 2849–2854 (2011).
12. Koga, R. & Moran, N. A. Swapping symbionts in spittlebugs: evolutionary replacement of a reduced genome symbiont. *The ISME Journal* **8**, 1237–1246 (2014).
13. Salem, H. *et al.* Vitamin supplementation by gut symbionts ensures metabolic homeostasis in an insect host. *Proc. R. Soc. B.* **281**, 20141838 (2014).
14. Bell-Roberts, L., Douglas, A. E. & Werner, G. D. A. Match and mismatch between dietary switches and microbial partners in plant sap-feeding insects. *Proc. R. Soc. B.* **286**, 20190065 (2019).
15. Nikoh, N. *et al.* Evolutionary origin of insect– *Wolbachia* nutritional mutualism. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 10257–10262 (2014).
16. Peterson, B. F., Stewart, H. L. & Scharf, M. E. Quantification of symbiotic contributions to lower termite lignocellulose digestion using antimicrobial treatments. *Insect Biochemistry and Molecular Biology* **59**, 80–88 (2015).
17. Zhao, P. *et al.* Fungal pathogen promotes caterpillar feeding and weight gain using a host-like trehalase. *Current Biology* **35**, 3601-3610.e3 (2025).
18. He, Z. *et al.* Effects of Wolbachia infection on the postmating response in Drosophila melanogaster. *Behav Ecol Sociobiol* **72**, 146 (2018).

19. Vigneron, A. *et al.* Insects Recycle Endosymbionts when the Benefit Is Over. *Current Biology* **24**, 2267–2273 (2014).
20. Himler, A. G. *et al.* Rapid Spread of a Bacterial Symbiont in an Invasive Whitefly Is Driven by Fitness Benefits and Female Bias. *Science* **332**, 254–256 (2011).
21. Jaenike, J., Unckless, R., Cockburn, S. N., Boelio, L. M. & Perlman, S. J. Adaptation via Symbiosis: Recent Spread of a *Drosophila* Defensive Symbiont. *Science* **329**, 212–215 (2010).
22. Katlav, A., Cook, J. M. & Riegler, M. Common endosymbionts affect host fitness and sex allocation via egg size provisioning. *Proc. R. Soc. B.* **289**, 20212582 (2022).
23. Chen, S.-J., Lu, F., Cheng, J.-A., Jiang, M.-X. & Way, M. O. Identification and Biological Role of the Endosymbionts *Wolbachia* in Rice Water Weevil (Coleoptera: Curculionidae). *Environ Entomol* **41**, 469–477 (2012).
24. Sumida, Y., Katsuki, M., Okada, K., Okayama, K. & Lewis, Z. Wolbachia induces costs to life-history and reproductive traits in the moth, Ephestia kuehniella. *Journal of Stored Products Research* **71**, 93–98 (2017).
25. Whitcomb, R. F., Tully, J. G., Bové, J. M. & Saglio, P. Spiroplasmas and Acholeplasmas: Multiplication in Insects. *Science* **182**, 1251–1253 (1973).
26. Herren, J. K. *et al.* Insect endosymbiont proliferation is limited by lipid availability. *eLife* **3**, e02964 (2014).
27. Chiel, E. *et al.* Assessments of Fitness Effects by the Facultative Symbiont *Rickettsia* in the Sweetpotato Whitefly (Hemiptera: Aleyrodidae). *Annals of the Entomological Society of America* **102**, 413–418 (2009).

28. Rodriguero, M. S., Confalonieri, V. A., Guedes, J. V. C. & Lanteri, A. A. *Wolbachia* infection in the tribe Naupactini (Coleoptera, Curculionidae): association between thelytokous parthenogenesis and infection status. *Insect Molecular Biology* **19**, 631–640 (2010).
29. Elias-Costa, A. J., Confalonieri, V. A., Lanteri, A. A. & Rodriguero, M. S. Game of clones: Is Wolbachia inducing speciation in a weevil with a mixed reproductive mode? *Molecular Phylogenetics and Evolution* **133**, 42–53 (2019).
30. Giorgini, M., Bernardo, U., Monti, M. M., Nappo, A. G. & Gebiola, M. *Rickettsia* Symbionts Cause Parthenogenetic Reproduction in the Parasitoid Wasp *Pnigalio soemius* (Hymenoptera: Eulophidae). *Appl Environ Microbiol* **76**, 2589–2599 (2010).
31. Gotoh, T., Noda, H. & Ito, S. Cardinium symbionts cause cytoplasmic incompatibility in spider mites. *Heredity* **98**, 13–20 (2007).
32. White, J. A., Kelly, S. E., Perlman, S. J. & Hunter, M. S. Cytoplasmic incompatibility in the parasitic wasp Encarsia inaron: disentangling the roles of Cardinium and Wolbachia symbionts. *Heredity* **102**, 483–489 (2009).
33. Werren, J. H. *et al.* Rickettsial relative associated with male killing in the ladybird beetle (Adalia bipunctata). *J Bacteriol* **176**, 388–394 (1994).
34. Hurst, G. D. D., Purvis, E. L. & Sloggett, J. J. The effect of infection with male-killing Rickettsia on the demography of female Ada/ia bipunctata L. (two spot ladybird).
35. Jiggins, Hurst, Dolman, & Majerus. High-prevalence male-killing *Wolbachia* in the butterfly *Acraea encedana*. *Journal of Evolutionary Biology* **13**, 495–501 (2000).

36. Harumoto, T. & Lemaitre, B. Male-killing toxin in a bacterial symbiont of Drosophila. *Nature* **557**, 252–255 (2018).

37. McClanan, M. E., Luckhart, S. & Pfeiffer, D. G. Wolbachia Strains Associated with Univoltine and Multivoltine Plum Curculios (Coleoptera: Curculionidae). *Journal of Entomological Science* **39**, 132–135 (2004).

38. Clavé, C. *et al.* Physiological costs of facultative endosymbionts in aphids assessed from energy metabolism. *Functional Ecology* **36**, 2580–2592 (2022).

39. Salem, H., Kreutzer, E., Sudakaran, S. & Kaltenpoth, M. Actinobacteria as essential symbionts in firebugs and cotton stainers (Hemiptera, Pyrrhocoridae). *Environmental Microbiology* **15**, 1956–1968 (2013).

40. Engl, T. *et al.* Ancient symbiosis confers desiccation resistance to stored grain pest beetles. *Molecular Ecology* **27**, 2095–2108 (2018).

41. Ross, P. A. *et al.* Developing Wolbachia-based disease interventions for an extreme environment. *PLoS Pathog* **19**, e1011117 (2023).

42. Yue, H. *et al.* Diversity and saline-alkali resistance of Coleoptera endosymbiont bacteria in arid and semi-arid climate. *Microbiol Spectr* **12**, e00232-24 (2024).

43. Zhang, B., Leonard, S. P., Li, Y. & Moran, N. A. Obligate bacterial endosymbionts limit thermal tolerance of insect host species. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 24712–24718 (2019).

44. Truitt, A. M., Kapun, M., Kaur, R. & Miller, W. J. *Wolbachia* modifies thermal preference in *Drosophila melanogaster*. *Environmental Microbiology* **21**, 3259–3268 (2019).

45. Kontsedalov, S. *et al*. The presence of *Rickettsia* is associated with increased susceptibility of *Bemisia tabaci* (Homoptera: Aleyrodidae) to insecticides. *Pest Management Science* **64**, 789–792 (2008).
46. Ghanim, M. & Kontsedalov, S. Susceptibility to insecticides in the Q biotype of *Bemisia tabaci* is correlated with bacterial symbiont densities. *Pest Management Science* **65**, 939–942 (2009).
47. Cai, T. *et al*. Insecticide susceptibility in a planthopper pest increases following inoculation with cultured *Arsenophonus*. *The ISME Journal* **18**, wrae194 (2024).
48. Borm, S. V., Buschinger, A., Boomsma, J. J. & Billen, J. *Tetraponera* ants have gut symbionts related to nitrogen–fixing root–nodule bacteria. *Proc. R. Soc. Lond. B* **269**, 2023–2027 (2002).
49. Kaltenpoth, M., Göttler, W., Herzner, G. & Strohm, E. Symbiotic Bacteria Protect Wasp Larvae from Fungal Infestation. *Current Biology* **15**, 475–479 (2005).
50. Teixeira, L., Ferreira, Á. & Ashburner, M. The Bacterial Symbiont Wolbachia Induces Resistance to RNA Viral Infections in Drosophila melanogaster. *PLoS Biol* **6**, e1000002 (2008).
51. Xie, J., Butler, S., Sanchez, G. & Mateos, M. Male killing Spiroplasma protects Drosophila melanogaster against two parasitoid wasps. *Heredity* **112**, 399–408 (2014).
52. Bi, J., Sehgal, A., Williams, J. A. & Wang, Y.-F. Wolbachia affects sleep behavior in Drosophila melanogaster. *Journal of Insect Physiology* **107**, 81–88 (2018).

53. Horn, C. J., Yuli, S., Berry, J. A. & Luong, L. T. A male-killing Spiroplasma endosymbiont has age-mediated impacts on Drosophila endurance and sleep. *Journal of Insect Physiology* **161**, 104723 (2025).
54. Hosokawa, T., Kikuchi, Y., Shimada, M. & Fukatsu, T. Symbiont acquisition alters behaviour of stinkbug nymphs. *Biol. Lett.* **4**, 45–48 (2008).
55. Zhao, T., Axelsson, K., Krokene, P. & Borg-Karlson, A.-K. Fungal Symbionts of the Spruce Bark Beetle Synthesize the Beetle Aggregation Pheromone 2-Methyl-3-buten-2-ol. *J Chem Ecol* **41**, 848–852 (2015).
56. Dillon, R. J., Vennard, C. T. & Charnley, A. K. A Note: Gut bacteria produce components of a locust cohesion pheromone. *J Appl Microbiol* **92**, 759–763 (2002).
57. Woodbury, N. & Gries, G. Firebrats, *Thermobia domestica* , aggregate in response to the microbes *Enterobacter cloacae* and *Mycotypha microspora*. *Entomologia Exp Applicata* **147**, 154–159 (2013).
58. Grenier, A. M., Nardon, C. & Nardon, P. The role of symbiotes in flight activity of *Sitophilus* weevils. *Entomologia Exp Applicata* **70**, 201–208 (1994).
59. Goodacre, S. L. *et al.* Microbial modification of host long-distance dispersal capacity. *BMC Biol* **7**, 32 (2009).
60. Hamidović, A. *et al.* Gut symbiont Serratia symbiotica influences feeding and locomotor behavior of aphid host Aphis fabae. *Arthropod-Plant Interactions* **19**, 87 (2025).
61. Dillard, J. & Benbow, M. E. From Symbionts to Societies: How Wood Resources Have Shaped Insect Sociality. *Front. Ecol. Evol.* **8**, 173 (2020).

62. Dion, E., Polin, S. E., Simon, J.-C. & Outreman, Y. Symbiont infection affects aphid defensive behaviours. *Biol. Lett.* **7**, 743–746 (2011).

63. Lewis, Z., Heys, C., Prescott, M. & Lizé, A. You are what you eat: Gut microbiota determines kin recognition in *Drosophila*. *Gut Microbes* **5**, 541–543 (2014).