

The Risk of Avian Influenza in the Southern Ocean

A practical guide for operators interacting with wildlife

Advice from Avian Influenza experts suggests that there is a **high risk that Highly Pathogenic Avian Influenza will arrive in the Southern Ocean during the 2022/23-2024/25 austral summers.**

Summary

Since the beginning of 2022, the increasing intensity of highly pathogenic avian influenza (HPAI) H5N1 outbreaks has resulted in the death of hundreds of thousands of seabirds in the Northern Hemisphere, around the Atlantic and Pacific Oceans and southern Africa. The SCAR Antarctic Wildlife Health Working Group (AWHWG) is highly concerned about the likely arrival and subsequent impact HPAI H5N1 might have on Southern Ocean wildlife. Due to the heightened risk of HPAI being introduced to Antarctica during the 2022/23 Austral summer by migrating seabirds, the AWHWG recommends that:

- People working with or close to wildlife should assume that HPAI will arrive in the sub-Antarctic and Antarctica and take precautions to protect themselves when working around wildlife (including appropriate Personal Protective Equipment (PPE)) and maintain the highest biosecurity to prevent transmission between wildlife aggregations.
- All National Programmes (NPs), tourism and fishery operators should monitor colonies for signs of H5N1 before approaching, especially in migratory species such as skuas, gulls and giant petrels. Visitors from tourism and fishery operations should not enter colonies and high wildlife density areas with suspected HPAI and NPs should conduct risk analysis as to which activities need to continue.
- A detailed protocol on how to assess wildlife aggregations for HPAI prior to a visit and what to do if HPAI is detected should be provided to all stakeholders physically present in Antarctica this season.
- If you detect signs of HPAI, do not enter colonies and report these to your permit issuer. Videos of affected animals (collected from a distance) are very helpful for experts to help determine whether or not this is HPAI.

- Operators should refresh themselves with and review all biosecurity and any response guidelines to unusual/mass mortality events.

This document aims to:

1. Outline the likely risk to Southern Ocean taxa (we are working on a separate document).
2. Suggest which risks can be mitigated in light of human activity, transmission into and out of Antarctica and the sub-Antarctic regions through all operators as well as movements between sites within the Southern Ocean (primarily for science and tourism).
3. Start discussions with National Programmes about ongoing monitoring for disease and consequences.

What is Avian Influenza?

Avian influenza virus (AIV) comprises the avian strains of Influenza A virus (species *Alphanfluenzavirus influenzae*, family *Orthomyxoviridae*). The AIVs maintained in wild birds are low pathogenic avian influenza (LPAI) viruses, which cause no clinical signs of disease in wild birds. Of concern are subtypes H5 and H7, which may become highly pathogenic in poultry and then escape into wild populations. (Horimoto and Kawaokade 2005; Monne et al 2014; Bruin et al 2022). These highly pathogenic avian influenza viruses (HPAI) can range from causing no clinical disease to 100% mortality. The current outbreaks of global concern are caused by HPAI H5N1 viruses.

What is the risk to humans?

There have been 863 human cases of HPAI H5N1 with 456 deaths in 2003–2021. This translates to a low risk of infection in humans, but high risk of mortality if infected (~50%). Human cases caused by the current circulating lineage of HPAI H5 (lineage 2.3.4.4) have been reported in China, Laos, Russia, Nigeria, the UK and USA, with no examples of human-to-human transmission detected (Wille and Barr 2022). To date, all human infections with HPAI have been in people closely interacting directly for prolonged periods of time with birds. If NPs decide to work at affected sites, we suggest that only appropriately trained field staff in appropriate Personal Protective Equipment (PPE) enter colonies with suspected HPAI and **do not handle sick, dying, or dead birds unless specifically trained and permitted for HPAI detection.**

How is it spread?

Transmission and pathogenicity of avian influenza varies by strain and which species it infects. In wild birds, LPAI is transmitted via the faecal-oral route and environmental contamination (mostly water). Direct contact is also believed to be a mechanism of transmission and may be important in the spread of HPAI, in addition to transmission through respiratory droplets.

Colonial nesting seabirds may be at greater risk of disease spread due to the proximity of individuals and nests over a long period of time, increasing direct contact rates among individuals, or with infected faeces and water (Vittecoq et al. 2017).

AIV in seabirds and the risk to Southern Ocean wildlife

The natural reservoir for avian influenza virus comprises aquatic birds, including Anseriformes (ducks, geese, swans) and Charadriiformes (shorebirds, gulls, terns, skuas, auks). Since 2022, a substantial increase in outbreaks of HPAI has led to considerable mortality in seabirds in the Northern Hemisphere, southern Africa and the Atlantic and Pacific Oceans. Outbreaks in seabird colonies of southern Africa began ~2017, and repeated detections in South African and Namibian seabirds include African Penguins, Common Terns, Swift Terns, and Cape Cormorants (Khomenko et al. 2018; Mollini et al. 2020), all taxonomically close to Southern Ocean seabirds. The current HPAI H5N1 has been linked to the death of Harbour Seals (*Phoca vitulina*) and Grey Seals (*Halichoerus grypus*) in the United States and Canada, in addition to a range of other mammals, indicating that the disease is not limited to seabirds (NOAA Fisheries, 2022; Ann 2022).

LPAI strains have been detected in seabirds on the Antarctic Peninsula and South Shetland Islands, comprising Adélie (*Pygoscelis adeliae*), Gentoo (*P. papua*), and Chinstrap (*P. antarcticus*) Penguins (Sanfilippo 2010; Hurt et al. 2014, 2016; Barriga et al. 2016), Southern Giant Petrel (*Macronectes giganteus*) (de Souza Petersen et al. 2017), Brown Skua (*Stercorarius antarcticus*) (Seixas 2014), and Snowy Sheathbill (*Chionis albus*) (Hurt et al. 2016). Antibodies against AIV have also been detected in a range of seabird species and localities in Antarctica and sub-Antarctic islands (Lang et al. 2016; Smeele et al. 2018).

As of August 2022, no HPAI outbreak has been recorded in the Southern Ocean, however, the recent outbreaks elsewhere suggest high risk of introduction to the region, given several species currently impacted in the Northern Hemisphere are Antarctic migrants, such as skuas and terns, and other migratory species (such as gull species) that mix with Antarctic migrants. Migratory species arriving from outside the Southern Ocean (mostly between September and November) that could transmit

the virus to the Antarctic and sub-Antarctic coincides with the arrival of penguins and other flying seabirds for breeding, thus increasing the risk amongst susceptible species.

Monitoring for early detection of HPAI

HPAI could most likely be introduced and transmitted to wildlife 'naturally' in Antarctica and sub-Antarctic islands via long-distance migratory species for which HPAI does not preclude long-distance flight. It could also be introduced by humans, via contaminated clothing and equipment.

Continued monitoring of HPAI along migratory pathways will enable all operators to assess the risk and likely arrival of HPAI into sub-Antarctic and Antarctica. Currently there is no central reporting agency, so we recommend communicating with your permitting authority. We recommend working with seabird biologists to identify whether HPAI is already present, the important migratory species that may be vectors, to monitor flyways and the movement of HPAI towards the Southern Ocean.

Detecting and responding to possible outbreaks

Mass mortality events occur normally in times of food stress. These are hard to distinguish from mortality due to disease or contaminants. Most diseased birds will die at sea, making it difficult to notice the impacts of an infectious agent such as HPAI. Ongoing disease monitoring programmes aim to resolve background and 'natural' diseases versus introduced and emerging diseases that pose a risk to populations. Despite this, where mass mortality occurs, it is necessary to react in a precautionary manner because of the risk of HPAI to visitors and wildlife alike and the high risk of spreading it to other colonies. If unusually large numbers or clusters of dead animals are observed, it is best to assume a mass mortality event due to disease and close the site until further investigation. As a rule, you should alert the person who authorises your visit or issues your permit, which may differ between Antarctica and the sub-Antarctic islands.

We strongly advise people against entering possibly infected colonies, touching sick or dying birds, or taking any samples unless they (1) have been trained to do so, (2) are wearing appropriate PPE and taking appropriate biosecurity precautions, (3) have a clear need to do so in light of an outbreak, and (4) have permits to collect diseased samples.

To identify potential presence of HPAI in wildlife, some of the most **common behavioural signs** observed in the current outbreak in the Northern Hemisphere include;

- **Neurological issues such as loss of coordination and balance,**

- **Trembling head and body,**
- **Sudden and rapid increase in the number of birds found dead between visits,**
- **Lethargy and depression, unresponsiveness, lying down, drooping wings, dragging legs,**
- Swollen head,
- Closed and excessively watery eyes, possibly with opaque cornea or darkened iris (new symptom associated with current outbreak),
- Twisting of the head and neck,
- Haemorrhages on shanks of the legs and under the skin of the neck,
- Respiratory distress such as gaping (mouth breathing), nasal snicking (coughing sound), sneezing, gurgling, or rattling, and
- Discoloured or loose watery droppings, bright green in some species.

Bold points are practical observations that could be made via binoculars prior to landing or approaching the colony. In the event of an outbreak, it would be expected that multiple of these behavioural signs are manifested by multiple individuals. If in doubt, it is best to refrain from approaching the colony and reporting for someone else to confirm.

Examples of behaviours observed in infected seabirds in recent outbreak:

- <https://twitter.com/i/status/1542423370511302656>
- <https://twitter.com/gannetrocks/status/1549641794165538817>
- <https://photos.app.goo.gl/u9f3jpuy4SKQr1Lw9>

One of the most practical changes that visitors can make prior to approaching a colony is a site assessment looking for behavioural signs (listed above). Experienced researchers, guides and naturalists will be especially good at spotting atypical behaviour and spasms, which could be integrated into a site evaluation prior to landing. If one is unsure if the behaviour is typical of HPAI, video footage could be taken using a mobile phone and sent to the Antarctic Wildlife Health Working Group for verification by an expert team (contact details below). In case of an abnormal number of dead individuals, panoramic pictures should be taken. **Do not approach colonies if there is a suspected disease outbreak.**

Prevention and Control

Given that Antarctica and sub-Antarctic islands are at high risk for the 2022/23 Austral summer, personnel from all operators living, working, and visiting these regions should **act as if HPAI will arrive in Antarctica and sub-Antarctic Islands** and should take appropriate precautions. Responses may be stepped up if cases are confirmed. Suggested prevention and control measures that operators can immediately implement to prevent the spread of HPAI are outlined below:

Before leaving for Antarctica and the sub-Antarctic:

1. All personnel including, but not limited to, expeditioners, guides and crew working in other wildlife areas (e.g., seabird colonies, the Arctic, Alaska, etc.) should thoroughly remove any soiled material (i.e., soil, faeces) from their boots and disinfect all clothing, footwear, and any equipment before leaving for Antarctica and the sub-Antarctic to prevent introduction of HPAI from other wildlife areas. It is recommended to use new clothes, footwear, and equipment when possible.
2. All personnel, including tourists, should remove all soiled material and disinfect footwear before boarding a vessel or aircraft, especially if they have been near wildlife colonies.
3. The use of Virkon™ S, F10 or soap followed by 10% bleach is recommended for disinfection of boots and outer clothing.

Before visiting a colony or wildlife aggregation:

1. In general, regional specific advice/regulation may exist, so we suggest adopting whichever is the more precautionary.
2. To identify if a colony is potentially infected with HPAI, an appropriately trained or experienced guide/research team should be sent via small boat, avoid landing if possible (or via high vantage point at remote southern colonies), and observe the colony using binoculars to look for carcasses and behaviour of live birds (listed above) before entering the site or sending tourists/parties to shore. In case of terrestrial approaching, observations using binoculars should be made 150 m from the closest animal aggregation.
3. If there are signs of an unusual/mass mortality event or behavioural signs of HPAI within a colony, the visit should be aborted. **All equipment and clothing should be completely disinfected as soon as possible** and the observations reported to your permit issuer/national authority immediately.

There may be specific reporting forms or procedures, particularly for sub-Antarctic Islands.

4. In general, all equipment (including boots, backpacks, hiking poles, tripods, camping equipment and cones or route markers etc.) should be cleaned of any soiled material (i.e., soil, faeces) and disinfected before disembarking the boat/leaving base and every time after visiting a colony.

During a visit:

1. Unless specifically permitted to do otherwise, visitors to wildlife colonies should always keep **a minimum distance of 5 metres from wildlife**, in addition to strict adherence to ATCM general guidelines, IAATO guidelines or any other local requirements. (https://documents.ats.aq/recatt/att483_e.pdf) given a greater distance may be required for different animal species and age cohorts. If an animal spontaneously approaches an individual or a group of visitors, the visitors should retreat to ensure this minimum distance is rigorously adhered to unless specifically studying or sampling animals and being permitted to do so.
2. People should not sit on or lie on the ground close to animal activity or faecal matter (within 10 metres of nests, haul-out sites or pathways). Freshwater pools should also be avoided.
3. To minimise animal exposure to equipment/potential fomite transmission (i.e. transmission from objects or equipment carrying infection), field equipment should not be left unattended and should be kept away from wildlife. If necessary, consider grouping separate items onto a disinfected ground sheet or similar.
4. **Visitors must adhere to strict recommendations for personal hygiene at all times** (frequent hand washing for the appropriate time and regular disinfection).
5. **Visitors must never touch birds, dead or alive unless they have a permit that, in this case, specifically includes birds with suspected disease.**

Post-visit:

1. All equipment (including boots, tripods and anything that may have been in contact with the ground) should be also cleaned of any soiled material (i.e., soil, faeces) and disinfected upon returning to the ship or a base after a shore visit. Wear a mask to prevent inhalation during cleaning.

2. Field equipment should be thoroughly disinfected prior to and after shore visits.

General Recommendations:

1. All logistics providers and expeditions should be on high alert for HPAI from September through to April, **assume HPAI will arrive in the Southern Ocean** and take appropriate precautions. In addition:
 - a. Wear appropriate PPE including face masks and gloves when handling seabirds in the absence of disease or even working close to wildlife. If disease is suspected, PPE should be increased to include, disposable overalls/aprons and safety glasses/shield,
 - b. Be appropriately trained in working with potentially infectious animals, and correctly wear/remove/dispose of PPE. The World Health Organization has online training for this.
<https://openwho.org/courses/IPC-PPE-EN>
2. Anyone responsible for groups visiting/transiting through wildlife colonies should be briefed on identifying signs of mass mortality events and HPAI behavioural signs, and abort visits if signs of mass mortality are seen.
3. Minimise movements between wildlife aggregations without biosecurity (e.g., fieldworkers walking between aggregations within a single day).
4. To avoid transmission through fomites, disinfect outer clothing using soap and warm water every day prior to and after work with animals and in between colonies (if researchers visit more than one colony per day).
5. Field equipment should be thoroughly disinfected prior to and after animal handling, especially when working in different areas or colonies. Wherever possible, it is recommended that field/sampling equipment not be shared between locations if appropriate disinfection cannot be achieved. The use of 70% ethanol, Virkon™ S, F10 or 10% bleach (sodium hypochlorite) and soap are recommended.
6. Poultry, meat and eggs must be kept in a strictly controlled space and their remains in the vessels and stations and waste products be incinerated.

Response to Suspected Avian Influenza Outbreak

1. Abort a visit if signs of HPAI are seen.
 - a. Do not enter colonies with suspected HPAI

- b. **Do not touch dead, sick or dying birds.** Advice from vets is not to remove carcasses to minimise the likelihood of infection.
 - a. **Immediately report the event following your organisation's reporting procedures,** collect evidence such as videos from a distance.
2. Sampling requires appropriate authorisation/permitting and, if done without training, PPE and containment, could do more harm than good as well as being a breach of permits or the law.
3. The suspected event should be **reported immediately to the following:**
 - a. The National Competent Authority which permits/authorises the operation;
 - b. National Programs and Authorities are required to report the event to the World Organisation for Animal Health (WOAH) as HPAI is a globally reportable disease

All operators to the Antarctic and sub-Antarctic are encouraged to share information of suspected outbreak(s) and their locations with all other programmes and operators in the region, and with organisations such as CCAMLR and IAATO. Relevant contacts are listed below.

Helpful information to collect/provide:

- OrganismID
- Date;
- Time in UTC
- Location including name of site and coordinates (use a GPS if available),
- Name of observer, organisation and vessel (if applicable)
- Area affected,
- Species (*Scientific name*) present, age class and sex ratio, and which one(s) seem(s) affected by the outbreak (if available),
- Indication of the number of animals involved - if possible, count the dead and dying and estimate the percentage of each among the colony,
- Any behavioural signs noted, and

- Photographs and/or videos of unusual behaviours, but also of groups of apparently healthy and unhealthy/dead individuals.

If a landing was conducted and later the presence of HPAI suspected, all clothing and footwear should be cleaned and disinfected using 70% ethanol, Virkon S, F10 or soap and then 10% commercial bleach (sodium hypochlorite).

4. If a National Competent Authority permits/authorises an investigation team, they must be adequately trained and aware of all risks. The team must follow guidelines on health and safety and ensure adequate personal protection. The team will not visit any unaffected colonies after visiting the affected colony. Specific guidelines for fieldworkers and investigation teams are attached.

Contacts

- Antarctic National Competent Authorities:

<https://www.ats.aq/devAS/Ats/NationalCompetentAuthorities?lang=e>

- The International Association of Antarctica Tour Operators

operations@iaato.org; +1 401 608 2090; www.iaato.org

- The World Organisation for Animal Health (WOAH)

As HPAI is a globally notifiable disease, the World Organisation for Animal Health (WOAH) **needs to be notified if there is any suspected outbreak.**

Submission of a suspected outbreak can additionally be reported to the PROMED website using the following link: <https://promedmail.org/submitinfo>

Authors:

Meagan Dewar^{1, 13*}, Michelle Wille^{2, 13}, Amandine Gamble^{3, 4, 13}, Ralph E. T. Vanstreels^{5, 13}, Thierry Bouliner^{6, 13}, Adrian Smith^{7, 13}, Arvind Varsani^{8, 13}, Norman Ratcliffe⁹, Jennifer Black¹⁰, Amanda Lynnes¹¹, Andrés Barbosa^{12, 14} and Tom Hart^{13, 14*}

1. Future Regions Research Centre, Federation University Australia, Berwick Australia
2. Sydney Institute for Infectious Diseases, School of Life and Environmental Sciences and School of Medical Sciences, The University of Sydney, New South Wales, Australia
3. School of Biodiversity, One Health and Veterinary Medicine, University of Glasgow, United Kingdom
4. Department of Public and Ecosystem Health, Cornell University, New York, United States

5. Latin America Program, Karen C. Drayer Wildlife Health Center, University of California - Davis, São Paulo, Brazil
6. Centre d'Ecologie Fonctionnelle et Evolutive, CNRS, Université de Montpellier,
7. Department of Biology, Oxford University
8. The Biodesign Center for Fundamental and Applied Microbiomics, School of Life Sciences, Center for Evolution and Medicine, Arizona State University, Tempe, Arizona, AZ 85287, US
9. British Antarctic Survey, Cambridge, United Kingdom
10. C/o Foreign, Commonwealth and Development Office, United Kingdom
11. International Association of Antarctica Tour Operators
12. Dept. Evolutionary Ecology, National Museum of Natural Sciences, Spanish Research Council
13. Penguin Watch, Department of Biology, Oxford University
14. Antarctic Wildlife Health Working Group. Expert Group for Birds and Marine Mammals. Scientific Committee for Antarctic Research

*Corresponding Authors: Meagan Dewar m.dewar@federation.edu.au; Tom Hart tom.hart@biology.ox.ac.uk

References

Ann, V. 2022. Death of seals in Quebec linked to highly pathogenic avian flu. The Canadian Press.

<https://www.theglobeandmail.com/canada/article-death-of-seals-in-quebec-linked-to-highly-pathogenic-avian-flu/>

Barriga, G.P., Boric-Bargetto, D., Cortez-San Martin, M., Neira, V., van Bakel, H., Thompson, M., Tapia, R., Toro-Ascuy, D., Moreno, L., Vasquez, Y., Sallaberry, M., Torres-Perez, F., Gonzalez- Acuna, D., Medina, R.A. 2016. Avian influenza virus H5 strain with North American and Eurasian lineage genes in an Antarctic penguin. *Emerging infectious diseases*. 22(12) 2221 - 2223.

de Bruin ACM, Funk M, Spronken MI, Gulyaev AP, Fouchier RAM, Richard M. 2022. Hemagglutinin Subtype Specificity and Mechanisms of Highly Pathogenic Avian Influenza Virus Genesis. *Viruses*. 14(7):1566. <https://doi.org/10.3390/v14071566>

de Souza Petersen, E., de Araujo, J., Krüger, L., Seixas, M. M., Ometto, T., Thomazelli, L. M., Walker, D., Durigon, E. L., & Petry, M. V. (2017). First detection of avian influenza virus (H4N7) in Giant Petrel monitored by geolocators in the Antarctic region. *Marine Biology*, 164(4). <https://doi.org/10.1007/s00227-017-3086-0>

Horimoto T, Kawaoka Y. 2005. Influenza: lessons from past pandemics, warnings from current incidents. *Nat Rev Microbiol*. (8):591-600.

Hurt, A., Vjaykrishna, D., Butler, J., Bass, C., Mauer-Stroh, S., Silva-de-la-Fuente, C.M., Medina-Vogel, G., Olsen, B., Kelso, A, Barr, I.G., Gonzalez-Acuna, D. 2014. Detection of evolutionary distinct avian influenza A virus in Antarctica. *mBio* 5(3)e01098-14.

Hurt, A.C, Su, Y.C.F, Aban, M, Peck, H, Lau, H, Baas, C, Deng, Y-M, Spirason, N, Ellström, P, Hernandez, J, Olsen, B, Barr, I.G, Vijaykrishna, D, Gonzalez-Acuna, D. 2016. Evidence for the Introduction, Reassortment, and Persistence of Diverse Influenza A Viruses in Antarctica. *Journal of Virology* 90 (21) pp. 9674 –9682.

International Association of Antarctica Tour Operators (IAATO). IAATO Procedures upon discovery of a high Mortality events. ATCM XXXIX CEP XIX, Santiago (2016), www.ats.aq.

Khomenko, Abolnik, Roberts, Waller, Shaw, Monne, Taylor, Dhingra, Pittiglio, Mugyeom, Roche, Fredrick, Kamata, Okuthe, Kone, Wiersma, Von Dobschuetz, Soumare, Makonnen, Morzaria, Lubroth 2018. 2016–2018 Spread of H5N8 highly pathogenic avian influenza (HPAI) in sub-Saharan Africa (No. 12). Food and Agriculture Organization of the United Nations.
<https://www.fao.org/3/ca1209en/CA1209EN.pdf>

Lang, A. S., Lebarbenchon, C., Ramey, A. M., Robertson, G. J., Waldenström, J., & Wille, M. (2016). Assessing the role of seabirds in the ecology of influenza A viruses. *Avian diseases*, 60(1s), 378-386.

Monne, I. Fusaro, A. Nelson, M.I. Bonfanti, L. Mulatti, P. Hughes, J. Murcia, P.R. Schivo, A. Valastro, V. Moreno, A. Holmes, E.C. Cattoli, G. 2014. Emergence of a Highly Pathogenic Avian Influenza Virus from a Low-Pathogenic Progenitor. *Journal of Virology*. 88 (8): 4375-4388

Molini, U, Aikukutu, G, Roux, J.P, Kemper, J, Ntahonshikira, C, Marruchella, G, Khaiseb, S, Cattoli, G, Dundon, W.G. 2020. Avian Influenza H5N8 Outbreak in African Penguins (*Spheniscus demersus*), Namibia, 2019. *Journal of Wildlife Disease* 56(1) pp. 214-218.

NOAA Fisheries 2022. NOAA Fisheries Confirms Highly Pathogenic Avian Influenza Linked to Recent Increase in Seal Deaths in Maine. 6th July 2022.

Sanfilippo, L.F. 2010. Epidemiologia e caracterização molecular do vírus da influenza em quatro espécies de pinguins na região Antártica. Doctoral thesis, University of São Paulo, São Paulo, Brazil.

<https://www.teses.usp.br/teses/disponiveis/42/42132/tde-11082011-105843/pt-br.php>

Seixas M. M. M. 2014. Estudo do vírus Influenza em aves marinhas da região subantártica. Masters dissertation, University of São Paulo, São Paulo, Brazil.
<https://www.teses.usp.br/teses/disponiveis/42/42132/tde-12082016-115854/pt-br.php>

Smeele, Z. E., Ainley, D. G., & Varsani, A. (2018). Viruses associated with Antarctic wildlife: From serology based detection to identification of genomes using high throughput sequencing. *Virus Research*, 243, 91-105.

USDA 2022 Detections of Highly Pathogenic Avian Influenza.
<https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/avian/avian-influenza/2022-hpai>

Verhagen, J.H.; Fouchier, R.A.M.; Lewis, N. Highly Pathogenic Avian Influenza Viruses at the Wild–Domestic Bird Interface in Europe: Future Directions for Research and Surveillance. *Viruses* 2021, 13, 212.

Vittecoq, M., Gauduin, H., Oudart, T., Bertrand, O., Roche, B., Guillemain, M., & Boutron, O. (2017). Modelling the spread of avian influenza viruses in aquatic reservoirs: A novel hydrodynamic approach applied to the Rhône delta (southern France). *Science of The Total Environment*, 595, 787–800.
<https://doi.org/10.1016/j.scitotenv.2017.03.165>

Wille, M., Barr, I.G. 2022. Resurgence of avian influenza virus. *Science* 376 (6592): 459-460