

Stability in numbers: a positive link between honeybee colony size and thermoregulatory efficiency around the brood

Ugoline Godeau¹, Maryline Pioz¹, Olivier Martin², Charlotte Rüger³, Didier Crauser¹, Yves Le Conte¹, Mickael Henry¹, Cedric Alaux¹

5

¹ INRAE, Abeilles et Environnement, 84914 Avignon, France

² INRAE, Biostatistique et processus SPatiaux (BioSP), 84914 Avignon, France

³ ANSES, Epidémiologie et appui à la surveillance (EAS), 69364 Lyon Cedex 07, France

10 Corresponding author:

Ugoline Godeau, godeau.ugoline@gmail.com

Abstract

To ensure the optimal development of brood, a honeybee colony needs to maintain its
15 temperature within a certain range of values (thermoregulation), regardless of environmental
changes in biotic and abiotic factors. While the set of behavioural and physiological responses
implemented by honeybees to regulate the brood temperature has been well studied, less is
known about the factors that may influence the efficiency of this thermoregulation. Based on the
response threshold model of task allocation, increased colony homeostasis should be driven by
20 increases in group size. We therefore determined whether colony size (number of adult bees and
amount of brood), positively influenced the efficiency of thermoregulation that we measured via
two criteria: (i) the precision of the temperature close to its brood optimum, and (ii) the stability
of the temperature around this optimum value. Finally, within the applied perspective of
honeybee colony monitoring, we assessed whether the efficiency of thermoregulation could be
25 used as a proxy of colony size.

For that purpose, we followed 29 honeybee colonies over two years, measured both brood and
adult population size regularly over the beekeeping season, and monitored the in-hive
temperature over the 24 hours preceding the inspections of these colonies. We then studied the
effect of the size of the colony (number of adult bees and number of brood cells), as well as
30 meteorological variables, on the efficiency of thermoregulation (mean and stability of brood
temperature, *i.e.* between 32 and 36°C).

In addition to a clear link with meteorological conditions, we found that the mean brood
temperature and the stability of this temperature were both positively linked to the size of
colonies. The mean brood temperature was more dependent on the amount of the brood, while its

35 stability was more dependent on the number of bees. However, these relationships between colony size and thermoregulation were too weak for clearly discriminating colony population size based solely on the brood thermoregulatory efficiency. This demonstrates an extremely high flexibility and efficiency of honeybee colonies to thermoregulate the brood regardless of the amount of brood and the group size.

40 **Introduction**

Homeostasis denotes the ability of living organisms to actively maintain steady internal conditions necessary for survival. A classic example of organismal homeostasis is the regulation of body temperature within certain boundaries, even when environmental temperatures change. Such a phenomenon is also found in social insects, like the honeybee *Apis mellifera*, for which 45 the maintenance of nest conditions within a certain range of values (homeostasis), regardless of environmental changes in biotic and abiotic factors, is crucial for their colony development and survival.

The maintenance of inner hive conditions is one of the most crucial functions of honeybee colonies. While adult bees are rather eurytherms (*i.e.* can live under a wide range of 50 temperatures), with a minimum of 18°C for normal muscle function (Esch & Bastian, 1968) and a maximum for survival above 50°C (Coelho, 1991; Kovac et al., 2014), the brood is stenothermic (*i.e.* only able to survive and develop within a narrow temperature range) (Seeley, 1985). Accurate temperature regulation is therefore essential for proper development, with brood temperature strictly controlled within a temperature range of 32 to 36°C (Seeley, 1985) with 55 regulation even more precise during the pupal period (35±0.5°C, Jones et al. 2004; Kronenberg & Heller 1982; Stabentheiner et al. 2010, 2021). Maintaining this optimal temperature window is

crucial for the colony. Indeed, extended deviations are known to increase mortality (Koeniger, 1978; Wang et al., 2016), cause morphological defects (Fukuda & Sakagami, 1968; Himmer, 1932; Winston, 1987), disrupt synaptic organization in the brain of adult bees (Groh et al., 2004) and affect behavioural performances (Becher et al., 2009; Jones et al., 2005; Tautz et al., 2003).

The colony, through the cooperation and coordination between individuals, therefore implements a set of behavioural and physiological responses to ensure proper temperature regulation of the hive (Jones & Oldroyd, 2006). When temperature is perceived as being too high, workers regulate it by fanning hot air out of the nest with their wings and may simultaneously spread water to induce evaporative cooling (Prange, 1996). At a finer scale, young workers can passively absorb heat by placing themselves between the heat source and the brood cells. This behaviour is called heat shielding, and it is usually carried out by placing the ventral side against the hot surface (Bonoan et al., 2014; Siegel et al., 2005; Starks et al., 2005; Starks & Gilley, 1999). When temperature is perceived as being too low, workers can contract their thoracic muscles to produce heat (Esch et al., 1991; Heinrich, 1980, 1985, 1993; Heinrich & Esch, 1994). Another efficient heating strategy consists of entering an empty cell to warm the adjacent cells containing brood (Bujok et al., 2002; Kleinhenz et al., 2003). Finally, during longer periods of cold, workers can cluster together and generate metabolic heat (Kronenberg & Heller, 1982; Meikle et al., 2016; Seeley & Heinrich, 1981; Stabentheiner et al., 2010).

The thermoregulatory mechanisms within the hive are therefore numerous, of different natures (behavioural, physiological or passive), flexible and interlaced, resulting in an effective brood temperature homeostasis (Kronenberg & Heller, 1982; Stabentheiner et al., 2021), even in extreme ambient conditions (*e.g.* Himmer, 1932 and Lindauer, 1955). However, temperature can fluctuate around its optimal value (Stabentheiner et al., 2021). Within the goal of maintaining

80 temperature homeostasis, the efficiency of thermoregulation can be gauged through two criteria:
(i) the precision of the temperature close to its brood optimum, and (ii) the stability of the
temperature around this optimum value. Many studies have investigated how bees perform
thermoregulation (see above), but little is known about the factors that can influence the
efficiency of this thermoregulation. Based on the response threshold model of task allocation
85 (Beshers & Fewell, 2001), the probability that an individual bee will engage in thermoregulation
will depend on the level of the task stimulus and her threshold for that stimulus, *i.e.* the
likelihood of reacting to the task-associated stimuli. A greater between-individual variability and
within-individual consistency (specialisation) in task performance is therefore expected to
increase behavioural homeostasis within the colony (Ulrich et al., 2018). This was confirmed by
90 an increased stability in temperature changes within colonies composed of genetically diverse
worker bees as compared to colonies with a low level of genetic diversity (Jones et al., 2004). A
more recent study showed in ants that increased colony behavioral homeostasis is also driven by
increases in group size (number of adult ants), likely via a stabilization of task performance
frequency and a decrease in task neglect (Ulrich et al., 2018). We could therefore expect a
95 similar influence of colony size on thermoregulatory efficiency in honeybees. A link between
temperature regulation and the number of honeybees has been suggested (Seeley & Heinrich
1981; e.g. Southwick, 1985), as well as the role of the stimulus intensity (here brood amount),
but this remains to be investigated and characterized.

We therefore investigated in this paper whether the thermoregulatory efficiency around the brood
100 was related to colony size (number of adult bees and brood amount). For that purpose, we
monitored outside meteorological conditions, inner hive temperature and bee population level of
several colonies over two years. We then investigated the influence of colony size on the

thermoregulatory level (mean brood temperature) and stability (fluctuations around the mean brood temperature). Finally, by using the relationship between the thermoregulatory efficiency and colony size, we investigated whether the ability to regulate temperature around the brood
105 could be used to estimate the colony size (for instance, whether high variability in thermoregulatory capacities could be an indicator of a relatively weak colony, and vice versa), without needing more data such as climate data. Indeed, within the context of severe colony losses observed around the world over the past years (Ellis et al., 2010; Neumann & Carreck,
110 2010; Potts et al., 2010), there is a clear need for surveillance networks and beekeeping operations to identify simple and non-intrusive proxies of colony state for monitoring and assessing their development and potential decline (López-Uribe et al., 2020). Such proxies could be extremely useful given that connected hives now allow us to monitor real-time data on physical variables such as weight, temperature, humidity and respiratory gases (Marchal et al.,
115 2020; Meikle & Holst, 2015; Zacepins et al., 2011, 2012).

Methods

I. Experimental setup and colony monitoring

Experiments were performed at INRAE (Avignon, France, 43°540N-4°-520E) with honeybee colonies (*Apis mellifera*). A total of 28 ten-frame colonies were randomly selected from our local
120 apiary in 2018 plus a new colony in 2019, for a total of 29 different colonies over the two years. Each colony was equipped with a temperature sensor (SHT35-DIS-B2.5KS, Sensirion AG) measuring in-hive temperature every 5 min. with a precision of 0.1°C within a temperature range of 20 to 60°C. The sensor precision was verified and validated beforehand using a climatic chamber. Colony strength was found to be more related to temperature data from sensors nearest

125 to the geometric centre of the hive (Cook et al., 2022). Therefore, the sensor was placed between
the two central frames 5 and 6 and at mid-height, in order to be as close as possible to the brood,
which generally occupy the central place in the hive.

The sensor was wired to a STM32 microcontroller (STMicroelectronics) and data were stored on
a memory card (SanDisk Ultra SDHC 16 Go).

130 Colonies were inspected six times in 2018 (*i.e.* every two to three weeks between July and
October) and five times in 2019 (*i.e.* approximately every three weeks between April and July)
to estimate three parameters: the number of open and closed brood cells and the number of adult
bees. During colony visits, each side of each frame was visually inspected and the area covered
by each of these parameters was reported as a percentage (one full side = 100%). Considering
135 that a full side of a Dadant Hoffmann frame has a surface of 9.03 dm² and contains in theory
1,100 bees and 3,100 brood cells, percentages were ultimately converted into number of open
brood cells, number of closed brood cells, and number of adult bees inside the hive (Alaux et al.,
2018; Hernandez et al., 2020). Initial population size was different for each colony, and ranged
from 4,536 to 40,131 adult bees, 15,200 to 44,250 open and closed brood cells in July 2018, and
140 from 5,292 to 40,950 adult bees and 0 to 37,600 open and closed brood cells in April 2019
(Appendix S2 Figure S1, Appendix S3 Table S3).

II. Data analysis

1. Link between thermoregulatory efficiency, meteorological conditions and colony size

Thermoregulatory variables

145 Using temperature sensor data, we calculated the mean of in-hive temperatures over the 24 hours preceding the day of colony evaluation (hereafter MeanT – Appendix S2 Figure S2). We chose a 24-hour time period since brood population can rapidly evolve over days (e.g. adult emergence). We therefore minimized the risk of having brood population changes between the temperature and population monitoring. We then discarded observations for which this mean (MeanT) was
150 outside the range 32-36°C, since it is unlikely that brood develop near the sensor at such temperatures (Seeley, 1985), resulting in a total number of 236 observations for the 29 colonies. The mean brood temperature in the dataset was 34.3°C (min = 32.15, Q1 = 33.9, Median = 34.4, Q3 = 34.81, max = 35.85), and rarely exceeded 35°C (n = 29 events for 236 observations). We then calculated the coefficient of variation (CV; *i.e.* standard deviation expressed as a percentage
155 of MeanT) within this same 24-hour period, to obtain a dimensionless variable representing the variation of temperature as a percentage of the mean temperature. The final response variables are therefore (i) the in-hive mean temperature (MeanT), representing the thermoregulatory precision and (ii) the coefficient of variation of the in-hive temperature (CV), representing the thermoregulatory stability.

160 *Predictor variables*

Both colony size and environmental meteorological conditions can potentially influence the regulation of the in-hive temperature (Stabentheiner et al., 2021). Regarding colony size, we studied two predictor variables: (i) the number of adult bees (Nbees), and (ii) the total number of brood cells (Nbrood - as the sum of the number of capped and uncapped brood cells). Regarding
165 meteorological conditions, we retrieved data from a local INRAE weather station and investigated the three following variables as relevant indicators of environmental conditions: (i) the daily mean external temperature over the 24 hours preceding the day of colony evaluation

(temperature mean TM, in degree Celsius), (ii) the daily global radiation (GR, in joule/cm², Burrill & Dietz 1981) and (iii) the daily precipitation (rainfall rate RR, in mm). In addition, to
170 take into account a possible effect of phenological advancement of the colony, we have adapted
the cumulative growing degree-day, usually used to estimate the growth and development of
plants, to the foraging activity of bees, which largely contributes to colony development. The
cumulative growing degree-day (GDDcum) was calculated as the sum of mean daily
temperatures (TM) above 12.5°C, from the beginning of each year, *i.e.* 2018 and 2019
175 (Appendix S2 Figure S3). A temperature of 12.5°C corresponds to the minimum temperature at
which honeybee foraging activity starts (Vicens & Bosch, 2000). For a given year the GDDcum
was calculated as follows (where t=1 corresponds to the 1st of January):

$$GDDcum_t = \sum_{i=1}^t (\max(TM_i - 12.5, 0))$$

Finally, colony replicate was included as a random effect to take into account potential variation
180 in thermoregulatory capacity inherent to the colony (such as colony genetics, Jones et al. 2004).

The various meteorological predictors (TM, GR, RR and GDDcum) were tested in addition to
the strength of the colony (via Nbrood and Nbees variables) to explain the thermoregulatory
efficiency criteria: the mean temperature around the brood (MeanT) and the variability of
temperatures around the brood (CV). For the later, we also integrated MeanT as a predictor to
185 assess its potential influence on thermoregulatory stability.

Statistical analysis and model selection

We conducted a model-averaging analysis in order to study in detail the predictors explaining the
variations of MeanT and CV and their contributions. For this purpose, we first used generalized

linear mixed models to model the relationship between the two response temperature variables
190 (MeanT and CV) and the predictors. For MeanT, we specified a Gaussian distribution and the
Identity link function. For CV, since this variable was continuous and severely skewed, we
specified an inverse-Gaussian distribution with a Log link function. Because variables are
measured in different units, we centred and scaled (by dividing by the standard deviation) the
numerical variables when used as predictors (hereafter with an “S” at the end of their names).
195 The two models are written as follows:

$$g(\mu_{ij}) = \beta_0 + \sum_{h=1}^p \beta_h x_{hij} + \alpha_i$$

Where

μ_{ij} is the expectation of the variable Y_{ij}

y_{ij} is the j th observation of the i th colony (either MeanT or CV),

200 β_0 is the intercept,

β_h is the regression coefficient for the h th predictor,

x_{hij} is the j th value of the i th colony for the h th of p fixed-effect predictors,

α_i is the colony-specific effect and $\alpha_i \sim \text{Gaussian}(0, \sigma^2)$,

g is the link function (identity for the normal distribution and log for the inverse gaussian
205 distribution).

We fitted the generalized linear models using the "glm" function (from the "stats" package - R
Core Team 2021) for fixed effects models (without colony random effect), or the "glmer"

function (from the "lme4" packages – Bates et al. 2015) for mixed models (with colony random effect).

210 In order to avoid multicollinearity, which is highly problematic in the case of model averaging (Banner & Higgs, 2017; Cade, 2015), we have excluded the possibility for the models to integrate simultaneously variables that introduce multicollinearity (for multicollinearity detection method see Appendix S1). In the end, the following pairs were not integrated into the same model: NBeeS and NBroodS, GRS and RRS, GRS and TMS, and GDDcumS and MeanTS.

215 A model selection procedure was applied by using the corrected Akaike Information Criterion (AICc). This procedure was done with the dredge function in the “MuMIn” package (Bartoń, 2020). Finally, we conducted model averaging based on AICc, a multimodel inference approach that allows one to derive inference from a subset of closely related best models, and not just from a single best model. Regarding the choice of the subset, we included models with a ΔAICc of
220 less than seven points from the best model, grouping models that are likely to be the best models and that should all be used when making further inferences ($\Delta\text{AICc} < 2$) and models that are unlikely to be best models but that should not be discounted ($\Delta\text{AICc} \in [4,7]$, Burnham & Anderson, 2002). Having previously forced some predictors not to be included in certain models, in order to avoid multicollinearities and a strong bias of underestimation of coefficients, we
225 preferred the results of the conditional average. This model only averages over the models where the parameter appears, ignoring the cases where the model does not include the predictors when calculating the coefficients (unlike the full average for which the coefficients are set to 0.0 if the predictors are not included in the model). We did not consider any interaction between predictors because we had no *a priori* biological reason for doing so, and integrating these interactions, in
230 particular with the random effects, was too ambitious in relation to the quantity of data available.

Based on the average models of MeanT and CV, for each predictor, we extracted its regression coefficient (to study effect size) and its P-value (to evaluate if its relationship with the studied variable is statistically significant). We also assessed the relative importance of each predictor by summing the Akaike weight across all the models in the set in which the predictor appeared. The
235 closer the sum is to 1.0, the more important the predictor is in the set of fitted models (Burnham & Anderson, 2004).

2. Thermoregulatory efficiency across colony size categories

Within the applied perspective of honeybee colony monitoring, we tested whether data on thermoregulatory capacities on their own (*i.e.* without the climatic information), could provide
240 information on colony size. For that purpose, we transformed the two quantitative variables representing the strength of the colony (Nbees and Nbrood), into ordinal variables of four categories (respectively; catBees and catBrood) based on the quartiles (balanced in terms of number; Appendix S3 Table S3). We then assessed whether colony size categories were associated with specific in-hive temperature variables (MeanT and CV) by comparing the later
245 across the different categories representing the strength of the colony. Since the MeanT and CV data were not normally distributed (neither globally nor by categories), we applied the nonparametric test of Kruskal-Wallis to test whether the medians of the thermoregulatory variables (MeanT or CV) differ across colony-size categories. We also looked at the effect size of this Kruskal-Wallis test (as being the *eta* squared based on the H-statistic, with $\epsilon^2 < 0.01$: very
250 small effect, $0.01 < \epsilon^2 < 0.08$: small effect, $0.08 < \epsilon^2 < 0.26$: medium effect and $\epsilon^2 \geq 0.26$: large effect, Cohen, 1988). In the case of significant results ($P < 0.05$), we applied a Dunn post-hoc test to investigate multiple pairwise comparisons. These tests were carried out using the package “rstatix” (Kassambara, 2021).

255 Finally, we conducted ordinal logistic regression (with “clm” function from the “ordinal”
package - Christensen 2019) in order to try to predict colony-size category (brood or number of
bees) based on the in-hive temperature (with MeanT scaled). We compared predictions of the
models with the real observations and extracted accuracy (the proportion of all correctly
classified validation points) and Cohen’s Kappa statistics ($\kappa = (\text{Observed Accuracy} -$
260 $\text{Expected Accuracy}) / (1 - \text{Expected Accuracy})$), which evaluate classification performance,
taking into account the possibility of the agreement occurring by chance.


Statistical analyses were conducted on the filtered data (temperature of brood thermoregulation:
MeanT $\in [32,36^\circ\text{C}]$), and therefore the sample size of each population category was no longer
balanced (see Appendix S3 Table S8 & Appendix S3 Table S9 for the effective categories sizes).

265 **Results**

I. Link between the mean temperature around the brood, colony size and meteorological conditions


270 After setting aside models prone to multicollinearity, a total of 13 models remained (Appendix
S3 Table S4) from which the average model was estimated (Appendix S3 Table S5). The
combined outputs of the average model suggest that MeanT deviance was better explained by the
scaled cumulative growing degree-day (GDDcumS), the scaled global radiation (GRS), the
scaled mean external temperature (TMS), the scaled total number of brood cells (NbroodS), the
scaled total number of adult bees (NBeesS), the scaled precipitation levels (RRS), and a colony
random effect (sum of weights=0.20). NbroodS, GRS and TMS had a significant positive effect

275 on MeanT, contrary to GDDcumS that had a significant negative effect on MeanT (Figure 1). NbeesS and RRS had no significant effect on MeanT (Figure 1). When looking at the sum of Akaike weights, GRS, NBroodS and particularly GDDcumS, had high relative importance. The model was not very efficient in predicting the observed data, with weak MeanT strongly overestimated and strong MeanT heavily underestimated (Figure 2).



280

Figure 1: Left: Predictor estimates (β), and P-values (P) of fixed-effect parameters for the in-hive mean temperature (MeanT) average model. Bars represent the 95% confidence intervals of predictor estimates. Red points and bars are for predictors with significant effect at a level of 5%. Right: Sum of weights across all models in the set where the variable occurred.



285

Figure 2: Mean in-hive temperature (MeanT) predicted by the average model as a function of the observed MeanT, with first bisector in blue ($\text{Predicted MeanT} = \text{Observed MeanT}$), deviations from this line in grey and regression line of the point cloud as a dotted red line.


290 **II. Link between the variability of the temperature around the brood, colony size and meteorological conditions**

290

After setting aside models prone to multicollinearity, a total of 3 models remained (Appendix S3 Table S6) from which the average model was estimated (Appendix S3 Table S7). The combined outputs of the average model suggest that CV deviance was better explained by the scaled mean in-hive temperature (MeanTS), the scaled precipitation levels (RRS), the scaled external mean temperature (TMS), the scaled number of adult bees (NBeesS), the scaled total number of brood

295


cells (NbroodS) and a colony random effect. All models composing the average model included the colony random effect, which had a significant effect on CV. TMS had a significant positive effect on CV, contrary to MeanTS, RRS, and NBeesS which had a significant negative effect (Figure 3). NBroodS had no significant effect on CV (Figure 3). When looking at the sum of 300 Akaike weights, TMS, MeanTS, RRS and, to a lesser extent, NbeeS, had high relative importance. The model was again not very efficient in predicting the observed data, with weak CVs slightly overestimated and the few strong CVs heavily underestimated, meaning that the model better explains weak CVs than strong CVs (Figure 4).



305

Figure 3: Left: Predictor estimates (β), and P-values (P) of fixed-effect parameters for the in-hive temperature CV average model. Bars represent the 95% confidence intervals of predictor

estimates. Red points and bars are for predictors with significant effect at a level of 5%. Right: Sum of weights across all models in the set where the variable occurred.



310

Figure 4: Coefficient of variation of in-hive temperature (CV) predicted by the average model as a function of the observed CV, with first bisector in blue (Predicted CV=Observed CV), deviations from this line in grey and regression line of the point cloud as a dotted red line.

III. Link between thermoregulatory efficiency and categorized colony sizes

315 We then assessed whether thermoregulatory efficiency (precision MeanT and stability CV) differed between colony size categories (based on quartiles of brood amount: catBrood and of bee number: catBees). We found significant variations across colony size categories with moderate magnitude for both MeanT (catBrood: P-value < 0.001, $\epsilon^2 = 0.112$ and catBees: P-

value < 0.001, $\varepsilon^2 = 0.0796$, Kruskal-Wallis tests) and CV (catBrood: P-value < 0.001, $\varepsilon^2 = 0.1010$ and catBees: P-value < 0.001, $\varepsilon^2 = 0.0834$). Thermoregulatory efficiency of middle-size colonies (cat2 and cat3 of catBrood and catBees) did not consistently differ from thermoregulatory efficiency of the smallest and largest colonies and from each other (Figure 5). However, the MeanT and the CV were always significantly different between colonies of the smallest and largest size category (for both catBees and catBrood).

325 The effect of Nbrood or Nbees on the precision or stability of thermoregulation was not distinct enough to clearly discriminate different potential colony sizes. However, a low temperature (MeanT $\approx 33.8^\circ\text{C}$) combined with poor stability of this temperature (CV ≈ 1.67) was somewhat associated with a weak colony at least in terms of number of bees (catBees cat1 $\leq 13,419$ bees, cf. Appendix S3 Table S8). In the same way, a temperature near the optimal brood temperature 330 (MeanT $\approx 34.7^\circ\text{C}$) combined with a good stability of this temperature (CV ≈ 0.962), seemed to indicate a strong colony, at least in terms of the amount of brood (catBrood cat4 $> 27,225$ brood cells, cf. Appendix S3 Table S9). The analysis of the density plot (Appendix S2 Figure S1**Error! Reference source not found.**) showed that, around a MeanT of 35°C , colonies were unlikely to belong to the first category of NBrood. Conversely, below a MeanT of 34°C , colonies were 335 unlikely to belong to the fourth category of NBrood. Regarding the CV, we did not observe any clear discrimination of categories (NBrood and NBees, Figure 5 and Appendix S2 Figure S4).




Figure 5: Thermoregulatory efficiency (MeanT and CV) across categories of number of adult bees (catBees) and brood cells (catBrood). ns: not significant, stars indicate significant differences between colony categories (post-hoc Dunn test). *** P-value < 0.005, ** P-value < 0.01, * P-value < 0.05, ns P-value \geq 0.05. For category ranges see Appendix S3 Table S1.

340

345

Finally, the ordinal logistic regression showed that MeanT and CV were not significantly correlated to categories of number of bees, catBees (MeanT: P-value = 0.151; CV: P-value = 0.075), and the predictions of the model based on the two predictors were predominantly incorrect in comparison with the observations (accuracy = 0.3136, κ = 0.0313; Table 1). On the other hand, MeanT had a significant effect on Nbrood categories (catBrood) with ordinal logistic

models (MeanT: P-value < 0.001; CV: P-value = 0.601), but the model predictions were still not correct (accuracy = 0.3432, κ = 0.0884; Table 1).

350 *Table 1 : Confusion table for both ordinal logistic models predicting catBees or catBrood based on MeanT and CV. Grey highlights indicate true positives. For category ranges see Appendix S3 Table S2.*

Variable studied	Data	Model predictions			
	observations	cat1	cat2	cat3	cat4
catBees	cat1	10	10	26	1
	cat2	5	5	50	1
	cat3	5	8	58	2
	cat4	3	3	48	1
catBrood	cat1	8	15	21	1
	cat2	10	15	35	4
	cat3	5	13	47	7
	cat4	1	5	38	11

Discussion

355 I. Link between the mean temperature around the brood and colony size

In order for the brood to develop normally, honeybee colonies need to regulate the brood temperature between 32°C and 36°C, and optimally at 35°C (Seeley, 1985). Brood temperature is therefore regulated within a narrow range of temperatures but according to the response

threshold model of division of labour, we still expected that an increase in group size would
360 generate a higher level of social homeostasis and therefore increased capacity of reaching
optimal nest conditions, due to higher variability and task specialization between individuals
(Ulrich et al., 2018). We did not find that brood temperature significantly increases with the
number of adult bees within colonies, the link seems present but not strong enough in our data to
be significant. However, the increase in thermoregulation was significantly and positively related
365 to the amount of brood. Social homeostasis, and thus the ability to thermoregulate, is not only
due to the likelihood of individuals to react to a stimulus but also to the intensity of the stimulus
and whether it exceeds the individual response threshold (Theraulaz et al., 1998). Under this last
scenario, it is possible that the stimulus intensity of the thermoregulatory tasks (brood amount)
was high enough to surpass the threshold response of many individual bees, regardless of their
370 respective thresholds. However, a non-mutually exclusive hypothesis is that the greater the
quantity of brood, the greater the chance there is that the sensor is well surrounded by brood, and
therefore to record optimal temperatures. By selecting temperatures within the brood
thermoregulatory range, we expected to always have brood in the vicinity of the sensor, which
was confirmed during colony inspection, but a large amount of brood would exclude situations in
375 which the temperature sensor would be located at the periphery of the brood patch where the
brood temperature would be lower.

II. Link between the variability of the temperature around the brood and colony size

The variability of brood temperature was mainly linked to the mean temperature around which
380 this variability was calculated: as the value of the brood temperature increased, approaching

35°C, the temperature variability decreased. This confirms that a finer-tuned temperature regulation occurs around the temperature of 35°C, which is the optimal temperature for pupal development (Jones et al., 2004; Kronenberg & Heller, 1982).

Variation in brood temperature was also correlated to the colony size (number of adult bees and, 385 to a lesser extent, amount of brood), with larger colonies exhibiting reduced changes in brood temperature. This could be expected given that we first found that brood temperature increased with colony size (amount of brood, see above), and brood temperature increase was associated with lower temperature variability. However, when analysing the contribution of different environmental and colony variables, the ability to keep brood temperature stable seemed to be 390 more linked to the number of adult bees (significant) than to the amount of brood (selected by models but not significant). This suggests that an increase in group size allows honeybee colony responses to better buffer against environmental fluctuation and therefore regulate hive temperature. This phenomenon could be attributed to differences among individuals, which generally increases with colony size in social insects (Ulrich et al., 2018). Inter-individual 395 behavioural variation was notably found to favour the collective control of nest climate in bumblebees (*Bombus terrestris*, Weidenmüller 2004). However, the contribution of group size to the stability of brood temperature was of low magnitude and needs to be confirmed in future studies with perhaps a greater range of colony sizes. More generally, we also found important inter-colony variation in the ability of keeping brood temperature stable. These differences 400 among colonies could be linked to various underlying reasons, for example, the exact location of the hive (more or less shaded) or the genetics of the bees (Graham et al., 2006; Jones et al., 2004).

III. Link between thermoregulatory efficiency and environmental conditions

This efficiency of thermoregulation also depends on environmental conditions. The effect of the
405 environmental temperature on the hive temperature has been highlighted previously. Stabentheiner et al. (2010) notably showed that the environmental temperatures have a non-negligible impact on the temperature regulation capacity inside the hive, in particular at the level of the brood. We also highlighted an influence of these environmental conditions on the regulation of brood temperature.

410 The external temperature had a non-negligible influence on both the mean brood temperature and the colony's ability to maintain a stable brood temperature. Indeed, the hotter the environmental temperature, the hotter mean brood temperature was, and the more difficult it was for the colony to stabilise the brood temperature. This might be explained by the mechanisms used by colonies to compensate for high environmental temperatures, which consist of the collection and
415 evaporation of water above the brood. Notably, in response to a simulated heat wave at 37°C (2°C above the optimal temperature), a 70% increase in forager traffic to sustain water needs was previously observed (Bordier et al., 2017). However, the efficacy of water collection not only depends on the foraging capacity but also on water availability in the environment (distance from the hive, water amount), which might lead to some degree of fluctuation in the regulation of
420 brood temperature as compared to the more "passive" response to cold (changes in bee density and endothermy), especially in our experimental site characterized by high summer temperatures (between 35 and 40°C).

The cumulative growing degree-day, which gives an indication of phenological advancement of the colony, was negatively associated with the mean brood temperature: the further we advanced

425 in the beekeeping season, the lower the brood temperature was. Such an association could be
easily explained by the link between brood temperature and brood size, which declined between
the spring and fall. More interestingly, even though precipitation was not significantly associated
with an increase in the brood temperature, it seemed to substantially influence the maintenance
of stable brood temperature, as the occurrence of precipitation was associated with a decrease in
430 temperature variability. This observation confirms the influence of group size (number of bees)
on the stability of brood temperature since higher numbers of bees are expected in the hive due
to reduced or no foraging activity on rainy days. Finally, solar radiation seemed to influence the
mean brood temperature, with an increase in radiation associated with an increase in mean
temperature, likely due to the direct heating of the hive by the sun's rays on sunnier days.

435 **IV. From thermoregulatory data to colony size evaluation**

Measurements of in-hive temperatures have already been suggested and used for monitoring
honeybee colony populations. It was notably found that the adult and brood mass of colonies
were positively correlated with the in-hive temperatures (Cook et al., 2022; Meikle et al., 2016,
2017). Similarly, brood mass (but not adult mass) was inversely related to the amplitude of in-
440 hive temperatures (Meikle et al., 2017). These results were obtained by including all in-hive
temperatures (no pre-selection of brood temperatures ranging between 32 and 36°C). As a
consequence, strict control of temperatures (low temperature variation) was indicative of
colonies with brood and large temperature amplitudes were indicative of colonies with little or
no brood (Meikle et al., 2017). The fact that we obtained similar results but on brood
445 thermoregulation is promising within the goal of estimating colony population size. Indeed, in-
hive temperatures could be used as a first filter to discriminate colonies with brood from colonies

with little or no brood. Then, analysis of brood thermoregulatory efficiency could be used in a second step to evaluate in more detail the state of colonies with a relatively high amount of brood. The greater the brood temperature homeostasis, the larger the colony would be (as indicated by our results). By splitting colonies into size categories, we effectively found that large colonies had significantly better brood thermoregulation than small colonies. However, when looking at the boxplot and density plot of colony size categories according to their thermoregulatory levels (Figure 5, Appendix S2 Figure S4), it was only possible to state that a temperature below 33.8°C was not indicative of large colonies in our dataset ($\geq 27,225$ adult bees or $\geq 27,225$ brood cells). While in-hive temperatures can easily discriminate colonies with brood from colonies with almost no brood, a higher level of colony-size discrimination was not possible when focusing on brood temperatures.

V. Conclusion

Brood thermoregulatory efficiency was associated with colony size, and the propensity to reach 35°C was more related to the amount of brood near the sensor and the temperature stability more related to the number of adult bees. This highlights the importance of increased group size notably for maintaining stable temperature conditions within the hive. However, when considering the size effect (magnitude), the influence of colony size on thermoregulation was relatively marginal, which indicates a very high efficiency of the honeybee colony to thermoregulate whatever the amount of brood and the group size. As a consequence, the discrimination of colony population level based on brood thermoregulatory data was rather difficult. Nevertheless, within the applied perspective of honeybee colony monitoring, it would be useful to analyse a larger range of colony size variation, including depopulated or collapsing

colonies, to fully conclude on the potential of brood temperature as a proxy for colony size
470 estimation.

References

- Alaux, C., Soubeyrand, S., Prado, A., Peruzzi, M., Maisonnasse, A., Vallon, J., Hernandez, J.,
Jourdan, P., & Conte, Y. L. (2018). Measuring biological age to assess colony
demographics in honeybees. *PLOS ONE*, *13*(12), e0209192.
475 <https://doi.org/10.1371/journal.pone.0209192>
- Banner, K. M., & Higgs, M. D. (2017). Considerations for assessing model averaging of
regression coefficients. *Ecological Applications*, *27*(1), 78-93.
<https://doi.org/10.1002/eap.1419>
- Bartoń, K. (2020). *MuMIn : Multi-Model Inference*. R package version 1.43.17. [https://CRAN.R-](https://CRAN.R-project.org/package=MuMIn)
480 [project.org/package=MuMIn](https://CRAN.R-project.org/package=MuMIn)
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models
Using lme4. *Journal of Statistical Software*, *67*, 1-48.
<https://doi.org/10.18637/jss.v067.i01>
- Becher, M. A., Scharpenberg, H., & Moritz, R. F. A. (2009). Pupal developmental temperature
485 and behavioral specialization of honeybee workers (*Apis mellifera* L.). *Journal of*
Comparative Physiology A, *195*(7), 673-679. <https://doi.org/10.1007/s00359-009-0442-7>
- Beshers, S. N., & Fewell, J. H. (2001). Models of Division of Labor in Social Insects. *Annual*
Review of Entomology, *46*(1), 413-440. <https://doi.org/10.1146/annurev.ento.46.1.413>
- Bonoan, R. E., Goldman, R. R., Wong, P. Y., & Starks, P. T. (2014). Vasculature of the hive :
490 Heat dissipation in the honey bee (*Apis mellifera*) hive. *Naturwissenschaften*, *101*(6),
459-465. <https://doi.org/10.1007/s00114-014-1174-2>

- Bordier, C., Dechatre, H., Suchail, S., Peruzzi, M., Soubeyrand, S., Pioz, M., Pélissier, M., Crauser, D., Conte, Y. L., & Alaux, C. (2017). Colony adaptive response to simulated heat waves and consequences at the individual level in honeybees (*Apis mellifera*). *Scientific Reports*, 7(1), 3760. <https://doi.org/10.1038/s41598-017-03944-x>
- 495
- Bujok, B., Kleinhenz, M., Fuchs, S., & Tautz, J. (2002). Hot spots in the bee hive. *Naturwissenschaften*, 89(7), 299-301. <https://doi.org/10.1007/s00114-002-0338-7>
- Burnham, K. P., & Anderson, D. R. (2002). *Model selection and multimodel inference: A practical information-theoretic approach* (Second Ed.). Springer.
- 500
- Burnham, K. P., & Anderson, D. R. (2004). Multimodel Inference : Understanding AIC and BIC in Model Selection. *Sociological Methods & Research*, 33(2), 261-304. <https://doi.org/10.1177/0049124104268644>
- Burrill, R. M., & Dietz, A. (1981). The response of honey bees to variations in solar radiation and temperature. *Apidologie*, 12(4), 319-328. <https://doi.org/10.1051/apido:19810402>
- 505
- Cade, B. S. (2015). Model averaging and muddled multimodel inferences. *Ecology*, 96(9), 2370-2382. <https://doi.org/10.1890/14-1639.1>
- Christensen, R. H. B. (2019). *ordinal—Regression Models for Ordinal Data*. R package version 2019.12-10. <https://CRAN.R-project.org/package=ordinal>
- Coelho, J. R. (1991). Heat Transfer and Body Temperature in Honey Bee (Hymenoptera : Apidae) Drones and Workers. *Environmental Entomology*, 20(6), 1627-1635. <https://doi.org/10.1093/ee/20.6.1627>
- 510
- Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences* (Second ed.). Routledge. <https://doi.org/10.4324/9780203771587>

- 515 Cook, D., Tarlinton, B., McGree, J. M., Blackler, A., & Hauxwell, C. (2022). Temperature Sensing and Honey Bee Colony Strength. *Journal of Economic Entomology*, toac034. <https://doi.org/10.1093/jee/toac034>
- Ellis, J. D., Evans, J. D., & Pettis, J. (2010). Colony losses, managed colony population decline, and Colony Collapse Disorder in the United States. *Journal of Apicultural Research*, 49(1), 134-136. <https://doi.org/10.3896/IBRA.1.49.1.30>
- 520 Esch, H., & Bastian, J. (1968). Mechanical and electrical activity in the indirect flight muscles of the honey bee. *Zeitschrift Für Vergleichende Physiologie*, 58(4), 429-440. <https://doi.org/10.1007/BF00343515>
- Esch, H., Goller, F., & Heinrich, B. (1991). How do bees shiver? *The Science of Nature*, 78(7), 325-328. <https://doi.org/10.1007/BF01221422>
- 525 Fukuda, H., & Sakagami, S. F. (1968). Worker brood survival in honeybees. *Population Ecology*, 10(1), 31-39. <https://doi.org/10.1007/BF02514731>
- Graham, S., Myerscough, M. R., Jones, J. C., & Oldroyd, B. P. (2006). Modelling the role of intracolony genetic diversity on regulation of brood temperature in honey bee (*Apis mellifera* L.) colonies. *Insectes Sociaux*, 53(2), 226-232. <https://doi.org/10.1007/s00040-005-0862-5>
- 530 Groh, C., Tautz, J., & Rössler, W. (2004). Synaptic organization in the adult honey bee brain is influenced by brood-temperature control during pupal development. *Proceedings of the National Academy of Sciences of the United States of America*, 101(12), 4268-4273. <https://doi.org/10.1073/pnas.0400773101>

- 535 Heinrich, B. (1980). Mechanisms of Body-Temperature Regulation in Honeybees, *Apis Mellifera* : II. Regulation of Thoracic Temperature at High Air Temperatures. *Journal of Experimental Biology*, 85(1), 73-87. <https://doi.org/10.1242/jeb.85.1.73>
- Heinrich, B. (1985). The social physiology of temperature regulation in honeybees. In B. Holldobler & M. Lindauer (Éds.), *Experimental behavioral ecology and sociobiology* (p. 393-406). Sinauer Associates.
- 540 Heinrich, B. (1993). *The Hot-Blooded Insects : Strategies and Mechanisms of Thermoregulation*. Springer.
- Heinrich, B., & Esch, H. (1994). Thermoregulation in Bees. *American Scientist*, 82(2), 164-170.
- Hernandez, J., Maisonnasse, A., Cousin, M., Beri, C., Le Quintrec, C., Bouetard, A., Castex, D., 545 Decante, D., Servel, E., Buchwalder, G., Brunet, F., Feschet-Destrella, E., de Bellescize, K., Kairo, G., Frontero, L., Pédehontaa-Hiaa, M., Buisson, R., Pouderoux, T., Aebi, A., & Kretzschmar, A. (2020). ColEval : Honeybee COLony Structure EVALuation for Field Surveys. *Insects*, 11(1), 41. <https://doi.org/10.3390/insects11010041>
- Himmer, A. (1932). Die Temperaturverhältnisse bei den sozialen Hymenopteren. *Biological 550 Reviews*, 7(3), 224-253. <https://doi.org/10.1111/j.1469-185X.1962.tb01042.x>
- Jones, J. C., Helliwell, P., Beekman, M., Maleszka, R., & Oldroyd, B. P. (2005). The effects of rearing temperature on developmental stability and learning and memory in the honey bee, *Apis mellifera*. *Journal of Comparative Physiology. A, Neuroethology, Sensory, Neural, and Behavioral Physiology*, 191(12), 1121-1129. [https://doi.org/10.1007/s00359-](https://doi.org/10.1007/s00359-005-0035-z)
- 555 005-0035-z

- Jones, J. C., Myerscough, M. R., Graham, S., & Oldroyd, B. P. (2004). Honey Bee Nest Thermoregulation : Diversity Promotes Stability. *Science*, 305(5682), 402-404. <https://doi.org/10.1126/science.1096340>
- Jones, J. C., & Oldroyd, B. P. (2006). Nest Thermoregulation in Social Insects. In *Advances in Insect Physiology* (Vol. 33, p. 153-191). Elsevier. [https://doi.org/10.1016/S0065-2806\(06\)33003-2](https://doi.org/10.1016/S0065-2806(06)33003-2)
- Kassambara, A. (2021). *rstatix : Pipe-Friendly Framework for Basic Statistical Tests*. R package version 0.7.0. <https://CRAN.R-project.org/package=rstatix>
- Kleinhenz, M., Bujok, B., Fuchs, S., & Tautz, J. (2003). Hot bees in empty broodnest cells : Heating from within. *Journal of Experimental Biology*, 206(23), 4217-4231. <https://doi.org/10.1242/jeb.00680>
- Koeniger, N. (1978). Das Wärmes der Brut bei der Honigbiene (*Apis mellifera* L.). *Apidologie*, 9(4), 305-320. <https://doi.org/10.1051/apido:19780404>
- Kovac, H., Käfer, H., Stabentheiner, A., & Costa, C. (2014). Metabolism and upper thermal limits of *Apis mellifera carnica* and *A. m. Ligustica*. *Apidologie*, 45(6), 664-677. <https://doi.org/10.1007/s13592-014-0284-3>
- Kronenberg, F., & Heller, H. C. (1982). Colonial thermoregulation in honey bees (*Apis mellifera*). *Journal of Comparative Physiology*, 148(1), 65-76. <https://doi.org/10.1007/BF00688889>
- Lindauer, M. (1955). The Water Economy and Temperature Regulation of the Honeybee Colony. *Bee World*, 36(5), 81-92. <https://doi.org/10.1080/0005772X.1955.11094876>
- López-Urbe, M. M., Ricigliano, V. A., & Simone-Finstrom, M. (2020). Defining Pollinator Health : A Holistic Approach Based on Ecological, Genetic, and Physiological Factors.

- 580 *Annual Review of Animal Biosciences*, 8(1), 269-294. <https://doi.org/10.1146/annurev-animal-020518-115045>
- Marchal, P., Buatois, A., Kraus, S., Klein, S., Gomez-Moracho, T., & Lihoreau, M. (2020). Automated monitoring of bee behaviour using connected hives: Towards a computational apidology. *Apidologie*, 51(3), 356-368. <https://doi.org/10.1007/s13592-019-00714-8>
- 585 Meikle, W. G., & Holst, N. (2015). Application of continuous monitoring of honeybee colonies. *Apidologie*, 46(1), 10-22. <https://doi.org/10.1007/s13592-014-0298-x>
- Meikle, W. G., Weiss, M., Maes, P. W., Fitz, W., Snyder, L. A., Sheehan, T., Mott, B. M., & Anderson, K. E. (2017). Internal hive temperature as a means of monitoring honey bee colony health in a migratory beekeeping operation before and during winter. *Apidologie*, 590 48(5), 666-680. <https://doi.org/10.1007/s13592-017-0512-8>
- Meikle, W. G., Weiss, M., & Stilwell, A. R. (2016). Monitoring colony phenology using within-day variability in continuous weight and temperature of honey bee hives. *Apidologie*, 47(1), 1-14. <https://doi.org/10.1007/s13592-015-0370-1>
- Mela, C. F., & Kopalle, P. K. (2002). The impact of collinearity on regression analysis: The 595 asymmetric effect of negative and positive correlations. *Applied Economics*, 34(6), 667-677. <https://doi.org/10.1080/00036840110058482>
- Neumann, P., & Carreck, N. L. (2010). Honey bee colony losses. *Journal of Apicultural Research*, 49(1), 1-6. <https://doi.org/10.3896/IBRA.1.49.1.01>
- Potts, S. G., Roberts, S. P. M., Dean, R., Marris, G., Brown, M. A., Jones, R., Neumann, P., & 600 Settele, J. (2010). Declines of managed honey bees and beekeepers in Europe. *Journal of Apicultural Research*, 49(1), 15-22. <https://doi.org/10.3896/IBRA.1.49.1.02>

- Prange, H. D. (1996). Evaporative cooling in insects. *Journal of Insect Physiology*, 42(5), 493-499. [https://doi.org/10.1016/0022-1910\(95\)00126-3](https://doi.org/10.1016/0022-1910(95)00126-3)
- R Core Team. (2021). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
- 605
- Seeley, T. D. (1985). *Honeybee ecology: A study of adaptation in social life*. Princeton University Press.
- Seeley, T. D., & Heinrich, B. (1981). Regulation of temperature in the nests of social insects. In B. Heinrich (Éd.), *Insect thermoregulation* (p. 160-234). Wiley.
- 610 Siegel, A. J., Hui, J., Johnson, R. N., & Starks, P. T. (2005). Honey bee workers as mobile insulating units. *Insectes Sociaux*, 52(3), 242-246. <https://doi.org/10.1007/s00040-005-0805-1>
- Southwick, E. E. (1985). Allometric relations, metabolism and heart conductance in clusters of honey bees at cool temperatures. *Journal of Comparative Physiology B*, 156(1), 143-149.
- 615 <https://doi.org/10.1007/BF00692937>
- Stabentheiner, A., Kovac, H., & Brodschneider, R. (2010). Honeybee Colony Thermoregulation – Regulatory Mechanisms and Contribution of Individuals in Dependence on Age, Location and Thermal Stress. *PLOS ONE*, 5(1), e8967. <https://doi.org/10.1371/journal.pone.0008967>
- 620 Stabentheiner, A., Kovac, H., Mandl, M., & Käfer, H. (2021). Coping with the cold and fighting the heat: Thermal homeostasis of a superorganism, the honeybee colony. *Journal of Comparative Physiology A*, 207(3), 337-351. [https://doi.org/10.1007/s00359-021-01464-](https://doi.org/10.1007/s00359-021-01464-8)

- 625 Starks, P. T., & Gilley, D. C. (1999). Heat Shielding : A Novel Method of Colonial
Thermoregulation in Honey Bees. *Naturwissenschaften*, 86(9), 438-440.
<https://doi.org/10.1007/s001140050648>
- Starks, P. T., Johnson, R. N., Siegel, A. J., & Decelle, M. M. (2005). Heat shielding : A task for
youngsters. *Behavioral Ecology*, 16(1), 128-132. <https://doi.org/10.1093/beheco/arh124>
- 630 Tautz, J., Maier, S., Groh, C., Rössler, W., & Brockmann, A. (2003). Behavioral performance in
adult honey bees is influenced by the temperature experienced during their pupal
development. *Proceedings of the National Academy of Sciences*, 100(12), 7343-7347.
<https://doi.org/10.1073/pnas.1232346100>
- Theraulaz, G., Bonabeau, E., & Deneubourg, J.-N. (1998). Response threshold reinforcements
and division of labour in insect societies. *Proceedings of the Royal Society of London.
635 Series B: Biological Sciences*, 265(1393), 327-332.
<https://doi.org/10.1098/rspb.1998.0299>
- Ulrich, Y., Saragosti, J., Tokita, C. K., Tarnita, C. E., & Kronauer, D. J. C. (2018). Fitness
benefits and emergent division of labour at the onset of group living. *Nature*, 560(7720),
635-638. <https://doi.org/10.1038/s41586-018-0422-6>
- 640 Vicens, N., & Bosch, J. (2000). Weather-Dependent Pollinator Activity in an Apple Orchard,
with Special Reference to *Osmia cornuta* and *Apis mellifera* (Hymenoptera :
Megachilidae and Apidae). *Environmental Entomology*, 29(3), 413-420.
<https://doi.org/10.1603/0046-225X-29.3.413>
- 645 Wang, Q., Xu, X., Zhu, X., Chen, L., Zhou, S., Huang, Z. Y., & Zhou, B. (2016). Low-
Temperature Stress during Capped Brood Stage Increases Pupal Mortality,

Misorientation and Adult Mortality in Honey Bees. *PLOS ONE*, 11(5), e0154547.
<https://doi.org/10.1371/journal.pone.0154547>

650 Weidenmüller, A. (2004). The control of nest climate in bumblebee (*Bombus terrestris*)
colonies: Interindividual variability and self reinforcement in fanning response.
Behavioral Ecology, 15(1), 120-128. <https://doi.org/10.1093/beheco/arg101>

Winston, M. L. (1987). *The Biology of the Honey Bee*. Harvard University Press.

655 Zacepins, A., Meitalovs, J., Komasilovs, V., & Stalidzans, E. (2011). Temperature sensor
network for prediction of possible start of brood rearing by indoor wintered honey bees.
12th International Carpathian Control Conference (ICCC).
<https://doi.org/10.1109/CARPATIANCC.2011.5945901>

Zacepins, A., Stalidzans, E., & Meitalovs, J. (2012). *Application of information technologies in
precision apiculture*.

Acknowledgements

660 This project received funding from France Agrimer (ToolBee, 17-03-R) (MP, CR, DC, MH, YLC, CA). U. Godeau was supported by the European Horizon 2020 research and innovation program under grant agreement no. 817622 (B-GOOD). We are extremely grateful to Hervé Mugnier, Jean-Camille Lebreton, Aurelien Neveux and Sully Vitry for their technical assistance with the hive sensors.