

1 **Title:** Telomere length in house sparrows increases in early-life and can be paternally inherited

2

3 Sophie Bennett<sup>1,2\*</sup>, Antje Girndt<sup>1,3,4</sup>, Alfredo Sánchez-Tójar<sup>1,3,4</sup>, Terry Burke<sup>5</sup>, Mirre Simons<sup>5</sup>, Julia  
4 Schroeder<sup>1</sup>

5

6 <sup>1</sup> Division of Biology, Imperial College London, Silwood Park, United Kingdom

7 <sup>2</sup> UK Centre for Ecology & Hydrology, Bush Estate, Penicuik, Midlothian EH26 0QB, UK

8 <sup>3</sup> Department of Evolutionary Biology, Max Planck Institute for Ornithology, Seewiesen, Germany

9 <sup>4</sup> Department of Evolutionary Biology, Bielefeld University, Germany

10 <sup>5</sup> Department of Animal and Plant Sciences, University of Sheffield, Sheffield, United Kingdom

11

12 **\*Corresponding author contact:** bennett.i.sophie@gmail.com

13 **Abstract:**

14 Offspring of older parents in many species display decreased longevity, a faster ageing rate and  
15 lower fecundity than offspring born to younger parents. Biomarkers, such as telomeres, that tend to  
16 shorten as individual age, may provide insight into the mechanisms of parental age effects. Parental  
17 age could determine telomere length either through inheritance of shortened telomeres or through  
18 indirect effects, such as variation in parental care with parent ages, which in turn might lead to  
19 variation in offspring telomere length. There is no current consensus as to the heritability of  
20 telomere length, and the direction and extent of parental age effects however. To address this, here  
21 we experimentally investigate how parental age is associated with telomere length at two time  
22 points in early life in a captive population of house sparrows (*Passer domesticus*). We  
23 experimentally separated parental age from sex effects by allowing the parent birds to only mate  
24 with young, or old partners. We found that telomere length of the offspring increased between the  
25 age of 0.5 and 3 months at the group and individual level, which has been reported previously  
26 predominantly in non-avian taxa. We further show that older fathers produced daughters with a  
27 greater early-life increase in telomere length, supporting sex-specific inheritance, and or sex-  
28 specific non-genetic effects. Overall, our results highlight the need for more studies testing early-  
29 life telomere dynamics and sex-specific heritability of telomere length.

30 **Key words:** telomere dynamics, ageing, inter-generational effects, z-linked inheritance,  
31 transgenerational effects, Lansing effect

32 ***Introduction***

33 Parent age at conception is often associated with their offspring's' life-history, with offspring of  
34 older parents commonly having reduced reproductive success and longevity (Heidinger et al., 2016;  
35 Priest et al., 2002; Schroeder et al., 2015). Moreover, in some species, offspring of older parents  
36 experience higher rates of senescence, cellular ageing, and decreased longevity compared to their  
37 older siblings (Bouwhuis et al., 2010; Broer et al., 2013; Torres et al., 2011). While some studies do  
38 not find such effects (Froy et al., 2017; Unryn et al., 2005), the associations are reported across a  
39 wide range of taxa from rotifers (King, 1983) and insects (Priest et al., 2002) to birds and mammals  
40 (Bize et al., 2009; Haussmann et al., 2003b), and is termed the Lansing effect (Lansing, 1947).

41 The relative length of telomeres, the chromosome capping structures consisting of TTAGGG base  
42 pair repeats, is associated with biological age and longevity (Heidinger et al., 2012; Mather et al.,  
43 2011; Vedder et al., 2021). Telomeres partly function to prevent DNA damage from reactive  
44 oxygen species (Aubert and Lansdorp, 2008). The activity levels of telomerase, the RNA-protein  
45 complex responsible for ligating TTAGGG repeats, decline rapidly in early life and are tissue  
46 specific (Taylor and Delany, 2000). Together this leads to a gradual telomere shortening over an  
47 individual's lifetime (Aubert and Lansdorp, 2008; Finkel and Holbrook, 2000), which is why  
48 telomere length is often used as a biomarker for biological age (Mather et al., 2011; Zglinicki and  
49 Martin-Ruiz, 2005). However, whether there is a direct causal link between telomere length and an  
50 individual's age remains unclear (Boonekamp et al., 2013; Simons, 2015).

51 In birds, telomere loss is fastest in early-life and an initially longer telomere length is associated  
52 with longer subsequent lifespans in captive (Reichert et al., 2013; Wilbourn et al., 2018) and wild  
53 (Haussmann et al., 2003a; Heidinger et al., 2016; Reed et al., 2008; Richardson et al., 2001;  
54 Salomons et al., 2009; Vedder et al. 2021) bird populations. There is evidence for telomere length  
55 being heritable in birds (Vedder et al., 2021), and telomere dynamics have been associated with sex-  
56 specific parental age and telomere length (Asghar et al., 2015; Horn et al., 2011; Reichert et al.,  
57 2015; Salomons et al., 2009). This suggests that indeed, some Lansing-type effects may be inherited  
58 via telomere length. However, the direction of the association between telomere length, and  
59 maternal and paternal age varies even within bird species (Dugdale and Richardson, 2018). In birds,  
60 the offspring of older mothers may have shorter telomeres and a faster attrition rate, especially in  
61 early development (Asghar et al., 2015; Salomons et al., 2009). Conversely, negative associations  
62 between paternal age and offspring telomere length have been observed in the absence of maternal  
63 correlation (Horn et al., 2011).

64 Between taxa, studies on the heritability of telomere length are conflicting. The heritability of  
65 telomere length can be sex-specific and is often larger in the heterogametic sex; suggesting some  
66 degree of maternal inheritance in birds (Asghar et al., 2015; Horn et al., 2011; Reichert et al., 2015)  
67 and paternal inheritance in humans (Eisenberg et al., 2017; Njajou et al., 2007; Nordfjäll et al.,  
68 2009). However, homogametic inheritance of telomere length has also been found in humans (Broer  
69 et al., 2013), in some bird species (Bauch et al., 2019; Bouwhuis et al., 2018), and in lizards (Olsson  
70 et al., 2011). A sex-specific lack of heritability has also been found in several bird species (Atema et  
71 al., 2015; Heidinger et al., 2012; Kucera, 2018). Overall, parental age effects on offspring telomere  
72 length, dynamics and heritability are complex, and vary in extent and direction of impact within and  
73 between taxa.

74 Here, we test for sex-specific, age-related parental effects on offspring telomere dynamics in  
75 captive house sparrows *Passer domesticus*. By pairing different age categories of parent birds, we  
76 experimentally test the hypothesis that offspring of older parents have shorter telomeres and faster  
77 telomere attrition rates than offspring from younger parents.

## 78 **Methods:**

### 79 ***Study species and experimental design:***

80 We used captive house sparrows at the Max Planck Institute for Ornithology, Seewiesen, Germany,  
81 during the breeding season of 2014. We used 42 pairs of male and female sparrows, which were  
82 assigned to four treatments, each with an equal sex ratio and a uniform distribution of ages across  
83 both sexes to control for age-assortative mating. We experimentally bred pairs in one of four age  
84 combinations: old-female/ old-male (OO, n=8 pairs), old-female/ young-male (OM, n=11 pairs),  
85 young-female/ old-male (YO, n=13 pairs), and young-female/ young-male (YY, n=10 pairs). Young  
86 birds hatched the preceding summer. Old (O) was defined as sparrows aged 4 years and older,  
87 although most individuals were 7 years or older (Males: 8 years = 2, 9 years = 21; Females: 4 years  
88 = 1, 7 years= 10, 8 years= 4, and 9 years= 1). The difference in age distribution between females  
89 and males corresponded to that observed in the wild, where females live shorter than  
90 males (Schroeder et al., 2012). We did not use the middle aged groups because in wild house  
91 sparrows, reproductive senescence may start at 3 years for females (Schroeder et al., 2012), or 5  
92 years in males (Hsu et al. 2017). Each treatment group was replicated twice in two separate  
93 breeding groups located in separate aviaries. Each replicate aviary contained 15.3 (s.d.= 4.9) males  
94 and 14.6±2.4 females of the respective age class. Bird husbandry is described in Girndt et al.  
95 (2017).

96 Each replicate aviary was equipped with one more nest box than breeding pairs to reduce male-male  
97 competition for nest boxes. Sparrows were then allowed to naturally display, form pair bonds,  
98 choose a mate restricted by the age class present, and raise their young (Girndt et al, 2018). We  
99 systematically monitored breeding and identified the parents attending each nest box by observing  
100 the individual birds' colour ring combinations.

#### 101 ***Blood sample collection:***

102 We took blood samples from chicks 0.5 months after they hatched (n= 75). After fledging, offspring  
103 remained in the same aviary as their parents and siblings, and 2.5 months later were blood sampled  
104 again (n=59). Blood samples were collected from the brachial vein of offspring using 1mm  
105 capillary tubes and stored in 1ml of 96% ethanol. We collected samples of 56 individuals at both  
106 0.5 and 3 months to test for within-individual changes.

#### 107 ***DNA extraction and quantification:***

108 Following standard DNA extraction (Richardson et al., 2001), we measured the DNA concentration  
109 of the samples using a ThermoScientific NanoDrop8000 Spectrophotometer and standardised the  
110 concentration in our samples to 20-30ng/ml to ensure equal amplification of samples during qPCR.  
111 Where necessary, samples were diluted with T10E0.1 (10mM Tris-HCl, pH 8.0, 0.1mM EDTA, pH  
112 8.0) or concentrated using a ThermoScientific Savant DNA SpeedVac Concentrator.

#### 113 ***Estimation of telomere length:***

114 We used multiplex qPCR to determine relative telomere length. We determined 'T' as the number  
115 of telomere repeats and 'S' as the number of control gene repeats. We then used the T/S ratio as a  
116 proxy for telomere length. The four DNA primers we used are described in Criscuolo et al. (2009).  
117 We used DNA from house sparrows not included in this analysis as standards at five DNA  
118 concentrations of 80, 20, 5, 1.25 and 0.31ng/ml, on each plate. We then used these standards to  
119 produce a standard curve for all analysed samples. In each well we added 1.5µl of DNA sample,  
120 0.9µl of each primer, 10µl of Sybr®Select Master Mix and 4.9µl ddH<sub>2</sub>O. We ran each plate with an  
121 equal number of 0.5 and 3 months sample pairs from the same individual to account for any  
122 potential sample and plate effects when comparing within-individual changes in telomere length.  
123 We ran 42 samples, the five standards and a negative (with all components except a DNA sample)  
124 in duplicate on each 96-well plate. We ran the qPCR cycling conditions using QuantStudio 12kFlex  
125 Software v1.2.2 following the cycle timings given in Cawthon (2009). We analysed the software  
126 output to calculate the T/S ratio in each sample (Appendix 1.1). We altered the thresholds for the  
127 standard curve of the telomere and GAPDH primers for each plate to optimise amplification  
128 efficiencies to between a standard of 95-110. Efficiencies for each plate were between 99.3-99.7 for

129 GAPDH and 99.3-105.8 for telc and telg. The standard curve for each plate had an  $R^2$  of 0.99 and  
130 the intra- and inter-plate variation coefficients all met adequate levels (Cawthon, 2009). We also ran  
131 a melt curve to examine whether the expected two products were generated in the reaction.  
132 Additionally, we checked all plate amplification curves to see if DNA was present in the control, as  
133 this would indicate contamination. In all plates DNA was absent, or present only in very low levels  
134 in negatives, apart from very late amplification due to primer dimerization. We repeated any sample  
135 duplicates that had a standard deviation of  $>0.05$  following thresholding and used the mean T/S  
136 ratio of duplicates in our analysis. T/S ratios of offspring at 0.5 months old are referred to as T/S<sub>0.5</sub>  
137 and samples at 3 months old T/S<sub>3</sub> in our analyses. We then calculated the difference between the  
138 two measurements as  $\Delta T/S$ . All samples were analysed for telomere length at the same time and had  
139 a similar shelf time (Lieshout et al., 2020). All reagents and equipment were produced by Thermo  
140 Fisher Scientific, Waltham, Massachusetts, US.

141 ***Ethical Note:***

142 The Government of Upper Bavaria, Germany, approved the care, handling and husbandry of all  
143 birds in this study and granted a license for animal experiments to JS (Nr311.5–5682.1/1-2014-  
144 024).

145 ***Statistical Analysis:***

146 Next, we tested for a change in telomere length over the 2.5 months period by running a linear  
147 mixed-effects model (LMM) with T/S as response variable, time of sampling (0.5 months or 3  
148 months) as an explanatory fixed factor, and individual chick ID as a random effect on the intercept.  
149 Then we tested whether telomere lengths in offspring were more variable at either 0.5, or 3 months  
150 using a two-tailed F-test. Next, we ran two further LMMs with the response variable T/S<sub>0.5</sub> and  
151 T/S<sub>3</sub>, respectively. For each model we tested the fixed effects of the paternal and maternal age  
152 categories (either ‘young’ or ‘old’). To test for sex-specific parental effects, we included offspring  
153 sex as a categorical variable (with ‘male’ as the reference level) and an interaction of chick sex with  
154 parental age in the T/S<sub>0.5</sub> model. Because not all chicks were sampled at exactly 3 months after  
155 hatching (mean= 100.8 days, s.d.= 8.4), we also tested for an effect of the exact age in days of  
156 offspring in T/S<sub>3</sub> with a LMM with T/S<sub>3</sub> as the response variable, and ‘sample age’ as an  
157 explanatory covariate. We found that ‘sample age’ did not have a statistically significant effect on  
158 T/S<sub>3</sub> (posterior mode= -0.001, 95% credible interval= -0.01, 0.001, pMCMC=0.809). Still, to  
159 account for any potential bias we retained ‘sample age’ as a fixed effect in the T/S<sub>3</sub> model.

160 As the 0.5 months samples were a mix of newly-, and already-extracted DNA samples, we also  
161 tested whether time of extraction had any effect on the calculated T/S ratio as a result of DNA

162 degradation (Madisen et al., 1987) (n samples newly-extracted= 10 out of 75). We fitted a LMM  
163 with T/S<sub>0.5</sub> as the response and the time of extraction as a fixed factor, either ‘newly-’ or ‘already-  
164 extracted’. We found no statistically significant difference between newly-, and already-extracted  
165 samples (posterior mode= -0.06, 95% credible interval= -0.20, 0.08, pMCMC=0.389).

166 We included the nest box ID and aviary ID in which chicks were born as random effects on the  
167 intercept in all models to account for variance between broods and aviaries. We also included the  
168 random term of qPCR plate ID in all models to account for between-plate variance on the intercept.  
169 All models were run using the Markov chain Monte Carlo (MCMC) method in the R package  
170 MCMCglmm v.2.29 (Hadfield, 2010).

### 171 ***Model validation:***

172 As we used a Bayesian modelling approach, we deemed fixed terms to be statistically significant if  
173 their 95% credible intervals (95CI) did not span zero, and we also report MCMC-p-values  
174 (pMCMC) (Hadfield, 2010). All terms were retained in models irrespective of their statistical  
175 significance. We directly assessed model autocorrelation for fixed and random effects to ensure the  
176 risk of type I errors was not inflated. We also inspected iteration and density plots to ensure that  
177 effects showed equal variation around a constant mode and demonstrated convergence (Gelman and  
178 Hill, 2006; Hadfield, 2010). We examined collinearity of fixed effects, as collinearity could distort  
179 model results, which did not exceed 0.7 (Dormann et al., 2013). We ran all models for 100,000  
180 iterations with a thinning interval of 10 and used default priors. All statistical analyses were carried  
181 out in R v.3.6.1 (R Core Team, 2019)

### 182 ***Results:***

183 Unexpectedly, the telomere length for offspring increased within 80% of individuals between 0.5  
184 and 3 months of age (n = 45/56) for those where both measurements were available. On average,  
185 the difference between T/S<sub>0.5</sub> and T/S<sub>3</sub> was statistically significantly positive (Figure 1 and Table 1).  
186 Further, as chicks aged, they varied more in their telomere lengths; there was greater variance in  
187 T/S<sub>3</sub> than in T/S<sub>0.5</sub> (coefficient of variance (CV)<sub>±s.e.</sub>: 0.5 months: 0.22±0.02, n=75, 3 months:  
188 0.27±0.02, n=59; F-test: F=0.43, p<0.01).

189

190

191


192

193

194 **Table 1:** Results from a Bayesian MCMC linear mixed-effects model testing the difference  
 195 between telomere length in house sparrow chicks at 0.5 and 3 months of age.

| <i>Parameter</i>        | <i>Estimate</i> | <i>95% confidence intervals</i> | <i>p<sub>MCMC</sub></i> |
|-------------------------|-----------------|---------------------------------|-------------------------|
| <b><i>Intercept</i></b> | <b>0.93</b>     | <b>0.84 - 0.99</b>              | <b>&lt;0.001</b>        |
| <b><i>Chick age</i></b> | <b>0.19</b>     | <b>0.12 - 0.26</b>              | <b>&lt;0.001</b>        |
| <i>Random effects</i>   |                 |                                 |                         |
| <i>Chick ID</i>         | 0.00            | 0.00 - 0.01                     |                         |
| <i>Nest box</i>         | 0.02            | 0.01 - 0.04                     |                         |
| <i>Aviary</i>           | 0.00            | 0.00 - 0.00                     |                         |
| <i>qPCR plate ID</i>    | 0.00            | 0.00 - 0.01                     |                         |
| <i>Residual</i>         | 0.04            | 0.02 - 0.05                     |                         |

202 Chick age was modelled as a binary variable of either 0.5 months or 3 months, with 0.5 months as a reference level. 0.5  
 203 months: n=75 chicks, 3 months: n=59. Estimates shown are posterior modes. Significant effects are shown in bold.



204 **Figure 1:** Change in telomere length (log(T/S Ratio)) within house sparrow chicks at 0.5 and 3  
 205 months of age. A) Individuals are connected by a line (n offspring with samples at 0.5 months=75,  
 206 at 3 months=59). B) Boxplots show the mean (central line) and 25<sup>th</sup> and 75<sup>th</sup> percentiles (lower and  
 207 upper box bounds respectively) of the log(T/S Ratio) within age group of the chicks' parents (Y =  
 208 <2 years old, O > 3 years old for females and >7 years old for males). T/S Ratio is presented on a  
 209 log scale to aid visualisation. YO = young mothers, old fathers (n=19, 12). OO = both parents old  
 210 (n=18, 19). OY = old mothers, young fathers (n=17, 18). YY = both parents young (n=15, 10).

211 We did not find a statistically significant effect of parental age class on T/S<sub>0.5</sub>, which is shortly  
 212 before sparrows gain independence and fledge from their nest (Table 2). However, the T/S<sub>3</sub> model  
 213 detected statistically significant effects of paternal age, and the interaction between these two

214 variables. This means that daughters of young fathers had shorter telomeres than daughters of old  
 215 fathers (Table 2, Fig. 2).


216 **Table 2:** Results from two Bayesian MCMC general linear mixed-effects models with telomere  
 217 length of house sparrow chicks at age 0.5 months and 3 months as response variables, respectively.

| <i>Parameter</i>                       | <i>T/S<sub>0.5</sub></i> |                    |                         | <i>T/S<sub>3</sub></i> |                     |                         |
|----------------------------------------|--------------------------|--------------------|-------------------------|------------------------|---------------------|-------------------------|
|                                        | <i>Estimate</i>          | <i>95% CI</i>      | <i>p<sub>MCMC</sub></i> | <i>Estimate</i>        | <i>95% CI</i>       | <i>p<sub>MCMC</sub></i> |
| <b><i>Intercept</i></b>                | <b>0.97</b>              | <b>0.84 - 1.10</b> | <b>&lt;0.001</b>        | <b>1.57</b>            | <b>0.32 - 2.71</b>  | <b>0.022</b>            |
| <i>Chick sex</i>                       | -0.08                    | -0.22 - 0.11       | 0.469                   | -0.23                  | -0.41 - 0.61        | 0.121                   |
| <i>Maternal age</i>                    | -0.07                    | -0.23 - 0.08       | 0.346                   | 0.06                   | -0.32 - 0.21        | 0.711                   |
| <i>Maternal age x Chick sex</i>        | 0.06                     | -0.15 - 0.25       | 0.597                   | 0.09                   | -0.21 - 0.43        | 0.573                   |
| <b><i>Paternal age</i></b>             | -0.03                    | -0.20 - 0.10       | 0.481                   | <b>-0.27</b>           | <b>-0.52 - 0.00</b> | <b>0.047</b>            |
| <b><i>Paternal age x Chick sex</i></b> | 0.6                      | -0.07 - 0.34       | 0.168                   | <b>-0.40</b>           | <b>0.09 - 0.71</b>  | <b>0.016</b>            |
| <i>Sample day</i>                      |                          |                    |                         | 0.00                   | -0.01 - 0.01        | 0.694                   |
| <b><i>Random effects</i></b>           |                          |                    |                         |                        |                     |                         |
| <i>Nest box</i>                        | 0.01                     | 0.00 - 0.03        |                         | 0.01                   | 0.00 - 0.02         |                         |
| <i>Aviary</i>                          | 0.00                     | 0.00 - 0.01        |                         | 0.01                   | 0.00 - 0.01         |                         |
| <i>qPCR plate ID</i>                   | 0.00                     | 0.00 - 0.01        |                         | 0.00                   | 0.00 - 0.01         |                         |
| <i>Residual</i>                        | 0.08                     | 0.03 - 0.11        |                         | 0.08                   | 0.03 - 0.11         |                         |

218 Maternal and paternal age were modelled as a binary variable of either young or old; young was <2 years old, and old  
 219 was determined as >3 years old for females and >7 years for males). 0.5 months: n=69 chicks, 3 months: n=59. The  
 220 reference level for parental ages was ‘old’, and the ‘female’ was the reference level for chick sex. Estimates shown are  
 221 posterior modes. Significant effects are shown in bold.

222

223



224

225 **Figure 2:** Post-hoc effect size plot from a linear mixed-effects model testing the relationship  
 226 between T/S<sub>3</sub>, father age, and sex of chicks (Table 1). Fathers were assigned an age category of  
 227 young, ‘Y’, or old, ‘O’. A young father was <2 years old, and an old father was determined as >7  
 228 years old. Chick sex is indicated as either female, ‘red’, or male, ‘blue’. The number of offspring in  
 229 each category; Y, and female=12, male=16, O, and female=16, male=15. Squares represent the  
 230 model estimates effect sizes of T/S ratio for each paternal age x chick sex, associated lines represent  
 231 95% credible intervals.

232 **Discussion:**

233 Individual chick telomere length increased between 0.5 and 3 months of age. This increase  
 234 disagrees with much of the published literature, which generally find a decrease in telomere length  
 235 in early-life (Boonekamp et al., 2014; Cerchiara et al., 2017; De Meyer et al., 2007; Hoelzl et al.,  
 236 2016; Salomons et al., 2009). While a population level increase in telomere length has previously  
 237 been found in some long-lived bird species (Hausmann et al., 2007; Pauliny et al., 2012), other  
 238 studies have found that telomeres elongation for a proportion of chicks is more common in smaller,  
 239 shorter-lived species (Brown et al., 2021; Eisenberg, 2019). For example, a study on jackdaws  
 240 *Corvus monedula* found that between 5 and 30 days post-hatching, telomere lengths increased for  
 241 25% of sampled offspring (Grasman et al., 2011). An increase in early-life telomere length has also  
 242 been observed in non-avian taxa, including water pythons *Liasis fuscus* (Ujvari and Madsen, 2009)  
 243 and European badgers *Meles meles* (van Lieshout et al., 2019). A lack of comparable published  
 244 research exploring a change in telomere length using multiple time points in early life may, in part,  
 245 explain the surprising nature of our observed increase in telomere length in early-life.

246 An increase in telomere length can have methodological and/or biological explanations. First, it  
247 could be due to DNA in samples degrading over time (Madisen et al., 1987; but see Seutin et al.,  
248 1991). Since we used pre-extracted DNA for the majority of 0.5 month samples, we investigated  
249 whether differential telomere degradation rates between extracted DNA and blood sample types  
250 could be a cause for the observed increase. However, we found no statistically significant difference  
251 between the telomere lengths of newly- and already-extracted samples and thus, telomere  
252 degradation is an unlikely explanation for our results.

253 Second, qPCR plates contained both 0.5 and 3 months samples, and between-plate variance was  
254 negligible in all our models, highlighting that this element of our methodology had little impact on  
255 our results. Overall, we monitored procedural efficiency throughout data collection and did not  
256 identify any other potential methodological sources of variation, and so, we are convinced that the  
257 increase in telomere length observed in our study has a biological explanation. For example,  
258 telomerase activity might have been maintained in the offspring after the first sample was taken.  
259 Indeed, two studies have shown that telomerase activity can be maintained up to five weeks post-  
260 hatching in zebra finches *Taeniopygia guttata* (Hausmann et al., 2007) and chickens *Gallus gallus*  
261 (Taylor and Delany, 2000). Yet, neither of these studies assessed telomerase activity at multiple  
262 time points in the same individual's early-life post-hatching, which remains as an interesting future  
263 avenue for the field.

264 While we expected that old parents would produce offspring with shorter telomeres, as found in  
265 other short-lived bird species (Bauch et al., 2019; Criscuolo et al., 2017; Sparks et al., 2020), our  
266 experimental approach found that old fathers produced daughters with longer telomeres, but only 3  
267 months after hatching, indicating an environmental effect. Similar positive effects of parental age  
268 have also previously been found in long and short-lived bird species (Dupont et al., 2018, Asghar et  
269 al., 2015; Becker et al., 2015). Positive effects of parental age on offspring telomere length may  
270 arise from a potentially improved parental care that older individuals may be able to provide  
271 compared to inexperienced, young breeders. Again though, previous studies have found a negative  
272 effect of parental age resulting from the poorer condition of these old individuals (Bouwhuis et al.,  
273 2018; Criscuolo et al., 2017), or a lack of an effect of parental age on parental care (Nakagawa et  
274 al., 2007). Further, in some studies testing parent sex-specific effects, offspring telomere length was  
275 found to correlate only with maternal age, and only relatively soon after hatching (ten days:  
276 Reichert et al. (2015); nine days: Asghar et al. (2015)). As we found no influence of maternal age  
277 in our study, an influence of maternal age on offspring telomere length may well have been present,  
278 but already diminished below detectable levels 0.5 months after hatching.

279 While telomere lengths in offspring have been shown to be affected by an offspring's environment  
280 (Dugdale and Richardson, 2018; Lieshout et al., 2021), effects of paternal age in birds have been  
281 found to be independent of this (Bauch et al., 2019; Boonekamp et al., 2014). As such, overall, there  
282 is growing support for at least contributory paternal inheritance of telomere length in some species  
283 of birds (Bouwhuis et al., 2018; Olsson et al., 2011; this study). The combined positive effect of  
284 having an older father has been theorised to result from an upregulation of telomerase activity in  
285 sperm and a subsequent increase in gamete telomere length as males age (De Meyer et al., 2007); as  
286 such a positive association of paternal telomere length with age has also been found in humans  
287 (Kimura et al., 2008; Unryn et al., 2005). Therefore, a combination of telomerase activity in sperm  
288 in fathers, a form of Z-linked inheritance, and potential parental care benefits discussed above may  
289 explain the positive effect of increasing father age on offspring telomere lengths, with larger effects  
290 seen on daughters compared to sons as observed here.

291 However, we did not detect an effect of parental age on offspring telomere length at 0.5 months  
292 after hatching. Heidinger et al. (2016) similarly found no effect of parental age on offspring  
293 telomere length in very early-life at 25 days after hatching in European shags *Phalacrocorax*  
294 *aristotelis*. Further, variation in pre-fledging telomere length may in part be explained by brood-  
295 specific additive genetic effects (Voillemot et al., 2012). As such, it may be that at later time points  
296 effects of parent age and post-fledging environmental factors appear to be more important than  
297 brood-specific effects in determining offspring telomere length. Again, there is a need for more  
298 studies investigating the relationship between paternal age and telomere dynamics to detect when  
299 and how patterns of telomere dynamics are driven.

300 In sum, our results indicate that paternal age effects are more influential on offspring telomere  
301 length than maternal age effects in our population of house sparrows, with the daughters of older  
302 fathers having longer telomeres. Future analyses of telomerase activity levels in both the sperm of  
303 adult males and the somatic tissues of offspring would yield further insights into the drivers of  
304 parental age effects on offspring telomere dynamics in early-life.

305 **Acknowledgements:**

306 The authors would like to thank to Natalie dos Remedios for her help and support with processing  
307 samples, and Marta Precioso for helpful discussions on the methodology. We thank Annemarie  
308 Grötsch and Natalie Fischer for care of the captive population.

309 **Competing interests' statement:**

310 No competing interests declared.

311 **Author contributions:**

312 Conceptualization: SB, JS; Methodology: SB, JS, MJPS; Validation: SB, MJPS; Formal analysis:  
313 SB, JS; Investigation: SB, AG, JS, AST; Resources: JS, MJPS, TB; Data curation: SB, JS, TB;  
314 Writing- original draft: SB; Writing- review & editing: SB, JS, AG, AST, MJPS; Visualization: SB,  
315 JS, AG, AST; Supervision: JS; Project administration: JS, TB; Funding acquisition: JS, TB.

316 **Funding:**

317 AG was supported by the Bielefeld Young Researcher's Fund; AST was funded by the German  
318 Research Foundation (DFG) as part of the SFB 592 TRR 212 (NC<sup>3</sup>; project numbers 316099922,  
319 396782608); MJPS is a Wellcome Sir Henry Dale Fellow [MJPS]. This work was funded by the  
320 Volkswagen Foundation and a Grant CIG from the European Union [PCIG12-GA-2012-333096 to  
321 JS].

322 **References:**

- 323 Asghar, M., Bensch, S., Tarka, M., Hansson, B., Hasselquist, D., 2015. Maternal and genetic factors  
324 determine early life telomere length. *Proc. R. Soc. Lond. B Biol. Sci.* 282, 20142263.  
325 <https://doi.org/10.1098/rspb.2014.2263>
- 326 Atema, E., Mulder, E., Dugdale, H.L., Briga, M., Noordwijk, A.J. van, Verhulst, S., 2015. Heritability of  
327 telomere length in the Zebra Finch. *J. Ornithol.* 156, 1113–1123. [https://doi.org/10.1007/s10336-](https://doi.org/10.1007/s10336-015-1212-7)  
328 [015-1212-7](https://doi.org/10.1007/s10336-015-1212-7)
- 329 Aubert, G., Lansdorp, P.M., 2008. Telomeres and Aging. *Physiol. Rev.* 88, 557–579.  
330 <https://doi.org/10.1152/physrev.00026.2007>
- 331 Bauch, C., Boonekamp, J.J., Korsten, P., Mulder, E., Verhulst, S., 2019. Epigenetic inheritance of telomere  
332 length in wild birds. *PLOS Genet.* 15, e1007827. <https://doi.org/10.1371/journal.pgen.1007827>
- 333 Becker, P.J.J., Reichert, S., Zahn, S., Hegelbach, J., Massemin, S., Keller, L.F., Postma, E., Criscuolo, F., 2015.  
334 Mother–offspring and nest-mate resemblance but no heritability in early-life telomere length in  
335 white-throated dippers. *Proc. R. Soc. B Biol. Sci.* 282, 20142924.  
336 <https://doi.org/10.1098/rspb.2014.2924>
- 337 Bize, P., Criscuolo, F., Metcalfe, N.B., Nasir, L., Monaghan, P., 2009. Telomere dynamics rather than age  
338 predict life expectancy in the wild. *Proc. R. Soc. Lond. B Biol. Sci.* 276, 1679–1683.  
339 <https://doi.org/10.1098/rspb.2008.1817>
- 340 Boonekamp, J.J., Mulder, G.A., Salomons, H.M., Dijkstra, C., Verhulst, S., 2014. Nestling telomere  
341 shortening, but not telomere length, reflects developmental stress and predicts survival in wild  
342 birds. *Proc. R. Soc. Lond. B Biol. Sci.* 281, 20133287. <https://doi.org/10.1098/rspb.2013.3287>
- 343 Boonekamp, J.J., Simons, M.J.P., Hemerik, L., Verhulst, S., 2013. Telomere length behaves as biomarker of  
344 somatic redundancy rather than biological age. *Aging Cell* 12, 330–332.  
345 <https://doi.org/10.1111/accel.12050>
- 346 Bouwhuis, S., Charmantier, A., Verhulst, S., Sheldon, B.C., 2010. Trans-generational effects on ageing in a  
347 wild bird population. *J. Evol. Biol.* 23, 636–642. <https://doi.org/10.1111/j.1420-9101.2009.01929.x>
- 348 Bouwhuis, S., Verhulst, S., Bauch, C., Vedder, O., 2018. Reduced telomere length in offspring of old fathers  
349 in a long-lived seabird. *Biol. Lett.* 14, 20180213. <https://doi.org/10.1098/rsbl.2018.0213>
- 350 Broer, L., Codd, V., Nyholt, D.R., Deelen, J., Mangino, M., Willemsen, G., Albrecht, E., Amin, N., Beekman,  
351 M., de Geus, E.J.C., Henders, A., Nelson, C.P., Steves, C.J., Wright, M.J., de Craen, A.J.M., Isaacs, A.,  
352 Matthews, M., Moayyeri, A., Montgomery, G.W., Oostra, B.A., Vink, J.M., Spector, T.D., Slagboom,  
353 P.E., Martin, N.G., Samani, N.J., van Duijn, C.M., Boomsma, D.I., 2013. Meta-analysis of telomere  
354 length in 19 713 subjects reveals high heritability, stronger maternal inheritance and a paternal age  
355 effect. *Eur. J. Hum. Genet.* 21, 1163–1168. <https://doi.org/10.1038/ejhg.2012.303>
- 356 Brown, thomas, Dugdale, H., Spurgin, L., Komdeur, J., Burke, T., Richardson, D., 2021. Causes and  
357 Consequences of Telomere Lengthening in a Wild Vertebrate Population (preprint). Authorea.  
358 <https://doi.org/10.22541/au.161408541.15345829/v1>
- 359 Cawthon, R.M., 2009. Telomere length measurement by a novel monochrome multiplex quantitative PCR  
360 method. *Nucleic Acids Res.* 37, e21–e21. <https://doi.org/10.1093/nar/gkn1027>
- 361 Cerchiara, J.A., Risques, R.A., Prunkard, D., Smith, J.R., Kane, O.J., Dee Boersma, P., 2017. Telomeres  
362 shorten and then lengthen before fledging in Magellanic penguins (*Spheniscus magellanicus*). *Aging*  
363 9, 487–493. <https://doi.org/10.18632/aging.101172>
- 364 Criscuolo, F., Bize, P., Nasir, L., Metcalfe, N.B., Foote, C.G., Griffiths, K., Gault, E.A., Monaghan, P., 2009.  
365 Real-time quantitative PCR assay for measurement of avian telomeres. *J. Avian Biol.* 40, 342–347.  
366 <https://doi.org/10.1111/j.1600-048X.2008.04623.x>
- 367 Criscuolo, F., Zahn, S., Bize, P., 2017. Offspring telomere length in the long lived Alpine swift is negatively  
368 related to the age of their biological father and foster mother. *Biol. Lett.* 13, 20170188.  
369 <https://doi.org/10.1098/rsbl.2017.0188>
- 370 De Meyer, T., Rietzschel, E.R., Buyzere, D., L, M., De Bacquer, D., Van Criekinge, W., Backer, D., G, G.,  
371 Gillebert, T.C., Van Oostveldt, P., Bekaert, S., 2007. Paternal age at birth is an important  
372 determinant of offspring telomere length. *Hum. Mol. Genet.* 16, 3097–3102.  
373 <https://doi.org/10.1093/hmg/ddm271>

374 Dormann, C.F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J.R.G., Gruber, B.,  
375 Lafourcade, B., Leitão, P.J., Münkemüller, T., McClean, C., Osborne, P.E., Reineking, B., Schröder, B.,  
376 Skidmore, A.K., Zurell, D., Lautenbach, S., 2013. Collinearity: a review of methods to deal with it and  
377 a simulation study evaluating their performance. *Ecography* 36, 27–46.  
378 <https://doi.org/10.1111/j.1600-0587.2012.07348.x>

379 Dugdale, H.L., Richardson, D.S., 2018. Heritability of telomere variation: it is all about the environment!  
380 *Philos. Trans. R. Soc. B Biol. Sci.* 373, 20160450. <https://doi.org/10.1098/rstb.2016.0450>

381 Dupont, S.M., Barbraud, C., Chastel, O., Delord, K., Ruault, S., Weimerskirch, H., Angelier, F., 2018. Young  
382 parents produce offspring with short telomeres: A study in a long-lived bird, the Black-browed  
383 Albatross (*Thalassarche melanophrys*). *PLOS ONE* 13, e0193526.  
384 <https://doi.org/10.1371/journal.pone.0193526>

385 Eisenberg, D.T.A., 2019. Paternal age at conception effects on offspring telomere length across species—  
386 What explains the variability? *PLOS Genet.* 15, e1007946.  
387 <https://doi.org/10.1371/journal.pgen.1007946>

388 Eisenberg, D.T.A., Tackney, J., Cawthon, R.M., Cloutier, C.T., Hawkes, K., 2017. Paternal and grandpaternal  
389 ages at conception and descendant telomere lengths in chimpanzees and humans. *Am. J. Phys.*  
390 *Anthropol.* 162, 201–207. <https://doi.org/10.1002/ajpa.23109>

391 Finkel, T., Holbrook, N.J., 2000. Oxidants, oxidative stress and the biology of ageing. *Nature* 408, 239–247.  
392 <https://doi.org/10.1038/35041687>

393 Froy, H., Bird, E.J., Wilbourn, R.V., Fairlie, J., Underwood, S.L., Salvo-Chirnside, E., Pilkington, J.G., Béréños,  
394 C., Pemberton, J.M., Nussey, D.H., 2017. No evidence for parental age effects on offspring  
395 leukocyte telomere length in free-living Soay sheep. *Sci. Rep.* 7, 9991.  
396 <https://doi.org/10.1038/s41598-017-09861-3>

397 Gelman, A., Hill, J., 2006. *Data Analysis Using Regression and Multilevel/Hierarchical Models*. Cambridge  
398 University Press.

399 Girndt, A., Chng, C.W.T., Burke, T., Schroeder, J., 2018. Male age is associated with extra-pair paternity, but  
400 not with extra-pair mating behaviour. *Sci. Rep.* 8, 8378. [https://doi.org/10.1038/s41598-018-](https://doi.org/10.1038/s41598-018-26649-1)  
401 [26649-1](https://doi.org/10.1038/s41598-018-26649-1)

402 Girndt, A., Cockburn, G., Sánchez-Tójar, A., Løvlie, H., Schroeder, J., 2017. Method matters: Experimental  
403 evidence for shorter avian sperm in faecal compared to abdominal massage samples. *PLOS ONE* 12,  
404 e0182853. <https://doi.org/10.1371/journal.pone.0182853>

405 Grasman, J., Salomons, H.M., Verhulst, S., 2011. Stochastic modeling of length-dependent telomere  
406 shortening in *Corvus monedula*. *J. Theor. Biol.* 282, 1–6. <https://doi.org/10.1016/j.jtbi.2011.04.026>

407 Hadfield, J., 2010. MCMC methods for Multi-response Generalised Linear Mixed Models: The MCMCglmm R  
408 package. *J. Stat. Softw.* 33, 1–22.

409 Haussmann, M.F., Vleck, C.M., Nisbet, I.C.T., 2003a. Calibrating the telomere clock in common terns, *Sterna*  
410 *hirundo*. *Exp. Gerontol.*, Proceedings of the 2nd Symposium on Organisms with Slow Aging (SOSA-  
411 2) 38, 787–789. [https://doi.org/10.1016/S0531-5565\(03\)00109-8](https://doi.org/10.1016/S0531-5565(03)00109-8)

412 Haussmann, M.F., Winkler, D.W., Huntington, C.E., Nisbet, I.C.T., Vleck, C.M., 2007. Telomerase activity is  
413 maintained throughout the lifespan of long-lived birds. *Exp. Gerontol.* 42, 610–618.  
414 <https://doi.org/10.1016/j.exger.2007.03.004>

415 Haussmann, M.F., Winkler, D.W., O'Reilly, K.M., Huntington, C.E., Nisbet, I.C.T., Vleck, C.M., 2003b.  
416 Telomeres shorten more slowly in long-lived birds and mammals than in short-lived ones. *Proc. R.*  
417 *Soc. Lond. B Biol. Sci.* 270, 1387–1392. <https://doi.org/10.1098/rspb.2003.2385>

418 Heidinger, B.J., Blount, J.D., Boner, W., Griffiths, K., Metcalfe, N.B., Monaghan, P., 2012. Telomere length in  
419 early life predicts lifespan. *Proc. Natl. Acad. Sci.* 109, 1743–1748.  
420 <https://doi.org/10.1073/pnas.1113306109>

421 Heidinger, B.J., Herborn, K.A., Granroth-Wilding, H.M.V., Boner, W., Burthe, S., Newell, M., Wanless, S.,  
422 Daunt, F., Monaghan, P., 2016. Parental age influences offspring telomere loss. *Funct. Ecol.* 30,  
423 1531–1538. <https://doi.org/10.1111/1365-2435.12630>

424 Hoelzl, F., Smith, S., Cornils, J.S., Aydinonat, D., Bieber, C., Ruf, T., 2016. Telomeres are elongated in older  
425 individuals in a hibernating rodent, the edible dormouse (*Glis glis*). *Sci. Rep.* 6, 36856.  
426 <https://doi.org/10.1038/srep36856>

427 Horn, T., Robertson, B.C., Will, M., Eason, D.K., Elliott, G.P., Gemmell, N.J., 2011. Inheritance of Telomere  
428 Length in a Bird. *PLOS ONE* 6, e17199. <https://doi.org/10.1371/journal.pone.0017199>

429 Kimura, M., Cherkas, L.F., Kato, B.S., Demissie, S., Hjelmberg, J.B., Brimacombe, M., Cupples, A., Hunkin,  
430 J.L., Gardner, J.P., Lu, X., Cao, X., Sastrasinh, M., Province, M.A., Hunt, S.C., Christensen, K., Levy, D.,  
431 Spector, T.D., Aviv, A., 2008. Offspring's Leukocyte Telomere Length, Paternal Age, and Telomere  
432 Elongation in Sperm. *PLOS Genet.* 4, e37. <https://doi.org/10.1371/journal.pgen.0040037>

433 King, C.E., 1983. A re-examination of the Lansing effect. *Hydrobiologia* 104, 135–139.  
434 <https://doi.org/10.1007/BF00045959>

435 Kucera, A., 2018. Sperm Telomere Dynamics: Natural Variation and Sensitivity to Environmental Influences  
436 in House Sparrows (*Passer domesticus*) (PhD). North Dakota State University.

437 Lansing, A.I., 1947. A Transmissible, Cumulative, and Reversible Factor in Aging. *J. Gerontol.* 2, 228–239.  
438 <https://doi.org/10.1093/geronj/2.3.228>

439 Lieshout, S.H.J. van, Froy, H., Schroeder, J., Burke, T., Simons, M.J.P., Dugdale, H.L., 2020. Slicing: A  
440 sustainable approach to structuring samples for analysis in long-term studies. *Methods Ecol. Evol.*  
441 11, 418–430. <https://doi.org/10.1111/2041-210X.13352>

442 Lieshout, S.H.J. van, Sparks, A.M., Bretman, A., Newman, C., Buesching, C.D., Burke, T., Macdonald, D.W.,  
443 Dugdale, H.L., 2021. Estimation of environmental, genetic and parental age at conception effects on  
444 telomere length in a wild mammal. *J. Evol. Biol.* 34, 296–308. <https://doi.org/10.1111/jeb.13728>

445 Madisen, L., Hoar, D.I., Holroyd, C.D., Crisp, M., Hodes, M.E., Reynolds, J.F., 1987. The effects of storage of  
446 blood and isolated DNA on the integrity of DNA. *Am. J. Med. Genet.* 27, 379–390.  
447 <https://doi.org/10.1002/ajmg.1320270216>

448 Mather, K.A., Jorm, A.F., Parslow, R.A., Christensen, H., 2011. Is Telomere Length a Biomarker of Aging? A  
449 Review. *J. Gerontol. Ser. A* 66A, 202–213. <https://doi.org/10.1093/gerona/glq180>

450 Nakagawa, S., Ockendon, N., Gillespie, D.O.S., Hatchwell, B.J., Burke, T., 2007. Does the badge of status  
451 influence parental care and investment in house sparrows? An experimental test. *Oecologia* 153,  
452 749–760. <https://doi.org/10.1007/s00442-007-0765-4>

453 Njajou, O.T., Cawthon, R.M., Damcott, C.M., Wu, S.-H., Ott, S., Garant, M.J., Blackburn, E.H., Mitchell, B.D.,  
454 Shuldiner, A.R., Hsueh, W.-C., 2007. Telomere length is paternally inherited and is associated with  
455 parental lifespan. *Proc. Natl. Acad. Sci.* 104, 12135–12139.  
456 <https://doi.org/10.1073/pnas.0702703104>

457 Nordfjäll, K., Svenson, U., Norrback, K.-F., Adolfsson, R., Roos, G., 2009. Large-scale parent–child  
458 comparison confirms a strong paternal influence on telomere length. *Eur. J. Hum. Genet.* 18, 385–  
459 389. <https://doi.org/10.1038/ejhg.2009.178>

460 Olsson, M., Pauliny, A., Wapstra, E., Uller, T., Schwartz, T., Blomqvist, D., 2011. Sex Differences in Sand  
461 Lizard Telomere Inheritance: Paternal Epigenetic Effects Increases Telomere Heritability and  
462 Offspring Survival. *PLOS ONE* 6, e17473. <https://doi.org/10.1371/journal.pone.0017473>

463 Pauliny, A., Larsson, K., Blomqvist, D., 2012. Telomere dynamics in a long-lived bird, the barnacle goose.  
464 *ResearchGate* 12, 257. <https://doi.org/10.1186/1471-2148-12-257>

465 Priest, N.K., Mackowiak, B., Promislow, D.E.L., 2002. The role of parental age effects on the evolution of  
466 aging. *Evolution* 56, 927–935. [https://doi.org/10.1554/0014-3820\(2002\)056\[0927:TROPAE\]2.0.CO;2](https://doi.org/10.1554/0014-3820(2002)056[0927:TROPAE]2.0.CO;2)

467 R Core Team, 2019. R: A Language and Environment for Statistical Computing. R Found. Stat. Comput.  
468 Vienna Austria Version 3.6.1.

469 Reed, T.E., Kruuk, L.E.B., Wanless, S., Frederiksen, M., Cunningham, E.J.A., Harris, M.P., 2008. Reproductive  
470 senescence in a long-lived seabird: rates of decline in late-life performance are associated with  
471 varying costs of early reproduction. *Am. Nat.* 171, E89–E101. <https://doi.org/10.1086/524957>

472 Reichert, S., Criscuolo, F., Verinaud, E., Zahn, S., Massemin, S., 2013. Telomere Length Correlations among  
473 Somatic Tissues in Adult Zebra Finches. *PLOS ONE* 8, e81496.  
474 <https://doi.org/10.1371/journal.pone.0081496>

475 Reichert, S., Rojas, E.R., Zahn, S., Robin, J.-P., Criscuolo, F., Massemin, S., 2015. Maternal telomere length  
476 inheritance in the king penguin. *Heredity* 114, 10–16. <https://doi.org/10.1038/hdy.2014.60>

477 Richardson, D.S., Jury, F.L., Blaakmeer, K., Komdeur, J., Burke, T., 2001. Parentage assignment and extra-  
478 group paternity in a cooperative breeder: the Seychelles warbler (*Acrocephalus sechellensis*). *Mol.*  
479 *Ecol.* 10, 2263–2273. <https://doi.org/10.1046/j.0962-1083.2001.01355.x>

480 Salomons, H.M., Mulder, G.A., Zande, L. van de, Haussmann, M.F., Linskens, M.H.K., Verhulst, S., 2009.  
481 Telomere shortening and survival in free-living corvids. *Proc. R. Soc. Lond. B Biol. Sci.* 276, 3157–  
482 3165. <https://doi.org/10.1098/rspb.2009.0517>  
483 Schroeder, J., Burke, T., Mannarelli, M.-E., Dawson, D.A., Nakagawa, S., 2012. Maternal effects and  
484 heritability of annual productivity. *J. Evol. Biol.* 25, 149–156. <https://doi.org/10.1111/j.1420-9101.2011.02412.x>  
485  
486 Schroeder, J., Nakagawa, S., Rees, M., Mannarelli, M.-E., Burke, T., 2015. Reduced fitness in progeny from  
487 old parents in a natural population. *Proc. Natl. Acad. Sci.* 112, 4021–4025.  
488 <https://doi.org/10.1073/pnas.1422715112>  
489 Simons, M.J.P., 2015. Questioning causal involvement of telomeres in aging. *Ageing Res. Rev.* 24, 191–196.  
490 <https://doi.org/10.1016/j.arr.2015.08.002>  
491 Sparks, A.M., Spurgin, L.G., Velde, M. van der, Fairfield, E.A., Komdeur, J., Burke, T., Richardson, D.S.,  
492 Dugdale, H., 2020. Telomere heritability and parental age at conception effects in a wild avian  
493 population. *EcoEvoRxiv*. <https://doi.org/10.32942/osf.io/eq2af>  
494 Taylor, H.A., Delany, M.E., 2000. Ontogeny of telomerase in chicken: Impact of downregulation on pre- and  
495 postnatal telomere length in vivo. *Dev. Growth Differ.* 42, 613–621. <https://doi.org/10.1046/j.1440-169x.2000.00540.x>  
496  
497 Torres, R., Drummond, H., Velando, A., 2011. Parental age and lifespan influence offspring recruitment: a  
498 long-term study in a seabird., Parental Age and Lifespan Influence Offspring Recruitment: A Long-  
499 Term Study in a Seabird. *PLoS One* PLoS ONE 6, 6, e27245–e27245.  
500 <https://doi.org/10.1371/journal.pone.0027245>, 10.1371/journal.pone.0027245  
501 Ujvari, B., Madsen, T., 2009. Short Telomeres in Hatchling Snakes: Erythrocyte Telomere Dynamics and  
502 Longevity in Tropical Pythons. *PLOS ONE* 4, e7493. <https://doi.org/10.1371/journal.pone.0007493>  
503 Unryn, B.M., Cook, L.S., Riabowol, K.T., 2005. Paternal age is positively linked to telomere length of  
504 children. *Ageing Cell* 4, 97–101. <https://doi.org/10.1111/j.1474-9728.2005.00144.x>  
505 van Lieshout, S.H.J., Bretman, A., Newman, C., Buesching, C.D., Macdonald, D.W., Dugdale, H.L., 2019.  
506 Individual variation in early-life telomere length and survival in a wild mammal. *Mol. Ecol.* 28,  
507 4152–4165. <https://doi.org/10.1111/mec.15212>  
508 Vedder, O., Moiron, M., Bichet, C., Bauch, C., Verhulst, S., Becker, P.H., Bouwhuis, S., 2021. Telomere length  
509 is heritable and genetically correlated with lifespan in a wild bird. *Mol. Ecol. Early-view*.  
510 <https://doi.org/10.1111/mec.15807>  
511 Voillemot, M., Hine, K., Zahn, S., Criscuolo, F., Gustafsson, L., Doligez, B., Bize, P., 2012. Effects of brood size  
512 manipulation and common origin on phenotype and telomere length in nestling collared  
513 flycatchers. *BMC Ecol.* 12, 17. <https://doi.org/10.1186/1472-6785-12-17>  
514 Wilbourn, R.V., Moatt, J.P., Froy, H., Walling, C.A., Nussey, D.H., Boonekamp, J.J., 2018. The relationship  
515 between telomere length and mortality risk in non-model vertebrate systems: a meta-analysis.  
516 *Philos. Trans. R. Soc. B Biol. Sci.* 373, 20160447. <https://doi.org/10.1098/rstb.2016.0447>  
517 Zglinicki, T. v, Martin-Ruiz, C.M., 2005. Telomeres as Biomarkers for Ageing and Age-Related Diseases. *Curr.*  
518 *Mol. Med.* 5, 197–203. <https://doi.org/info:doi/10.2174/1566524053586545>  
519