

1 Validating morphological condition indices and their relationship
2 with reproductive success in great-tailed grackles

3 Berens JM¹ Folsom M² Sevchik A¹ Bergeron L³ Logan CJ²
4 McCune KB^{3*}

5 2021-07-25

6 **Affiliations:**

- 7 1) Arizona State University School of Life Sciences
8 2) Max Planck Institute for Evolutionary Anthropology
9 3) University of California Santa Barbara

10 *Corresponding author: KB McCune (kelseybmccune@gmail.com)

11 **Cite as:** Berens JM, Logan CJ, Folsom M, Sevchik A, Bergeron L, McCune KB. Submitted to PCI Ecology
12 Nov 2020. Validating morphological condition indices and their relationship with reproductive success in
13 great-tailed grackles.

14 **This preregistration has been pre-study peer reviewed and received an In Principle Acceptance
15 by:**

16 Marcos Mendez (2019 In Principle Acceptance) Are condition indices positively related to each other and to
17 fitness?: a test with grackles. *Peer Community in Ecology*, 100035. 10.24072/pci.ecology.100035

- 18 • Reviewers: Javier Seoane and Isabel López-Rull

19 **ABSTRACT**

20 Morphological and physiological variation among individuals has the potential to influence multiple life
21 history characteristics such as dispersal, migration, reproductive success, and survival. Individuals that are
22 in better “condition” can disperse or migrate further or more successfully, have greater reproductive success,
23 and survive longer, particularly in years where environmental conditions are harsh. Condition is defined in
24 various ways, but is most often measured using an individual’s energetic state. These traits are difficult to
25 measure directly, therefore a variety of morphological proxies to quantify energetic condition are used instead,
26 including fat score, weight, ratio of weight to tarsus length, and a scaled mass index. However, there is mixed
27 support regarding whether these energetic condition indices relate to life history characteristics, and whether
28 the relationship is linear. Additionally, although some investigations use multiple morphological proxies for
29 energetic condition, rarely have there been direct comparisons among proxies to validate that they measure
30 the same trait. In this investigation, we define condition as an energetic state and we attempt to measure
31 it by comparing two morphological indices (fat score and the scaled mass index) to validate whether they
32 measure the same trait and whether they correlate with measures of reproductive success in our study system,
33 the great-tailed grackle (*Quiscalus mexicanus*). We found that the morphological proxies did not correlate
34 with each other, indicating that they do not measure the same trait. Further, neither proxy significantly
35 correlated with reproductive success in males, measured as whether a male held a territory containing nests

36 or not. We found that females with a high scaled mass index had a significantly lower probability that
37 their nest would survive on any given day. However, there was no relationship between female fat score and
38 nest survival. These results indicate that morphological measures of energetic condition should be validated
39 before relying on their use as a condition proxy in grackles and birds in general. Future research should
40 investigate behavioral mechanisms underlying our result that higher scaled mass index correlated with lower
41 nest survival to better understand the importance of energetic condition for reproductive success - a necessary
42 component for selection to act.

43 INTRODUCTION

44 Morphological and physiological variation among individuals has the potential to influence multiple life
45 history characteristics such as dispersal, migration, reproductive fitness, and survival (Wilder et al., 2016).
46 One trait that might be particularly likely to influence these life history characteristics is energetic condition.
47 Individuals that are in better “condition” can disperse or migrate further or more successfully, have greater
48 reproductive success, and survive longer (Heidinger et al., 2010; Liao et al., 2011; Wilder et al., 2016),
49 particularly in years where environmental conditions are harsh (Milenkaya et al., 2015). For example, a
50 study conducted on vipers showed that while the level of fat reserves in males was not related to their sexual
51 activity, females with low fat reserves engaged in sexual interactions less frequently than those with higher
52 fat reserves (Aubret et al., 2002). In contrast, mantids showed conflicting results regarding the relationship
53 between fat reserves and reproductive success (Barry & Wilder, 2013). Female mantids were fed either a
54 high protein, low lipid diet, or a high lipid, low protein diet. The females that received the high lipid diet
55 had higher lipid content in most parts of their body compared to that of their high protein diet counterparts.
56 However, they were not able to produce even half as many eggs as the females fed the high protein, low
57 lipid diet. This led to lower male attraction, measured by the number of copulation events, thus negatively
58 impacting further reproductive success.

59 A variety of morphological proxies have been used to quantify energetic condition [i.e., fat score, weight,
60 ratio of mass to structural size, residuals from a linear regression of mass as a function of structural body
61 size; Labocha et al. (2014); Jacobs et al. (2012)]. However, there is mixed support regarding whether and
62 how these proxies relate to life history characteristics (Labocha et al., 2014; Wilder et al., 2016). A review
63 conducted by Barnett (2015) shows that, while mass or body size measures of energetic condition are often
64 assumed to have a positive linear relationship with fitness, this is not always the case, and the relationship
65 should first be empirically validated before being used as a proxy (Barnett et al., 2015). In some instances, the
66 proxy might relate to life history characteristics, but in a non-linear way. For example, theoretical simulations
67 of small birds show that survival does not increase linearly with energy (i.e., fat) reserves (McNamara et
68 al., 2005). If the reserves are too low, the individual is at risk of starvation. However, once the reserves get
69 too high, the individual is at an increased risk of predation (McNamara et al., 2005). Thus, fat reserves can
70 relate to a life history variable (survival), but in a U-shaped relationship rather than a linear one.

71 Although some studies use multiple morphological proxies for energetic condition (e.g., Warnock & Bishop,
72 1998), rarely are these variables directly compared. Multiple proxies should correlate with each other if
73 they measure the same trait (energetic condition). Furthermore, there is still confusion about what trait
74 some proxies actually measure. For example, a study conducted on two species of crickets showed that
75 three estimates of energetic condition based on fat content or on the relationship between body mass and
76 body length (scaled mass index or ordinary least squares regression) did not correlate with each other (Kelly
77 et al., 2014). A similar lack of a relationship was found in flying animals such as birds (A. G. Gosler
78 et al., 1995; Jacobs et al., 2012) and bats (McGuire et al., 2018) thus indicating cross-taxa support that
79 morphological proxy measures do not always measure the same trait. This is an example of the jingle fallacy
80 (Block, 1995; Carter et al., 2013), where a single trait label (“energetic condition”) actually encompasses
81 more than one distinct trait. In this case, two investigations using different proxies can be conducted on the
82 same research question, using the same species, but may end up with different results. This is problematic
83 because inconsistency in results among researchers can result in potentially misleading interpretations of the
84 impact of variation in morphology in relation to life history and population variables (Stevenson & Woods
85 Jr, 2006).

86 Here we compare two indices (fat score and the scaled mass index) of an individual’s energetic state to
87 validate whether they correlate with each other, which would indicate that they both measure energetic
88 condition. Fat score, as described by Kaiser (1993), is a numerical estimate of the amount of fat visible
89 under the skin (Fig. 1). The score ranges from 0 to 8 depending on the size and appearance of the fat
90 located in the individual’s abdomen and interclavicular depression, with 0 indicating no visible fat and 8
91 indicating extensive fat covering the ventral surface such that no muscle tissue is visible. For example, a
92 score of 1 corresponds to sparse traces of fat visible in the interclavicular depression and abdomen. This
93 measure is frequently used in birds (Cornelius Ruhs et al., 2019; Erciyas et al., 2010; Merilä & Svensson,
94 1997), and is a straightforward, non-invasive method for estimating energetic condition. However, previous
95 research found that it does not always positively relate with life history variables. For example, Haas (1998)
96 found no difference between fat scores in individuals that had successful or failed nests in American robins
97 and brown thrashers, indicating that fat score may not explain much of the variation in nest success in
98 some species. Further research is needed to understand the relationship between fat score measures and life
99 history characteristics.

100 In contrast, the scaled mass index (SMI) is more difficult to calculate than the fat score, but it has become
101 the predominant ratio method for quantifying energetic condition within and among populations (Delciellos
102 et al., 2018; English et al., 2018; Maceda-Veiga et al., 2014). The SMI is an individual’s mass scaled
103 by skeletal body size (Peig & Green, 2009). Unlike the common alternative which uses a simple ratio of
104 tarsus (lower leg) length to body mass, the SMI accounts for the tendency towards allometric scaling where
105 the relationship between body mass and structural size increases by a power law (Huxley, 1932). When
106 individuals with different structural body sizes can be standardized to the population average structural
107 body size, then energetic condition (the amount of mass not explained by structural body size) can be more
108 directly compared within and across populations. That is, the SMI calculates the energetic condition as
109 the mass of an individual relative to the population by first computing the mass that the individual would
110 have at the population average of a specific body measurement (e.g., tarsus length). Next, structural body
111 size of the individual is standardized by scaling the individual’s structural body length by the population
112 average of that body measurement, which accounts for population differences. The SMI is calculated as:

113 $Mass_i \left[\frac{AvgLength_p}{Length_i} \right]^{slope_p}$ where $Mass_i$ is each individual’s weight in grams, $Length_i$ is the value of the
114 chosen measure of structural body length for each bird, $AvgLength_p$ is the average structural body length in
115 the population, and $slope_p$ is calculated from the standard major axis regression (which is used to compare
116 variables that were both directly measured and thus have residual error) of a structural body size measure,
117 like tarsus length on mass (Peig & Green, 2009), and is interpreted as the expected change in structural
118 length for a one unit increase in mass. Therefore, individuals with superior energetic condition (larger weight
119 for their structural body size) will have a higher SMI compared to individuals that with inferior energetic
120 condition. Studies across taxa found that the SMI relates positively to reproductive success and survival. For
121 example, mallards with a lower SMI had lower rates of survival compared to their higher SMI counterparts
122 (Champagnon et al., 2012), while in crimson finches SMI was positively related to the number of young that
123 survived to independence (Milenkaya et al., 2015).

124 Our research will determine whether these two indices of energetic condition measure the same trait, and
125 whether this trait relates to an important life history characteristic: reproductive success. Measuring repro-
126 ductive success in birds involves finding and monitoring nests (Mayfield, 1961). However, nests are usually
127 built in cryptic locations and parents behave secretly (Gill, 1995), thus making it difficult to quantify the
128 number of eggs and nestlings inside the nest over time. Additionally, it is difficult and time-consuming to
129 track the survival of offspring once they leave the nest. Therefore, we will use the predominant method in
130 this field for quantifying reproductive success: whether a nest fledged offspring (Mayfield, 1961).

131 Our study system is a population of great-tailed grackles (*Quiscalus mexicanus*), hereafter “grackles,” in
132 Tempe, Arizona. This system is ideal for this investigation because grackles are native to the tropical
133 climates of Central America (Johnson & Peer, 2001), but have rapidly expanded their geographic range into
134 new areas (Wehtje, 2003). Because grackles are a water-associated species, the desert habitat of Tempe
135 presents physiological challenges that could lead to an increased likelihood of a tradeoff between survival
136 and reproductive attempts (Henderson et al., 2017). Deserts are characterized by a scarcity of water and
137 extreme temperature fluctuations, which require behavioral and physiological adaptations (Costa, 2012).

138 Wide variation in energetic condition and reproductive success is possible if grackle physiology requires more
139 water than is present in the environment, and some individuals may cope with physiological stress, or find
140 hidden sources of water, better than others (Henderson et al., 2017).

141 **Hypotheses** We measured two morphological proxy variables of energetic condition and observed repro-
142 ductive success in grackles to test two hypotheses. The first examined the relationship between two morpho-
143 logical proxies of energetic condition to validate that they measure the same inherent trait. Secondly, we
144 hypothesized that energetic condition, as measured by either or both of the morphological proxy variables,
145 would relate to reproductive success in male and female grackles.

146 METHODS

147 The methods below are based on the preregistration, with small changes summarized in the Deviations from
148 the planned methods section and further explained in the preregistration (in italics).

149 **Preregistration details** The preregistration used secondary data that were collected as part of other
150 ongoing investigations (tarsus length in http://corinalogan.com/Preregistrations/g_flexgenes.html; tarsus
151 length, body weight, number of fledglings, and whether a male holds a territory in http://corinalogan.com/Preregistrations/g_withinpop.html; and tarsus length in http://corinalogan.com/Preregistrations/g_expansion.html). The preregistration, containing the hypotheses, methods, and analysis plan, was written
152 (July 2019) and submitted to Peer Community In Ecology for pre-study peer review (August 2019) before any
153 analyses were conducted. We revised according to reviewer comments and received in principle acceptance
154 by PCI Ecology of the version on 8 Nov 2019. After that, we conducted the analyses in the preregistration.
155 Our final methods, results, and discussion, including all data and code, are listed below.
156
157



158 Summary of methods

159 Figure 1: A male grackle showing the yellow/orange tint of fat under the skin in the intraclavicular depression
160 (left); and a female grackle showing *no* fat under the skin of the intraclavicular region, but significant fat
161 deposits under the skin of the abdomen (right).

162 Great-tailed grackles are caught year-round in the wild in Tempe, Arizona using a variety of methods
163 (e.g., walk-in trap, bownet, mist net). After capture we immediately processed birds by attaching colored
164 leg bands in unique combinations for individual identification, conducted morphological measurements of
165 weight, tarsus length, flattened wing length, tail length, skull length, bill length and fat score (Fig. 1, the
166 amount of visible fat under the skin in the clavicle and abdomen as in Kaiser, 1993). Most grackles were
167 released after completion of color band marking, measurements, and acquiring a blood sample. A subset
168 of grackles were held in aviaries for up to 6 months for behavioral testing, and then released back to the
169 wild at their location of capture. We tested the repeatability of our structural size measures on this subset

170 of individuals by measuring them again before release. The second measures were collected by the same
171 experimenter in 11 out of 17 females and 10 out of the 18 males that were repeatedly sampled.

172 From March - August, we monitor the behavior of all color-marked grackles to determine their nesting status.
173 We follow females carrying nesting materials to find their nest. We determine whether the male territory
174 owner is color-marked as well. Then we check each nest approximately every day to determine the status
175 based on the female's behavior (building, incubation, feeding nestlings, feeding fledglings, failed).

176 Individuals included in our sample were those for which we have measures of energetic condition when they
177 were adults. We did not include individuals whose data were collected as juveniles. We also excluded data
178 that was collected from the grackles when they were released from the aviaries to avoid any confounds due
179 to their time in the aviary (e.g., perhaps unlimited nutritious food in the aviaries affected their fat score).
180 However, to validate that our measures of structural body size (tarsus length or wing length) are precise
181 and accurate, we measured twice the subset of grackles brought into aviaries - once when they were initially
182 caught, and again up to 6 months later when we released them. We calculated the repeatability of these
183 multiple measures. All other data included in this study came from wild-caught grackles (including the data
184 from the birds that were brought into the aviaries on their first capture).

185 We first used logistic mixed-effect models to determine whether SMI and fat score are correlated. We also
186 tested whether SMI and fat score varied by season because grackles are difficult to catch such that we were
187 unable to structure our data collection to coincide with the breeding season and instead caught and measured
188 grackles as often as possible. Previous research found a non-linear relationship between reproductive success
189 and energetic condition variables (Milenkaya et al., 2015). To check whether this is occurring in our data,
190 we visually examined our raw data to determine if we need to include a non-linear energetic condition
191 independent variable into our models (i.e. FatScore²). Then we used we used two types of logistic mixed-
192 effect models to determine the relationship between energetic condition and reproductive success. Both types
193 are supported in the literature, but are slightly different in the way in which the link function is specified.
194 First, we modeled the effect of energetic condition on reproductive success using a generalized linear mixed
195 model framework with a logit link function (i.e. Milenkaya et al., 2015). We then also used a logistic exposure
196 model that has a link function which accounts for the time interval between nest checks when estimating
197 the probability of daily nest survival (Bolker, 2014; Shaffer, 2004).

198 **After pre-study peer review: Deviations from the planned methods**

199 1) We realized that the sexual dimorphism of male and female body sizes necessitates separate analyses.
200 Therefore, we calculated SMI for males and females separately, and ran separate models for each sex
201 for the repeatability analysis (P1 and P2).

202 2) Fat score data were distributed such that the majority of scores were 0, with some 1's and very few
203 higher numbers. Specifically, of the 21 males, 15 had fat scores at 0, 5 scored 1, and a single male had
204 a fat score of 2. Out of 47 females, 26 scored 0, 18 scored 1, 2 scored 2, and a single female scored 3.
205 This lack of variance in the response variable led to problems when we ran the models: it was difficult
206 to fit models using an ordinal regression. The function "simulateResiduals," which we used to check
207 our data, does not work with data in the ordinal family. Consequently, we modified the model to use
208 a logistic regression where the dependent variable FatScore is categorized as individuals that showed
209 no visible fat ($y = 0$), or some fat was present ($y = 1$) where we combined all individuals that had fat
210 score values of 1 or greater. Subsequent data checking indicated that these data were not zero-inflated
211 or overdispersed.

212 **Deviations when testing hypothesis 1: correlation between SMI and Fat score**

213 3) Warning messages occurred during the repeatability analysis using the "rptR" package in R (Stoffel et
214 al., 2017) indicating that the fit was singular, likely because the variance for the Experimenter random
215 effect in the model for both female and male wing length was 0.001. We thus conducted an unregistered

216 analysis where we confirmed that our repeatability values from the repeatability models were valid,
217 despite the warning, by hand calculating repeatability following Nakagawa & Schielzeth (2010). The
218 hand-calculated repeatabilities were nearly identical (female $R = 0.5$, male $R = 0.71$) to the output
219 from the rpt function.

220 4) Despite the data checking which indicated our model was not overdispersed or zero inflated, we could
221 not get the fixed effects or random effect to converge using the Bayesian package in R “MCMCglmm.”
222 We found no improvement in model fit by tweaking the priors or iterations/burnin/thin options. There-
223 fore, we fit these models using the function glmer, a frequentist framework.

224 5) The Season variable only includes 2 males in the breeding season category, thus we do not have a
225 large enough sample to produce reliable estimates. We removed the Season variable from the model
226 for males.

227 **Deviations when testing hypothesis 2: energetic condition and reproductive success**

228 6) Only two females had reproductive success data from more than one year in our study (2019 and 2020).
229 Consequently, there were very few repeated measures in this sample and our random effect of bird ID
230 accounted for zero variance. This led to a warning that our model fit was singular. Therefore, we
231 removed the data for these females for 2020 so we could remove ID as a random effect from the model,
232 which resulted in the model running without warnings. We removed the 2020 data for these females
233 because their energetic condition data was collected in 2019 and these measures were more likely to
234 relate to their 2019 reproductive success data than to their reproductive success in 2020.

235 7) The fit of the model analyzing the relationship between energetic condition and male reproductive
236 success (ability to hold a territory containing female nests) was singular. The Year random effect
237 accounted for zero variance in the data, so we removed it. The fit was still singular, but we retained
238 the ID random effect (although it also explained zero variance) to account for repeated measures in
239 this sample.

240 8) The model fit was again singular in our logistic exposure model because the Year random effect ex-
241 plained zero variance in the data. We removed this random effect from the analysis.

242 **RESULTS**

243 **Prediction 1: correlation between SMI and Fat Score**

244 We were able to calculate SMI for 24 males and 62 females, and fat score values were available for 22 males
245 and 47 females (Table 1).

246 We found that wing length was more tightly correlated with body mass than tarsus length in both sexes,
247 therefore we used wing length in our SMI calculations (female $n = 62$, $r = 0.26$, $p = 0.03$; male $n = 24$, $r =$
248 0.35 , $p = 0.08$). This allows us to account for as much variation in body mass as possible that is associated
249 with skeletal body size because leftover variation in body mass is more likely to relate to energetic condition.
250 Consequently, we used wing length in our calculation of SMI.

251 To validate that we were measuring structural body size consistently across experimenters, we analyzed the
252 repeatability of wing length in the birds in our sample that were measured more than once. We found that
253 average wing length was repeatable ($n = 17$ females, Repeatability \pm standard error = 0.53 ± 0.18 ; $n = 18$
254 males, Repeatability \pm SE = 0.75 ± 0.11). Data permutations and a likelihood ratio test both confirmed
255 that these repeatability values were statistically significant at $p < 0.01$.

256 In females, we found that for every one unit increase in SMI, the bird is 1.3 times more likely to have some
257 fat (a 30% increase in the odds of having fat), which is not a statistically significant relationship (female p
258 = 0.81; Table 2). In males, a one unit increase in SMI corresponds to an odds ratio of 1.6, or a 60% increase

Table 1: Table 1. Sample sizes for P1 and P2. The *Breeding* and *Non-breeding season* categories refer to the number of individuals measured in each season. The *Reprod. success* category represents the total number of individuals in each year observed engaging in breeding behaviors. Note that the 2019 and 2020 reproductive success sample sizes include some of the same individuals that were observed in both years. Whereas, the *Prop. successful* category represents the proportion of the total individuals observed engaging in breeding behaviors in each year that held a territory containing nests (males) or fledged young (females).

Category	Males	Females
Breeding Season Fat	2	12
Non-breeding fat	20	35
Breeding season SMI	6	24
Non-breeding SMI	18	38
Aviaries	16	9
Repro. success 2019	8	9
Repro. success 2020	17	13
Prop. successful 2019	0.63	0.22
Prop. successful 2020	0.47	0.54

Table 2: Table 2. Results from the logistic mixed-effect regression for 47 females and fixed-effect regression for 21 males to determine whether fat score and scaled mass index (SMI) are correlated. Estimates are presented with the standard error in parentheses. Our sample size was too small to test for a season effect in males.


Parameter	Females		Males	
	Estimate (SE)	p-value	Estimate (SE)	p-value
Intercept	-0.20 (0.74)	0.79	-0.82 (0.64)	0.21
SMI	0.07 (0.30)	0.81	0.46 (0.62)	0.46
Season	0.27 (0.71)	0.70	NA	NA

259 in the odds of having some fat, which is also not a statistically significant relationship ($p = 0.50$; Table 2).
 260 Together, this indicates that SMI and fat score are not equally measuring energetic condition. There was
 261 also no relationship between season (breeding or non-breeding) and female fat score ($p = 0.71$). Only 2 males
 262 were measured during the breeding season, therefore we omitted season as an independent variable in the
 263 male model (Table 1).

264 **Prediction 2: energetic condition and reproductive success**


265 Our sample size for P2, where individuals had measures of reproductive success, SMI, and fat scores, was 20
 266 for females and 20 for males.

267 To determine whether we should include any non-linear effects of SMI in our models (A. G. Gosler et al.,
 268 1995; Milenkaya et al., 2015), we visually evaluated whether individuals in any of 5 categories, ranging from
 269 low to high SMI, were more likely to be reproductively successful (Fig. 2). We found no visual evidence for
 270 a non-linear relationship between reproductive success and SMI for males or females (Fig. 3). Consequently,
 271 we did not include non-linear terms in subsequent models.



272

273 Figure 2: Frequency histogram of the SMI scores, illustrating the SMI categories, for the 33 males and 31
 274 females for which we also had reproductive success data. The mean SMI value is indicated by a red vertical
 275 line. We created SMI category bins (indicated with vertical blue lines) in 1 standard deviation increments,
 276 centered on the mean. Category 3 indicates the SMI value is close to the population mean value. Categories
 277 1 and 2 are individuals that have SMI scores that are low, and moderately low, respectively, compared to
 278 the population mean value. Similarly, categories 4 and 5 contain individuals that have SMI scores that are
 279 moderately high and high, respectively, compared to the population mean value.




280

281 Figure 3: The proportion of individuals that successfully fledged nests (females: left) or held a territory
 282 (males: right) in low (1), moderately low (2), moderate (3), moderately high (4) and high (5) scaled mass

283 index (SMI) categories. Dots are sized according to the number (n) of individuals in that category. There is
284 no evidence of a non-linear relationship.

285 We used linear models to determine whether season would be important to include in our models testing
286 whether body condition relates to reproductive success. We found that SMI did not differ by season for
287 females (Estimate (SE): $\beta = -0.30$ (0.26), $p = 0.26$) or males ($\beta = -0.65$ (0.43), $p = 0.15$). Similarly, fat
288 score for females ($\beta = 0.28$ (0.68), $p = 0.68$) and males ($\beta = 17.08$ (2797.4), $p = 0.99$) did not differ by
289 season (Fig. 4). Although we note that, as stated above and indicated in the standard error value, we lack
290 sufficient fat score data from males in the breeding season so these results should be interpreted with caution.
291 Consequently, we did not include season as an independent variable in our subsequent models testing the
292 relationship between our body condition proxies and reproductive success.



293

294 Figure 4: Scaled mass index (SMI) was not significantly different between the breeding and non-breeding
295 seasons for either sex.

296 Because fat score and SMI did not correlate, we included both as independent variables in our models testing
297 prediction 2. For both males and females, we found no statistically significant relationships between either
298 proxy of energetic condition and reproductive success (Table 3). Of note, the inconsistent direction of the
299 effects for the parameter estimates further supports that SMI and fat score do not measure the same trait.

300 For females, our SMI parameter estimate of -0.92 (exponentiated to get the log odds = 0.40) indicates that
301 a one unit increase in SMI corresponded to a 60% decrease in the odds a female would fledge an offspring
302 ($p = 0.13$). Whereas an increase from no visible fat to showing some fat corresponded to a 16% increase
303 in the odds a female would fledge an offspring (log odds = 1.16, $p = 0.82$). There was also no evidence of
304 a significant relationship between the ability of a female to produce fledglings and having previously spent
305 time in the aviaries (log odds = 0.25, $p = 0.22$), where the odds that a female would fledge an offspring were
306 75% lower if females spent time in the aviaries.

307 For males, there was also no statistically significant support for a relationship between whether a male
308 defended a territory and SMI (log odds = 3.25, $p = 0.13$). Nevertheless, this relationship may be biologically
309 important because a one unit increase in SMI corresponded to a more than 300% increase in the odds a
310 male will hold a territory containing nests. Fat score was also statistically unrelated to male reproductive
311 success where an increase from showing no visible fat to showing some fat corresponded to a 28% decrease
312 in territory holding (log odds = 0.72, $p = 0.76$). Lastly, we found that those males who spent time in the

Table 3: Table 3. Results from the logistic regression for 20 females and 20 males to test whether reproductive success relates to condition. Estimates are presented with the standard error in parentheses.

Parameter	Females		Males	
	Estimate (SE)	p-value	Estimate (SE)	p-value
Intercept	-0.02 (0.73)	0.98	3.05 (1.40)	0.03
FatScore	0.15 (1.02)	0.89	-0.33 (1.10)	0.77
SMI	-0.92 (0.61)	0.13	1.18 (0.78)	0.13
Aviary	-1.38 (1.14)	0.23	-3.62 (1.56)	0.02*

Table 4: Table 4. Results of the logistic exposure model showing the relationship between the probability of daily nest survival and scaled mass index (SMI), fat score, the amount of time spent in the aviaries, and the day of the year. Parameter estimates are presented with the standard error in parentheses. Odds ratios (OR) represent the exponentiated estimates, and are presented to increase interpretability with 95 percent confidence intervals in parentheses.


Parameter	Estimate (SE)	OR (CI)	p-value
Intercept	1.99 (0.40)	7.32 (3.3-16.0)	<0.001
Fat score	0.91 (0.49)	0.50 (0.27-0.92)	0.06
SMI	-0.69 (0.31)	2.48 (0.95-6.49)	0.03*
Day of year	-0.21 (0.15)	0.63 (0.19-2.10)	0.16
Aviary	-0.47 (0.61)	0.81 (0.60-1.10)	0.44

313 aviaries were statistically less likely (97% decrease in the odds) to hold a territory compared with males who
 314 were never in the aviaries (log odds = 0.03, $p = 0.02$). However, we stress that our sample size was relatively
 315 small (20 males), and we did not have a balanced sample because there were no males that did not defend a
 316 territory and were never in the aviaries. Additionally, only five males had data from more than one breeding
 317 season, which resulted in our model fit being singular because the random effect for bird ID accounted for
 318 essentially zero variance. However, we kept ID in the model to account for the repeated samples.

319 **Prediction 2: energetic condition and probability of daily nest survival**

320 Logistic regression analyses to determine reproductive success from nests discovered in different stages will
 321 be systematically biased (Shaffer, 2004). Nests discovered at a more progressed stage (i.e., nestling stage
 322 compared to building stage) are statistically more likely to succeed and nests with frequent and prolonged
 323 adult visits (such as those that occur when nests survive longer) are more likely to be discovered. Therefore,
 324 nests that fail early are less likely to be detected (Shaffer, 2004). Consequently, we analyzed female repro-
 325 ductive success using a logistic exposure model (Bolker, 2014), which uses survival analysis to determine the
 326 factors affecting the probability of daily nest survival, while accounting for incomplete nest observations.

327 We found that the probability of daily nest survival was significantly negatively related to SMI (log odds
 328 = 0.50, $p = 0.03$; Table 4), where, for every unit increase in SMI, the odds of daily nest survival decreased
 329 by half. This indicates that a female with a larger SMI (more mass for her structural body size) was less
 330 likely to have her nest survive each day (Fig. 5). There was no statistically significant relationship between
 331 the probability of daily nest survival and fat score (log odds = 2.48, $p = 0.06$), day of the year (log odds =
 332 0.81, $p = 0.16$), or time spent in the aviaries (log odds = 0.63, $p = 0.44$, Table 4). Although not statistically
 333 significant, the effect size for the relationship between fat score and daily nest survival is large (Fig. 5) and
 334 potentially biologically meaningful. The odds of a nest surviving on a given day are almost 2.5 times greater
 335 (248%) for birds with some fat (a score of 1) compared to no fat (a score of 0).



336

337 Figure 5: Odds ratios for independent variables affecting the probability of a nest surviving a given day. The
 338 dots and corresponding values represent the odds ratio values, and lines represent the confidence intervals
 339 around the odds ratio value. The vertical line at $x = 1$ delineates the odds ratio value for no relationship
 340 between the estimates and the probability of daily nest survival. The asterisk indicates an odds ratio value
 341 that is statistically significant.

342 DISCUSSION

343 Energetic condition is not directly observable, but variation can affect life history characteristics (Barnett
 344 et al., 2015; Labocha et al., 2014). Consequently, a large corpus of research attempts to measure energetic
 345 condition using various proxy measures (Labocha et al., 2014) and largely assumes that the chosen proxy
 346 accurately reflects energetic condition as a singular trait. Although it is often implicitly assumed that all
 347 proxy measures for energetic condition reflect the same inherent trait, it is rare for one study to compare
 348 multiple proxies. However, if all proxy measures are affected similarly by a singular energetic condition
 349 phenotype, then multiple proxy measures should produce correlated results. The aim of the current study
 350 was therefore to test the idea that multiple commonly used morphological proxies equally measure energetic
 351 condition (by correlating with each other), and that these measures can explain variation in reproductive
 352 success.

353 Here we found that two morphological proxies of energetic condition, fat score and SMI, did not correlate
 354 with each other in the great-tailed grackle, regardless of whether it was the breeding or non-breeding season.
 355 While both proxies are well supported in previous research as measures of energetic condition, our results
 356 indicate that they may not be measuring the same trait. This has also been found in studies on bats
 357 (McGuire et al., 2018), which are species that similarly experience distinct demands on body structure to
 358 facilitate flight. There are several potential reasons why grackle fat score and SMI did not correlate. First,

359 it is possible that we were unable to accurately measure the amount of fat the birds actually stored. In
360 addition to storing fat under their skin, birds may also store fat intraperitoneally (Musacchia, 1953), which
361 would not have been detected with our fat score measure. Second, SMI and fat score may measure different
362 components of energetic condition because variation in mass among grackles could be attributable to muscle
363 or body water content, whereas fat score only accounts for subcutaneous fat (Labocha & Hayes, 2012).
364 Research shows that stored fat is the primary source of energy in many taxa (Walsberg, 1988), especially in
365 birds (Blem, 1990; Pond, 1981) because the energy per ounce from fat is much higher than from proteins or
366 carbohydrates (Gessaman, 1999). However, because desert birds, such as the grackles in our investigation,
367 have inconsistent access to water sources, variation in body water content may obscure variation in lipid
368 content. Measuring muscle content often requires destructive methods [i.e. sacrificing the birds; Zhang et
369 al. (2015)] or less objective assessments such as keel prominence or breast muscle shape (Abolins-Abols &
370 Ketterson, 2017; A. Gosler, 1991), which was beyond the scope of the current research program. Third,
371 it is possible that fat score and SMI did not correlate due to temporal variation at a fine scale that we
372 were unable to capture. Although we found no evidence that SMI or fat score varied by season, there is
373 evidence from other studies that avian mass changes with time of day (Nip et al., 2019) and stage of breeding
374 (Milenkaya et al., 2013). It was logistically impossible in our project (and in many avian research programs)
375 to capture birds multiple times within a season or at several times per day, therefore temporal variation in
376 data collection could obscure the correlation between these two proxies, if such a correlation exists. However,
377 the stage of breeding is unlikely to introduce additional variance to our study because we did not catch any
378 females that were actively engaged in any stage of the breeding process. Finally, our sample sizes might have
379 been too small to detect an effect, but the effect size for the relationship between fat score and SMI was
380 essentially zero (0.001), therefore it is unlikely that a larger sample size would find a biologically informative
381 relationship between these two proxies.

382 Energetic condition can have a large impact on reproductive success in birds (Drent & Daan, 1980; Montreuil-
383 Spencer, 2017) and in flying mammals (Welbergen, 2011). For example, female chickadees with higher
384 winter fat scores are more likely to lay eggs earlier in the subsequent breeding season, as well as go on
385 to feed those offspring more frequently (Montreuil-Spencer, 2017). Energetic condition is likely a factor in
386 reproductive success in our system because previous research in great-tailed grackles found that larger and
387 heavier males were more likely to hold territories, have more social mates, and sire more offspring (Johnson
388 et al., 2000). Our study additionally considered female morphology and reproductive success, subcutaneous
389 fat, and controlled for the impact of structural body size on mass. However, we found reproductive success,
390 measured as the ability to produce fledglings (females) or to hold a territory containing nests (males), did
391 not significantly correlate with fat score or SMI. Although our results were not statistically significant, in
392 some cases the parameter estimates revealed log-odds that may be large enough to be biologically significant.
393 Notably, a one unit increase in SMI corresponded to a more than 300% increase in the odds a male will hold
394 a territory containing nests, but a 60% decrease in the odds a female would fledge an offspring.

395 We additionally used logistic exposure models to determine whether the energetic condition of females
396 related to the probability of daily nest survival. We only included females in this analysis because males
397 were never observed contributing to nest building, incubation, or feeding nestlings in our population and so
398 will not have a direct effect on daily nest survival. We found a negative relationship between female SMI
399 and the likelihood of daily nest survival. This could be due to larger females actually carrying proportionally
400 smaller energetic reserves than their smaller female counterparts (Jacobs et al., 2012), as seen in red-winged
401 blackbirds (Langston et al., 1990). In some species, females with smaller body sizes are able to initiate
402 breeding earlier because they can allocate more resources to reproduction compared to larger individuals
403 that have higher bodily energy demands and therefore fewer excess energetic resources (Barbraud et al.,
404 2000; Langston et al., 1990; Murphy, 1986). This indirectly affects reproductive success because nesting
405 earlier increases the probability of nesting success and multiple nesting attempts (Johnson & Peer, 2001;
406 Perrins, 1970). Yet, in our study we found no relationship between the probability of daily nest survival and
407 day of the year, therefore this is unlikely to explain the negative relationship between SMI and nest survival.
408 Alternatively, it is possible that larger females are unable to build a more concealed nest in the most dense
409 vegetation, or that larger females are unable to build nests in delicate vegetation structure that is more likely
410 to be inaccessible to predators. Moreover, the parameter estimate for the relationship between fat score and
411 the daily probability of nest survival indicates that females with some visible fat are more than twice as

412 likely to have a nest survive a given day. Because the direction of this effect is opposite to the relationship
413 between SMI and nest survival, this is further evidence that these two proxies represent different traits.

414 Great-tailed grackles are an interesting system to study energetic condition and reproductive success because
415 they recently expanded their range into Arizona, where the climate and habitat are distinct from that in
416 Central America where the species originally evolved (Wehtje, 2003). The increase in temperature variation
417 and decrease in available water at our desert study site are both environmental stressors that have previously
418 been found to negatively affect energetic condition (Pendlebury et al., 2004). Although our study spanned
419 only two years, our data are likely representative of reproductive success in this environment because the
420 temperatures during our study were in line with those from the previous three years (National Climatic
421 Data Center, 2020). Reproductive success is vital to species persistence and abundance in novel environ-
422 ments (Maspons et al., 2019). Therefore, an understanding of energetic condition and its relationship with
423 reproductive success in grackles outside of their original range could broadly inform conservation research in
424 invasive and non-native species. While reproductive success of certain avian species may be easier to monitor
425 at a more fine scale (i.e. cavity nesters), the predominant measure of reproductive success currently used
426 by avian ecologists is the ability of adults to fledge offspring (since foundational work by Mayfield, 1961)
427 because it is financially and logistically accessible to more researchers. Therefore, we believe our measure of
428 reproductive success in grackles is informative, and that research that spans taxa with diverse reproductive
429 strategies is important for understanding general trends in energetic condition and the appropriate proxies.

430 The results of this study highlight the need to better understand proxy measures of energetic condition,
431 not only in grackles, but for birds in general. Most studies on avian energetic condition only use one proxy
432 variable, but because energetic condition is difficult to measure directly, it is important to compare mul-
433 tiple proxy variables to determine whether the proxy is measuring the intended trait (Block, 1995; Carter
434 et al., 2013). If financially and logistically feasible, future research could measure total body composition
435 and relative mass of fat using the relatively new and promising method of quantitative magnetic resonance
436 (Guglielmo et al., 2011), or researchers could incorporate additional physiological methods to measure en-
437 ergetic condition, for example, blood hematocrit levels (Dawson & Bortolotti, 1997; but see Fair et al.,
438 2007). Additionally, studying traits that could relate to variation in energy stores, such as dispersal (Ellers
439 et al., 1998), migratory endurance (Deppe et al., 2015), or survival (Liao et al., 2011) would allow us to
440 disentangle whether morphological proxies like fat score and SMI are poor proxy measures for energetic
441 condition, or whether fat score and SMI do not affect reproductive success but may be associated with other
442 life history characteristics. Because SMI can perform poorly in birds with low lipid mass, future research
443 should also compare several mass by structural body size equations to determine the most appropriate proxy
444 for a specific study system (Jacobs et al., 2012). Lastly, future research would benefit from using logistic
445 exposure models to examine the relationship between energetic condition and reproductive success because
446 these models control for the bias that arises when early nest failures are not detected, which is not possible
447 in logistic regression models, and it is more sensitive to changes in a bird's nest status (Shaffer, 2004).

DETAILED HYPOTHESES AND METHODS FROM THE PRE-REGISTRATION

HYPOTHESES

We measured two morphological proxy variables of energetic condition and observed reproductive success in grackles to test two hypotheses:

H1 - There is a relationship between two different morphological indices of energetic condition: fat score and the scaled mass index.

Prediction 1: Fat score and the scaled mass index will be positively correlated. This would indicate that these two indices measure the same trait, and it is likely they both are proxies for fat content.

Prediction 1 alternative 1: There is a negative correlation between fat score and the scaled mass index. This would indicate that there may be a tradeoff between the two indices where a larger value of the scaled mass index may measure muscle content rather than fat, and individuals with more muscle have less visible fat.

Prediction 1 alternative 2: There is no correlation between fat score and the scaled mass index. This indicates that these two variables do not measure the same trait. Fat score may not adequately capture a bird's energetic condition because birds may be selected to only store the minimal fat necessary to prevent starvation, while also minimizing the weight gain that would make them easier targets for predators (Barnett et al., 2015). Similarly, the scaled mass index could be heavily influenced by body size, therefore reflecting structural size rather than fat storage (Labocha & Hayes, 2012).

H2 - Energetic condition (as measured by fat score and the scaled mass index) relates to reproductive success (measured as a binary variable of whether a female had one or more fledglings (1) or not (0), and whether a male defended a territory containing nests (1) or not (0)).

Prediction 2: Morphological indices of energetic condition (fat score and the scaled mass index) will correlate positively with reproductive success. This would indicate that individuals with more fat, and therefore higher energy reserves, are better able to acquire the resources necessary for reproduction.

Prediction 2 alternative 1: Morphological indices of energetic condition (fat score and the scaled mass index) will correlate negatively with reproductive success. This indicates that individuals may make trade offs, with some acquiring more food and increasing their energy reserves, and others prioritizing reproductive activities over increasing energy reserves.

Prediction 2 alternative 2: Morphological indices of energetic condition (fat score and the scaled mass index) do not correlate with reproductive success. This indicates that other, potentially non-morphological, individual characteristics relate to reproductive success (i.e., cognition, nest site selection, breeding experience, predator vigilance, etc.).

METHODS

The methods below are based on the preregistration, with small changes as described in the Deviations from the planned methods section above.

Planned Sample Great-tailed grackles are caught in the wild in Tempe, Arizona using a variety of methods (e.g., walk-in trap, bownet, mist net). After capture we immediately process birds by attaching colored leg bands in unique combinations for individual identification, conducting morphological measurements of weight, tarsus length, flattened wing length, tail length, skull length, bill length and fat score (the amount of visible fat under the skin in the clavicle and abdomen as in Kaiser, 1993). Most grackles are released after completion of color band marking, measurements, and acquiring a blood sample. A subset of grackles

491 are held in aviaries for up to 6 months for behavioral testing, and then released back to the wild at their
492 location of capture.

493 From March - August, we monitor the behavior of all color-marked grackles to determine their nesting status.
494 We follow females carrying nesting materials to find their nest. We determine whether the male territory
495 owner is color-marked as well. Then we check each nest approximately every day to determine the status
496 based on the female's behavior (building, incubation, feeding nestlings, feeding fledglings, failed).

497 Individuals included in this sample will be those for which we have measures of energetic condition when they
498 were adults. We will not include individuals whose data were collected as juveniles. As of 30 July 2019, we
499 have fledgling data for 14 females that exhibited breeding behavior (5 had 1+ fledgling, 9 had no fledglings)
500 and breeding territory status for 10 males (7 territory holders, 3 non-territory holders, 2 not observed so not
501 part of this sample). Therefore, the minimum sample size for H2 will be 24. The minimum sample size for
502 H1 will be 72, because that is how many marked individuals we have biometric data for so far. However,
503 we expect to be able to add to the sample size for both H1 and H2 before the end of this investigation in
504 Tempe, Arizona. *UPDATE Oct 2020: In the second breeding season we had 20 females and 20 males with*
505 *reproductive success and energetic condition data.*

506 **Sample size rationale** We will continue to color mark as many grackles as possible, and collect biometric
507 data and fat scores. Our current sample of reproductive success is small because the grackles in Tempe
508 nest in very tall palms, making it difficult to determine nest status. However, we plan to collect additional
509 reproductive success data during the breeding season in summer 2020. *UPDATE Oct 2020: In the second*
510 *breeding season we had 20 females and 20 males with reproductive success and energetic condition data.*

511 **Data collection stopping rule** We will stop collecting data for this project in early August 2020 when
512 research at the Tempe, Arizona field site will be finished.

513 **Open materials** Biometric measurement protocol: [https://gitlab.com/corinalogan/the-grackle-project/
514 blob/master/protocolBiometrics.pdf](https://gitlab.com/corinalogan/the-grackle-project/blob/master/protocolBiometrics.pdf)

515 Nest check protocol: [https://gitlab.com/corinalogan/the-grackle-project/blob/master/protocolNestCheck.
516 pdf](https://gitlab.com/corinalogan/the-grackle-project/blob/master/protocolNestCheck.pdf)

517 **Open data** All data (Berens et al., 2020) are available at [https://knb.ecoinformatics.org/view/doi:10.
518 5063/F1NZ862D](https://knb.ecoinformatics.org/view/doi:10.5063/F1NZ862D) and at github (the provided code will load these files directly from github).

519 **Randomization and counterbalancing** There is no randomization or counterbalancing in this investi-
520 gation.

521 **Blinding of conditions during analysis** No blinding is involved in this investigation.

522 **Dependent Variables**

523 **P1: correlation between fat and the scaled mass index**

524 1) Fat score [the amount of visible fat under the skin in the clavicle and abdomen reported as a score
525 from 0 (no fat) to 8 (fat completely covers muscles and underside of the bird); Kaiser (1993)] *UPDATE*
526 *Oct 2020: Fat score was heavily 0 skewed with few scores greater than one. To increase model fit we*
527 *used a binomial response variable instead, where 0 is no fat and 1 is some fat observed under the skin.*

528 **P2: energetic condition and reproductive success**

- 529 1) Female had one or more fledglings (yes, no)
530 2) Male held a territory consisting of 1 to 3 clumped palms containing nests (yes, no)

531 Independent Variables

532 P1: correlation between fat and the scaled mass index

- 533 1) Scaled mass index using measures of body weight and tarsus length or flattened wing length (average
534 of left and right as in Bleeker et al., 2005). We will choose the measure that is most correlated with
535 body weight (Peig & Green, 2009).
- 536 2) Season (non-breeding [Sep-Feb], breeding [Mar-Aug]). *UPDATE Oct 2020: The Season variable only*
537 *includes 2 males in the breeding season category, thus we do not have a large enough sample to produce*
538 *reliable estimates. We removed the Season variable from the model for males.*
- 539 3) Random effect: Experimenter (because several different experimenters measure dependent variables on
540 multiple different birds)

541 P2: energetic condition and reproductive success

- 542 1) Fat score
- 543 • Note 1: if the fat score and the scaled mass index are positively correlated, then we will use only
544 fat score in the model for P2. If they are not positively correlated, then we will add the scaled
545 mass index as an independent variable in the P2 analysis
 - 546 • Note 2: if fat score and/or the scaled mass index vary by season (breeding or non-breeding), then
547 we will only use the data from the breeding season to ensure that less time has elapsed between
548 the collection of energetic condition and reproductive success variables
- 549 2) Temporarily held in aviaries for behavioral testing at any point during this study, because this may
550 affect breeding behaviors (yes, no)
- 551 3) Random effect: Year (to determine whether conditions in a given breeding season similarly affected all
552 grackle behavior and nest success)
- 553 4) Random effect: Bird ID (because there may be multiple measures of reproductive success for each bird)

554 ANALYSIS PLAN

555 *UPDATE Oct 2020:*

- 556 1) *We realized that the sexual dimorphism of male and female body sizes necessitates separate analyses.*
557 *Therefore, we calculated SMI for males and females separately, ran separate models for each sex for the*
558 *repeatability analysis, P1 and P2.*
- 559 2) *Fat score data were distributed such that the majority of scores were 0, with some 1's and very few higher*
560 *numbers. This made it difficult to fit models using an ordinal regression. The function simulateResiduals,*
561 *which we used to check our data, does not work with data in the ordinal family. Consequently, we used*
562 *logistic regression where the dependent variable FatScore represents no fat (score = 0), or some fat (score =*
563 *1)*
- 564 3) *Despite the data checking which indicated our model was not overdispersed or zero inflated, we could not*
565 *get the fixed effects or random effect to converge using the Bayesian MCMCglmm. We found no improvement*
566 *in model fit by tweaking the priors or iterations/burnin/thin options. Therefore, we fit these models using*
567 *the function glmer, a frequentist framework.*

568 4) *The Season variable only includes 2 males in the breeding season category, thus we do not have a large*
569 *enough sample to produce reliable estimates. We removed the Season variable from the model for males.*

570 We will **exclude** data that was collected from the grackles when they were released from the aviaries to
571 avoid any confounds due to their time in the aviary (e.g., perhaps unlimited nutritious food in the aviaries
572 affected their fat score). However, to validate that our measures of structural body size (tarsus length or
573 wing length) are precise and accurate, we will measure twice a subset of grackles brought into aviaries -
574 once when they are initially caught, and again up to 6 months later when we release them. We will then
575 calculate the repeatability of these multiple measures. All other data included in this study will come only
576 from wild-caught grackles (including the birds that were brought into the aviaries on their first capture).
577 When **missing data** occur, the existing data for that individual will be included in the analyses for which
578 their data exist. Analyses will be conducted in R [current version 4.0.5; R Core Team (2017)].

579 **Ability to detect actual effects** To begin to understand what kinds of effect sizes we will be able to
580 detect given our sample size limitations, we used G*Power Faul et al. (2009) to conduct power analyses
581 based on confidence intervals. G*Power uses pre-set drop down menus and we chose the options that were
582 as close to our analysis methods as possible (listed in each analysis below). Note that there were no explicit
583 options for GLMMs, thus the power analyses are only an approximation of the kinds of effect sizes we can
584 detect. We realize that these power analyses are not fully aligned with our study design and that these
585 kinds of analyses are not appropriate for Bayesian statistics (e.g., our MCMCglmm below), however we are
586 unaware of better options at this time. Additionally, it is difficult to run power analyses because it is unclear
587 what kinds of effect sizes we should expect due to the lack of data on this species for these particular research
588 questions.

589 **Data checking** The data will be checked for overdispersion, underdispersion, zero-inflation, and het-
590 eroscedasticity with the DHARMA R package (Hartig, 2019) following methods by Hartig.

591 *P1 analysis: correlation between fat and the scaled mass index*

592 We will calculate the scaled mass index as described by Peig & Green (2009) using either tarsus or flattened
593 wing length - whichever measure is most correlated with body weight (Peig & Green, 2009).

594 We use a Generalized Linear Mixed Model (GLMM; MCMCglmm function, MCMCglmm package; (Hadfield
595 2010)) with an ordinal distribution (for categorical variables in MCMCglmm) and probit link using 130,000
596 iterations with a thinning interval of 10, a burnin of 30,000, and minimal priors (V=1, nu=0) (Hadfield,
597 2014). We will ensure the GLMM shows acceptable convergence [lag time autocorrelation values <0.01;
598 Hadfield (2010)], and adjust parameters if necessary to meet this criterion. We will determine whether an
599 independent variable had an effect or not using the Estimate in the full model.

600 Where we have multiple measures of tarsus or flattened wing length, we will check that our measurements
601 are repeatable using the rptR package (Stoffel et al., 2017).

602 To roughly estimate our ability to detect actual effects (because these power analyses are designed for
603 frequentist statistics, not Bayesian statistics), we ran a power analysis in G*Power with the following settings:
604 test family=F tests, statistical test=linear multiple regression: Fixed model (R² deviation from zero), type
605 of power analysis=a priori, alpha error probability=0.05. We changed the power and the effect size until we
606 reached an output that we project our sample size will be (n=90). The number of predictor variables was
607 restricted to only the fixed effects because this test was not designed for mixed models. The protocol of the
608 power analysis is here:

609 *Input:*

610 Effect size $f^2 = 0.15$

611 err prob = 0.05

612 Power (1- err prob) = 0.86

613 Number of predictors = 3

614 *Output:*

615 Noncentrality parameter = 13.3500000

616 Critical F = 2.7119214

617 Numerator df = 3

618 Denominator df = 85

619 Total sample size = 89

620 Actual power = 0.8635760

621 This means that, with a sample size of 89, we would have an 86% chance of detecting a medium effect
622 (approximated at $f^2=0.15$ by Cohen, 1988).

623 *code shown in .rmd*

624 *P2 analysis: energetic condition and reproductive success*

625 To model the effect of energetic condition on reproductive success, we will use two types of logistic mixed-
626 effect models. Both types are supported in the literature, but are slightly different in the way in which
627 the link function is specified. First, we will model reproductive success using a generalized linear mixed
628 model framework with a logit link function (i.e. Milenkaya et al., 2015). We will also use a logistic exposure
629 model that has a link function which accounts for the time interval between nest checks when estimating
630 the probability of daily nest survival (Bolker, 2014; Shaffer, 2004). If fat score and the scaled mass index
631 are positively correlated in P1, then we will use only fat score as the independent variable in this GLMM. If
632 they are not positively correlated, we will include both as independent variables.

633 Previous research found a non-linear relationship between reproductive success and energetic condition vari-
634 ables (Milenkaya et al., 2015). To check whether this is occurring in our data, we will first plot our raw
635 data to determine if we need to include a non-linear energetic condition independent variable into our model
636 (i.e. FatScore²). Our dependent variable is binary, so to more clearly see the trends in the data, on the x-axis
637 we will bin our energetic condition scores into 5 categories based on standard deviations (sd) around the
638 mean (low = < 2 sd, moderately low = -2 sd to -1 sd, moderate = -1 sd to +1 sd, moderately high = +1 sd
639 to +2 sd, high = > 2 sd). Then on the y-axis we will use the proportion of individuals in each category that
640 had successful nests. *UPDATE Oct 2020: Because most individuals fell within the medium category when*
641 *we grouped data using 1 standard deviation around the mean, we switched to using half standard deviation*
642 *increments around the mean.*

643 A power analysis was conducted as above for P1 and the protocol reported here:

644 *Input:*

645 Effect size $f^2 = 0.15$

646 err prob = 0.05

647 Power (1- err prob) = 0.90

648 Number of predictors = 2

649 *Output:*

650 Noncentrality parameter = 13.2000000

651 Critical F = 3.1038387

652 Numerator df = 2

653 Denominator df = 85

654 Total sample size = 88

655 Actual power = 0.9020264

656 This means that, with a sample size of 88, we would have a 90% chance of detecting a medium effect
657 (approximated at $f^2=0.15$ by Cohen, 1988).

658 *code shown in .rmd*

659 **Do energetic condition variables vary by season?** *code shown in .rmd*

660 **Does energetic condition relate to reproductive success?** *code shown in .rmd*

661 **Does female energetic condition relate to the probability of daily nest survival?** Our measure
662 of female nest success could be systematically biased against nests that failed early (Shaffer, 2004). Conse-
663 quently, we also analyzed female reproductive success using a logistic exposure model. This type of model
664 determines the factors affecting daily nest survival probability.

665 *code shown in .rmd*

666 ETHICS

667 This research is carried out in accordance with permits from the:

- 668 1) US Fish and Wildlife Service (scientific collecting permit number MB76700A-0,1,2)
- 669 2) US Geological Survey Bird Banding Laboratory (federal bird banding permit number 23872)
- 670 3) Arizona Game and Fish Department (scientific collecting license number SP594338 [2017], SP606267
671 [2018], and SP639866 [2019])
- 672 4) Institutional Animal Care and Use Committee at Arizona State University (protocol number 17-1594R)

673 AUTHOR CONTRIBUTIONS

674 **Berens:** Hypothesis development, data collection, revising/editing.

675 **Logan:** Study design, write up, revising/editing, materials/funding.

676 **Folsom:** Data collection, revising/editing.

677 **Sevchik** Data collection, revising/editing.

678 **Bergeron:** Data collection, revising/editing.

679 **McCune:** Hypothesis development, data collection, data analysis, write up, revising/editing.

680 FUNDING

681 This research is funded by the Department of Human Behavior, Ecology and Culture at the Max Planck
682 Institute for Evolutionary Anthropology.

ACKNOWLEDGEMENTS

We thank Aaron Blackwell and Ken Kosik for being the UCSB sponsors of the Cooperation Agreement with the Max Planck Institute for Evolutionary Anthropology; and our research assistants for help with trapping the grackles and collecting the biometric and nest/territory data: Aelin Mayer, Nancy Rodriguez, Brianna Thomas, Aldora Messinger, Elysia Mamola, Michael Guillen, Rita Barakat, Adriana Boderash, Olateju Ojekunle, August Sevchik, Justin Huynh, Amanda Overholt, and Michael Pickett.

REFERENCES

- Abolins-Abols, M., & Ketterson, E. D. (2017). Condition explains individual variation in mobbing behavior. *Ethology*, *123*(8), 495–502.
- Aubret, F., Bonnet, X., Shine, R., & Lourdaïs, O. (2002). Fat is sexy for females but not males: The influence of body reserves on reproduction in snakes (*vipera aspis*). *Hormones and Behavior*, *42*(2), 135–147.
- Barbraud, C., Lormée, H., & LeNevé, A. (2000). Body size and determinants of laying date variation in the snow petrel *pagodroma nivea*. *Journal of Avian Biology*, *31*(3), 295–302.
- Barnett, C. A., Suzuki, T. N., Sakaluk, S. K., & Thompson, C. F. (2015). Mass-based condition measures and their relationship with fitness: In what condition is condition? *Journal of Zoology*, *296*(1), 1–5.
- Barry, K. L., & Wilder, S. M. (2013). Macronutrient intake affects reproduction of a predatory insect. *Oikos*, *122*(7), 1058–1064.
- Berens, J., Logan, C., Folsom, M., Sevchik, A., Bergeron, L., & McCune, K. (2020). Validating morphological condition indices and their relationship with reproductive success in great-tailed grackles. *Knowledge Network for Biocomplexity, Data package*. <https://doi.org/10.5063/7P8WSM>
- Bleeker, M., Kingma, S. A., Szentirmai, I., Székely, T., & Komdeur, J. (2005). Body condition and clutch desertion in penduline tit *remiz pendulinus*. *Behaviour*, *142*, 1465–1478.
- Blem, C. (1990). Avian energy storage. *Curr Ornithol*, *7*, 59–113.
- Block, J. (1995). A contrarian view of the five-factor approach to personality description. *Psychological Bulletin*, *117*(2), 187.
- Bolker, B. (2014). Logistic regression, accounting for differences in exposure. *Version 09.30. 2014. R Pubs*.
- Carter, A. J., Feeney, W. E., Marshall, H. H., Cowlshaw, G., & Heinsohn, R. (2013). Animal personality: What are behavioural ecologists measuring? *Biological Reviews*, *88*(2), 465–475.
- Champagnon, J., Guillemain, M., Elmberg, J., Massez, G., Cavallo, F., & Gauthier-Clerc, M. (2012). Low survival after release into the wild: Assessing “the burden of captivity” on mallard physiology and behaviour. *European Journal of Wildlife Research*, *58*(1), 255–267.
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences 2nd edn*. Erlbaum Associates, Hillsdale.
- Cornelius Ruhs, E., Vézina, F., & Karasov, W. H. (2019). Physiological and immune responses of free-living temperate birds provided a gradient of food supplementation. *Physiological and Biochemical Zoology*, *92*(1), 106–114.
- Costa, G. (2012). *Behavioural adaptations of desert animals*. Springer Science & Business Media.
- Dawson, R. D., & Bortolotti, G. R. (1997). Are avian hematocrits indicative of condition? American kestrels as a model. *The Journal of Wildlife Management*, 1297–1306.
- Delciellos, A. C., Barros, C. dos S. de, Prevedello, J. A., Ferreira, M. S., Cerqueira, R., & Vieira, M. V. (2018). Habitat fragmentation affects individual condition: Evidence from small mammals of the brazilian atlantic forest. *Journal of Mammalogy*, *99*(4), 936–945.

- 725 Deppe, J. L., Ward, M. P., Bolus, R. T., Diehl, R. H., Celis-Murillo, A., Zenzal, T. J., Moore, F. R., Benson,
726 T. J., Smolinsky, J. A., Schofield, L. N., & others. (2015). Fat, weather, and date affect migratory
727 songbirds' departure decisions, routes, and time it takes to cross the gulf of mexico. *Proceedings of the*
728 *National Academy of Sciences*, 112(46), E6331–E6338.
- 729 Drent, R., & Daan, S. (1980). The prudent parent: Energetic adjustments in avian breeding 1. *Ardea*,
730 55(1–2), 225–252.
- 731 Eillers, J., Van Alphen, J. J., & Sevenster, J. G. (1998). A field study of size–fitness relationships in the
732 parasitoid *asobara tabida*. *Journal of Animal Ecology*, 67(2), 318–324.
- 733 English, M. D., Robertson, G. J., Peck, L. E., Pirie-Hay, D., Roul, S., & Mallory, M. L. (2018). Body
734 condition of american black ducks (*anas rubripes*) wintering in atlantic canada using carcass composition
735 and a scaled mass index. *Canadian Journal of Zoology*, 96(10), 1137–1144.
- 736 Erciyas, K., Gürsoy, A., Özsemir, A., & Barış, Y. (2010). Body mass and fat score changes in recaptured
737 birds during the autumn migration at the cernek ringing station in turkey. *The Ring*, 32(1-2), 3–15.
- 738 Fair, J., Whitaker, S., & Pearson, B. (2007). Sources of variation in haematocrit in birds. *Ibis*, 149(3),
739 535–552.
- 740 Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using g* power 3.1:
741 Tests for correlation and regression analyses. *Behavior Research Methods*, 41(4), 1149–1160.
- 742 Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G* power 3: A flexible statistical power analysis
743 program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39(2), 175–191.
- 744 Gessaman, J. (1999). Evaluation of some nonlethal methods of estimating avian body fat and lean mass.
745 *Proceedings of the 22nd International Ornithological Congress. University of Natal Press, Durban*, 2–16.
- 746 Gill, F. (1995). *Ornithology*. Freeman; Company.
- 747 Gosler, A. (1991). On the use of greater covert moult and pectoral muscle as measures of condition in
748 passerines with data for the great tit *parus major*. *Bird Study*, 38(1), 1–9.
- 749 Gosler, A. G., Greenwood, J. J., & Perrins, C. (1995). Predation risk and the cost of being fat. *Nature*,
750 377(6550), 621–623.
- 751 Guglielmo, C. G., McGuire, L. P., Gerson, A. R., & Seewagen, C. L. (2011). Simple, rapid, and non-invasive
752 measurement of fat, lean, and total water masses of live birds using quantitative magnetic resonance.
753 *Journal of Ornithology*, 152(1), 75.
- 754 Haas, C. A. (1998). Effects of prior nesting success on site fidelity and breeding dispersal: An experimental
755 approach. *The Auk*, 115(4), 929–936.
- 756 Hadfield, J. (2010). MCMC methods for multi-response generalized linear mixed models: The MCMCglmm
757 r package. *Journal of Statistical Software*, 33(2), 1–22.
- 758 Hadfield, J. (2014). *MCMCglmm course notes*. [http://cran.r-project.org/web/packages/MCMCglmm/
759 vignettes/CourseNotes.pdf](http://cran.r-project.org/web/packages/MCMCglmm/vignettes/CourseNotes.pdf)
- 760 Hartig, F. (2019). *DHARMA: Residual diagnostics for hierarchical (multi-level / mixed) regression models*.
761 <http://florianhartig.github.io/DHARMA/>
- 762 Heidinger, I. M. M., Hein, S., & Bonte, D. (2010). Patch connectivity and sand dynamics affect dispersal-
763 related morphology of the blue-winged grasshopper *oedipoda caerulescens* in coastal grey dunes. *Insect*
764 *Conservation and Diversity*, 3(3), 205–212.
- 765 Henderson, L., Evans, N., Heidinger, B., Herborn, K., & Arnold, K. (2017). Do glucocorticoids predict fit-
766 ness? Linking environmental conditions, corticosterone and reproductive success in the blue tit, *cyanistes*
767 *caeruleus*. *Royal Society Open Science*, 4(10), 170875.
- 768 Huxley, J. (1932). *Problems of relative growth*. Dover Publications.

- 769 Jacobs, S. R., Elliott, K., Guigueno, M. F., Gaston, A. J., Redman, P., Speakman, J. R., & Weber, J.-M.
770 (2012). Determining seabird body condition using nonlethal measures. *Physiological and Biochemical*
771 *Zoology*, 85(1), 85–95.
- 772 Johnson, K., DuVal, E., Kiehl, M., & Hughes, C. (2000). Male mating strategies and the mating system of
773 great-tailed grackles. *Behavioral Ecology*, 11(2), 132–141.
- 774 Johnson, K., & Peer, B. D. (2001). *Great-tailed grackle: Quiscalus mexicanus*. Birds of North America,
775 Incorporated.
- 776 Kaiser, A. (1993). A new multi-category classification of subcutaneous fat deposits of songbirds (una nueva
777 clasificación, con multi-categorías, para los depósitos de grasa en aves canoras). *Journal of Field Or-*
778 *mithology*, 246–255.
- 779 Kelly, C. D., Tawes, B. R., & Worthington, A. M. (2014). Evaluating indices of body condition in two cricket
780 species. *Ecology and Evolution*, 4(23), 4476–4487.
- 781 Labocha, M. K., & Hayes, J. P. (2012). Morphometric indices of body condition in birds: A review. *Journal*
782 *of Ornithology*, 153(1), 1–22.
- 783 Labocha, M. K., Schutz, H., & Hayes, J. P. (2014). Which body condition index is best? *Oikos*, 123(1),
784 111–119.
- 785 Langston, N. E., Freeman, S., Rohwer, S., & Gori, D. (1990). The evolution of female body size in red-winged
786 blackbirds: The effects of timing of breeding, social competition, and reproductive energetics. *Evolution*,
787 44(7), 1764–1779.
- 788 Liao, C.-Y., Rikke, B. A., Johnson, T. E., Gelfond, J. A., Diaz, V., & Nelson, J. F. (2011). Fat maintenance
789 is a predictor of the murine lifespan response to dietary restriction. *Aging Cell*, 10(4), 629–639.
- 790 Maceda-Veiga, A., Green, A. J., & De Sostoa, A. (2014). Scaled body-mass index shows how habitat quality
791 influences the condition of four fish taxa in north-eastern Spain and provides a novel indicator of ecosystem
792 health. *Freshwater Biology*, 59(6), 1145–1160.
- 793 Maspons, J., Molowny-Horas, R., & Sol, D. (2019). Behaviour, life history and persistence in novel environ-
794 ments. *Philosophical Transactions of the Royal Society B*, 374(1781), 20180056.
- 795 Mayfield, H. (1961). Nesting success calculated from exposure. *The Wilson Bulletin*, 255–261.
- 796 McGuire, L. P., Kelly, L. A., Baloun, D. E., Boyle, W. A., Cheng, T. L., Clerc, J., Fuller, N. W., Gerson,
797 A. R., Jonasson, K. A., Rogers, E. J., & others. (2018). Common condition indices are no more effective
798 than body mass for estimating fat stores in insectivorous bats. *Journal of Mammalogy*, 99(5), 1065–1071.
- 799 McNamara, J. M., Barta, Z., Houston, A. I., & Race, P. (2005). A theoretical investigation of the effect
800 of predators on foraging behaviour and energy reserves. *Proceedings of the Royal Society B: Biological*
801 *Sciences*, 272(1566), 929–934.
- 802 Merilä, J., & Svensson, E. (1997). Are fat reserves in migratory birds affected by condition in early life?
803 *Journal of Avian Biology*, 279–286.
- 804 Milenkaya, O., Catlin, D. H., Legge, S., & Walters, J. R. (2015). Body condition indices predict reproductive
805 success but not survival in a sedentary, tropical bird. *PLoS One*, 10(8), e0136582.
- 806 Milenkaya, O., Weinstein, N., Legge, S., & Walters, J. R. (2013). Variation in body condition indices of
807 crimson finches by sex, breeding stage, age, time of day, and year. *Conservation Physiology*, 1(1).
- 808 Montreuil-Spencer, C. (2017). *Relationships between winter energetic condition and reproductive investment*
809 *in a wild bird* [PhD thesis].
- 810 Murphy, M. T. (1986). Body size and condition, timing of breeding, and aspects of egg production in eastern
811 kingbirds. *The Auk*, 103(3), 465–476.
- 812 Musacchia, X. (1953). A study of the lipids in arctic migratory birds. *The Condor*, 55(6), 305–312.

- 813 Nakagawa, S., & Schielzeth, H. (2010). Repeatability for gaussian and non-gaussian data: A practical guide
814 for biologists. *Biological Reviews*, 85(4), 935–956.
- 815 National Climatic Data Center, N. (2020). *Climate data online global summary of the month*. <https://www.ncdc.noaa.gov/cdo-web/>
816
- 817 Nip, E. J., Frei, B., & Elliott, K. H. (2019). Seasonal and temporal variation in scaled mass index of
818 black-capped chickadees (*poecile atricapillus*). *The Canadian Field-Naturalist*, 132(4), 368–377.
- 819 Peig, J., & Green, A. J. (2009). New perspectives for estimating body condition from mass/length data:
820 The scaled mass index as an alternative method. *Oikos*, 118(12), 1883–1891.
- 821 Pendlebury, C., MacLeod, M., & Bryant, D. (2004). Variation in temperature increases the cost of living in
822 birds. *Journal of Experimental Biology*, 207(12), 2065–2070.
- 823 Perrins, C. (1970). The timing of birds ‘breeding seasons. *Ibis*, 112(2), 242–255.
- 824 Pond, C. (1981). Storage. *Physiological Ecology*, 190–219.
- 825 R Core Team. (2017). *R: A language and environment for statistical computing*. R Foundation for Statistical
826 Computing. <https://www.R-project.org>
- 827 Shaffer, T. L. (2004). A unified approach to analyzing nest success. *The Auk*, 121(2), 526–540.
- 828 Stevenson, R., & Woods Jr, W. A. (2006). Condition indices for conservation: New uses for evolving tools.
829 *Integrative and Comparative Biology*, 46(6), 1169–1190.
- 830 Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2017). rptR: Repeatability estimation and variance decom-
831 position by generalized linear mixed-effects models. *Methods in Ecology and Evolution*, 8(11), 1639–1644.
- 832 Walsberg, G. E. (1988). Evaluation of a nondestructive method for determining fat stores in small birds and
833 mammals. *Physiological Zoology*, 61(2), 153–159.
- 834 Warnock, N., & Bishop, M. A. (1998). Spring stopover ecology of migrant western sandpipers. *The Condor*,
835 100(3), 456–467.
- 836 Wehtje, W. (2003). The range expansion of the great-tailed grackle (*quiscalus mexicanus gmelin*) in north
837 america since 1880. *Journal of Biogeography*, 30(10), 1593–1607.
- 838 Welbergen, J. A. (2011). Fit females and fat polygynous males: Seasonal body mass changes in the grey-
839 headed flying fox. *Oecologia*, 165(3), 629–637.
- 840 Wilder, S. M., Raubenheimer, D., & Simpson, S. J. (2016). Moving beyond body condition indices as an
841 estimate of fitness in ecological and evolutionary studies. *Functional Ecology*, 30(1), 108–115.
- 842 Zhang, Y., Eyster, K., Liu, J.-S., & Swanson, D. L. (2015). Cross-training in birds: Cold and exercise
843 training produce similar changes in maximal metabolic output, muscle masses and myostatin expression
844 in house sparrows (*passer domesticus*). *Journal of Experimental Biology*, 218(14), 2190–2200.