

# **A systematic quantitative review of ecological responses to variation in seasonal snow cover**

Rachel A. Slatyer\*, Pieter A. Arnold

Division of Ecology & Evolution, Research School of Biology, The Australian National University,  
Canberra, ACT, Australia

\* Corresponding author: Rachel A. Slatyer

Address: 46 Sullivans Creek Rd, Acton, ACT 2600, Australia

Email: [rachel.slatyer@anu.edu.au](mailto:rachel.slatyer@anu.edu.au)

**Keywords:** Climate change, phenology, snow manipulation, subnivean, winter

# 1 **A systematic review of ecological responses to variation in seasonal**

## 2 **snow cover**

### 3 **Abstract**

4 Seasonal snow is among the most important factors governing the ecology of many terrestrial  
5 ecosystems, but rising global temperatures are changing snow regimes and driving widespread  
6 declines in the depth and duration of snow cover. Loss of the insulating snow layer will  
7 fundamentally change the environment – far more than incremental temperature increases.  
8 Understanding how individuals, populations, and communities respond to different snow  
9 conditions is thus essential for predicting and managing future ecosystem change. We conducted  
10 a systematic review to synthesize 365 studies that have examined ecological responses to  
11 variation in winter snow conditions. This substantial body of research encompasses a broad  
12 range of research methods (experimental manipulations, natural snow gradients, and long-term  
13 monitoring approaches), locations (35 countries), study organisms (plants, mammals, arthropods,  
14 birds, fish, lichen, and fungi), and response measures. Although we found trends towards  
15 advanced phenology with earlier snowmelt and greater mortality and/or physical injury with  
16 shallower snow, the mechanisms behind these responses remain largely unknown. With 96% of  
17 studies from the northern hemisphere, the generality of these trends across ecosystems is also  
18 unclear. We identified substantial research gaps and highlight five key areas for future research  
19 to better understand the mechanisms underlying responses to changing snow conditions and the  
20 consequences of those responses for seasonally snow-covered ecosystems.

## 21 **Introduction**

22 The presence of seasonal snow, covering the ground for weeks to months each year, is a feature  
23 of many temperate and montane ecosystems with up to a third of the Earth's terrestrial surface  
24 covered by seasonal snow at any time (Vaughan et al. 2013). Snow is one of the most important  
25 factors governing the ecology of these ecosystems due to its influence on the timing and length  
26 of the growing season, local and regional hydrology, and soil nutrient influxes (Billings &  
27 Mooney 1968; Körner 2003; Vavrus 2007; Blankinship & Hart 2012). Yet snow conditions are  
28 changing in many parts of the world, altering winter and growing season conditions for both  
29 plants and animals and with the potential to drive significant biodiversity loss (Vaughan et al.  
30 2013; Niittynen et al. 2018).

31 Mean global temperatures have increased by 0.7°C over the last 50 years (Stocker et al. 2013),  
32 with global snow cover decreasing by 10% over the same period (Walther et al. 2002). The most  
33 rapid and consistent losses of snow, in terms of both depth and duration, are mid-elevation areas  
34 (e.g. sub-alpine zone) and those with Mediterranean/maritime climates (e.g. Australian alpine  
35 region), where mean air temperatures are close to freezing and snow is primarily temperature-  
36 limited (Brown & Mote 2009; Steger et al. 2013; Vaughan et al. 2013). While shifts in regional  
37 and global atmospheric circulation patterns are driving elevated snowfall in areas where snow is  
38 limited by precipitation (e.g. high northern latitudes), these regions are still likely to experience  
39 reduced spring snow and a shorter growing season over the next 50 years (Räisänen 2008;  
40 Brown & Mote 2009; Vaughan et al. 2013).

41 Changes in winter precipitation and temperature regimes in seasonally snow-covered  
42 environments are mediated by the snowpack – the layer of accumulated snow – and changes to  
43 the snowpack will have diverse ecological effects (Geiger et al. 1995; Fig. 1). Experimental field  
44 manipulations that artificially advance snowmelt consistently induce earlier phenology in plants

45 (Wipf & Rixen 2010). With earlier spring snowmelt forecast to be among the most prominent  
46 effects of global warming on seasonal snowpacks (Déry & Brown 2007; Räisänen 2008), an  
47 overall advancement in spring phenology is expected. How organisms adapted to a seasonally  
48 snow-covered environment will respond to changes in snow conditions more generally –  
49 including the extent of cover, depth, density, and the timing of snowmelt – is, however, unclear.

50 The extent of snow cover determines the availability of snow-associated habitat. Both mammals  
51 and arthropods, can be active on the snow surface during winter. Small arthropods such as  
52 springtails and mites can also inhabit the snowpack itself, moving through air pockets between  
53 ice crystals and using thermal gradients within the snowpack to regulate their microclimate  
54 (Leinaas 1981; Hågvar 2010). Finally, a narrow space between the ground and the base of the  
55 snowpack – the subnivean space, or subnivium (Pauli et al. 2013) – provides a physically  
56 sheltered and thermally stable overwinter refuge for plants and animals (both active and  
57 hibernating/diapause) (Pauli et al. 2013). Changes to the extent of snow cover will have a  
58 direct impact on the availability of these habitats (Fig. 1).

59 Snow depth and density determine the degree of thermal insulation offered by the snowpack  
60 (Pruitt 1970). Snow has a low thermal conductivity and, depending on density, 20 cm of snow is  
61 generally sufficient to effectively insulate the subnivean space from diel fluctuations in ambient  
62 air temperature (Pruitt 1970). This buffering effect means that subnivean organisms are expected  
63 to experience the coldest temperatures during early autumn and late spring – not during winter –  
64 in contrast to ecosystems without seasonal snow cover. Groffman et al. (2001) suggested that  
65 seasonally snow-covered ecosystems might thus experience “colder soils in a warmer world”,  
66 with snowpack decline exposing soils and organisms to air temperatures up to 15°C colder than  
67 those in a snow-buffered airspace (Mölders & Walsh 2004). A shallower snowpack will also  
68 increase ground temperature fluctuations, which are thus more likely to cross critical

69 physiological thresholds for subnivean organisms (Geiser & Broome 1993; Marshall & Sinclair  
70 2012; Williams et al. 2015).

71 The duration of snow cover directly determines growing season length for plants, with little  
72 growth and development under the snow (Körner 2003). While a longer growing season could  
73 increase productivity (e.g. Billings & Bliss 1959), snowmelt timing determines the conditions to  
74 which plants are exposed when they emerge from snow, and earlier snowmelt can increase  
75 exposure to damaging frost and extreme temperatures (Steltzer et al. 2009; Gezon et al. 2016).  
76 The timing of snowmelt influences water availability during the growing season, which is  
77 gradually released from melting snow, and late-season moisture limitation is a risk from an early  
78 snowmelt (Litaor et al. 2008; Berdanier & Klein 2011). Changes to snowmelt timing are  
79 particularly relevant for plants because they are unable to track the presence (or absence) of the  
80 snowpack, and for interactions between plants and other organisms such as pollinators and  
81 herbivores (e.g. Forrest & Thomson 2011).

82 The ecological responses of organisms to changes in snow conditions can be measured using  
83 both experimental and observational approaches. Experimental methods that manipulate specific  
84 aspects of the snowpack (e.g. snow depth) allow a targeted assessment of biotic responses but are  
85 often (necessarily) limited in spatial scale. Observational approaches include both natural snow  
86 gradients and multi-year monitoring. Snow gradients typically describe long-term responses of  
87 populations, species, and communities to spatial variation in snow conditions (e.g. adaptive  
88 differences in cold tolerance among populations: Briceño et al. 2014). By contrast, studies that  
89 monitor ecological responses across years with varying snow conditions generally describe  
90 shorter-term effects (e.g. body mass following years with low/high snow: Hendrichsen & Tyler  
91 2014). Experimental, gradient, and monitoring methods provide complementary approaches for

92 examining ecological responses to changes in snow conditions but are not often considered  
93 together.

94 In this review, we synthesize studies that have explored ecological responses to spatial and  
95 temporal variation in snow conditions using a systematic review approach (Pullin & Stewart  
96 2006; Lortie 2014; Pickering & Byrne 2014). In particular, we explore (a) the geographic  
97 locations of research, (b) what has been measured and how, (c) whether any conclusions can be  
98 made about general responses to snow conditions, and (d) gaps in current knowledge and future  
99 research directions.

## 100 **Methods**

### 101 *Search procedure and inclusion criteria*

102 The systematic review approach provides reproducible protocols and transparent reporting for  
103 searching, screening, and extracting data from the literature to give an overview of a field  
104 (Koricheva & Gurevitch 2013; Lortie 2014). We used the Preferred Reporting Items in  
105 Systematic Reviews and Meta-Analyses (PRISMA) framework (Moher et al. 2009) to compile a  
106 database of studies that measured ecological responses to variation in snow conditions.

107 To identify relevant literature, we searched three databases with the term “snow” in combination  
108 with any one of the following: “manipulation”, “experimental warming”, “climate change”,  
109 “ecology”, “long-term monitoring”, “long term monitoring”, “ploughing”, “gradient”,  
110 “grooming”, “snowpatch”, “phenology”, “winter warming”, (“winter” and “climate change”).  
111 These terms were used within “Topic” in the Web of Science database, within “Abstract, title,  
112 author, keywords” in the Scopus database, and within “Keywords” in the Science Direct  
113 database, limiting results to studies in English-language journals. These searches were initially  
114 conducted in May 2016 and repeated in May 2019 to update the database, which produced 9,047

115 unique results (Fig. 2). To supplement this topic-based search, 24 reviews on related topics were  
116 identified that have been published since 1999 (Appendix S1). All studies citing or cited by these  
117 reviews were retrieved in May 2016, returning an additional 860 unique studies (Fig. 2).  
118 Unpublished data and “grey” literature, such as protected area management plans, were not  
119 included as much of this literature is not publicly available and is challenging to search  
120 systematically via electronic databases (Côté et al. 2013).

121 All studies were screened for eligibility, based on the following criteria: (1) the study was  
122 original research, not a review, and published in an English-language academic journal; (2) the  
123 study was carried out at a site where there is seasonal snow cover; (3) the study measured some  
124 form of biotic response; (4) the study measured responses to changes in snow cover. For  
125 criterion 2, we excluded studies from polar regions and permanently snow-covered areas. Cooper  
126 (2014) reviewed the effects of winter climate change on arctic ecosystems and the effects of  
127 snow regime change in permanently snow-covered ecosystems are likely to differ from those in  
128 seasonal environments, where plants and animals are adapted to snow for only part of the year.  
129 For criterion 3, we considered any form of response measured in an animal or plant but excluded  
130 studies on soil microbes. For criterion 4, we included studies that experimentally manipulated  
131 snow cover in the field (“manipulation”), those that measured responses along a snowmelt  
132 gradient (“gradient”), and those that recorded responses over multiple years across which snow  
133 conditions differed (“monitoring”). Several experimental methods can be used to reduce snow  
134 cover, including manual snow removal (e.g. Bombonato & Gerdol 2012), external heating (e.g.  
135 Adler et al. 2007), soil heating (e.g. Bokhorst et al. 2012), the addition of material that increases  
136 albedo and facilitates snowmelt (e.g. Steltzer et al. 2009), and physical covering to prevent snow  
137 accumulation (e.g. Drescher & Thomas 2013). Natural snowmelt gradients and long-term studies  
138 offer a complementary field-based approach to assess the effects of snow depth, duration, and  
139 structure on organisms. In contrast to experimental snow removal, which is limited in scale,

140 these studies allow assessments of larger-scale and longer-term effects of growing season  
141 duration and winter snow conditions on community composition, individual behaviors, and  
142 functional traits. Studies were only included if they measured snow cover directly. A total of 365  
143 studies met all inclusion criteria (Fig. 2; Appendix S2).

#### 144 *Data extraction*

145 For each study, the following information was extracted: (1) the location (hemisphere, continent,  
146 country(ies), and study site(s)); (2) the focal taxonomic group(s); (3) the methodology, including  
147 the type of study, length of study and, for experimental studies, the form of manipulation; and (4)  
148 the type of measures made, including when responses were recorded, whether they were  
149 recorded for individuals, populations, or communities, and the type of response recorded (e.g.  
150 phenology, growth, survival, behavior).

151 For responses recorded during winter, we additionally recorded which snow habitat (on-snow:  
152 “snow surface”, in-snow: “intranivean”, under-snow: “subnivean”) was being used by the study  
153 organism. Data were analyzed using descriptive methods to reveal patterns in the literature and  
154 identify research gaps. Note that the numbers given in the results do not always sum to the total  
155 number of studies (365) because individual studies often included results in several categories.

156 In addition to the data above, which were extracted directly from each paper, we determined the  
157 general snow conditions for each study (or each site when a study included multiple sites). For  
158 each study, the latitude and longitude of the study site(s) was obtained either directly from the  
159 paper or by georeferencing named locations. For studies conducted over a large geographic area,  
160 we used an approximately central point of the study area. Data on seasonal snow classification  
161 (Sturm et al. 1995; Liston & Sturm 1998) were obtained from the Atlas of the Cryosphere, at a  
162  $0.5^{\circ} \times 0.5^{\circ}$  spatial resolution (Maurer 2007). Sturm et al. (1995)’s seasonal snow classification  
163 defines six classes of snow (tundra, taiga, alpine, maritime, prairie, ephemeral) based on the



164 stratigraphy, thickness, density, crystal morphology, and thermal gradient of the snowpack, and  
165 their spatial and temporal variability. Although this classification may not apply to all areas with  
166 seasonal snow (e.g. Sanecki et al. 2006a), it is a useful standard for comparisons. Snow  
167 classification was extracted for each study/site using RASTER 2.5-8 (Hijmans 2016), RGDAL 1.2-5  
168 (Bivand et al. 2016), and SP (Pebesma & Bivand 2005) packages in the R environment for  
169 statistical computing v3.3.0 (R Core Team 2016). The ephemeral snow classification (< 2  
170 months snow) covers large areas across the world that do not typically have seasonal snow,  
171 therefore it was not represented on the world map. Maps were plotted using GGMAP 3.0.0 (Kahle  
172 & Wickham 2013) and GGPLOT2 3.2.1 (Wickham 2016).

173 To summarize the main results, we tallied studies that had shown positive, negative, nil, or  
174 mixed responses to variation in snow conditions. Although such vote-counting methods are  
175 generally unsuitable as a formal statistical technique for research syntheses (Koricheva &  
176 Gurevitch 2013), they are valuable as a summary tool and highlight areas where formal meta-  
177 analysis might be warranted in the future. This approach was chosen because sample sizes were  
178 small for most response variables and many different types of response variable were measured,  
179 limiting the utility of quantitative analysis. Responses were summarized for plants, mammals,  
180 and arthropods – groups for which there were at least 20 studies. Twelve response variables were  
181 identified that were comparable across taxonomic groups. These are presented in relation to  
182 changes in snow depth and snowmelt timing (the most common aspects of snow variation  
183 measured). Summaries of results for all response variables measured across taxa are provided in  
184 Appendix S3.

## 185 **Results**

### 186 *Time and place*

187 There were 365 studies on ecological responses to variation in snow conditions that met all  
188 inclusion criteria. These studies were published between 1959 and 2019 with a median study  
189 duration of 2 years (range 1 – 60 years). While studies have been conducted in 35 countries,  
190 most of the research was from the USA (118 studies, 32%), Sweden (41 studies, 11%) and  
191 Canada (33 studies, 9%), and nearly all (349, 96%) from the northern hemisphere (Fig. 3,  
192 Table 1). Studies were conducted in alpine/montane (218 studies), temperate forest or grassland  
193 (94 studies), and sub-arctic environments (112 studies) (Table 1). Two locations featured  
194 prominently: Abisko in northern Sweden (27 studies) and the Rocky Mountain Biological  
195 Laboratory in Colorado, USA (20 studies).

196 The study sites cover a variety of snow conditions and, in the northern hemisphere, all snow  
197 types were represented: maritime ( $n = 193$  studies/sites), alpine ( $n = 86$ ), prairie ( $n = 63$ ), tundra  
198 ( $n = 79$ ), and taiga ( $n = 31$ ), with 38 studies lacking a snow classification. Note that a single  
199 study could have multiple sites. The predominance of studies on alpine (cold, deep snow cover)  
200 and maritime snow (warm, deep snow cover) does not correspond to the relative frequencies of  
201 these two snow types across the landscape: each are <10% of snow-covered land area in the  
202 northern hemisphere. In the southern hemisphere, maritime snow was the only snow type  
203 represented, although there were 15 sites that lacked a snow classification. This is likely due to  
204 the snow classification system being developed for northern hemisphere snow conditions, which  
205 are different to those in the southern hemisphere (Sanecki et al. 2006a).

### 206 *Organisms*

207 The impacts of seasonal snow cover have been assessed, in some way, for a broad range of plant  
208 and animal groups (Table 1). For plants (66% of all studies), this includes research on small

209 vascular plants ( $n = 158$ ), shrubs ( $n = 72$ ), trees ( $n = 40$ ), and bryophytes ( $n = 21$ ). For animals,  
210 most of the research has focused on mammals ( $n = 76$ ) or arthropods ( $n = 23$ ), with few studies  
211 for birds ( $n = 16$ ), fish ( $n = 2$ ), reptiles ( $n = 1$ ), or amphibians ( $n = 1$ ). Finally, a few studies  
212 included lichens ( $n = 13$ ) or fungi ( $n = 7$ ). Considering only the southern hemisphere, however,  
213 there was only one study of arthropods, four studies of mammals, and 13 studies of plants.

#### 214 *Study approach*

215 Research on ecological responses to variation in snow conditions has used experimental  
216 ( $n = 164$ ) and observational ( $n = 212$ ) methods (Table 2). This is true for research in both  
217 hemispheres, and all climatic zones. Observational studies included research using natural snow  
218 cover or snowmelt gradients ( $n = 119$ ) and year-to-year variation in snow conditions ( $n = 113$ ). A  
219 few studies used multiple methods: experimental manipulations with measures across snowmelt  
220 gradients ( $n = 7$ ) or through time ( $n = 5$ ), or long-term monitoring across snowmelt gradients  
221 ( $n = 20$ ).

222 Experimental manipulations of snow depth tested the effects of both more snow (increased  
223 depth:  $n = 62$ ; increased persistence:  $n = 47$ ), less snow (decreased depth:  $n = 68$ ; decreased  
224 persistence:  $n = 46$ ), and the effects of unusual weather events (e.g. mid-winter snowmelt:  
225  $n = 14$ ). However, more than half of the studies that altered snow depth also altered snowmelt  
226 timing (and *vice versa*), meaning that these effects are frequently confounded. Studies that  
227 altered snow duration almost always did so by manipulating the timing of spring snowmelt, with  
228 only three studies changing the timing of snow accumulation. Experimental manipulations of the  
229 effects of snow density ( $n = 17$ ) and snow chemistry ( $n = 4$ ) were most often related to  
230 anthropogenic use of snow: compaction from oversnow vehicles or skiing, and changes to  
231 chemistry or density due to artificial snowmaking.

232 Experimental approaches were common in physiological studies, community composition,  
233 chemistry, and overwinter survival, as well as the response of arthropods ( $n = 31$ ) and plants  
234 ( $n = 84$ ). By contrast, gradient and monitoring studies provide most of the evidence for effects of  
235 snow conditions on animal movements ( $n = 28$  and  $n = 18$ , respectively) and plant phenology  
236 ( $n = 33$  and  $n = 44$ , respectively).

### 237 *Timing of measurement*

238 Experimental studies nearly always measured responses in the subsequent growing season (93%  
239 of studies), while 20% of monitoring studies and 24% of gradient studies included winter  
240 measurements (Table 2). In contrast to all other taxa, more studies measured mammal responses  
241 during winter ( $n = 49$ ) than during the subsequent snow-free period ( $n = 32$ ), with these studies  
242 primarily exploring activity (e.g. home range size, habitat use) in relation to snow characteristics.  
243 There were 154 studies that measured the responses of small vascular plants during the growing  
244 season, but only five (Bell & Bliss 1979; Blume-Werry et al. 2016; Saarinen et al. 2016; Tessier  
245 2017; Mo et al. 2018) included measurements of winter response. In total, only 71 (19%) studies,  
246 of which only 22 were studies on non-mammalian organisms, included winter measurements.

### 247 *Ecological responses to snow variation*

248 We recorded 214 different response variables measured, across all studies (Appendix S3).  
249 Taking 12 response variables that are comparable to some extent between plants, mammals, and  
250 arthropods (Fig. 4), three results stand out. First, earlier snowmelt was consistently associated  
251 with earlier spring phenology across all groups (Fig. 4). Second, reduced snow depth was  
252 frequently associated with higher mortality and/or damage in plants; this effect was not clear for  
253 either arthropods or mammals, nor was there a clear association with snowmelt timing. Third,  
254 there seemed to be no clear directional effect of changes in either snow depth or snowmelt timing  
255 on body size (for animals) or total biomass (for plants), or on abundance/relative cover overall

256 (Appendix S3). In addition, variation in snow conditions was often (37 of 49 studies) associated  
257 with differences in plant and arthropod community composition in experimental, gradient, and  
258 monitoring studies.

## 259 **Discussion**

260 There is a substantial body of research on ecological responses to changes in snow conditions,  
261 ranging from studies of habitat use by large mammals during winter, to those testing the effects  
262 of shallow snow cover on plant physiology. These studies encompass many locations, study  
263 organisms, research methods and response variables, reflecting the widespread ecological  
264 importance of snow. Nevertheless, the large number of studies belies a thin research coverage for  
265 many taxa, locations, and research questions, and there are several notable gaps in the current  
266 literature, including geographic gaps and the measurement of winter responses.

### 267 *Research on seasonal snow cover is geographically skewed*

268 Snow occurs on every continent, but this is only partly reflected in the locations of research on  
269 seasonal snow, which is strongly biased towards European and North American mountain  
270 systems. Notably absent were studies from South America, Africa, and the sub-Antarctic. The  
271 Andes range in South America maintains both permanent and seasonal snowpacks, with the latter  
272 covering approximately 61,000 km<sup>2</sup> and accounting for 98% of snow-covered area on the  
273 continent (Saavedra et al. 2017). In the Chilean Andes, Cavieres and Arroyo (2000) found that  
274 seeds of the herb *Phacelia secunda* from higher elevations needed longer periods of cold prior to  
275 germination than those from lower elevation populations, however this study did not specifically  
276 examine effects of snow conditions. In Africa there are mountain systems with seasonal snow,  
277 including the Atlas Mountains in Algeria, Morocco and Tunisia, mountains in the Eastern Cape  
278 region of South Africa, and the Maluti Mountains in Lesotho. Studies from low-elevation forest

279 and grassland, many of which have “prairie”, “tundra” or “taiga” snow types, were also globally  
280 scarce, perhaps reflecting relatively higher human land-use across these areas.

281 Large regional differences in snow regimes, the drivers of snow cover variation (e.g.  
282 temperature- vs precipitation-limitation), and the likely impacts of climate change on snow  
283 conditions (Räisänen 2008) mean that expanding studies to include these underrepresented areas  
284 is important. The type and nature of the biota present also differs among regions and ecosystems  
285 (e.g. Sinclair & Chown 2005; Bannister 2007). In Australia, for example, snow-covered  
286 environments have many scleromorphic shrubs and no large mammals (Green & Osborne 2012).  
287 This ecosystem is likely to have fundamentally different responses to changes in snow conditions  
288 compared to, for example, a northern boreal forest with many large mammals.

289 While the ecological effects of snow might vary geographically due to differences in the  
290 underlying nature of the biota, variation might also be driven by differences in effects of  
291 changing snow conditions on the abiotic environment. When mean ambient air temperatures are  
292 above freezing, loss of the insulating snowpack should tend to increase, rather than decrease  
293 near-ground temperatures (Slatyer et al. 2017). By contrast, ambient winter air temperatures are  
294 well below freezing in many seasonally snow-covered ecosystems, driving a decrease in near-  
295 ground temperature with a reduced snowpack (e.g. Groffman et al. 2001; Decker et al. 2003; Tan  
296 et al. 2014; Petty et al. 2015). If the physical effects of reduced snow cover vary among regions,  
297 then inferences regarding ecological impacts will necessarily be region-specific. It is thus critical  
298 that studies of snow ecology measure and consider these differences. Conducting further studies  
299 in seasonally snow-covered areas of the southern hemisphere and other regions where the abiotic  
300 and/or ecological effects of reduced snow might be unique is essential to better understand which  
301 ecological impacts could be generalizable and which are specific to particular regions or snow  
302 conditions.

303 *Winter responses: the black box*

304 Fifteen years since Campbell et al. (2005) highlighted a paucity of ecological studies during  
305 winter, measurements of winter responses to variable snow conditions remain limited. Only 71  
306 of the 365 studies included in this review measured responses during the winter. Two studies of  
307 arthropods exemplify the value in measuring winter responses: following experimentally reduced  
308 snow cover, abundances of small arthropods were lower during the winter but, at the scale of the  
309 experiment, these effects did not carry over into the growing season (Sulkava & Huhta 2003;  
310 Bokhorst et al. 2013). For grazing mammals such as the Svalbard reindeer, climate warming has  
311 potential to lead to opposing effects on body mass in summer and winter. While warming during  
312 the growing season might increase plant growth and forage availability, winter warming that  
313 increases rain-on-snow events that in turn cause icing will restrict winter grazing, thereby  
314 reducing body mass (Albon et al. 2017). In these examples, measurements made only during the  
315 growing season would have failed to detect or explain the effects of changing snow conditions.

316 Among the studies included here, most of the winter measurements were on habitat use and  
317 activity patterns of mammals moving on the snow surface, with a tendency for individuals to  
318 favor areas with shallower snow than surrounding habitat (e.g. Mermod & Liberek 2002; Kolbe  
319 et al. 2007; Matthews 2010). An additional three studies examined how snow conditions affected  
320 habitat use and overwinter survival for subnivean animals. Artificially expanding the subnivean  
321 space increased winter activity and improved the overwinter survival of voles in Norway  
322 (Korslund & Steen 2006), while reducing the subnivean space lowered detection of small  
323 mammals in Australia (Sanecki et al. 2006b). However, winter survival of bank voles was not  
324 related to subnivean temperature or snow depth (Johnsen et al. 2018), with food availability more  
325 important than subnivean habitat structure for abundance and overwinter survival (Johnsen et al.  
326 2017). Shallow snow, and the associated increase in temperature fluctuations, can also increase  
327 the energy expenditure of hibernating subnivean mammals and dormant arthropods (e.g. Geiser

328 & Broome 1993; Irwin & Lee 2003). Taken together, these studies suggest contrasting effects of  
329 reduced snow on snow-surface and subnivean fauna.

330 The ecology of the subnivean environment remains elusive. With the exception of a detailed  
331 series of studies in Canada (e.g. Aitchison 1979a, b, c, 1984) and a couple of others (Berzitis et  
332 al. 2017; Slatyer et al. 2017), there are few surveys of subnivean arthropods and, although many  
333 mammals, reptiles, and amphibians are known or assumed to overwinter beneath the snow, their  
334 winter ecology is generally not well known. A second ecological unknown is the snow layer  
335 itself. Despite extensive searching of the literature, we found no studies that examined how  
336 changes in snow conditions might affect the intranivean fauna – small arthropods such as mites  
337 and springtails living within the snow layer itself. One might expect these organisms to be  
338 affected by the depth, density, and/or crystal structure of the snowpack, which affect the  
339 temperature gradient within the snowpack and the size of the spaces through which animals can  
340 move (Marchand 2013). While Leinaas (1981) studied movements of intranivean arthropods  
341 within the snowpack and (Hågvar 2010) provided an overview of intranivean taxa, we are  
342 unaware of further work on this topic.

### 343 *Conclusions*

344 The results of our systematic review provide a tantalizing glimpse into possible effects of snow  
345 conditions on organisms during winter, with individual studies showing that physiology, patterns  
346 of activity, habitat use, and foraging behavior can each be influenced by snow conditions.  
347 Winter-time frost hardiness, for example, has been shown to be greater for shrubs under deeper  
348 snow in both experimental (Taulavuori et al. 2011) and gradient studies (Palacio et al. 2015).  
349 These results hint at both short-term (plastic) and long-term (adaptive) physiological adjustments  
350 to variation in snow conditions. How such effects translate into growing season responses



351 remains unclear, but linking winter and growing season effects (e.g. Korslund & Steen 2006)  
352 will be an important avenue for future research.

353 By evaluating the current literature on ecological effects of changing snow conditions in  
354 seasonally snow-covered environments, this review provides an outline of where, how, and what  
355 research has been published, highlighting several knowledge gaps. Although many studies have  
356 examined ecological effects of changes in seasonal snow, when studies are divided by taxonomic  
357 group, location or climate zone, these numbers rapidly attenuate. There is ample scope for future  
358 research that both broadens the current literature and adds depth and detail to what already exists.

359 We highlight five key areas for future research:

- 360 1. Additional studies in underrepresented snow-covered areas, including the Andes  
361 mountain range in South America, the southern hemisphere generally, and low-elevation  
362 grassland and forest. These studies should be accompanied by measures of microclimate,  
363 so that observed ecological responses can be compared with studies from other regions.
- 364 2. Integration of natural snowmelt gradients with experimental manipulations or long-term  
365 monitoring (e.g. Dunne et al. 2004; Cornelius et al. 2013). Understanding how changing  
366 snow conditions will affect species and communities adapted to different snow conditions  
367 will require integrated approaches. Variation in, for example, physiological tolerances  
368 (e.g. Vrba 2012; Briceño et al. 2014), developmental temperatures (e.g. Forrest &  
369 Thomson 2011), or species interactions (e.g. Callaway et al. 2002) in areas with naturally  
370 high or low snow cover could affect responses to changing snow conditions.
- 371 3. Exploring the effects of changing snow conditions on species interactions. Only 14 of the  
372 studies in this review explicitly tested species interactions (but see also Nystuen et al.  
373 2014; Penczykowski et al. 2017). Early snowmelt could have large impacts on plant-

374 pollinator and plant-herbivore interactions by generating phenological mismatches (e.g.  
375 Kudo & Ida 2013).

376 4. Investigations into the mechanisms underlying higher mortality/injury with reduced  
377 snow/early snowmelt for plants. For example, is mortality caused by an accumulation of  
378 sub-lethal injuries or a single extreme event? Injury could similarly be caused by many  
379 factors such as species interactions (e.g. herbivory: Roy et al. 2004; fungal attack: Graae  
380 et al. 2008), physical damage from ice formation (e.g. Briceño et al. 2014), and  
381 physiological stress (e.g. Bokhorst et al. 2010). While similar mechanisms might be  
382 expected to affect mortality/injury in arthropods (e.g. ice encasement: Coulson et al.  
383 2000; food availability: Konestabo et al. 2007; crossing physiological thresholds:  
384 Marshall & Sinclair 2012), further studies testing both responses to changing snow  
385 conditions and the mechanisms behind these are needed.

386 5. Tests of the effects of early snowmelt on recruitment (e.g. seed germination and seedling  
387 establishment in plants Milbau et al. 2013; and hatching success in arthropods).  
388 Phenological shifts induced by early snowmelt are likely to cause decoupling between  
389 life stage and the climatic conditions to which that life stage has historically been  
390 exposed. Effects on recruitment, which typically manifest early in the growing season,  
391 will potentially have larger impacts at the ecosystem level than effects on adult growth.

392  
393 Seasonal snow is a central feature in the ecology of many terrestrial ecosystems. With continued  
394 climate change altering snow regimes worldwide, an understanding of how individuals,  
395 populations, species, and communities respond to different snow conditions is essential for  
396 predicting and managing future ecosystem change. There is great scope for more research on  
397 ecological effects of snow, both in terms of geographic and taxonomic breadth, and uncovering  
398 the most salient effects that might themselves be region- or species-specific.

399 **Acknowledgements:** Many thanks to Verónica Briceño for the fun and thought-provoking  
400 discussions that led to this review, and for providing comments on the manuscript. Thanks also  
401 to Bente Graae, Adrienne Nicotra, Kate Umbers, Catherine Pickering, and Kimberley Thompson  
402 for many constructive comments. Rachel Slatyer was supported by an American Australia  
403 Association fellowship while completing this work.

404 **Supporting dataset:** Available from the figshare repository: 10.6084/m9.figshare.4977998.

## 405 **Literature cited**

406 Adler LS, De Valpine P, Harte J, Call J. 2007. Effects of long-term experimental warming on  
407 aphid density in the field. *Journal of the Kansas Entomological Society* **80**:156-168.

408 Aitchison CW. 1979a. Winter-active subnivean invertebrates in southern Canada. I. Collembola.  
409 *Pedobiologia* **19**:113-120.

410 Aitchison CW. 1979b. Winter-active subnivean invertebrates in southern Canada. II. Coleoptera.  
411 *Pedobiologia* **19**:121-1979.

412 Aitchison CW. 1979c. Winter-active subnivean invertebrates in southern Canada. III. Acari.  
413 *Pedobiologia* **19**:153-160.

414 Aitchison CW. 1984. The ecology of winter-active collembolans and spiders. PhD thesis.  
415 University of Manitoba, Canada, University of Manitoba.

416 Albon SD, et al. 2017. Contrasting effects of summer and winter warming on body mass explain  
417 population dynamics in a food-limited Arctic herbivore. *Global Change Biology*  
418 **23**:1374-1389.

419 Bannister P. 2007. A touch of frost? Cold hardiness of plants in the Southern Hemisphere. *New*  
420 *Zealand Journal of Botany* **45**:1-33.

421 Bell KL, Bliss LC. 1979. Autecology of *Kobresia bellardii*: Why winter snow accumulation  
422 limits local distribution. *Ecological Monographs* **49**:377-402.

- 423 Berdanier AB, Klein JA. 2011. Growing season length and soil moisture interactively constrain  
424 high elevation aboveground net primary production. *Ecosystems* **14**:963-974.
- 425 Berzitis EA, Hager HA, Sinclair BJ, Hallett RH, Newman JA. 2017. Winter warming effects on  
426 overwinter survival, energy use, and spring emergence of *Cerotoma trifurcata*  
427 (Coleoptera: Chrysomelidae). *Agricultural and Forest Entomology* **19**:163-170.
- 428 Billings WD, Bliss LC. 1959. An alpine snowbank environment and its effects on vegetation,  
429 plant development, and productivity. *Ecology* **40**:388-397.
- 430 Billings WD, Mooney HA. 1968. The ecology of arctic and alpine plants. *Biological Reviews*  
431 **43**:481-529.
- 432 Bivand R, Keitt T, Rowlingson B. 2016. rgdal: bindings for the geospatial data abstraction  
433 library. R package version 1.2-5. <https://CRAN.R-project.org/package=rgdal>.
- 434 Blankinship JC, Hart SC. 2012. Consequences of manipulated snow cover on soil gaseous  
435 emission and N retention in the growing season: a meta-analysis. *Ecosphere* **31**:1-20.
- 436 Blume-Werry G, Kreyling J, Laudon H, Milbau A. 2016. Short-term climate change  
437 manipulation effects do not scale up to long-term legacies: effects of an absent snow  
438 cover on boreal forest plants. *Journal of Ecology* **104**:1638-1648.
- 439 Bokhorst S, Bjerke JW, Davey MP, Taulavuori K, Taulavuori E, Laine K, Callaghan TV,  
440 Phoenix GK. 2010. Impacts of extreme winter warming events on plant physiology in a  
441 sub-Arctic heath community. *Physiologia Plantarum* **140**:128-140.
- 442 Bokhorst S, Metcalfe DB, Wardle DA. 2013. Reduction in snow depth negatively affects  
443 decomposers but impact on decomposition rates is substrate dependent. *Soil Biology and*  
444 *Biochemistry* **62**:157-164.
- 445 Bokhorst S, Phoenix GK, Bjerke JW, Callaghan TV, Huyer-Brugman F, Berg MP. 2012.  
446 Extreme winter warming events more negatively impact small rather than large soil  
447 fauna: shift in community composition explained by traits not taxa. *Global Change*  
448 *Biology* **18**:1152-1162.
- 449 Bombonato L, Gerdol R. 2012. Manipulating snow cover in an alpine bog: effects on ecosystem  
450 respiration and nutrient content in soil and microbes. *Climatic Change* **114**:261-272.

- 451 Briceño VF, Harris-Pascal D, Nicotra AB, Williams E, Ball MC. 2014. Variation in snow cover  
452 drives differences in frost resistance in seedlings of the alpine herb *Aciphylla glacialis*.  
453 *Environmental and Experimental Botany* **106**:174-181.
- 454 Brown RD, Mote PW. 2009. The response of northern hemisphere snow cover to a changing  
455 climate. *Journal of Climate* **22**:2124-2145.
- 456 Callaway RM, et al. 2002. Positive interactions among alpine plants increase with stress. *Nature*  
457 **417**:844-848.
- 458 Campbell JL, Mitchell MJ, Groffman PM, Christenson LM, Hardy JP. 2005. Winter in  
459 northeastern North America: A critical period for ecological processes. *Frontiers in*  
460 *Ecology and the Environment* **3**:314-322.
- 461 Cavieres LA, Arroyo MTK. 2000. Seed germination response to cold stratification period and  
462 thermal regime in *Phacelia secunda* (Hydrophyllaceae). *Plant Ecology* **149**:1-8.
- 463 Cooper EJ. 2014. Warmer shorter winters disrupt arctic terrestrial ecosystems. *Annual Review of*  
464 *Ecology, Evolution, and Systematics* **45**:271-295.
- 465 Cornelius C, Leingärtner A, Hoiss B, Krauss J, Steffan-Dewenter I, Menzel A. 2013.  
466 Phenological response of grassland species to manipulative snowmelt and drought along  
467 an altitudinal gradient. *Journal of Experimental Botany* **64**:241-251.
- 468 Côté I, Curtis PS, Rothstein HR, Stewart GB. 2013. Gathering data: searching literature and  
469 selection criteria. Pages 37-51 in Koricheva J, Gurevitch J, and Mengersen K, editors.  
470 *Handbook of meta-analysis in ecology and evolution*. Princeton University Press,  
471 Princeton, NJ, USA.
- 472 Coulson SJ, Leinaas HP, Ims RA, Søvik G. 2000. Experimental manipulation of the winter  
473 surface ice layer: The effects on a High Arctic soil microarthropod community.  
474 *Ecography* **23**:299-306.
- 475 Decker KLM, Wang D, Waite C, Scherbatskoy T. 2003. Snow removal and ambient air  
476 temperature effects on forest soil temperatures in Northern Vermont. *Soil Science Society*  
477 *of America Journal* **67**:1234-1242.

- 478 Déry SJ, Brown RD. 2007. Recent Northern Hemisphere snow cover extent trends and  
479 implications for the snow-albedo feedback. *Geophysical Research Letters* **34**:L22504.
- 480 Drescher M, Thomas SC. 2013. Snow cover manipulations alter survival of early life stages of  
481 cold-temperate tree species. *Oikos* **122**:541-554.
- 482 Dunne JA, Saleska SR, Fischer ML, Harte J. 2004. Integrating experimental and gradient  
483 methods in ecological climate change research. *Ecology* **85**:904-916.
- 484 Forrest JRK, Thomson JD. 2011. An examination of synchrony between insect emergence and  
485 flowering in Rocky Mountain meadows. *Ecological Monographs* **81**:469-491.
- 486 Geiger R, Aron RA, Todhunter P 1995. *The Climate Near the Ground*. Harvard University Press,  
487 Cambridge, UK.
- 488 Geiser F, Broome LS. 1993. The effect of temperature on the pattern of torpor in a marsupial  
489 hibernator. *Journal of Comparative Physiology B* **163**:133-137.
- 490 Gezon ZJ, Inouye DW, Irwin RE. 2016. Phenological change in a spring ephemeral: Implications  
491 for pollination and plant reproduction. *Global Change Biology* **22**:1779-1793.
- 492 Graae BJ, Alsos IG, Erjnaes R. 2008. The impact of temperature regimes on development,  
493 dormancy breaking and germination of dwarf shrub seeds from arctic, alpine and boreal  
494 sites. *Plant Ecology* **1998**:275-284.
- 495 Green K, Osborne W 2012. *A Field Guide to the Wildlife of the Australian Snow Country*. New  
496 Holland Publishers, Wahroonga, NSW, Australia.
- 497 Groffman PM, Driscoll CT, Fahey TJ, Hardy JP, Fitzhugh RD, Tierney GL. 2001. Colder soils in  
498 a warmer world: A snow manipulation study in a northern hardwood forest ecosystem.  
499 *Biogeochemistry* **56**:135-150.
- 500 Hågvar S. 2010. A review of Fennoscandian arthropods living on and in snow. *European Journal*  
501 *of Entomology* **107**:281-298.
- 502 Hendrichsen DK, Tyler NJC. 2014. How the timing of weather events influences early  
503 development in a large mammal. *Ecology* **95**:1737-1745.

504 Hijmans RJ. 2016. raster: geographic data analysis and modeling. R package version 2.5-8.  
505 <https://CRAN.R-project.org/package=raster>.

506 Irwin JT, Lee RE. 2003. Cold winter microenvironments conserve energy and improve  
507 overwintering survival and potential fecundity of the goldenrod gall fly, *Eurosta*  
508 *solidaginis*. *Oikos* **100**:71-78.

509 Johnsen K, Boonstra R, Boutin S, Devineau O, Krebs CJ, Andreassen HP. 2017. Surviving  
510 winter: Food, but not habitat structure, prevents crashes in cyclic vole populations.  
511 *Ecology and Evolution* **7**:115-124.

512 Johnsen K, Devineau O, Andreassen HP. 2018. The effects of winter climate and intrinsic factors  
513 on survival of cyclic vole populations in southeastern Norway. *Annales Zoologici Fennici*  
514 **55**:173-185.

515 Kahle D, Wickham H. 2013. ggmap: spatial visualization with ggplot2. *The R Journal* **5**:144-  
516 161.

517 Kolbe JA, Squires JR, Pletscher DH, Ruggiero LF. 2007. The effect of snowmobile trails on  
518 coyote movements within lynx home ranges. *Journal of Wildlife Management* **71**:1409-  
519 1418.

520 Konestabo HS, Michelsen A, Holmstrup M. 2007. Responses of springtail and mite populations  
521 to prolonged periods of soil freeze-thaw cycles in a sub-arctic ecosystem. *Applied Soil*  
522 *Ecology* **36**:136-146.

523 Koricheva J, Gurevitch J. 2013. Place of meta-analysis among other methods of research  
524 synthesis. Pages 3-13 in Koricheva J, Gurevitch J, and Mengersen K, editors. *Handbook*  
525 *of Meta-Analysis in Ecology and Evolution*. Princeton University Press, Princeton, NJ,  
526 USA.

527 Körner C 2003. *Alpine plant life: functional plant ecology of high mountain ecosystems*.  
528 Springer-Verlag, Berlin, Germany.

529 Korslund L, Steen H. 2006. Small rodent winter survival: snow conditions limit access to food  
530 resources. *Journal of Animal Ecology* **75**:156-166.

- 531 Kudo G, Ida TY. 2013. Early onset of spring increases the phenological mismatch between  
532 plants and pollinators. *Ecology* **94**:2311-2320.
- 533 Leinaas HP. 1981. Activity of Arthropoda in snow within a coniferous forest, with special  
534 reference to Collembola. *Holarctic Ecology* **4**:127-138.
- 535 Liston GE, Sturm M 1998. Global Seasonal Snow Cover System. National Snow and Ice Data  
536 Center. Digital media, Boulder, Colorado, USA.
- 537 Litaor MI, Williams M, Seastedt TR. 2008. Topographic controls on snow distribution, soil  
538 moisture, and species diversity of herbaceous alpine vegetation, Niwot Ridge, Colorado.  
539 *Journal of Geophysical Research-Biogeosciences* **113**:G02008.
- 540 Lortie CJ. 2014. Formalized synthesis opportunities for ecology: systematic reviews and meta-  
541 analyses. *Oikos* **123**:897-902.
- 542 Marchand PJ 2013. Life in the cold: an introduction to winter ecology. University Press of New  
543 England, Hanova, NH, USA.
- 544 Marshall KE, Sinclair BJ. 2012. Threshold temperatures mediate the impact of reduced snow  
545 cover on overwintering freeze-tolerant caterpillars. *Naturwissenschaften* **99**:33-41.
- 546 Matthews A. 2010. Changes in fine-scale movement and foraging patterns of common wombats  
547 along a snow-depth gradient. *Wildlife Research* **37**:175-182.
- 548 Maurer J 2007. Atlas of the Cryosphere. National Snow and Ice Data Center. Digital media,  
549 Boulder, Colorado, USA.
- 550 Mermod CP, Liberek M. 2002. The role of snowcover for European wildcat in Switzerland.  
551 *Zeitschrift Fur Jagdwissenschaft* **48**:17-24.
- 552 Milbau A, Shevtsova A, Osler N, Mooshammer M, Graae BJ. 2013. Plant community type and  
553 small-scale disturbances, but not altitude, influence the invasibility in subarctic  
554 ecosystems. *New Phytologist* **197**:1002-1011.
- 555 Mo L, Luo P, Mou CX, Yang H, Wang J, Wang ZY, Li YJ, Luo C, Li T, Zuo DD. 2018. Winter  
556 plant phenology in the alpine meadow on the eastern Qinghai-Tibetan Plateau. *Annals of*  
557 *Botany* **122**:1033-1045.



558 Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. 2009. Preferred reporting  
559 items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Medicine*  
560 **6**:e1000097.

561 Mölders N, Walsh JE. 2004. Atmospheric response to soil-frost and snow in Alaska in March.  
562 *Theoretical and Applied Climatology* **77**:77-105.

563 Niittynen P, Heikkinen RK, Luoto M. 2018. Snow cover is a neglected driver of Arctic  
564 biodiversity loss. *Nature Climate Change* **8**:997-1001.

565 Nystuen KO, Evju M, Rusch GM, Graae BJ, Elde NE. 2014. Rodent population dynamics affect  
566 seedling recruitment in alpine habitats. *Journal of Vegetation Science* **25**:1004-1014.

567 Palacio S, Lenz A, Wipf S, Hoch G, Rixen C. 2015. Bud freezing resistance in alpine shrubs  
568 across snow depth gradients. *Environmental and Experimental Botany* **118**:95-101.

569 Pauli JN, Zuckerberg B, Whiteman JP, Porter W. 2013. The subnivium: a deteriorating seasonal  
570 refugium. *Frontiers in Ecology and the Environment* **11**:260-267.

571 Pebesma EJ, Bivand R. 2005. Classes and methods for spatial data in R. *R News* **5**:[https://cran.r-](https://cran.r-project.org/doc/Rnews)  
572 [project.org/doc/Rnews](https://cran.r-project.org/doc/Rnews).

573 Penczykowski RM, Connolly BM, Barton BT. 2017. Winter is changing: trophic interactions  
574 under altered snow regimes. *Food Webs* **13**:80-91.

575 Petty S, Zuckerberg B, Pauli JN. 2015. Winter conditions and land cover structure the  
576 subnivium, a seasonal refuge beneath the snow. *PLoS ONE* **10**:e0127613.

577 Pickering C, Byrne J. 2014. The benefits of publishing systematic quantitative literature reviews  
578 for PhD candidates and other early-career researchers. *Higher Education Research and*  
579 *Development* **33**:534-548.

580 Pruitt WO. 1970. Some ecological aspects of snow in Raup HM, editor. *Ecology of the subarctic*  
581 *regions*. UNESCO, Paris, France.

582 Pullin AS, Stewart GB. 2006. Guidelines for systematic review in conservation and  
583 environmental management. *Conservation Biology* **20**:1647-1656.

584 R Core Team 2016. R: A language and environment for statistical computing. R Foundation for  
585 Statistical Computing, Vienna, Austria. URL: <http://www.R-project.org/>.

586 Räisänen J. 2008. Warmer climate: less or more snow? *Climate Dynamics* **30**:307-319.

587 Roy BA, Güsewell S, Harte J. 2004. Response of plant pathogens and herbivores to a warming  
588 experiment. *Ecology* **85**:2570-2581.

589 Saarinen T, Rasmus S, Lundell R, Kauppinen OK, Hanninen H. 2016. Photosynthetic and  
590 phenological responses of dwarf shrubs to the depth and properties of snow. *Oikos*  
591 **125**:364-373.

592 Saavedra FA, Kampf SK, Fassnacht SR, Sibold JS. 2017. A snow climatology of the Andes  
593 Mountains from MODIS snow cover data. *International Journal of Climatology* **37**:1526-  
594 1539.

595 Sanecki G, Green K, Wood H, Lindenmayer D. 2006a. The characteristics and classification of  
596 Australian snow cover: an ecological perspective. *Arctic, Antarctic, and Alpine Research*  
597 **38**:429-435.

598 Sanecki G, Green K, Wood H, Lindenmayer D. 2006b. The implications of snow-based  
599 recreation for small mammals in the subnivean space in south-east Australia. *Biological*  
600 *Conservation* **129**:511-518.

601 Sinclair BJ, Chown SL. 2005. Climatic variability and hemispheric differences in insect cold  
602 tolerance: support from southern Africa. *Functional Ecology* **19**:214-221.

603 Slatyer RA, Nash MA, Hoffmann AA. 2017. Measuring the effects of reduced snow cover on  
604 Australia's alpine arthropods. *Austral Ecology* **42**:844-857.

605 Steger C, Kotlarski S, Jonas S, C. 2013. Alpine snow cover in a changing climate: a regional  
606 climate model perspective. *Climate Dynamics* **41**:735-754.

607 Steltzer H, Landry C, Painter TH, Anderson J, Ayers E. 2009. Biological consequences of earlier  
608 snowmelt from desert dust deposition in alpine landscapes. *Proceedings of the National*  
609 *Academy of Sciences* **106**:11629-11634.

- 610 Stocker TF, et al. 2013. Technical Summary in Stocker TF, Qin D, Plattner G-K, Tignor M,  
611 Allen SK, Boschung J, Nauels A, Xia Y, Bex V, and Midgley PM, editors. Climate  
612 Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth  
613 Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge  
614 University Press, Cambridge, UK.
- 615 Sturm M, Holmgren J, Liston GE. 1995. A seasonal snow classification system for local and  
616 global applications. *Journal of Climate* **8**:1261-1283.
- 617 Sulkava P, Huhta V. 2003. Effects of hard frost and freeze-thaw cycles on decomposer  
618 communities and N mineralisation in boreal forest soil. *Applied Soil Ecology* **22**:225-  
619 239.
- 620 Tan B, Wu FZ, Yang WQ, He XH. 2014. Snow removal alters soil microbial biomass and  
621 enzyme activity in a Tibetan alpine forest. *Applied Soil Ecology* **76**:34-41.
- 622 Taulavuori K, Bauer E, Taulavuori E. 2011. Overwintering stress of *Vaccinium vitis-idaea* in the  
623 absence of snow cover. *Environmental and Experimental Botany* **72**:397-403.
- 624 Tessier JT. 2017. Importance of depth in soil to corm survival in *Erythronium americanum*  
625 (Liliaceae). *Rhodora* **119**:33-43.
- 626 Vaughan DG, et al. 2013. Observations: Cryosphere in Stocker TF, Qin D, Plattner G-K, Tignor  
627 M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V, and Midgley PM, editors. Climate  
628 Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth  
629 Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge  
630 University Press, Cambridge, UK.
- 631 Vavrus S. 2007. The role of terrestrial snow cover in the climate system. *Climate Dynamics*  
632 **29**:73-88.
- 633 Vrba ES. 2012. Reverse altitudinal cline in cold hardiness among *Erebia* butterflies. *CryoLetters*  
634 **33**:251-258.
- 635 Walther GR, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC, Fromentin JM, Hoegh-  
636 Guldberg O, Bairlein F. 2002. Ecological response to recent climate change. *Nature*  
637 **416**:389-395.

- 638 Wickham H 2016. *ggplot2: elegant graphics for data analysis*. Springer-Verlag, NY, USA.
- 639 Williams CM, Henry HAL, Sinclair BJ. 2015. Cold truths: how winter drives responses of  
640 terrestrial organisms to climate change. *Biological Reviews* **90**:214-235.
- 641 Wipf S, Rixen C. 2010. A review of snow manipulation experiments in Arctic and alpine tundra  
642 ecosystems. *Polar Research* **29**:95-109.
- 643

644 **Supporting Information**

645 **Appendix S1.** Summary of studies identified via review searches.

646 **Appendix S2.** Citation details for all studies included in the review.

647 **Appendix S3.** Summary tables of responses of plants, mammals, and arthropods to reduced  
648 snow conditions.

649 **Data availability**

650 The full database of studies included in the review is available at: [10.6084/m9.figshare.4977998](https://doi.org/10.6084/m9.figshare.4977998)

651 **Tables**

652 **Table 1.** Summary of location and study organism information for original research papers  
 653 examining ecological effects of snow conditions. Percentages given are out of the total number  
 654 of studies (365) and do not always add up to 100 as some studies covered multiple categories.

Category	Total		Category	Total
All papers	365			
<b>Continent/region</b>			<b>Taxonomic/functional group</b>	
Europe	159 (44%)		Plant	241 (66%)
North America	149 (41%)		Small vascular plant	158 (43%)
Asia	40 (11%)		Shrub	72 (20%)
Australia	12 (3%)		Tree	40 (11%)
Oceania	6 (2%)		Bryophyte	21 (6%)
South America	0 (0%)		Animal	131 (36%)
Africa	0 (0%)		Mammal	76 (21%)
			Arthropod	37 (10%)
<b>Climate zone</b>			Bird	16 (4%)
Temperate alpine	157 (43%)		Fish	2 (1%)
Sub-arctic/boreal	112 (31%)		Reptile	1 (< 1%)
Temperate sub-alpine	61 (17%)		Amphibian	1 (< 1%)
Temperate forest	57 (16%)		Lichen	13 (4%)
Temperate grassland	37 (10%)		Fungi	7 (2%)
Sub-Antarctic	0 (0%)			
Tropical alpine	0 (0%)			

655

656 **Table 2.** Summary of methodological approaches used to study the ecological effects of snow  
 657 conditions on plants and animals. Percentages given are out of the total number of studies (365)  
 658 and do not always add up to 100 because some studies covered multiple categories.

<b>Category</b>	<b>Total</b>
<b>Type of study</b>	
Experimental	164 (45%)
Snow depth	114 (31%)
Snow duration	75 (21%)
Snow density	20 (5%)
Snow chemistry	4 (1%)
Observational	212 (58%)
Spatial variation	119 (33%)
Temporal variation	113 (31%)
<b>Timing of measurement</b>	
Summer	309 (85%)
Winter	71 (19%)
Snow-surface	59 (16%)
Intranivean	0 (0%)
Subnivean	17 (5%)

659

660 **Figure legends**

661 **Figure 1.** Some potential effects of changing snow conditions on organisms in seasonally snow-  
662 covered environments. Different colors are indicative of the type of effect (e.g. behavior,  
663 physiology, growth) that the change in snow condition might have.



664 **Figure 2.** Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA:  
665 Moher et al. 2009) flowchart, outlining the process followed to compile the dataset used in the  
666 literature review;  $n$  = number of original research papers (studies).

667 **Figure 3.** Distribution of study sites in relation to snow type and geography. Colors indicate  
668 different snow classifications according to Sturm et al. (1995) and studies included in the review  
669 are shown as orange circles. Snow classification data were obtained from the Atlas of the  
670 Cryosphere (Maurer 2007). Note some regions with seasonal snow, primarily in the southern  
671 hemisphere, do not have a classification according to the system of Sturm et al. (1995).



672 **Figure 4.** Summary of responses of plants, mammals, and arthropods to changes in snow depth  
673 and the timing of snowmelt, based on a simple vote-counting procedure (see Methods). Response  
674 variables are on the left and responses are shown in relation to (a) reduced snow depth and (b)  
675 earlier snowmelt; numbers indicate the number of studies. Light blue shading indicates a higher  
676 value or an earlier occurrence (for autumn/spring phenology) in > 50% of studies; dark blue  
677 shading indicates a lower value or a later occurrence in > 50% of studies. Grey shading indicates  
678 no clear directional response; this could be due to different studies showing results in opposite  
679 directions, individual studies showing mixed results, or individual studies showing no effect of  
680 snow variation on the response variable. Unfilled boxes indicate no studies.






Aspect of snow conditions




Abiotic link





Individual-level effects



Population- & community-level effects







Snow classification

- Tundra
- Taiga
- Maritime
- Prarie
- Alpine
- Ice

**(a) Reduced snow depth**

	Plant	Mammal	Arthropod
Diversity, species richness	14		2

	Plant	Mammal	Arthropod
Growing season density, abundance, relative abundance	14	2	7
Population growth rate	2	3	

	Plant	Mammal	Arthropod
Mortality, injury, damage	23	8	12
Germination/establishment/hatching success	6	1	3
Fecundity/number of progeny	17	3	1
Individual growth rate	17	1	2
Body mass, body size, total biomass	26	5	3

	Plant	Mammal	Arthropod
Spring phenology	22	1	2
Autumn phenology		1	
Phenological overlap (inter- or intra-specific)			
Duration of growing season activity	2		
Winter density, abundance, relative abundance		1	3

**(b) Earlier snowmelt**

	Plant	Mammal	Arthropod
Diversity, species richness	15		1

	Plant	Mammal	Arthropod
Growing season density, abundance, relative abundance	10	3	3
Population growth rate	1		1

	Plant	Mammal	Arthropod
Mortality, injury, damage	17	1	2
Germination/establishment/hatching success	7		
Fecundity/number of progeny	16	4	1
Individual growth rate	24	1	
Body mass, body size, total biomass	21	2	

	Plant	Mammal	Arthropod
Spring phenology	73	3	6
Autumn phenology	6		1
Phenological overlap (inter- or intra-specific)	4		2
Duration of growing season activity	13		1
Winter density, abundance, relative abundance			

Response	
	Increased/advanced
	Mixed
	Decreased/delayed
	No data

Community responses

Diversity, species richness

Population responses

Growing season density, abundance, relative abundance

Population growth rate

Mortality, recruitment, and growth

Mortality, injury, damage

Germination/establishment/hatching success

Fecundity/number of progeny

Individual growth rate

Body mass, body size, total biomass

Phenology

Spring phenology

Autumn phenology

Phenological overlap (inter- or intra-specific)

Duration of growing season activity

Winter density, abundance, relative abundance