

1 **Matters Arising: Response to Crossley et al. 2020**

2

3 **Title**

4 Meta-analyses of insect temporal trends must account for the complex sampling histories  
5 inherent to many long-term monitoring efforts

6

7 **Response to:**

8 Crossley *et al.* 2020. No net insect abundance and diversity declines across US Long Term  
9 Ecological Research sites. *Nature Ecology & Evolution* (2020) doi:10.1038/s41559-020-1269-4.

10

11 **Authors**

12 Ellen A. R. Welte<sup>1\*</sup>, Anthony Joern<sup>2</sup>, Aaron M. Ellison<sup>3</sup>, David C. Lightfoot<sup>4</sup>, Sydne Record<sup>5</sup>,  
13 Nicholas Rodenhouse<sup>6</sup>, Emily H. Stanley<sup>7</sup>, Michael Kaspari<sup>8</sup>

14

15 **Author Affiliations**

16 <sup>1</sup> Senckenberg Research Institute and Natural History Museum Frankfurt, Gelnhausen, Germany

17 <sup>2</sup> Division of Biology, Kansas State University, Manhattan, KS, USA

18 <sup>3</sup> Harvard Forest, Harvard University, Petersham, MA, USA

19 <sup>4</sup> Biology Department, University of New Mexico, Albuquerque, NM, USA

20 <sup>5</sup> Department of Biology, Bryn Mawr College, Bryn Mawr, PA, USA

21 <sup>6</sup> Biological Sciences, Wellesley College, Wellesley, MA, USA

22 <sup>7</sup> Center for Limnology, University of Wisconsin-Madison, Madison, WI, USA

23 <sup>8</sup> Geographical Ecology Group, Department of Biology, University of Oklahoma, Norman, OK,  
24 USA

25 \*corresponding author; email: ellen.welte@senckenberg.de

26

27 **Abstract**


28 In an article recently published in *Nature Ecology & Evolution* (Crossley et al. 2020 “No net  
29 insect abundance and diversity declines across US Long Term Ecological Research sites”),  
30 sampling effort within Long-Term Ecological Research (LTER) datasets was assumed to be  
31 consistent across years. Given the complex history of many long-term datasets at LTER sites,  
32 this assumption does not often hold and we believe this assumption led to errors in Crossley et  
33 al.’s analysis. Here we first use the Konza Prairie grasshopper dataset as an example of how  
34 changes in sampling locations and effort can cause errors when data are assumed to be collected  
35 with invariant sampling. Second, we describe similar and additional errors in data use from 7 of  
36 the 13 LTER sites included in Crossley et al. (2020)’s analysis.

37  
38 **Matters Arising**

39 Crossley et al. (2020)<sup>1</sup> conducted a meta-analysis to examine patterns of change in insect  
40 abundance and diversity across US Long-Term Ecological Research (LTER) sites, concluding “a  
41 lack of overall increase or decline”. This is notable if true, given mixed conclusions in the  
42 literature regarding the nature and ubiquity of insect declines across regions and insect  
43 taxonomic groups<sup>2-6</sup>. The data analyzed, downloaded from and collected by US LTER sites,  
44 represent unique time series of arthropod abundances. These long-term datasets often provide  
45 much needed insights, capturing both steady changes and responses to sudden unpredictable  
46 events. However, they are also rarely uniform in sampling effort across their full duration as a  
47 result of the changing goals and abilities of a research site to collect data<sup>7</sup>. We suggest that  
48 Crossley et al.’s results rely upon a key, but flawed, assumption, that sampling was collected “in  
49 a consistent way over time within each dataset”. We highlight how this incorrect assumption  
50 contributed to errors in their key finding for the Konza Prairie (KNZ) grasshopper dataset  
51 (CGR02), and we describe other errors in Crossley et al.’s data use, listing only errors from  
52 datasets of which either we ourselves are the PIs or we have been able to confirm with the  
53 corresponding LTER PIs and information managers.

54  
55 The KNZ CGR02 dataset documents grasshopper species abundances on 15 KNZ watersheds,  
56 and spans 1982-present (up to 2015 included in Crossley et al. 2020). Crossley et al. analyze  
57 time series of individual species from each dataset (increasing the number of “Time trends” in  
58 their Table 1). At the same time, regardless of variant sampling effort, they regularly sum all  
59 individuals within LTER datasets to yield a single value of abundance for a given species and  
60 year. This is the case for KNZ grasshoppers, and most other included LTER datasets (number of  
61 “Sites” in their Table 1). Importantly, sampling effort at KNZ and other LTER sites was not  
62 constant. At KNZ, the number of watersheds in which grasshoppers were collected, and the  
63 number of samples per watershed, both varied over time (Fig. 1). The number of watersheds  
64 sampled per year has varied from 6-14. Most notably, 6 bison-grazed watersheds were added to  
65 KNZ sampling in 2002. Changes in watersheds sampled over time are documented in the online  
66 metadata (<https://portal.edirepository.org/nis/metadataviewer?packageid=knb-lter-knz.29.12>).

67



68  
 69 **Figure 1. The complex history of sampling of the KNZ grasshopper dataset.** The KNZ  
 70 grasshopper dataset (CGR01) exhibits high variance both in number of watersheds sampled per  
 71 year (number of bars per year) and number of samples collected within each watershed each year  
 72 (depicted in color). Other complexities include the tragic loss of three years (1992-1995) of  
 73 sampling due to a freezer crash, changes in sampling month, changes in watershed burn  
 74 frequencies, and the reintroduction of bison in the 1990s to six of the later-sampled watersheds.

75  
 76 Accounting for KNZ’s complex sampling history matters. Bison-grazed watersheds support  
 77 higher grasshopper abundances and species richness<sup>8,9</sup>. In a recent analysis using the CGR02  
 78 dataset, to account for this change in sampling effort, data were combined only from watersheds  
 79 collected in the same years (e.g. by splitting samples from grazed watersheds into a separate time  
 80 series) and abundances within each watershed and year were divided by the number of samples.  
 81 Analysis of the data structured in this way showed a 2% annual decline in grasshopper  
 82 abundance, with only one common species increasing<sup>10</sup>. Crossley et al., in contrast, report no  
 83 such decline in grasshoppers, and instead report most grasshopper species increased in  
 84 abundance from 1982-2015. The authors of Crossley et al. (2020) note the discrepancy with both  
 85 this study<sup>10</sup> and another meta-analysis<sup>3</sup> and suggest it is “driven by falling numbers of just two  
 86 once-dominant species... whereas many other formerly rare species have become more abundant  
 87 and both evenness and species richness have increased”. However, we believe the discrepancy  
 88 arises because Crossley et al. did not account for variable sampling effort, including KNZ’s  
 89 incorporation of additional, more diverse grazed habitats midway in the time series, leading to  
 90 the perception of increased grasshopper abundances over the full time series record.

91  
 92 We have thus far been able to confirm issues with data used from 7 of the 13 LTER sites  
 93 included in Crossley et al. (2020). Very similar data misuse (i.e. where raw annual abundances  
 94 were summed irrespective of changes in sampling effort and location) was observed for data  
 95 from six additional LTER sites (Cedar Creek, Central Arizona-Phoenix, Harvard Forest,  
 96 Hubbard Brook, North Temperate Lakes, and Sevilleta). Other notable inaccuracies include: the  
 97 inappropriate use of experimental datasets with confounding treatment effects from Cedar Creek,  
 98 Harvard Forest, and North Temperate Lakes; the inclusion of taxa from the KNZ grasshopper  
 99 dataset for the full time series (1982-2015) which were only recorded in the dataset starting in

100 2013, not accounting for multi-year population cycles in time series known to have these  
101 dynamics such as Hubbard Brook Lepidoptera<sup>11</sup> and KNZ grasshoppers<sup>10</sup>, and non-inclusion of a  
102 dataset in the final analysis stated to be included (in Crossley et al. [2020]’s Table 1) from North  
103 Temperate Lakes due to the replacement of all abundance values with zeros. We provide details  
104 about these and other errors in the Supplementary Information.

105  
106 Given these mistakes, we urge skepticism regarding Crossley et al. (2020)’s general conclusion  
107 of no net decline in insect abundances at US LTER sites in recent decades. Although their goal is  
108 laudable, we believe that no conclusion can be reached at this point regarding general trends in  
109 US insect populations, in part due to flaws in their analyses. Failure to take into account  
110 sampling effort in long-term datasets at best will increase measurement error and bias toward a  
111 null result (as was the case with Crossley et al.’s overall conclusion). At worst, when a site’s  
112 sampling effort increases or decreases among years, it can generate even more erroneous  
113 conclusions regarding population change. Recently, a study reporting widespread collapse of  
114 rainforest insect populations at the LTER site Luquillo necessitated a similar correction<sup>5</sup>. We  
115 echo those authors, when they suggest that scientists can avoid such fixable errors by reading  
116 corresponding metadata and contacting the data providers in advance or even including the field  
117 biologists as authors. Like the ecology they document, it is important to take into account that  
118 long-term monitoring efforts by LTERs and similar institutions are themselves complex and full  
119 of history.

120

#### 121 **Author Contributions**

122 E.A.R.W., S. R., A.J., and M.K. conceived the idea for the paper. E.A.R.W. wrote the first draft.  
123 A.M.E., D.L., S.R., N.R., and E.S. identified further errors in the Crossley et al. online data. All  
124 authors significantly contributed to revisions.

125

#### 126 **Acknowledgements**

127 Stevan Earl, Susan Barrott, Elizabeth Borer, and Steve Pennings aided in identifying errors in  
128 Crossley et al.’s online data. Karl Roeder provided comments on an early draft of this  
129 manuscript. Jeff Taylor aided in identifying changes in Konza watershed names. We thank all  
130 LTER information managers and PIs who help keep online metadata updated. We thank Pam  
131 Montz, John Haarstad, Amanda Kuhl, Matthew Ayres, Richard Holmes, and the numerous others  
132 who did the hard work to generate these long-term datasets. NSF DEB-1556280 to MK and  
133 Konza Prairie LTER NSF DEB-1440484 supported this work.

134

#### 135 **Competing Interests**

136 The authors declare no competing interests.

137

#### 138 **Data Availability**

139 KNZ grasshopper abundance data are available from the Long-Term Ecological Research Data  
140 Portal (<https://doi.org/10.6073/pasta/7b2259dcb0e499447e0e11dfb562dc2f>). Links to  
141 additionally described LTER datasets are provided in the Supplementary Information.

142

#### 143 **References**

- 144 1. Crossley, M. S. *et al.* No net insect abundance and diversity declines across US Long  
145 Term Ecological Research sites. *Nature Ecology & Evolution* (2020)  
146 doi:10.1038/s41559-020-1269-4.
- 147 2. Hallmann, C. A. *et al.* More than 75 percent decline over 27 years in total flying insect  
148 biomass in protected areas. *PLoS One* **12**, e0185809 (2017).
- 149 3. van Klink, R. *et al.* Meta-analysis reveals declines in terrestrial but increases in  
150 freshwater insect abundances. *Science* **368**, 417–420 (2020).
- 151 4. Lister, B. C. & Garcia, A. Climate-driven declines in arthropod abundance restructure  
152 a rainforest food web. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E10397–E10406 (2018).
- 153 5. Willig, M. *et al.* Populations are not declining and food webs are not collapsing at the  
154 Luquillo Experimental Forest. *Proceedings of the National Academy of Sciences* **116**,  
155 12143–12144 (2019).
- 156 6. Pilotto, F. *et al.* Meta-analysis of multidecadal biodiversity trends in Europe. *Nature*  
157 *Communications* **11**, 3486 (2020).
- 158 7. Didham, R. K. *et al.* Interpreting insect declines: seven challenges and a way forward.  
159 *Insect Conservation and Diversity* **13**, 103–114 (2020).
- 160 8. Welty, E. A. R. *et al.* Fire, grazing and climate shape plant-grasshopper interactions in  
161 a tallgrass prairie. *Funct. Ecol.* **33**, 735–745 (2019).
- 162 9. Bruckerhoff, L. A. *et al.* Harmony on the prairie? Grassland plant and animal  
163 community responses to variation in climate across land-use gradients. *Ecology* **101**,  
164 e02986 (2020).
- 165 10. Welty, E. A. R., Roeder, K. A., de Beurs, K. M., Joern, A. & Kaspari, M. Nutrient  
166 dilution and climate cycles underlie declines in a dominant insect herbivore. *Proc.*  
167 *Natl. Acad. Sci. U. S. A.* **117**, 7271–7275 (2020).
- 168 11. Martinat, P. J. & Allen, D. C. Saddled prominent outbreaks in North America.  
169 *Northern Journal of Applied Forestry* **5**, 88–91 (1988).
- 170
- 171

172 **Supplementary Information for Matters Arising:**  
173 Meta-analyses of insect temporal trends must account for the complex sampling histories  
174 inherent to many long-term monitoring efforts

175  
176 **Authors:** Ellen A. R. Welte, Anthony Joern, Aaron M. Ellison, David C. Lightfoot, Sydne  
177 Record, Nicholas Rodenhouse, Emily H. Stanley, Michael Kaspari

178  
179

### 180 **Description of data use errors in Crossley et al. (2020)<sup>1</sup>**

181  
182 The errors noted here are all examples of issues that can arise from the use of publicly available  
183 online datasets, without reading metadata or first communicating with the Principal Investigators  
184 (PIs) whose research created the datasets. The National Science Foundation funded LTER  
185 Network has specific data access and user policies (<https://lternet.edu/data-access-policy/>). While  
186 the LTER network strives to make research data publicly available, LTER also urges users of  
187 LTER datasets to contact the PIs of datasets with questions about methodology, and encourages  
188 data users to collaborate with the data authors. None of the authors of datasets described here are  
189 aware of being contacted by the authors of Crossley et al. (2020). We ask readers to read  
190 metadata and communicate or even collaborate with the PIs of publically available datasets that  
191 you intend to use for meta-analysis publications. Following these guidelines improves our ability  
192 to conduct good solid science.

193

194 The most common error we have noted in Crossley et al. (2020)'s use of Long-Term  
195 Ecological Research Network (LTER) data was to use raw annual sums of individuals for entire  
196 LTER datasets, which, combined with variation in sampling effort and location, produced  
197 unreliable estimates of arthropod temporal trends. It is evident that Crossley et al. (2020)'s  
198 analysis did not account for sampling variation because **1)** they state that they considered all  
199 included datasets to have invariant sampling effort, **2)** complex datasets were considered one  
200 time series in their analyses, and **3)** for many datasets it is evident that no averaging occurred  
201 because the sum of all species abundances in their online data  
202 (<https://datadryad.org/stash/dataset/doi:10.5061/dryad.cc2fqz645>) totals the same number as the  
203 total individuals collected within entire LTER datasets.

204

205 We note that Crossley et al. did consider quantifying sampling effort as they include a  
206 column in their online data called "n.obs". However, the corresponding author of Crossley et al.  
207 (2020), Michael Crossley, informed us that n.obs was never used in their analyses. We further  
208 note that even if abundances had been divided by n.obs, this may not appropriately account for  
209 changes in sampling effort/location because: **1)** based on Crossley et al. (2020)'s code, n.obs  
210 does not always capture sample observations correctly (e.g. watershed is not included in the  
211 calculation of n.obs for the Konza grasshopper dataset and n.obs is incorrectly listed as "1" for  
212 all rows of Crossley et al.'s online data for both Central Arizona-Phoenix pitfall datasets, the  
213 Cedar Creek grasshopper dataset, the Hubbard Brook White Mountains Region caterpillar  
214 dataset, the Sevilleta grasshopper dataset, and the Sevilleta pitfall dataset), and **2)** as we have  
215 shown for the Konza grasshopper dataset, that if changes in sampling location correspond to the  
216 gain/loss of habitats that support different species and abundances, it is not appropriate to divide  
217 abundances by the number of sampling locations. In this case, combining sampling locations into

218 one time series is only appropriate if the same sampling locations are sampled for the same  
219 duration of time.

220

221 Here we describe where the assumption of invariant sampling error occurred in Crossley  
222 et al. (2020)'s use of LTER datasets and provide the raw numbers of individuals from each  
223 dataset to more transparently allow others to check our work. We additionally provide  
224 information on other documented errors. We include information only where either we ourselves  
225 are the PIs of these datasets or we have been able to confirm errors with PIs and information  
226 managers from corresponding LTERs. LTER sites are listed in alphabetical order and include  
227 Cedar Creek, Central Arizona-Phoenix, Harvard Forest, Hubbard Brook, Konza Prairie, North  
228 Temperate Lakes, and Sevilleta.

229

230 Cedar Creek

231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267

**1) Grasshopper dataset (ghe014)**

**<https://www.cedarcreek.umn.edu/research/data/dataset?ghe014>,**

- A) No correction was made for variation in sampling effort and changes in sampling locations which are documented in the metadata (<https://www.cedarcreek.umn.edu/research/data/methods?e014>). The sum of all individuals in the Crossley et al. (2020) online data time series (52,116 individuals) is the same as the total individuals collected from the entire ghe014 dataset. The “n.obs” (number of observations) in Crossley et al. (2020)’s online data is set to “1” for all rows.
- B) This dataset is not correctly linked in Supplementary Table 1 and incorrectly described.

**1) Arthropod “Sweep1” dataset (arce153)**

**<https://www.cedarcreek.umn.edu/research/data/dataset?arce153>**

- A) This dataset is not appropriate to answer questions about general insect trends, since it is an experiment including nitrogen addition treatments and herbivore exclosures.
- B) This dataset is not correctly linked in Crossley et al. (2020)’s Supplementary Table 1.

**2) Arthropod “Sweep2” dataset (aage120)**

**<https://www.cedarcreek.umn.edu/research/data/dataset?aage120>**

- A) Either no or an inappropriate correction was made for variation in sampling effort and changes in sampling locations which are documented in the metadata (<https://www.cedarcreek.umn.edu/research/data/methods?e120>). No correction was made if “n.obs”, Crossley et al. (2020)’s calculation of the number of observations, was not used in analyses as asserted by corresponding author Michael Crossley; an inappropriate correction was applied if n.obs was included in the analyses. The sum of all individuals in the Crossley et al. (2020) online data time series (151,227 individuals) is the same as the total individuals collected from the entire dataset (after subtracting the 44,027 unidentified “undet undet” individuals). The calculation of n.obs in Crossley et al. (2020)’s R code does not include plot number, only month and year of observation.
- B) This dataset is not appropriate to answer questions about general insect trends, since it is an experiment with treatments having different levels of plant diversity.
- C) This dataset is not correctly linked and incorrectly described in Crossley et al. (2020)’s Supplementary Table 1.



268 **Central Arizona-Phoenix**

269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303

**1) Arthropod sweep dataset (knb-lter-cap.652.2)**

**<https://doi.org/10.6073/pasta/0669ee6a71b24abb1ae3827f4ee77f6d>**

No correction was made for variation in sampling locations which are documented in the metadata (<https://data.sustainability.asu.edu/cap-portal/metadataviewer?packageid=knb-lter-cap.652.2>). The sum of all individuals in the Crossley et al. (2020) online data time series (34,316) is a similar number to the total individuals (34,323) in the entire dataset. There is 1 individual listed as unidentified but we cannot account for the discrepancy of the 6 remaining individuals. The calculation of n.obs in Crossley et al. (2020)'s R code does not include subsite, only sample date.

**2) Ground arthropod pitfall central Arizona-Phoenix dataset (knb-lter-cap.41.16)**  
**<https://data.sustainability.asu.edu/cap-portal/mapbrowse?packageid=knb-lter-cap.41.16>**

It is likely that no correction was made for variation in sampling effort and changes in sampling locations which are documented in the metadata (<https://data.sustainability.asu.edu/cap-portal/metadataviewer?packageid=knb-lter-cap.41.16>). While we cannot account for the discrepancy between the number of individuals in the full pitfall dataset (2,563,183 individuals) and the number in the Crossley et al. (2020) online data time series (2,529,604 individuals, 98% of those in the full dataset), considering the high variability in subsite number and location per year in this dataset we remain concerned that sampling effort and location were not accounted for. The “n.obs” (number of observations) in Crossley et al. (2020)'s online data is set to “1” for all rows.

**3) Ground arthropod pitfall McDowell dataset (knb-lter-cap.643.2)**

**<https://sustainability.asu.edu/cap/ter/data/view/knb-lter-cap.643.2/>**

No correction was made for variation in sampling effort and changes in sampling locations which are documented in the metadata (<https://data.sustainability.asu.edu/cap-portal/metadataviewer?packageid=knb-lter-cap.643.2>). The sum of all individuals in the Crossley et al. (2020) online data time series (22,360 individuals) is the same as the total individuals collected from the entire dataset from (after subtracting the 1 unidentified “Unknown” individual). The “n.obs” (number of observations) in Crossley et al. (2020)'s online data is set to “1” for all rows.

304 **Harvard Forest**

305

306 **1) Harvard Forest Hemlock Removal Experiment Ant dataset (knb-lter-hfr.118.30)**

307 **<https://doi.org/10.6073/pasta/7a6b956fb0960d7fe8bb048b1fe26956>**

308 **A)** Sampling effort differed among years for the Harvard Forest Hemlock Removal  
309 Experiment (HF-HeRE) dataset analyzed by Crossley et al. (2020). These ants were  
310 collected within a long-term experiment in which there were four plot types: two  
311 controls (intact hemlock and intact mixed hardwood) and two canopy manipulations  
312 (hemlocks girdled and logged).

313 **B)** In the Crossley et al. (2020) analysis this dataset is coded with Locales: “ants.pitfall”,  
314 “ants.bait”, “ants.hand”, and “ants.litter” that represent pitfall trapping, bait sampling  
315 with cookies and tuna fish, hand collections, and sieved litter samples of ants from  
316 2003-2015. Sampling effort differed among years for these different sampling  
317 methods coded as “Locales” in the Crossley et al. (2020) analysis. All four sampling  
318 methods (i.e., “Locales” equal to “ants.pitfall”, “ants.bait”, “ants.litter”, and  
319 “ants.hand” were sampled in June, July, and August from 2003-2005, in July and  
320 August in 2006, and in July only from 2007-2008. From 2009-2015, only pitfall traps  
321 were set within the HF-HeRE. The number of pitfall traps set from 2003-2012 was 25  
322 traps total (situated in a 10 m × 10 m array). In 2012, a deer and moose enclosure was  
323 set up within the experimental plots of the HF-HeRE and an additional 10 m × 10 m  
324 array of 25 pitfall traps was set up within the enclosure (i.e., pitfall trap sampling  
325 effort doubled from 2012-2015 relative to the number of pitfall traps from 2003-2011.  
326 We note that Crossley et al. (2020) do account for the different sampling methods and  
327 that the baits, litter, and hand samples were only collected from 2003-2008, but other  
328 differences in sampling effort were not accounted for. The “n.obs” (number of  
329 observations) in Crossley et al. (2020)’s online data is set to one for all rows in the  
330 dataset, so it does not account for differences in numbers of samples per year.

331 **C)** For the Harvard Forest ant data, Crossley et al. (2020) treat the number of ants  
332 collected by pitfall, bait, and litter samples as raw abundances, which may  
333 overestimate abundance of ants if they happen to occur nearby colonies with actively  
334 foraging workers<sup>2</sup>.

335

336 **2) Nantucket ant dataset (knb-lter-hfr.147.21)**

337 **<https://doi.org/10.6073/pasta/3493424abf9fc36eac7b62b732e4ea55>**

338 **(hf147-09-nantucket-sites-2004-09.csv)**

339 This dataset contains ants sampled with pitfall traps in two bogs and surrounding forests  
340 in 2000 combined with ants sampled from upland habitats from 2004–2009 by a variety  
341 of methods and at different intensities and sites. It also includes “velvet ants”, a group of  
342 ant-mimicking wasps, which were identified only to family (Mutillidae). These data were  
343 collected to assess relationships of ant diversity with habitat and management regime<sup>3</sup>  
344 and cannot be used to analyze temporal trends within a site. Either no or an inappropriate  
345 correction was made for this variation in sampling effort and changes in sampling  
346 locations, all of which are documented in the metadata

347 (<https://portal.lternet.edu/nis/metadataviewer?packageid=knb-lter-hfr.147.21>). No  
348 correction was made if “n.obs”, Crossley et al. (2020)’s calculation of the number of  
349 observations, was not used in analyses as asserted by corresponding author Michael

350 Crossley; an inappropriate correction was applied if n.obs was included in the analyses.  
351 The sum of all individuals in the Crossley et al. (2020) online data time series (32,146  
352 individuals) is the same as the total individuals collected from the entire dataset (after  
353 subtracting the 9 individuals with year listed as “NA” and 2 individuals with species code  
354 listed as “NA”). The calculation of n.obs in Crossley et al. (2020)’s R code does not  
355 include subsite (“site”) or collection method, only community type (habitat description),  
356 month, and year of observation.

### 3) Tick dataset (knb-lter-hfr.299.3)

357  
358 <https://doi.10.6073/pasta/b29a97941c11ddf45540ea30066fde35>

359 **A)** These data are collected with student time sheets for payroll to raise awareness of tick  
360 bites for students in the Harvard Forest Summer Research Program in Ecology. The  
361 tick survey is voluntary, has variable response rates each year depending on the group  
362 of students, and generally shows a decline in collection intensity during the summer  
363 as students increasingly fail to report weekly data. The summer of 2019 also had a  
364 much lower response rate because the program switched to using digital, rather than  
365 paper, time sheets. The calculation of n.obs (for this dataset coded as “n.y1”) in  
366 Crossley et al. (2020)’s R code is the sum of hours reported by the tick survey, which  
367 is the number of hours worked during the day when the student found a tick on their  
368 body. This number does not represent the response rate of the survey, which would  
369 need to be accounted for to address differences in samples per year.

370 **B)** For the tick data, Crossley et al. (2020) analyze 30 separate time series based on the  
371 locations of collection, but it is not clear how those locations were delineated. Many  
372 of the “location.names” from this dataset have overlap as they are filled in with text  
373 by students in the survey form. For instance, the “location.names” of “Harvard  
374 Forest” in the survey overlaps with many possible locations listed by students (e.g.,  
375 “greenhouse”, “Prospect Hill”, “Shaler Hall”). Thus, it is not appropriate to analyze  
376 these data as separate time series as they refer, in some instances, to the same general  
377 location.  
378

### 4) Carnivorous plant prey dataset (knb-lter-hfr.111.16)

379  
380 <https://doi.org/10.6073/pasta/cb95637eda0f96c3fdbd1a97e632c7b7>

381 These data were from a global review of arthropod prey spectra of carnivorous plants<sup>4</sup>.  
382 None of the data were collected at Harvard Forest (and most were collected on other  
383 continents), and for each carnivorous plant species, “year” indicates the year the data  
384 were published and no time-series (repeat collection) was observed or implied by the data  
385 or discussed in the review. Although these data were not included in the final analysis of  
386 Crossley et al. (2020), rows for these data are listed in Crossley et al. (2020)’s online data  
387 (<https://datadryad.org/stash/dataset/doi:10.5061/dryad.cc2fqz645>;  
388 External\_Database\_S1\_PerSpecies\_Abundance\_LTER\_annotated.csv) and all abundance  
389 values are listed as zero. The rows corresponding to this dataset (lines 28497 – 36898:  
390 8401 records) were inaccurately included in Crossley et al. (2020)’s count of 82,777  
391 observations (the number of rows in their online data:  
392 External\_Database\_S1\_PerSpecies\_Abundance\_LTER\_annotated.csv).  
393  
394

395 5) We also note more generally that for the Harvard Forest datasets that the environmental  
396 data are all for the Harvard Forest site in Petersham in central Massachusetts, but the  
397 Nantucket dataset should report different environmental data as it was collected from an  
398 island off eastern Massachusetts that has very different climate from central  
399 Massachusetts. Furthermore, the locations in the tick dataset, which are each given a  
400 different time series should also have location specific environmental data as locations of  
401 data collections were variable (e.g., most in western MA at Harvard Forest, but some in  
402 Connecticut; Cambridge, MA; etc.).  
403

404 **Hubbard Brook**

405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429

**1) Lepidoptera datasets (knb-lter-hbr.82.8)**

**<https://doi.org/10.6073/pasta/5d2a8c67c5a3278032b2b14d66c09a7f>**

- A) Sampling effort differed among plots and years for one of the two Hubbard Brook datasets analyzed by Crossley et al. (2020). These data, coded as Locale: “Lepidoptera1”, represent visual counts of caterpillars on one plot, 1986-2018. Consistent sampling effort occurred throughout this time series; however, sampling effort differed among both plots and years for the dataset coded as Locale “Lepidoptera2”. This second dataset spans 1986-1995 and represents three different plots in the White Mountains Region that are located outside of the Hubbard Brook valley. The sum of all individuals in the Crossley et al. (2020) online data time series (4,030 individuals) is the same as the total individuals collected from the entire White Mountains Region caterpillar dataset. The “n.obs” (number of observations) in Crossley et al. (2020)’s online data is set to “1” for all rows for the White Mountains Region dataset.
- B) Crossley et al. do not acknowledge that these data include Lepidoptera identified only to the family level as noted in the online metadata (<https://portal.edirepository.org/nis/mapbrowse?scope=knb-lter-hbr&identifier=82>).
- C) The caterpillar populations documented in these data exhibit outbreaks at long intervals (e.g., 10-13 years apart<sup>5</sup>), limiting the ability of linear trend analysis to detect meaningful trends with time series of shorter lengths (10-33 years for Hubbard Brook data used in Crossley et al.(2020)).
- D) Only one of the two Hubbard Brook datasets analyzed is described in Crossley et al. (2020)’s Supplementary Table 1.

430 **Konza Prairie**

431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466

**1) Grasshopper dataset (CGR02)**

**<https://doi.org/10.6073/pasta/7b2259dcb0e499447e0e11dfb562dc2f>**

- A) Either no or an inappropriate correction was made for variation in sampling effort and changes in sampling locations which are documented in the metadata (<https://portal.edirepository.org/nis/metadataviewer?packageid=knb-lter-knz.29.12>). No correction was made if “n.obs”, Crossley et al. (2020)’s calculation of the number of observations, was not used in analyses as was asserted by corresponding author Michael Crossley; an inappropriate correction was applied if n.obs was included in the analyses. The sum of all individuals in the Crossley et al. (2020) online data time series (121,229 individuals) is the same as the total individuals collected from the entire CGR02 dataset from 1982-2015 (after subtracting the 459 unidentified “unknown” individual grasshoppers). The calculation of n.obs in Crossley et al. (2020)’s R code does not include watershed, only month, day, and the replicate code within the watershed (“a” or “b”).
- B) Three taxa (Tettigoniidae, Oecanthinae, and Gryllidae) included in Crossley et. al. (2020)’s analysis for the full duration (1982-2015) were only recorded in the KNZ dataset starting in 2013.
- C) The grasshopper populations documented in these data exhibit cycles at ~5 year intervals, limiting the ability of linear trend analysis to detect meaningful trends with time series of shorter lengths<sup>6</sup>.

**2) Gall insects (CGP01)**

**<http://dx.doi.org/10.6073/pasta/b2ac9e918a66dbbb18c7a6b39dc1efab>**

Either no or an inappropriate correction was made for variation in sampling locations and plant species sampled which are documented in the metadata (<https://portal.edirepository.org/nis/metadataviewer?packageid=knb-lter-knz.27.11>). No correction was made if “n.obs”, Crossley et al. (2020)’s calculation of the number of observations, was not used in analyses as was asserted by corresponding author Michael Crossley; an inappropriate correction was applied if n.obs was included in the analyses. The sum of all individuals in the Crossley et al. (2020) online data time series (27,819 galled stems) is the same as the total galled stems in the entire CGP01 dataset. The calculation of n.obs (for this dataset coded as “n.y1”) in Crossley et al. (2020)’s R code does not include watershed or account for the different plant species sampled and only accounts for the number of sampled stems.

467 **North Temperate Lakes**

468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512

**1) Benthic macroinvertebrate dataset (knb-lter-ntl.11.34)**

<https://doi.org/10.6073/pasta/1bad728523ce4c39ade38fa666a59aee>

- A) Likely due to program R being case sensitive, the time series for Sparkling Lake which was coded both “SP” and “sp” was accidentally split into two time series with “sp” having non-zero values only in 2016-2017. However, Crossley et al. (2020) considered Locale “sp” a separate time series spanning 1981-2017.
- B) No correction was made for variation in sampling effort and changes in sampling locations if “n.obs”, Crossley et al. (2020)’s calculation of the number of observations, was not used in analyses as was asserted by corresponding author Michael Crossley. While we have not been able to identify why there is a discrepancy between the number of individuals in the Crossley et al. (2020) online data time series (126,041 individuals) and those in the full dataset (140,100 individuals), if n.obs was not included in trend calculation, sampling effort changes were not accounted for as the full time series (1981-2017) was included in Crossley et al. (2020)’s analysis, even though some lakes did not have sampling in all years.

**2) Pelagic macroinvertebrate dataset (knb-lter-ntl.14.30)**

<https://doi.org/10.6073/pasta/cc25694cdde49853271df465a15007fb>

While listed in Crossley et al. (2020)’s Table 1 and Supplemental Table 1, these data do not appear to be included in Crossley et al.’s online data, nor are they referenced in Crossley et al. (2020)’s online R code.

**3) Crayfish dataset (knb-lter-ntl.3.28)**

<https://doi.org/10.6073/pasta/61619e749daf99c71a289dcadafb795c>

While included in Crossley et al.’s online time series data (“External\_Database\_S1\_PerSpecies\_Abundance\_LTER\_annotated”), all abundance values are listed as zero. No entries from this dataset are listed in Crossley et al.’s online trend data (“External\_Database\_S2\_time\_trends\_arthropods\_relaxed”), thus these data were not included in Crossley et al. (2020)’s final analysis.

**4) Crayfish dataset (knb-lter-ntl.217.9)**

<https://doi.org/10.6073/pasta/4a22c4b3707f68ba5c03cc3ed70e98b6>

- A) This dataset has an incorrect link listed in Crossley et al. (2020)’s Supplementary Table 1. We were able to reconstruct which dataset was used by matching total sums between North Temperate Lakes crayfish datasets and finding identical yearly sums between the Crossley et al. (2020) online data and knb-lter-ntl.217.9 for 2001-2010 (both totaling 95,066 individuals for this duration).
- B) Crossley et al. (2020) online data for this dataset contains data from 2011, when none exists in the dataset.
- C) This dataset is not appropriate to answer questions about general arthropod trends, since it contains data on an experiment of crayfish removal.
- D) This dataset documents an invasive crayfish species, and therefore may not be appropriate for understanding general arthropod population trends.

513 **Sevilleta**

514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552

**1) Grasshopper dataset (sev-106)**

**<https://doi.org/10.6073/pasta/c1d40e9d0ec610bb74d02741e9d22576>**

No correction was made for changes in sampling locations (termination of the pinyon-juniper [Goat Draw] vegetation type sampling site, and the initiation of a new [Blue Grama] sampling site at SEV in 2002) which are documented in the metadata (<https://portal.edirepository.org/nis/metadataviewer?packageid=knb-lter-sev.106.152976>). The sum of all individuals in the Crossley et al. (2020) online data time series (36,634 individuals) is the same number as total individuals in the entire sev-106 dataset. The “n.obs” (number of observations) in Crossley et al. (2020)’s online data is set to “1” for all rows. The added Blue Grama site had considerably higher numbers of grasshoppers than the old pinyon-juniper site, and this change in sampling location likely inflated the numbers of grasshoppers in Crossley et al. (2020)’s calculation starting in 2002.

**2) Ground arthropod dataset (sev-29)**

**<https://doi.org/10.6073/pasta/9e7e6dc9c9d8f72e9e0bca07a1e76ccd>**

No correction was made for changes in sampling locations (termination of the pinyon-juniper [Goat Draw] vegetation type sampling site, and the initiation of a new [Blue Grama] sampling site at SEV in 2002) which are documented in the metadata (<https://portal.edirepository.org/nis/metadataviewer?packageid=knb-lter-sev.29.175390>). Collection of ground arthropods for this dataset also varied in number of traps per collection period/subsite. Some traps in each sample set of 3 subsample traps were often omitted from data tabulation due to individual traps being disturbed by precipitation runoff, or vertebrate animals. Summing omitted subsample traps (missing values, not zeros) would have reduced the sum counts for a line of 3 traps. While we cannot account for the discrepancy between the number of individuals in the Crossley et al. (2020) online data time series (39,926 individuals) and those in the full sev-29 dataset (52,188 individuals identified to genus level), this discrepancy appears to arise from Crossley et al. (2020)’s analysis removing species (they state 365 species/morphospecies were used in their Table 1) from the sev-29 full dataset (we count 433 species/morphospecies identified to genus). Crossley et al. did include some taxa identified to genus but not species in their analysis, so the reason for removing species/morphospecies is unclear. The “n.obs” (number of observations) in Crossley et al. (2020)’s online data is set to “1” for all rows. The added Blue Grama site had considerably higher numbers of ground arthropods than the old pinyon-juniper site, and this change in sampling location likely inflated the numbers of ground arthropods in Crossley et al. (2020)’s calculation starting in 2002.



553 **Supplementary Information References**

554

- 555 1. Crossley, M. S. *et al.* No net insect abundance and diversity declines across US Long Term  
556 Ecological Research sites. *Nature Ecology & Evolution* (2020) doi:10.1038/s41559-020-1269-  
557 4.
- 558 2. Gotelli, N. J., Ellison, A. M., Dunn, R. R. & Sanders, N. J. Counting ants (Hymenoptera:  
559 Formicidae): biodiversity sampling and statistical analysis for myrmecologists.  
560 *Myrmecological News* **15**, 13–19 (2011).
- 561 3. Ellison, A. M. The ants of Nantucket: Unexpectedly high biodiversity in an anthropogenic  
562 landscape. *Northeastern Naturalist* **19**, 43–66 (2012).
- 563 4. Ellison, A. M. & Gotelli, N. J. Energetics and the evolution of carnivorous plants—Darwin’s  
564 ‘most wonderful plants in the world’. *J Exp Bot* **60**, 19–42 (2009).
- 565 5. Martinat, P. J. & Allen, D. C. Saddled prominent outbreaks in North America. *Northern*  
566 *Journal of Applied Forestry* **5**, 88–91 (1988).
- 567 6. Welti, E. A. R., Roeder, K. A., de Beurs, K. M., Joern, A. & Kaspari, M. Nutrient dilution and  
568 climate cycles underlie declines in a dominant insect herbivore. *Proc. Natl. Acad. Sci. U. S. A.*  
569 **117**, 7271–7275 (2020).

570