

1 **Matters Arising: Response to Crossley et al. 2020**

2

3 **Title**

4 Meta-analyses of insect temporal trends must account for the complex sampling histories
5 inherent to many long-term monitoring efforts

6

7 **Response to:**

8 Crossley *et al.* 2020. No net insect abundance and diversity declines across US Long Term
9 Ecological Research sites. *Nature Ecology & Evolution* (2020) doi:10.1038/s41559-020-1269-4.

10

11 **Authors**

12 Ellen A. R. Welte^{1*}, Anthony Joern², Aaron M. Ellison³, David C. Lightfoot⁴, Sydne Record⁵,
13 Nicholas Rodenhouse⁶, Emily H. Stanley⁷, Michael Kaspari⁸

14

15 **Author Affiliations**

16 ¹ Senckenberg Research Institute and Natural History Museum Frankfurt, Gelnhausen, Germany

17 ² Division of Biology, Kansas State University, Manhattan, KS, USA

18 ³ Harvard Forest, Harvard University, Petersham, MA, USA

19 ⁴ Biology Department, University of New Mexico, Albuquerque, NM, USA

20 ⁵ Department of Biology, Bryn Mawr College, Bryn Mawr, PA, USA

21 ⁶ Biological Sciences, Wellesley College, Wellesley, MA, USA

22 ⁷ Center for Limnology, University of Wisconsin-Madison, Madison, WI, USA

23 ⁸ Geographical Ecology Group, Department of Biology, University of Oklahoma, Norman, OK,
24 USA

25 *corresponding author; email: ellen.welte@senckenberg.de

26

27 **Abstract**


28 In an article recently published in *Nature Ecology & Evolution* (Crossley et al. 2020 “No net
29 insect abundance and diversity declines across US Long Term Ecological Research sites”),
30 sampling effort within Long-Term Ecological Research (LTER) datasets was assumed to be
31 consistent across years. Given the complex history of many long-term datasets at LTER sites,
32 this assumption does not often hold and we believe this assumption led to errors in Crossley et
33 al.’s analysis. Here we first use the Konza Prairie grasshopper dataset as an example of how
34 changes in sampling locations and effort can cause errors when data are assumed to be collected
35 with invariant sampling. Second, we describe similar and additional errors in data use from 7 of
36 the 13 LTER sites included in Crossley et al. (2020)’s analysis.

37
38 **Matters Arising**

39 Crossley et al. (2020)¹ conducted a meta-analysis to examine patterns of change in insect
40 abundance and diversity across US Long-Term Ecological Research (LTER) sites, concluding “a
41 lack of overall increase or decline”. This is notable if true, given mixed conclusions in the
42 literature regarding the nature and ubiquity of insect declines across regions and insect
43 taxonomic groups²⁻⁶. The data analyzed, downloaded from and collected by US LTER sites,
44 represent unique time series of arthropod abundances. These long-term datasets often provide
45 much needed insights, capturing both steady changes and responses to sudden unpredictable
46 events. However, they are also rarely uniform in sampling effort across their full duration as a
47 result of the changing goals and abilities of a research site to collect data⁷. We suggest that
48 Crossley et al.’s results rely upon a key, but flawed, assumption, that sampling was collected “in
49 a consistent way over time within each dataset”. We highlight how this incorrect assumption
50 contributed to errors in their key finding for the Konza Prairie (KNZ) grasshopper dataset
51 (CGR02), and we describe other errors in Crossley et al.’s data use, listing only errors from
52 datasets of which either we ourselves are the PIs or we have been able to confirm with the
53 corresponding LTER PIs and information managers.

54
55 The KNZ CGR02 dataset documents grasshopper species abundances on 15 KNZ watersheds,
56 and spans 1982-present (up to 2015 included in Crossley et al. 2020). Crossley et al. analyze
57 time series of individual species from each dataset (increasing the number of “Time trends” in
58 their Table 1). At the same time, regardless of variant sampling effort, they regularly sum all
59 individuals within LTER datasets to yield a single value of abundance for a given species and
60 year. This is the case for KNZ grasshoppers, and most other included LTER datasets (number of
61 “Sites” in their Table 1). Importantly, sampling effort at KNZ and other LTER sites was not
62 constant. At KNZ, the number of watersheds in which grasshoppers were collected, and the
63 number of samples per watershed, both varied over time (Fig. 1). The number of watersheds
64 sampled per year has varied from 6-14. Most notably, 6 bison-grazed watersheds were added to
65 KNZ sampling in 2002. Changes in watersheds sampled over time are documented in the online
66 metadata (<https://portal.edirepository.org/nis/metadataviewer?packageid=knb-lter-knz.29.12>).

67



68
 69 **Figure 1. The complex history of sampling of the KNZ grasshopper dataset.** The KNZ
 70 grasshopper dataset (CGR01) exhibits high variance both in number of watersheds sampled per
 71 year (number of bars per year) and number of samples collected within each watershed each year
 72 (depicted in color). Other complexities include the tragic loss of three years (1992-1995) of
 73 sampling due to a freezer crash, changes in sampling month, changes in watershed burn
 74 frequencies, and the reintroduction of bison in the 1990s to six of the later-sampled watersheds.
 75

76 Accounting for KNZ’s complex sampling history matters. Bison-grazed watersheds support
 77 higher grasshopper abundances and species richness^{8,9}. In a recent analysis using the CGR02
 78 dataset, to account for this change in sampling effort, data were combined only from watersheds
 79 collected in the same years (e.g. by splitting samples from grazed watersheds into a separate time
 80 series) and abundances within each watershed and year were divided by the number of samples.
 81 Analysis of the data structured in this way showed a 2% annual decline in grasshopper
 82 abundance, with only one common species increasing¹⁰. Crossley et al., in contrast, report no
 83 such decline in grasshoppers, and instead report most grasshopper species increased in
 84 abundance from 1982-2015. The authors of Crossley et al. (2020) note the discrepancy with both
 85 this study¹⁰ and another meta-analysis³ and suggest it is “driven by falling numbers of just two
 86 once-dominant species... whereas many other formerly rare species have become more abundant
 87 and both evenness and species richness have increased”. However, we believe the discrepancy
 88 arises because Crossley et al. did not account for variable sampling effort, including KNZ’s
 89 incorporation of additional, more diverse grazed habitats midway in the time series, leading to
 90 the perception of increased grasshopper abundances over the full time series record.
 91

92 We have thus far been able to confirm issues with data used from 7 of the 13 LTER sites
 93 included in Crossley et al. (2020). Very similar data misuse (i.e. where raw annual abundances
 94 were summed irrespective of changes in sampling effort and location) was observed for data
 95 from six additional LTER sites (Cedar Creek, Central Arizona-Phoenix, Harvard Forest,
 96 Hubbard Brook, North Temperate Lakes, and Sevilleta). Other notable inaccuracies include: the
 97 inappropriate use of experimental datasets with confounding treatment effects from Cedar Creek,
 98 Harvard Forest, and North Temperate Lakes; the inclusion of taxa from the KNZ grasshopper
 99 dataset for the full time series (1982-2015) which were only recorded in the dataset starting in

100 2013, not accounting for multi-year population cycles in time series known to have these
101 dynamics such as Hubbard Brook Lepidoptera¹¹ and KNZ grasshoppers¹⁰, and non-inclusion of a
102 dataset in the final analysis stated to be included (in Crossley et al. [2020]’s Table 1) from North
103 Temperate Lakes due to the replacement of all abundance values with zeros. We provide details
104 about these and other errors in the Supplementary Information.

105
106 Given these mistakes, we urge skepticism regarding Crossley et al. (2020)’s general conclusion
107 of no net decline in insect abundances at US LTER sites in recent decades. Although their goal is
108 laudable, we believe that no conclusion can be reached at this point regarding general trends in
109 US insect populations, in part due to flaws in their analyses. Failure to take into account
110 sampling effort in long-term datasets at best will increase measurement error and bias toward a
111 null result (as was the case with Crossley et al.’s overall conclusion). At worst, when a site’s
112 sampling effort increases or decreases among years, it can generate even more erroneous
113 conclusions regarding population change. Recently, a study reporting widespread collapse of
114 rainforest insect populations at the LTER site Luquillo necessitated a similar correction⁵. We
115 echo those authors, when they suggest that scientists can avoid such fixable errors by reading
116 corresponding metadata and contacting in advance or even including as authors the data
117 providers. Like the ecology they document, it is important to take into account that long-term
118 monitoring efforts by LTERs and similar institutions are themselves complex and full of history.

119
120 **Author Contributions**
121 E.A.R.W., S. R., A.J., and M.K. conceived the idea for the paper. E.A.R.W. wrote the first draft.
122 A.M.E., D.L., S.R., N.R., and E.S. identified further errors in the Crossley et al. online data. All
123 authors significantly contributed to revisions.

124
125 **Acknowledgements**
126 Stevan Earl, Susan Barrott, Elizabeth Borer, and Steve Pennings aided in identifying errors in
127 Crossley et al.’s online data. Karl Roeder provided comments on an early draft of this
128 manuscript. Jeff Taylor aided in identifying changes in Konza watershed names. We thank all
129 LTER information managers and PIs who help keep online metadata updated. We thank Pam
130 Montz, John Haarstad, Amanda Kuhl, Matthew Ayres, Richard Holmes, and the numerous others
131 who did the hard work to generate these long-term datasets. NSF DEB-1556280 to MK and
132 Konza Prairie LTER NSF DEB-1440484 supported this work.

133
134 **Competing Interests**
135 The authors declare no competing interests.

136
137 **Data Availability**
138 KNZ grasshopper abundance data are available from the Long-Term Ecological Research Data
139 Portal (<https://doi.org/10.6073/pasta/7b2259dcb0e499447e0e11dfb562dc2f>). Links to
140 additionally described LTER datasets are provided in the Supplementary Information.

141
142 **References**
143 1. Crossley, M. S. *et al.* No net insect abundance and diversity declines across US Long
144 Term Ecological Research sites. *Nature Ecology & Evolution* (2020)
145 doi:10.1038/s41559-020-1269-4.

- 146 2. Hallmann, C. A. *et al.* More than 75 percent decline over 27 years in total flying insect
147 biomass in protected areas. *PLoS One* **12**, e0185809 (2017).
- 148 3. van Klink, R. *et al.* Meta-analysis reveals declines in terrestrial but increases in
149 freshwater insect abundances. *Science* **368**, 417–420 (2020).
- 150 4. Lister, B. C. & Garcia, A. Climate-driven declines in arthropod abundance restructure
151 a rainforest food web. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E10397–E10406 (2018).
- 152 5. Willig, M. *et al.* Populations are not declining and food webs are not collapsing at the
153 Luquillo Experimental Forest. *Proceedings of the National Academy of Sciences* **116**,
154 12143–12144 (2019).
- 155 6. Pilotto, F. *et al.* Meta-analysis of multidecadal biodiversity trends in Europe. *Nature*
156 *Communications* **11**, 3486 (2020).
- 157 7. Didham, R. K. *et al.* Interpreting insect declines: seven challenges and a way forward.
158 *Insect Conservation and Diversity* **13**, 103–114 (2020).
- 159 8. Welte, E. A. R. *et al.* Fire, grazing and climate shape plant-grasshopper interactions in
160 a tallgrass prairie. *Funct. Ecol.* **33**, 735–745 (2019).
- 161 9. Bruckerhoff, L. A. *et al.* Harmony on the prairie? Grassland plant and animal
162 community responses to variation in climate across land-use gradients. *Ecology* **101**,
163 e02986 (2020).
- 164 10. Welte, E. A. R., Roeder, K. A., de Beurs, K. M., Joern, A. & Kaspari, M. Nutrient
165 dilution and climate cycles underlie declines in a dominant insect herbivore. *Proc.*
166 *Natl. Acad. Sci. U. S. A.* **117**, 7271–7275 (2020).
- 167 11. Martinat, P. J. & Allen, D. C. Saddled prominent outbreaks in North America.
168 *Northern Journal of Applied Forestry* **5**, 88–91 (1988).
- 169
170

171 **Supplementary Information for Matters Arising:**
172 Meta-analyses of insect temporal trends must account for the complex sampling histories
173 inherent to many long-term monitoring efforts

174
175 **Authors:** Ellen A. R. Welti, Anthony Joern, Aaron M. Ellison, David C. Lightfoot, Sydne
176 Record, Nicholas Rodenhouse, Emily H. Stanley, Michael Kaspari

177
178

179 **Description of data use errors in Crossley et al. (2020)¹**

180
181 The errors noted here are all examples of issues that can arise from the use of publicly available
182 online datasets, without reading metadata or first communicating with the Principal Investigators
183 (PIs) whose research created the datasets. The National Science Foundation funded LTER
184 Network has specific data access and user policies (<https://lternet.edu/data-access-policy/>). While
185 the LTER network strives to make research data publicly available, LTER also urges users of
186 LTER datasets to contact the PIs of datasets with questions about methodology, and encourages
187 data users to collaborate with the data authors. None of the authors of datasets described here are
188 aware of being contacted by the authors of Crossley et al. (2020). We ask readers to read
189 metadata and communicate or even collaborate with the PIs of publically available datasets that
190 you intend to use for meta-analysis publications. Following these guidelines improves our ability
191 to conduct good solid science.

192

193 The most common error we have noted in Crossley et al. (2020)'s use of Long-Term
194 Ecological Research Network (LTER) data was to use raw annual sums of individuals for entire
195 LTER datasets, which, combined with variation in sampling effort and location, produced
196 unreliable estimates of arthropod temporal trends. It is evident that Crossley et al. (2020)'s
197 analysis did not account for sampling variation because **1)** they state that they considered all
198 included datasets to have invariant sampling effort, **2)** complex datasets were considered one
199 time series in their analyses, and **3)** for many datasets it is evident that no averaging occurred
200 because the sum of all species abundances in their online data
201 (<https://datadryad.org/stash/dataset/doi:10.5061/dryad.cc2fqz645>) totals the same number as the
202 total individuals collected within entire LTER datasets.

203

204 We note that Crossley et al. did consider quantifying sampling effort as they include a
205 column in their online data called "n.obs". However, the corresponding author of Crossley et al.
206 (2020), Michael Crossley, informed us that n.obs was never used in their analyses. We further
207 note that even if abundances had been divided by n.obs, this may not appropriately account for
208 changes in sampling effort/location because: **1)** based on Crossley et al. (2020)'s code, n.obs
209 does not always capture sample observations correctly (e.g. watershed is not included in the
210 calculation of n.obs for the Konza grasshopper dataset and n.obs is incorrectly listed as "1" for
211 all rows of Crossley et al.'s online data for both Central Arizona-Phoenix pitfall datasets, the
212 Cedar Creek grasshopper dataset, the Hubbard Brook White Mountains Region caterpillar
213 dataset, the Sevilleta grasshopper dataset, and the Sevilleta pitfall dataset), and **2)** as we have
214 shown for the Konza grasshopper dataset, that if changes in sampling location correspond to the
215 gain/loss of habitats that support different species and abundances, it is not appropriate to divide
216 abundances by the number of sampling locations. In this case, combining sampling locations into

217 one time series is only appropriate if the same sampling locations are sampled for the same
218 duration of time.

219

220 Here we describe where the assumption of invariant sampling error occurred in Crossley
221 et al. (2020)'s use of LTER datasets and provide the raw numbers of individuals from each
222 dataset to more transparently allow others to check our work. We additionally provide
223 information on other documented errors. We include information only where either we ourselves
224 are the PIs of these datasets or we have been able to confirm errors with PIs and information
225 managers from corresponding LTERs. LTER sites are listed in alphabetical order and include
226 Cedar Creek, Central Arizona-Phoenix, Harvard Forest, Hubbard Brook, Konza Prairie, North
227 Temperate Lakes, and Sevilleta.

228

229 Cedar Creek

230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266

1) Grasshopper dataset (ghe014)

<https://www.cedarcreek.umn.edu/research/data/dataset?ghe014>,

- A) No correction was made for variation in sampling effort and changes in sampling locations which are documented in the metadata (<https://www.cedarcreek.umn.edu/research/data/methods?e014>). The sum of all individuals in the Crossley et al. (2020) online data time series (52,116 individuals) is the same as the total individuals collected from the entire ghe014 dataset. The “n.obs” (number of observations) in Crossley et al. (2020)’s online data is set to “1” for all rows.
- B) This dataset is not correctly linked in Supplementary Table 1 and incorrectly described.

1) Arthropod “Sweep1” dataset (arce153)

<https://www.cedarcreek.umn.edu/research/data/dataset?arce153>

- A) This dataset is not appropriate to answer questions about general insect trends, since it is an experiment including nitrogen addition treatments and herbivore exclosures.
- B) This dataset is not correctly linked in Crossley et al. (2020)’s Supplementary Table 1.

2) Arthropod “Sweep2” dataset (aage120)

<https://www.cedarcreek.umn.edu/research/data/dataset?aage120>

- A) Either no or an inappropriate correction was made for variation in sampling effort and changes in sampling locations which are documented in the metadata (<https://www.cedarcreek.umn.edu/research/data/methods?e120>). No correction was made if “n.obs”, Crossley et al. (2020)’s calculation of the number of observations, was not used in analyses as asserted by corresponding author Michael Crossley; an inappropriate correction was applied if n.obs was included in the analyses. The sum of all individuals in the Crossley et al. (2020) online data time series (151,227 individuals) is the same as the total individuals collected from the entire dataset (after subtracting the 44,027 unidentified “undet undet” individuals). The calculation of n.obs in Crossley et al. (2020)’s R code does not include plot number, only month and year of observation.
- B) This dataset is not appropriate to answer questions about general insect trends, since it is an experiment with treatments having different levels of plant diversity.
- C) This dataset is not correctly linked and incorrectly described in Crossley et al. (2020)’s Supplementary Table 1.

267 **Central Arizona-Phoenix**

268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302

1) Arthropod sweep dataset (knb-lter-cap.652.2)

<https://doi.org/10.6073/pasta/0669ee6a71b24abb1ae3827f4ee77f6d>

No correction was made for variation in sampling locations which are documented in the metadata (<https://data.sustainability.asu.edu/cap-portal/metadataviewer?packageid=knb-lter-cap.652.2>). The sum of all individuals in the Crossley et al. (2020) online data time series (34,316) is a similar number to the total individuals (34,323) in the entire dataset. There is 1 individual listed as unidentified but we cannot account for the discrepancy of the 6 remaining individuals. The calculation of n.obs in Crossley et al. (2020)'s R code does not include subsite, only sample date.

2) Ground arthropod pitfall central Arizona-Phoenix dataset (knb-lter-cap.41.16)
<https://data.sustainability.asu.edu/cap-portal/mapbrowse?packageid=knb-lter-cap.41.16>

It is likely that no correction was made for variation in sampling effort and changes in sampling locations which are documented in the metadata (<https://data.sustainability.asu.edu/cap-portal/metadataviewer?packageid=knb-lter-cap.41.16>). While we cannot account for the discrepancy between the number of individuals in the full pitfall dataset (2,563,183 individuals) and the number in the Crossley et al. (2020) online data time series (2,529,604 individuals, 98% of those in the full dataset), considering the high variability in subsite number and location per year in this dataset we remain concerned that sampling effort and location were not accounted for. The “n.obs” (number of observations) in Crossley et al. (2020)'s online data is set to “1” for all rows.

3) Ground arthropod pitfall McDowell dataset (knb-lter-cap.643.2)

<https://sustainability.asu.edu/cap/ter/data/view/knb-lter-cap.643.2/>

No correction was made for variation in sampling effort and changes in sampling locations which are documented in the metadata (<https://data.sustainability.asu.edu/cap-portal/metadataviewer?packageid=knb-lter-cap.643.2>). The sum of all individuals in the Crossley et al. (2020) online data time series (22,360 individuals) is the same as the total individuals collected from the entire dataset from (after subtracting the 1 unidentified “Unknown” individual). The “n.obs” (number of observations) in Crossley et al. (2020)'s online data is set to “1” for all rows.

303 **Harvard Forest**

304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348

1) Harvard Forest Hemlock Removal Experiment Ant dataset (knb-lter-hfr.118.30)
<https://doi.org/10.6073/pasta/7a6b956fb0960d7fe8bb048b1fe26956>

- A) Sampling effort differed among years for the Harvard Forest Hemlock Removal Experiment (HF-HeRE) dataset analyzed by Crossley et al. (2020). These ants were collected within a long-term experiment in which there were four plot types: two controls (intact hemlock and intact mixed hardwood) and two canopy manipulations (hemlocks girdled and logged).
- B) In the Crossley et al. (2020) analysis this dataset is coded with Locales: “ants.pitfall”, “ants.bait”, “ants.hand”, and “ants.litter” that represent pitfall trapping, bait sampling with cookies and tuna fish, hand collections, and sieved litter samples of ants from 2003-2015. Sampling effort differed among years for these different sampling methods coded as “Locales” in the Crossley et al. (2020) analysis. All four sampling methods (i.e., “Locales” equal to “ants.pitfall”, “ants.bait”, “ants.litter”, and “ants.hand” were sampled in June, July, and August from 2003-2005, in July and August in 2006, and in July only from 2007-2008. From 2009-2015, only pitfall traps were set within the HF-HeRE. The number of pitfall traps set from 2003-2012 was 25 traps total (situated in a 10 m × 10 m array). In 2012, a deer and moose enclosure was set up within the experimental plots of the HF-HeRE and an additional 10 m × 10 m array of 25 pitfall traps was set up within the enclosure (i.e., pitfall trap sampling effort doubled from 2012-2015 relative to the number of pitfall traps from 2003-2011. We note that Crossley et al. (2020) do account for the different sampling methods and that the baits, litter, and hand samples were only collected from 2003-2008, but other differences in sampling effort were not accounted for. The “n.obs” (number of observations) in Crossley et al. (2020)’s online data is set to one for all rows in the dataset, so it does not account for differences in numbers of samples per year.
- C) For the Harvard Forest ant data, Crossley et al. (2020) treat the number of ants collected by pitfall, bait, and litter samples as raw abundances, which may overestimate abundance of ants if they happen to occur nearby colonies with actively foraging workers².

2) Nantucket ant dataset (knb-lter-hfr.147.21)
<https://doi.org/10.6073/pasta/3493424abf9fc36eac7b62b732e4ea55>
(hf147-09-nantucket-sites-2004-09.csv)

This dataset contains ants sampled with pitfall traps in two bogs and surrounding forests in 2000 combined with ants sampled from upland habitats from 2004–2009 by a variety of methods and at different intensities and sites. It also includes “velvet ants”, a group of ant-mimicking wasps, which were identified only to family (Mutillidae). These data were collected to assess relationships of ant diversity with habitat and management regime³ and cannot be used to analyze temporal trends within a site. Either no or an inappropriate correction was made for this variation in sampling effort and changes in sampling locations, all of which are documented in the metadata (<https://portal.lternet.edu/nis/metadataviewer?packageid=knb-lter-hfr.147.21>). No correction was made if “n.obs”, Crossley et al. (2020)’s calculation of the number of observations, was not used in analyses as asserted by corresponding author Michael

349 Crossley; an inappropriate correction was applied if n.obs was included in the analyses.
350 The sum of all individuals in the Crossley et al. (2020) online data time series (32,146
351 individuals) is the same as the total individuals collected from the entire dataset (after
352 subtracting the 9 individuals with year listed as “NA” and 2 individuals with species code
353 listed as “NA”). The calculation of n.obs in Crossley et al. (2020)’s R code does not
354 include subsite (“site”) or collection method, only community type (habitat description),
355 month, and year of observation.

356 **3) Tick dataset (knb-lter-hfr.299.3)**

357 **[https://doi.10.6073/pasta/ b29a97941c11ddf45540ea30066fde35](https://doi.10.6073/pasta/b29a97941c11ddf45540ea30066fde35)**

358 **A)** These data are collected with student time sheets for payroll to raise awareness of tick
359 bites for students in the Harvard Forest Summer Research Program in Ecology. The
360 tick survey is voluntary, has variable response rates each year depending on the group
361 of students, and generally shows a decline in collection intensity during the summer
362 as students increasingly fail to report weekly data. The summer of 2019 also had a
363 much lower response rate because the program switched to using digital, rather than
364 paper, time sheets. The calculation of n.obs (for this dataset coded as “n.y1”) in
365 Crossley et al. (2020)’s R code is the sum of hours reported by the tick survey, which
366 is the number of hours worked during the day when the student found a tick on their
367 body. This number does not represent the response rate of the survey, which would
368 need to be accounted for to address differences in samples per year.

369 **B)** For the tick data, Crossley et al. (2020) analyze 30 separate time series based on the
370 locations of collection, but it is not clear how those locations were delineated. Many
371 of the “location.names” from this dataset have overlap as they are filled in with text
372 by students in the survey form. For instance, the “location.names” of “Harvard
373 Forest” in the survey overlaps with many possible locations listed by students (e.g.,
374 “greenhouse”, “Prospect Hill”, “Shaler Hall”). Thus, it is not appropriate to analyze
375 these data as separate time series as they refer, in some instances, to the same general
376 location.
377

378 **4) Carnivorous plant prey dataset (knb-lter-hfr.111.16)**

379 **<https://doi.org/10.6073/pasta/cb95637eda0f96c3fdbd1a97e632c7b7>**

380 These data were from a global review of arthropod prey spectra of carnivorous plants⁴.
381 None of the data were collected at Harvard Forest (and most were collected on other
382 continents), and for each carnivorous plant species, “year” indicates the year the data
383 were published and no time-series (repeat collection) was observed or implied by the data
384 or discussed in the review. Although these data were not included in the final analysis of
385 Crossley et al. (2020), rows for these data are listed in Crossley et al. (2020)’s online data
386 (<https://datadryad.org/stash/dataset/doi:10.5061/dryad.cc2fqz645>;
387 External_Database_S1_PerSpecies_Abundance_LTER_annotated.csv) and all abundance
388 values are listed as zero. The rows corresponding to this dataset (lines 28497 – 36898:
389 8401 records) were inaccurately included in Crossley et al. (2020)’s count of 82,777
390 observations (the number of rows in their online data:
391 External_Database_S1_PerSpecies_Abundance_LTER_annotated.csv).

392

393

394 5) We also note more generally that for the Harvard Forest datasets that the environmental
395 data are all for the Harvard Forest site in Petersham in central Massachusetts, but the
396 Nantucket dataset should report different environmental data as it was collected from an
397 island off eastern Massachusetts that has very different climate from central
398 Massachusetts. Furthermore, the locations in the tick dataset, which are each given a
399 different time series should also have location specific environmental data as locations of
400 data collections were variable (e.g., most in western MA at Harvard Forest, but some in
401 Connecticut; Cambridge, MA; etc.).
402

403 **Hubbard Brook**

404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428

1) Lepidoptera datasets (knb-lter-hbr.82.8)

<https://doi.org/10.6073/pasta/5d2a8c67c5a3278032b2b14d66c09a7f>

- A) Sampling effort differed among plots and years for one of the two Hubbard Brook datasets analyzed by Crossley et al. (2020). These data, coded as Locale: “Lepidoptera1”, represent visual counts of caterpillars on one plot, 1986-2018. Consistent sampling effort occurred throughout this time series; however, sampling effort differed among both plots and years for the dataset coded as Locale “Lepidoptera2”. This second dataset spans 1986-1995 and represents three different plots in the White Mountains Region that are located outside of the Hubbard Brook valley. The sum of all individuals in the Crossley et al. (2020) online data time series (4,030 individuals) is the same as the total individuals collected from the entire White Mountains Region caterpillar dataset. The “n.obs” (number of observations) in Crossley et al. (2020)’s online data is set to “1” for all rows for the White Mountains Region dataset.
- B) Crossley et al. do not acknowledge that these data include Lepidoptera identified only to the family level as noted in the online metadata (<https://portal.edirepository.org/nis/mapbrowse?scope=knb-lter-hbr&identifier=82>).
- C) The caterpillar populations documented in these data exhibit outbreaks at long intervals (e.g., 10-13 years apart⁵), limiting the ability of linear trend analysis to detect meaningful trends with time series of shorter lengths (10-33 years for Hubbard Brook data used in Crossley et al.(2020)).
- D) Only one of the two Hubbard Brook datasets analyzed is described in Crossley et al. (2020)’s Supplementary Table 1.

429 **Konza Prairie**

430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465

1) Grasshopper dataset (CGR02)

<https://doi.org/10.6073/pasta/7b2259dcb0e499447e0e11dfb562dc2f>

- A) Either no or an inappropriate correction was made for variation in sampling effort and changes in sampling locations which are documented in the metadata (<https://portal.edirepository.org/nis/metadataviewer?packageid=knb-lter-knz.29.12>). No correction was made if “n.obs”, Crossley et al. (2020)’s calculation of the number of observations, was not used in analyses as was asserted by corresponding author Michael Crossley; an inappropriate correction was applied if n.obs was included in the analyses. The sum of all individuals in the Crossley et al. (2020) online data time series (121,229 individuals) is the same as the total individuals collected from the entire CGR02 dataset from 1982-2015 (after subtracting the 459 unidentified “unknown” individual grasshoppers). The calculation of n.obs in Crossley et al. (2020)’s R code does not include watershed, only month, day, and the replicate code within the watershed (“a” or “b”).
- B) Three taxa (Tettigoniidae, Oecanthinae, and Gryllidae) included in Crossley et. al. (2020)’s analysis for the full duration (1982-2015) were only recorded in the KNZ dataset starting in 2013.
- C) The grasshopper populations documented in these data exhibit cycles at ~5 year intervals, limiting the ability of linear trend analysis to detect meaningful trends with time series of shorter lengths⁶.

2) Gall insects (CGP01)

<http://dx.doi.org/10.6073/pasta/b2ac9e918a66dbbb18c7a6b39dc1efab>

Either no or an inappropriate correction was made for variation in sampling locations and plant species sampled which are documented in the metadata (<https://portal.edirepository.org/nis/metadataviewer?packageid=knb-lter-knz.27.11>). No correction was made if “n.obs”, Crossley et al. (2020)’s calculation of the number of observations, was not used in analyses as was asserted by corresponding author Michael Crossley; an inappropriate correction was applied if n.obs was included in the analyses. The sum of all individuals in the Crossley et al. (2020) online data time series (27,819 galled stems) is the same as the total galled stems in the entire CGP01 dataset. The calculation of n.obs (for this dataset coded as “n.y1”) in Crossley et al. (2020)’s R code does not include watershed or account for the different plant species sampled and only accounts for the number of sampled stems.

466 **North Temperate Lakes**

467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511

1) Benthic macroinvertebrate dataset (knb-lter-ntl.11.34)

<https://doi.org/10.6073/pasta/1bad728523ce4c39ade38fa666a59aee>

- A) Likely due to program R being case sensitive, the time series for Sparkling Lake which was coded both “SP” and “sp” was accidentally split into two time series with “sp” having non-zero values only in 2016-2017. However, Crossley et al. (2020) considered Locale “sp” a separate time series spanning 1981-2017.
- B) No correction was made for variation in sampling effort and changes in sampling locations if “n.obs”, Crossley et al. (2020)’s calculation of the number of observations, was not used in analyses as was asserted by corresponding author Michael Crossley. While we have not been able to identify why there is a discrepancy between the number of individuals in the Crossley et al. (2020) online data time series (126,041 individuals) and those in the full dataset (140,100 individuals), if n.obs was not included in trend calculation, sampling effort changes were not accounted for as the full time series (1981-2017) was included in Crossley et al. (2020)’s analysis, even though some lakes did not have sampling in all years.

2) Pelagic macroinvertebrate dataset (knb-lter-ntl.14.30)

<https://doi.org/10.6073/pasta/cc25694cdde49853271df465a15007fb>

While listed in Crossley et al. (2020)’s Table 1 and Supplemental Table 1, these data do not appear to be included in Crossley et al.’s online data, nor are they referenced in Crossley et al. (2020)’s online R code.

3) Crayfish dataset (knb-lter-ntl.3.28)

<https://doi.org/10.6073/pasta/61619e749daf99c71a289dcadafb795c>

While included in Crossley et al.’s online time series data (“External_Database_S1_PerSpecies_Abundance_LTER_annotated”), all abundance values are listed as zero. No entries from this dataset are listed in Crossley et al.’s online trend data (“External_Database_S2_time_trends_arthropods_relaxed”), thus these data were not included in Crossley et al. (2020)’s final analysis.

4) Crayfish dataset (knb-lter-ntl.217.9)

<https://doi.org/10.6073/pasta/4a22c4b3707f68ba5c03cc3ed70e98b6>

- A) This dataset has an incorrect link listed in Crossley et al. (2020)’s Supplementary Table 1. We were able to reconstruct which dataset was used by matching total sums between North Temperate Lakes crayfish datasets and finding identical yearly sums between the Crossley et al. (2020) online data and knb-lter-ntl.217.9 for 2001-2010 (both totaling 95,066 individuals for this duration).
- B) Crossley et al. (2020) online data for this dataset contains data from 2011, when none exists in the dataset.
- C) This dataset is not appropriate to answer questions about general arthropod trends, since it contains data on an experiment of crayfish removal.
- D) This dataset documents an invasive crayfish species, and therefore may not be appropriate for understanding general arthropod population trends.

512 **Sevilleta**

513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551

1) Grasshopper dataset (sev-106)

<https://doi.org/10.6073/pasta/c1d40e9d0ec610bb74d02741e9d22576>

No correction was made for changes in sampling locations (termination of the pinyon-juniper [Goat Draw] vegetation type sampling site, and the initiation of a new [Blue Grama] sampling site at SEV in 2002) which are documented in the metadata (<https://portal.edirepository.org/nis/metadataviewer?packageid=knb-lter-sev.106.152976>). The sum of all individuals in the Crossley et al. (2020) online data time series (36,634 individuals) is the same number as total individuals in the entire sev-106 dataset. The “n.obs” (number of observations) in Crossley et al. (2020)’s online data is set to “1” for all rows. The added Blue Grama site had considerably higher numbers of grasshoppers than the old pinyon-juniper site, and this change in sampling location likely inflated the numbers of grasshoppers in Crossley et al. (2020)’s calculation starting in 2002.

2) Ground arthropod dataset (sev-29)

<https://doi.org/10.6073/pasta/9e7e6dc9c9d8f72e9e0bca07a1e76ccd>

No correction was made for changes in sampling locations (termination of the pinyon-juniper [Goat Draw] vegetation type sampling site, and the initiation of a new [Blue Grama] sampling site at SEV in 2002) which are documented in the metadata (<https://portal.edirepository.org/nis/metadataviewer?packageid=knb-lter-sev.29.175390>). Collection of ground arthropods for this dataset also varied in number of traps per collection period/subsite. Some traps in each sample set of 3 subsample traps were often omitted from data tabulation due to individual traps being disturbed by precipitation runoff, or vertebrate animals. Summing omitted subsample traps (missing values, not zeros) would have reduced the sum counts for a line of 3 traps. While we cannot account for the discrepancy between the number of individuals in the Crossley et al. (2020) online data time series (39,926 individuals) and those in the full sev-29 dataset (52,188 individuals identified to genus level), this discrepancy appears to arise from Crossley et al. (2020)’s analysis removing species (they state 365 species/morphospecies were used in their Table 1) from the sev-29 full dataset (we count 433 species/morphospecies identified to genus). Crossley et al. did include some taxa identified to genus but not species in their analysis, so the reason for removing species/morphospecies is unclear. The “n.obs” (number of observations) in Crossley et al. (2020)’s online data is set to “1” for all rows. The added Blue Grama site had considerably higher numbers of ground arthropods than the old pinyon-juniper site, and this change in sampling location likely inflated the numbers of ground arthropods in Crossley et al. (2020)’s calculation starting in 2002.

552 **Supplementary Information References**

553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569

1. Crossley, M. S. *et al.* No net insect abundance and diversity declines across US Long Term Ecological Research sites. *Nature Ecology & Evolution* (2020) doi:10.1038/s41559-020-1269-4.
2. Gotelli, N. J., Ellison, A. M., Dunn, R. R. & Sanders, N. J. Counting ants (Hymenoptera: Formicidae): biodiversity sampling and statistical analysis for myrmecologists. *Myrmecological News* **15**, 13–19 (2011).
3. Ellison, A. M. The ants of Nantucket: Unexpectedly high biodiversity in an anthropogenic landscape. *Northeastern Naturalist* **19**, 43–66 (2012).
4. Ellison, A. M. & Gotelli, N. J. Energetics and the evolution of carnivorous plants—Darwin’s ‘most wonderful plants in the world’. *J Exp Bot* **60**, 19–42 (2009).
5. Martinat, P. J. & Allen, D. C. Saddled prominent outbreaks in North America. *Northern Journal of Applied Forestry* **5**, 88–91 (1988).
6. Welti, E. A. R., Roeder, K. A., de Beurs, K. M., Joern, A. & Kaspari, M. Nutrient dilution and climate cycles underlie declines in a dominant insect herbivore. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 7271–7275 (2020).