

1 **Hunters as citizen scientists: contributions to biodiversity monitoring**

2 **in Europe**

3 Benjamin Cretois¹; John DC Linnell²; Matthew Grainger²; Erlend B Nilsen²; Jan Ketil Rød¹

4 ¹Department of Geography, Norwegian University of Science and Technology, 7491 Trondheim,

5 Norway

6 ²Norwegian Institute for Nature Research, PO Box 5685, Torgard, 7485 Trondheim, Norway

7

8

9

10

11

12

13

14

15

16

17 Abstract

- 18 1. Monitoring biodiversity characteristics at large scales and with adequate resolution
19 requires considerable effort and resources. Overall, there is clearly a huge scope for
20 European hunters, a special and often overlooked group of citizen scientist, to
21 contribute even more to biodiversity monitoring, especially because of their presence
22 across the entire European landscape.
- 23 2. Using the Essential Biodiversity Variables (EBVs) framework we reviewed the published
24 and grey literature and contacted experts to provide a comprehensive overview of
25 hunters' contributions to biodiversity monitoring. We examined the methods used to
26 collect data in hunter-based monitoring, the geographic and taxonomic extent of such
27 contributions and the scientific output stemming from hunter-based monitoring data.
- 28 3. Our study suggests that hunter-based monitoring is widely distributed across Europe
29 and across taxa as 32 out of the 36 European countries included in our analysis involve
30 hunters in the monitoring of at least one species group with ungulates and small game
31 species groups which have the widest hunter-based monitoring coverage. We found
32 that it is possible to infer characteristics on Genetic composition, Species population,
33 Species traits and Community composition with data that are being routinely collected
34 by hunters in at least some countries. The main types of data provided are hunting bags
35 data, Biological samples including carcasses of shot animals and non-invasive samplings
36 and observations for counts and indices.
- 37 4. Hunters collect data on biodiversity in its key dimensions, collaborations between
38 hunters and scientists are fruitful and should be considered a standard partnership for

39 biodiversity conservation. To overcome the challenges in the use of hunters' data, more
40 rigorous protocols for sampling data should be implemented and improvements made
41 in data integration methods.

42 **Keywords:** citizen science; hunter; biodiversity monitoring; Europe; Essential
43 biodiversity variables (EBV); biodiversity conservation; mammal; birds; game species

44

45

46

47

48

49

50

51

52

53

54

55

56

57 1. Introduction

58 Global biodiversity is undergoing severe declines (Díaz et al., 2019). This situation has led the
59 international community to take action to alter this trend by setting policy frameworks and
60 objectives. For example, the Aichi Biodiversity Targets set by the Convention on Biological
61 Diversity (UN General Assembly, 1992) and the United Nations Sustainable Development Goals
62 (UN General Assembly, 2015) are globally accepted frameworks which set targets for progress
63 toward a more sustainable world. Biodiversity monitoring is an essential component of
64 measuring progress towards these goals. However, monitoring biodiversity at large scales and
65 with adequate resolution requires considerable effort and resources and represents a logistical
66 challenge for researchers. This is one driver behind the recent enthusiasm towards the
67 involvement of the public in the data collection process (Silvertown, 2009).

68 Citizen science, here defined as the involvement of citizens in scientific research and knowledge
69 production, has repeatedly demonstrated its ability to gather massive amounts of data at a
70 spatial scale unattainable by professional scientists alone (Silvertown, 2009). Even though some
71 citizen science projects are able to provide data the quality of which equals expert-based data,
72 most citizen science biodiversity programs focus solely on species abundance and distribution,
73 limiting its use for assessing international biodiversity targets (Kosmala et al., 2016; Chandler et
74 al., 2017). Other concerns of citizen science include observational biases such as ‘false
75 absences’ or misidentification or uneven spatial and temporal coverage. This raises concerns
76 when making inference using this kind of data, despite the fact that advances in modelling
77 based on such data are currently being made (Hochachka et al., 2012). The extent to which

78 citizen science dataset are biased depends on both the sampling regime used in citizen science
79 programs and the expertise of the recorder which can reduce some of the biases mentioned
80 (e.g. bird watchers data are less clustered around urban areas; Geldmann et al., 2016; Isaac &
81 Pocock, 2015).

82 Here we study the monitoring of biodiversity characteristics by volunteer citizen scientists
83 taking the special, and often overlooked, case of European hunters. For the purposes of this
84 review we focus on the monitoring activities that hunters engage in that are specific to their
85 hunting activity. We do not include other citizen science activities that they might engage in
86 outside of hunting. Hunters collect data during their activity both voluntarily through
87 cooperation with veterinary or other research institutes, and as a parts of compulsory programs
88 when countries' hunting regulations mandate such reporting through hunting statistics or the
89 collection of other data (see for instance <http://artemis-face.eu/> for an overview of the
90 European hunting bag regulations; Mörner et al., 2014). To a large extent hunter collected data
91 is formally institutionalised into wildlife management structures that are intended to support
92 sustainable harvest. Virtually the entire European landscape is utilised for some form of
93 hunting, and most hunting systems are tied to some form of property rights that ensure a broad
94 distribution of hunters across the whole landscape (Linnell et al., 2015). These factors combined
95 make Europe's estimated 7 million hunters a potentially valuable resource for citizen science
96 data collection (www.face.eu/).

97 In this study we use the Essential Biodiversity Variables (EBVs) framework to categorise the
98 different types of data coming from hunter-based monitoring and hence assess their

99 contribution to biodiversity monitoring. EBVs are a set of variables that aim to represent
100 biodiversity across its key dimensions (space, time and biological organisation) and that can
101 accurately document biodiversity change (Kissling et al., 2018). EBVs are being defined and
102 refined by GEO BON, a global biodiversity network that contributes to effective management
103 policies for biodiversity. They provide a first level of aggregation computed from the raw data
104 and can be used to compute more complex biodiversity indicators that can be used to measure
105 the achievement of policy goals (Pereira et al., 2013). GEO BON has divided the EBVs into 22
106 candidates grouped among 6 classes (i.e. genetic composition, species population, species
107 traits, community composition, ecosystem functioning and ecosystem structure,
108 www.geobon.org/ebvs).

109 We aim at providing a comprehensive overview of hunters' contributions to biodiversity
110 monitoring in Europe, and review the methods used to collect data in hunter-based monitoring.
111 We also examine the geographic and taxonomic extent of such contributions and the scientific
112 output stemming from hunter-based monitoring data.

113 2. Method

114 2.1. Systematic literature search

115 The first step of the review process was to define the scope of research that focuses on the
116 research question (Booth et al., 2016). In the present study we aimed at identifying which
117 Essential Biodiversity Variables are possible to derive using hunter-based monitoring. We
118 initially developed a list of keywords listing the actor (i.e. hunter or hunting team), the full list of

119 EBVs as defined by GEO BON, the taxonomic scope and the geographic scope. The refinement
120 of this list was done in an iterative fashion, running the list of keywords through Scopus and
121 Web of Science Core Collection and adding new keywords that emerged, and then re-running
122 the search until we reached a plateau in the number of papers returned by the databases (for
123 the full list of keywords see Document A1 in Appendix).

124 Before any screening the search string returned 1335 papers that we exported to create a
125 dataset. The dataset was initially reduced to 962 papers after screening for duplicates. The
126 search returned many papers that were outside the scope of our study and that concerned
127 anthropological studies on hunter gatherers, studies on hunter-based monitoring outside
128 Europe, or sociological studies on hunters such as hunters' perception of management
129 decisions or hunters' willingness to contribute to species monitoring. We excluded these
130 studies after screening for titles and abstracts and reduced our dataset to 493 papers. If doubts
131 remained regarding the potential contribution of a paper to our study, we kept it in our dataset
132 for final screening. We finally screened the entire papers and rejected studies in which hunters'
133 contribution was unclear such as if hunters were only mentioned in the acknowledgement of
134 the papers or if it was unclear how hunter-based monitoring was used to compute the EBV.
135 After this final step we retained a total of 277 papers (Figure A1 in Appendix). The screening
136 process was facilitated using the R package 'revtools' (Westgate, 2019).

137 2.2. Non-systematic literature search

138 The non-systematic search of our study was divided in two parts; a targeted search on Web of
139 Science Core Collection and Scopus and a search in the grey literature. Literature collection

140 databases such as Scopus and Web of Science only screen through the title, the abstract and
141 keywords. Because hunters' contribution to data collection is in some cases only mentioned in
142 the method section of peer reviewed papers, we expected the systematic literature search to
143 return incomplete information. To be able to include such sources of information, we used
144 'snow-ball sampling' (Goodman, 1961), whereby we sampled the references found in the
145 systematic literature search searching for certain authors or countries that we suspected were
146 commonly using hunters' as their main data providers. This added 89 additional unique papers
147 to our dataset.

148 Secondly, we manually accessed a sample of the proceedings of the International Union of
149 Game Biologists (IUGB). This sample was restricted to the volumes available from our own
150 institutional archives and, did not constitute the whole collection. The included volumes but
151 spanned a time period ranging from 1957 to 2011. In total we analyzed 14 out of the 35 existing
152 proceedings. Some of the articles were written in a language that no authors in this paper were
153 able to read (i.e. Russian or German) and were directly excluded. We selected papers based on
154 the inclusion and exclusion criteria previously defined and the search in the IUGB resulted in
155 the addition of 92 papers to the dataset.

156 2.3. Expert knowledge

157 To further complete our search of hunters' contribution to wildlife monitoring, we sent an
158 email query to experts, wildlife managers and national hunting associations in different
159 European countries. The list of informants was primarily based on the authors' professional
160 networks, but in some cases we were redirected to more competent contacts they personally

161 knew. We first sent a small questionnaire asking if they were aware of any hunter-based
162 monitoring programs or other monitoring schemes involving hunters and if they could provide
163 documentations (i.e. scientific papers or other documents such as official reports or links to
164 websites), for which species and which method was used (Document A2 in Appendix). If
165 answers were unclear, we contacted the informant and asked for more details. A total of 28
166 informants returned a total of 89 inputs including 23 published papers used to complete our
167 dataset.

168 2.4. Classification of EBVs

169 Following the recommendations given by Stewart et al. (2005) and based on the pre-defined
170 keywords, we compiled a database constituted of inputs found in different sources i) a
171 systematic search of the peer-reviewed literature, ii) a non-systematic gathering of literature,
172 including 'snow-ball' search of the scientific and technical grey-literature (which included our
173 own knowledge and libraries), and iii) a survey among our professional networks (i.e. experts in
174 wildlife management or wildlife research, and among national hunting associations).

175 For each input provided we documented the country, the species, the EBVs computed from the
176 data collected by hunters, the methods used by hunters to collect the data and the source of
177 the case. If there was any doubt about the EBV computed we referred to the 'Measurement
178 and scalability' section of each EBV candidate on the GEO-BON website (<https://geobon.org/>)
179 to find the EBV that most proximately the data collected by hunters (e.g. if jaw bone length
180 was measured, we deduced the EBV candidate 'morphology' was used, although with the
181 implicit understanding that this was ultimately used as a metric to monitor demography / life

182 history). To facilitate the analysis and interpretation we pooled species into five broad
183 functional groups: “ungulates”, “large carnivores”, “waterfowl”, “other birds” and “small game”
184 (including lagomorphs and medium sized carnivores, Table A1 in Appendix for the full
185 description of each group). We aggregated the diverse data collection methods into eight
186 categories based on the criteria shown in Table 1, namely “Bag”, “Camera trap”, “Carcasses”,
187 “Direct Observations”, “Indirect observations”, “Questionnaire”, “Ringing” and “Others”.

188 3. Results

189 3.1 Geographic and taxonomic extent of hunter-based monitoring

190 Our study suggests that hunter-based monitoring is widely distributed across Europe and across
191 taxa (Figure 1). We found that 32 out of 36 countries involve hunters in the monitoring of at
192 least one species group. With respect to the species group present in those countries we found
193 that 16 countries use hunter-based monitoring for all potential species groups (UK, Ireland, the
194 Netherlands and Iceland do not host any populations of large carnivores). In four countries
195 (Albania, Kosovo, Macedonia and Liechtenstein), we did not find any published evidence of an
196 organised hunter involvement in the monitoring of game species.

197 Based on our review, ungulates and small game are the species groups which have the widest
198 hunter-based monitoring coverage, as nearly all European countries (80% for ungulates and
199 86% for small game; Figure 1) involve hunters in the monitoring of these species. Even though
200 geographically widespread, we found that waterfowl and large carnivores are the groups which
201 receive less attention from hunter-based monitoring coverage as 63% and 66% respectively of

202 European countries use some sort of hunter-based monitoring for these groups (Table A2)). We
203 did not find any hunter-based monitoring scheme for large carnivores in countries such as
204 France or Lithuania, even though they have population of large carnivores (Linnell & Cretois,
205 2018).

206 3.2. Diversity of biodiversity characteristics recorded by hunter-based monitoring

207 Overall, we found that a wide range of biodiversity characteristics are being derived from
208 hunter-based monitoring programs (Figure 2). In fact, our study suggests that researchers and
209 wildlife managers infer characteristics on Genetic composition, Species population, Species
210 traits and Community composition with data that are being routinely collected by hunters. We
211 did not find any evidence of hunter-based monitoring schemes directly gathering information
212 on Ecosystem function and Ecosystem structure.

213 Other game birds and small game were the taxonomic groups for which hunter-based
214 monitoring was the most diverse, with 79% of the EBV candidates (excluding Ecosystem
215 function and Ecosystem structure) being monitored in at least some countries. Hunter based
216 monitoring for other groups of species was less diverse, with 64% of EBVs being recorded for
217 ungulates and 57% for large carnivores and waterfowl (Table A2).

218 For all species groups, all characteristics concerning their population (i.e. species distribution,
219 population abundance and structure) were recorded except for ungulates, for which we did not
220 find any inputs explicitly documenting the monitoring of their distribution. Nevertheless, it has
221 been argued that species distribution can be inferred from species abundance (Kéry & Royle,

222 2015), hence it is an implicit by-product of other monitoring. Characteristics at the individual
223 level were also very well monitored by hunters, and examples of monitoring of traits such as
224 physiology, morphology and reproduction were found for all species groups. Even traits that are
225 normally hard to obtain with traditional citizen science such as species phenology and
226 movement were monitored by hunters for 4 out of the 5 species groups.

227 Our results also suggest that studies use hunters' data for monitoring of genetic composition
228 through studies on allelic diversity and studies on co-ancestry for large carnivores, ungulates,
229 small game and other game birds.

230 3.3 Methods used to obtain species characteristics

231 Hunters contribute to the collection of data relevant for monitoring in many ways, which vary
232 greatly with respect to data volume, coverage and quality (Table 2). The main types of data
233 provided include;

234 **Hunting bags:** Information on the numbers of individual animals of different species that are
235 killed by hunters is recorded in most countries, although the spatial resolution of the
236 information varies. Under assumptions of more or less similar effort and similar quotas,
237 between year variation in the numbers of animals shot is being used to infer broad scale spatio-
238 temporal changes in abundance and thus species demographic attributes (e.g. Aebischer, 2019;
239 Massei et al., 2015), especially if combined with secondary data sources (Moleón et al., 2008;
240 2013; van der Jeugd & Kwak, 2017). As well as being used to follow single population trends,
241 the analysis of such data from multiple populations is used to map changes in distribution and
242 elucidate the relative impacts of multiple drivers of population change (Hagen et al., 2014;

243 Grøtan et al., 2005; Reimoser et al., 2014). We also found that hunting bag data are being used
244 to infer species interaction characteristics through studying fluctuations in small game hunting
245 bags (Smedshaug, 2018).

246 **Biological samples including carcasses of shot animals and non-invasive samplings:** Shot
247 animals are used to yield a wide variety of information relevant for monitoring. For mammals,
248 this demographic data is made even more valuable when animals can be aged from tooth
249 sectioning. Data on age and sex can be used to infer population structure and survival rates via
250 analyses like life-tables or population reconstruction (Nilsen et al., 2012; Solberg et al., 1999)
251 and for spatial population structure (Swenson et al., 1998; Kojola & Laitala, 2000). Data on
252 reproduction can be obtained from the analysis of reproductive organs. Body weights and
253 measurement of jawbones or femurs are used to infer body size and condition. Bird wings are
254 used to infer age and sex of animals killed (Pöysä & Väänänen, 2018). Tissue samples are also
255 collected for disease and parasite screening, ecotoxicology screening, or for genetical analysis
256 (Garbarino et al., 2017; Jelenko & Pokorny, 2010; Tallmon et al., 2004). The rise in non-invasive
257 DNA methods has opened a whole new avenue for collaboration as hunters can collect samples
258 such as faeces for use in population census. For example, the collection of bear scats for DNA
259 based census depends on hunters in Norway, Sweden, Slovenia and Croatia who annually
260 collect thousands of samples (Kindberg et al., 2011; Skrbinšek et al., 2019).

261 **Observations for counts and indices:** Hunters also observe many animals while hunting or
262 tending to their hunting areas. These observations, whether direct or indirect, are used to
263 obtain very valuable data on abundance, distribution and structure if there is a robust design
264 and analysis. For example, in Scandinavia the number of moose and bear observations per

265 hunter per hunting day during the hunting season constitutes a robust index of relative
266 abundance (Ericsson & Wallin, 1999; Kindberg et al., 2011; Solberg & Saether, 1999; Swenson et
267 al., 1994). French roe deer hunters report numbers of roe deer seen along transects (Vincent et
268 al., 1991). Hunters in much of Scandinavia and Finland also take part in structured distance
269 sampling-based surveys of abundance of ptarmigan and forest-living grouse (Lindén, 1996).
270 Bear hunters in Slovenia and Croatia record data on sex ratio and reproductive rates of bears
271 observed on feeding stations (Reljic et al., 2018) as well as using simultaneous observations to
272 produce relatively robust minimum counts of the size of the bear population (Bordjan et al.,
273 2019). Hunters all across the Nordic and Baltic countries submit records of lynx and wolf tracks
274 (and increasingly camera trap images) that are used to produce minimum counts of lynx and
275 wolf populations (Linnell et al., 2007; 2010).

276 **Other types of data:** Hunters also collect other type of data used to infer a wide range of
277 characteristics regarding their species of interest. Questionnaires and interviews are used for
278 disease detections through documentation of what hunters observe on the hunting ground
279 such as scabies infestation for red fox, hair loss in moose, or inferring species distribution or
280 abundance based on their experience and past observations (Gortázar et al., 1998; Llaneza &
281 Núñez-Quirós, 2009, (Madslie et al., 2011). Hunter cooperation with researchers and
282 management authorities also includes their willingness to help ring birds or tag mammals, and
283 return the carcass of shot ringed and tagged animals (Guzmán et al., 2017; Jensen, 1973).

284 3.4. Origin of the information

285 Overall, we found that more than 70% of the diversity in EBV monitoring was documented in
286 both the systematic and unstructured literature search for large carnivores, other game birds,
287 small game and ungulates (Fig. 3). The result is slightly more contrasted for waterfowl as this
288 number goes down to about 60%. Inputs from the unstructured search generally brought as
289 much or more information than the systematic search for large carnivores, small game,
290 ungulates and waterfowl (respectively contributing to 20, 18, 10 and 25%) and didn't add any
291 unique information for small game.

292 With regard to the geographic extent we found that the unstructured search provided a large
293 amount of unique information for other birds (40%), waterfowl (31%), small game (25%) and
294 large carnivores, for which other sources contributed to 50%. The use of a structured search
295 only yielded more information than the unstructured search for waterfowl and ungulates
296 (respectively 47 and 21% of total information).

297 4. Discussion

298 Our study highlights the taxonomic and geographical potential hunter-based monitoring holds,
299 with evidence of nearly all European countries (32 out of 36) using hunter-based monitoring for
300 at least one species group. Moreover, in about half of Europe (including the Scandinavian and
301 Baltic countries) there were examples of hunter-based monitoring to monitor all species
302 groups. We found that the overwhelming majority of hunter-based monitoring programs focus
303 on multiple aspects of the EBVs grouped under species characteristics (i.e. species population,
304 species traits, genetic composition and species community). However, we found that not all
305 countries use hunter-based monitoring for all species groups and some gaps remain. More

306 specifically, we did not find any evidence of hunter-based monitoring in most of the Balkan
307 countries. Concerning species groups, waterfowl is less monitored by hunters than any other
308 groups. This could possibly be due to the large quantity of bird occurrences collected by other
309 types of citizen scientists through for instance eBird, or other national Atlas surveys (Amano et
310 al., 2016a). A lack of literature about the use of hunter-based monitoring due to a lack of
311 acknowledgement of hunters' work from the scientific community or due to language barriers
312 could be two other reasons for these gaps given that some of our data has been found only
313 through experts' inputs. Up to 35% of the conservation literature is not written in English
314 (Amano et al., 2016b) and given the diversity of languages in Europe we did not expect to get
315 the full picture of the contribution of hunters in biodiversity monitoring through the systematic
316 search of the literature. There is almost certainly a major geographic and species-specific bias in
317 the extent to which hunter derived data is analyzed and published by scientists in English. In
318 fact, the email queries sent to wildlife managers and hunting associations added a significant
319 number of inputs in our dataset which were not documented in academic databases through
320 the form of scientific papers or grey literature. This result highlights that we would have
321 dramatically underestimated the extent to which hunters take part in monitoring across Europe
322 without the input of experts. It is almost certain that there are more examples that our search
323 was not able to uncover. The scale of our study (i.e. pooling species into groups and studying
324 hunter based monitoring at country level) might also hide certain fine scale particularities. Even
325 though there is monitoring of a certain EBV in ungulates in a given country, this does not mean
326 that all harvested ungulates are monitored, nor that the whole country is included. As such, our

327 review gives an overview of the potential and some examples, but it may not give a full picture
328 of how widespread the use is.

329 4.1. Particularities of hunter-based monitoring

330 Our results have highlighted the particularity of hunters as citizen scientists due to their access
331 to certain forms of data. Even though highly controversial in some countries in Europe (Fischer
332 et al., 2012), we have shown that hunting delivers data that can be beneficial to researchers
333 and management authorities through the submission of body parts (ovaries, jaw bones, femurs,
334 wings, tissues) from the carcass of harvested animals from which certain species characteristics
335 would have been unobtainable otherwise.

336 Hunting data can provide a unique time series in some countries and we have found studies
337 using roe deer antlers over a 67-year period to study change in environmental pollution
338 (Kierdorf & Kierdorf, 2000), or studies using 30 years of bag data (Jansson & Pehrson, 2007).

339 Even though the hunting season is normally limited to only certain periods (i.e. hunting
340 seasons, depending on the taxa and the countries' regulations), some programs make hunters
341 monitor species characteristics throughout the entire year. For instance, the Finnish wildlife
342 triangle is carried out once during summer and once during winter, allowing the creation of
343 time-series dataset useful for ecological research.

344 Hunting grounds are also widely spread across almost the entire European continent, and even
345 most protected areas are usually open to hunting (Linnell et al., 2015). Hunting is an
346 opportunity for monitoring the status of species population in these areas as hunter-based

347 monitoring has the potential to supplement traditional citizen science which is highly biased
348 towards environments with easy access such as human infrastructures or nature reserves
349 (Tiago et al., 2017).

350 Hunters monitor characteristics primarily on species they hunt and hence species they can
351 recognise easily (i.e. for most countries in Europe, getting a hunting licence requires passing
352 examinations, including assessments of species knowledge). This feature is especially important
353 when making inference from hunters' data as species misidentification is presumably less of an
354 issue than in other citizen science data. Hunters' data are also characterised by their
355 institutionalisation. Most European countries oblige hunters to report their harvest to estimate
356 the relative population of game species and to set quotas for the following year. This system
357 involves hunters directly in the management loop, motivating the data collection.

358 4.2. The challenges with hunter-based monitoring

359 There are however several challenges regarding the use of hunter-based monitoring.

360 Institutionalising the data can potentially lead to some extent of misreporting, where hunters
361 might report more than they really harvest or observe to artificially boost population numbers
362 and thus aim to increase the quota for the following year (Popescu et al., 2016). Even when
363 misreporting is not purposeful, the discrepancies between population indexes resulting from
364 hunting game bags and other more systematic methods can result in mistrust in data provided
365 by hunters from both institutions and other stakeholders, potentially increasing the negative
366 perception of hunters.

367 Moreover, because each country has its own reporting system, misreporting can be facilitated
368 depending on the system. For instance, some of the very structured monitoring programs are
369 able to report data that provide precise indices of the variation of game populations over large
370 areas (Ueno et al., 2014; for a selected sample of these programs see Box 1). However, many
371 programs across Europe are much less structured (i.e. they do not follow a robust sampling
372 scheme) and are based on hunter reports of total numbers of animals believed to occur on their
373 hunting grounds, which are then aggregated onto other administrative levels. These procedures
374 have poorly described methodology, no robust measures to prevent multiple counts of the
375 same individuals and are highly prone to misinterpretation or even potential abuse (Popescu et
376 al., 2016). At best they may provide a rough relative index of temporal change in abundance
377 (Bragina et al., 2018) and an indication of broad scale distribution. However, despite their
378 somewhat ad hoc nature their utility in guiding sustainable hunting practices over the last 50
379 years in many areas must be acknowledged. There is potential to add value to these systems if
380 the underlying concrete observations can be separated from the interpretation, and if some
381 transparent structure can be placed onto both the observation and interpretation processes
382 (e.g. ENETWILD consortium et al., 2018).

383 Finally, hunting based monitoring programs potentially encounter similar biases as any other
384 citizen science programs. For instance, hunter-killed birds rarely constitute a random subset of
385 a population as juveniles or older birds are more likely to be shot, resulting in age biases in
386 hunting bags (Madsen, 2010). Geographical biases also exist with hunting-based monitoring
387 because of different regional management practices, making some hunting areas more popular
388 than others which render abundance indices less reliable (Ranta et al., 2008). Nevertheless,

389 obtaining accurate measures of game characteristics is possible using rigorous methods and
390 standardised protocols such as the collection of ‘indicators of ecological change’ as done in
391 France for roe deer (Morellet et al., 2011). Recent statistical developments regarding the
392 integration of different data sources and types also allowed the researcher to combine hunters’
393 bag statistics with other sources to overcome some of the biases inherent to hunting bag data
394 (Isaac et al., 2019; Rutten et al., 2019).

395

396 **Box 1.** Examples of highly structured hunter-based monitoring schemes

397 **Norwegian moose monitoring program:** Besides reporting the harvest, Norwegian moose
398 hunters are asked to report all moose observed during the hunting season on a standardised
399 form (Solberg & Saether, 1999). This system was started in a few municipalities in the late
400 1960s and extended to cover the entire country during the 1980s. On a daily basis, the leader of
401 each moose hunting team records the number, sex (male, female, unknown) and age (calf,
402 adult, unknown) of all moose observed by the team members but removes individuals that with
403 certainty are observed by more than one member of the team. In addition, they record the
404 number of members that are hunting each day of the hunting season. Data are later reported
405 to the municipality wildlife board and the national deer register (www.hjorteviltregisteret.no/)
406 and used to generate various indices of moose population density and structure for use by the
407 wildlife management (Solberg & Saether, 1999). In addition, hunters in a selection of
408 monitoring sites are required to submit jawbones, ovaries and body weight information of the
409 animals harvested. These data go back to the 1980’s.

410

411 **Danish bird wing survey:** This survey consists of collecting wings from bird shots during the
412 hunting season and is based on voluntary contributions from hunters across Denmark. Every
413 year thousands of wings are forwarded and are used to infer survival, population abundance
414 and structure of the Danish game birds. More information can be found on the website of
415 Aarhus university (<http://fauna.au.dk/en/hunting-and-game-management/wing-survey/>)

416

417 **Finnish wildlife triangles:** The Finnish Wildlife Triangle scheme was developed by the Finnish
418 Game and Fisheries Research Institute in cooperation with the Hunters' central organization in
419 1988. It provides a wide range of information on species population distribution, abundance
420 and structure for 30 wildlife species. This scheme is highly structured and consists of equilateral
421 triangles with 4 km sides distributed across the whole Finnish landscape. These transects are
422 travelled in winter where tracks in the snow are counted, and during the summer when species
423 seen are counted. Annually the census is carried out for 800 to 1000 triangles and involve 7000
424 volunteers, mainly hunters (Pellikka et al., 2005).

425

426 5. Conclusion

427 With limited resources and requests from governments to monitor diverse biodiversity
428 characteristics at large scales, there is a growing need for scientists and wildlife management
429 authorities to use cost effective methods to collect data. Our study shows that collaborations
430 between hunters and scientists can contribute to biodiversity monitoring in nearly all its key

431 dimensions, with the exception of habitat indicators. Nevertheless, hunter based monitoring is
432 not a panacea as geographical and taxonomical gaps exist in the information brought by hunter
433 based monitoring in Europe, possibly due to the low acceptance for the use of hunter-based
434 monitoring within some conservation circles because of the societal and ethical challenges
435 hunting is now facing (Fischer et al., 2012).

436 Furthermore, apart from some very structured programs, the unsystematic nature of hunting-
437 based monitoring poses challenges concerning the use of these data. Statistical developments
438 in data integration as well as more rigorous protocols for data collection when using hunting-
439 based monitoring is needed to unlock further the potential that hunters' data holds.

440 6. Acknowledgements

441 BC was funded by a PhD scholarship from the Norwegian University of Science and Technology.
442 The involvement of JDCL, EBN and MG was funded by the Research Council of Norway (grant
443 251112). We thank David Scallan from FACE – The European Federation for Hunting and
444 Conservation for facilitating access to Europe's national hunting associations. We also thank the
445 wildlife managers and scientists who kindly answered our survey and gave us precious
446 information.

447 7. Authors' contribution

448 All authors participated in the conception and design of the study. BC collected and analysed
449 the data. BC and JDCL drafted the manuscript with the help of all authors. All authors gave final
450 approval for publication.

451 8. Data availability statement

452 The R script and the full dataset use to carry out the analysis are both made available to ensure
453 full reproducibility and can be found at DOI 10.17605/OSF.IO/GKAZM. The dataset available
454 contains all papers from the structured and unstructured search as well as the answers from
455 the questionnaires. All human participating in the questionnaire did so with informed
456 consensus and knowing that they could withdraw at any time without any explanation. The
457 distribution of the data is in line with the participants and respect the ethical guidelines from
458 the Norwegian University of Science and Technology.

459 9. References

- 460 Aebischer, N. J. (2019). Fifty-year trends in UK hunting bags of birds and mammals, and
461 calibrated estimation of national bag size, using GWCT's National Gamebag Census.
462 *European Journal of Wildlife Research*, 65(4). <https://doi.org/10.1007/s10344-019-1299-x>
- 463 Amano, T., Lamming, J. D. L., & Sutherland, W. J. (2016a). Spatial Gaps in Global Biodiversity
464 Information and the Role of Citizen Science. *BioScience*, 66(5), 393–400.
465 <https://doi.org/10.1093/biosci/biw022>
- 466 Amano, T., González-Varo, J. P., & Sutherland, W. J. (2016b). Languages are still a major barrier
467 to global science. *PLoS biology*, 14(12).
- 468 Booth, A., Sutton, A., & Papaioannou, D. (2016). *Systematic approaches to a successful*
469 *literature review*. Sage.
- 470 Bordjan, D., Javornik J., Jerina, K. (2019). *Cost benefit analysis of different monitoring*
471 *approaches and guidelines for optimized monitoring of brown bears. Report prepared*
472 *within the LIFE DINALP BEAR Project (LIFE13 NAT/SI/0005)*.
- 473 Bragina, E. V., Ives, A. R., Pidgeon, A. M., Balčiauskas, L., Csányi, S., Khojetskyy, P., ... Radeloff,
474 V. C. (2018). Wildlife population changes across Eastern Europe after the collapse of
475 socialism. *Frontiers in Ecology and the Environment*, 16(2), 77–81.
476 <https://doi.org/10.1002/fee.1770>
- 477 Chandler, M., See, L., Copas, K., Bonde, A. M. Z., López, B. C., Danielsen, F., ... Turak, E. (2017).
478 Contribution of citizen science towards international biodiversity monitoring. *Biological*
479 *Conservation*, 213, 280–294. <https://doi.org/10.1016/j.biocon.2016.09.004>

480 Díaz, S; Settele, J; Brondízio, E; Ngo, H; Guèze, M; Agard, J; Arneth, A; Balvanera, P; Brauman, K;
481 Butchart, S; Chan, K; Garibaldi, L; Ichii, K; Liu, J; Subrmanian, S; Midgley, G; Miloslavich, P;
482 Molnár, Z; Obura, D; Pfaff, A; Polasky, S; Purvis, A; Razz, C. (2019). *No TiSummary for*
483 *policymakers of the global assessment report on biodiversity and ecosystem services of the*
484 *Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services*.

485 ENETWILD consortium, Oliver Keuling, Marie Sange, Pelayo Acevedo, Tomasz Podgorski,
486 Graham Smith, Massimo Scandura, Marco Apollonio, E. F. and J. V. (2018). Guidance on
487 estimation of wild boar population abundance and density: methods, challenges,
488 possibilities. *EFSA Supporting Publications*, 15(7), 1449E.

489 Ericsson, G., & Wallin, K. (1999). Hunter observations as an index of moose
490 *Alces alces* population parameters. *Wildlife Biology*, 5(1), 177–185.
491 <https://doi.org/10.2981/wlb.1999.022>

492 Fischer, A., Kereži, V., Arroyo, B., Mateos-Delibes, M., Tadie, D., Lowassa, A., ... & Skogen, K.
493 (2013). (De) legitimising hunting—Discourses over the morality of hunting in Europe and
494 eastern Africa. *Land Use Policy*, 32, 261-270.

495 Garbarino, C., Interisano, M., Chiatante, A., Marucci, G., Merli, E., Arrigoni, N., ... Pozio, E.
496 (2017). *Trichinella spiralis* a new alien parasite in Italy and the increased risk of infection
497 for domestic and wild swine. *Veterinary Parasitology*, 246(July), 1–4.
498 <https://doi.org/10.1016/j.vetpar.2017.08.021>

499 Geldmann, J., Heilmann-Clausen, J., Holm, T. E., Levinsky, I., Markussen, B., Olsen, K., ... Tøttrup,
500 A. P. (2016). What determines spatial bias in citizen science? Exploring four recording
501 schemes with different proficiency requirements. *Diversity and Distributions*, 22(11),
502 1139–1149. <https://doi.org/10.1111/ddi.12477>

503 Goodman, L. A. (1961). Snowball sampling. *The annals of mathematical statistics*, 148-170.

504 Gortázar, C., Villafuerte, R., Blanco, J. C., & Fernández-De-Luco, D. (1998). Enzootic sarcoptic
505 mange in red foxes in Spain. *European Journal of Wildlife Research*, 44(4), 251–256.

506 Grøtan, V., Sæther, B. E., Engen, S., Solberg, E. J., Linnell, J. D. C., Andersen, R., ... Lund, E.
507 (2005). Climate causes large-scale spatial synchrony in population fluctuations of a
508 temperate herbivore. *Ecology*, 86(6), 1472–1482. <https://doi.org/10.1890/04-1502>

509 Guzmán, J. L., Caro, J., & Arroyo, B. (2017). Factors influencing mobility and survival of Eurasian
510 Woodcock wintering in Spain. *Avian Conservation and Ecology*, 12(2).
511 <https://doi.org/10.5751/ace-01096-120221>

512 Hagen, R., Heurich, M., Kröschel, M., & Herdtfelder, M. (2014). Synchrony in hunting bags:
513 Reaction on climatic and human induced changes? *Science of the Total Environment*, 468–
514 469, 140–146. <https://doi.org/10.1016/j.scitotenv.2013.08.022>

515 Hochachka, W. M., Fink, D., Hutchinson, R. A., Sheldon, D., Wong, W. K., & Kelling, S. (2012).
516 Data-intensive science applied to broad-scale citizen science. *Trends in Ecology and*
517 *Evolution*, 27(2), 130–137. <https://doi.org/10.1016/j.tree.2011.11.006>

- 518 Isaac, N. J., Jarzyna, M. A., Keil, P., Dambly, L. I., Boersch-Supan, P. H., Browning, E., ... & Jarvis,
519 S. (2019). Data integration for large-scale models of species distributions. *Trends in ecology*
520 *& evolution*.
- 521 Isaac, N. J. B., & Pocock, M. J. O. (2015). Bias and information in biological records. *Biological*
522 *Journal of the Linnean Society*, 115(3), 522–531. <https://doi.org/10.1111/bij.12532>
- 523 Jansson, G., & Pehrson, Å. (2007). The recent expansion of the brown hare (*Lepus europaeus*) in
524 Sweden with possible implications to the mountain hare (*L. timidus*). *European Journal of*
525 *Wildlife Research*, 53(2), 125–130. <https://doi.org/10.1007/s10344-007-0086-2>
- 526 Jelenko, I., & Pokorny, B. (2010). Historical biomonitoring of fluoride pollution by determining
527 fluoride contents in roe deer (*Capreolus capreolus* L.) antlers and mandibles in the vicinity
528 of the largest Slovene thermal power plant. *Science of the Total Environment*, 409(2), 430–
529 438. <https://doi.org/10.1016/j.scitotenv.2010.10.012>
- 530 Jensen, B. (1973). *Movements of the red fox (Vulpes vulpes L.) in Denmark investigated by*
531 *marking and recovery*. Vildtbiologisk Station.
- 532 Kéry, M. and Royle, A. J. (2015). *Applied Hierarchical Modeling in Ecology: Analysis of*
533 *distribution, abundance and species richness in R and BUGS: Volume 1: Prelude and Static*
534 *Models*. Academic Press.
- 535 Kierdorf, H., & Kierdorf, U. (2000). Roe deer antlers as monitoring units for assessing temporal
536 changes in environmental pollution by fluoride and lead in a German forest area over a 67-
537 year period. *Archives of Environmental Contamination and Toxicology*, 39(1), 1–6.
538 <https://doi.org/10.1007/s002440010072>
- 539 Kindberg, J., Swenson, J. E., Ericsson, G., Bellemain, E., Miquel, C., & Taberlet, P. (2011).
540 Estimating population size and trends of the Swedish brown bear *Ursus arctos* population.
541 *Wildlife Biology*, 17(2), 114–123. <https://doi.org/10.2981/10-100>
- 542 Kissling, W. D., Ahumada, J. A., Bowser, A., Fernandez, M., Fernández, N., García, E. A., ...
543 Hardisty, A. R. (2018). Building essential biodiversity variables (EBVs) of species distribution
544 and abundance at a global scale. *Biological Reviews*, 93(1), 600–625.
545 <https://doi.org/10.1111/brv.12359>
- 546 Kojola, I., & Laitala, H. M. (2000). Changes in the structure of an increasing brown bear
547 population with distance from core areas: Another example of presaturation female
548 dispersal? *Annales Zoologici Fennici*, 37(1), 59–64.
- 549 Kosmala, M., Wiggins, A., Swanson, A., & Simmons, B. (2016). Assessing data quality in citizen
550 science. *Frontiers in Ecology and the Environment*, 14(10), 551–560.
551 <https://doi.org/10.1002/fee.1436>
- 552 Lindén, H. (1996). Wildlife triangle scheme in Finland: methods and aims for monitoring wildlife
553 populations. *Finnish Game Research*, 49, 4–11.
- 554 Linnell, J.D.C. & Cretois, B. (2018). *Research for AGRI Committee-The Revival of Wolves and*

- 555 *Other Large Predators and Its Impact on Farmers and Their Livelihood in Rural Regions of*
556 *Europe.*
- 557 Linnell, J. D. C., Broseth, H., Odden, J., & Nilsen, E. B. (2010). Sustainably harvesting a large
558 carnivore? Development of eurasian lynx populations in Norway during 160 years of
559 shifting policy. *Environmental Management*, 45(5), 1142–1154.
560 <https://doi.org/10.1007/s00267-010-9455-9>
- 561 Linnell, J. D. C., Kaczensky, P., Wotschikowsky, U., Lescureux, N., & Boitani, L. (2015). Framing
562 the relationship between people and nature in the context of European conservation.
563 *Conservation Biology*, 29(4), 978–985. <https://doi.org/10.1111/cobi.12534>
- 564 Linnell, J. D. C., Odden, J., Andersen, R., Brøseth, H., Andrén, H., Liberg, O., ... Okarma, H. (2007).
565 Distance rules for minimum counts of Eurasian lynx Lynx lynx family groups under different
566 ecological conditions. *Wildlife Biology*, 13(4), 447–455. [https://doi.org/10.2981/0909-6396\(2007\)13\[447:drfmco\]2.0.co;2](https://doi.org/10.2981/0909-6396(2007)13[447:drfmco]2.0.co;2)
- 568 Llana, L., & Núñez-Quirós, P. (2009). Distribution of the iberian wolf (*Canis lupus signatus*) in
569 Galicia (NW Spain): Concordance between field sampling and questionnaires. *Wildlife*
570 *Biology in Practice*, 5(1), 23–32. <https://doi.org/10.2461/wbp.2009.5.5>
- 571 Madsen, J. (2010). Age bias in the bag of pink-footed geese *Anser brachyrhynchus*: influence of
572 flocking behaviour on vulnerability. *European Journal of Wildlife Research*, 56(4), 577–582.
- 573 Massei, G., Kindberg, J., Licoppe, A., Gačić, D., Šprem, N., Kamler, J., ... Náhlik, A. (2015). Wild
574 boar populations up, numbers of hunters down? A review of trends and implications for
575 Europe. *Pest Management Science*, 71(4), 492–500. <https://doi.org/10.1002/ps.3965>
- 576 Moleón, M., Almaraz, P., & Sánchez-Zapata, J. A. (2008). An emerging infectious disease
577 triggering large-scale hyperpredation. *PLoS ONE*, 3(6), 12–17.
578 <https://doi.org/10.1371/journal.pone.0002307>
- 579 Moleón, M., Almaraz, P., & Sánchez-Zapata, J. A. (2013). Inferring ecological mechanisms from
580 hunting bag data in wildlife management: A reply to Blanco-Aguilar et al. (2012). *European*
581 *Journal of Wildlife Research*, 59(4), 599–608. <https://doi.org/10.1007/s10344-013-0711-1>
- 582 Morellet, N., GAILLARD, J. M., Hewison, A. M., Ballon, P., Boscardin, Y. V. E. S., Duncan, P., ... &
583 Maillard, D. (2007). Indicators of ecological change: new tools for managing populations of
584 large herbivores. *Journal of Applied Ecology*, 44(3), 634–643.
- 585 Morner, T., Fischer, J., & Bengis, R. (2014). The value of increasing the role of private individuals
586 and organisations in One Health. *Scientific and Technical Review*, 33(3), 605–613.
- 587 Nilsen, E. B., Brøseth, H., Odden, J., & Linnell, J. D. C. (2012). Quota hunting of Eurasian lynx in
588 Norway: Patterns of hunter selection, hunter efficiency and monitoring accuracy. *European*
589 *Journal of Wildlife Research*, 58(1), 325–333. <https://doi.org/10.1007/s10344-011-0585-z>
- 590 Pellikka, J., Rita, H., & Lindén, H. (2005). Monitoring wildlife richness - Finnish applications
591 based on wildlife triangle censuses. *Annales Zoologici Fennici*, 42(2), 123–134.

- 592 Pereira, H. M., Ferrier, S., Walters, M., Geller, G. N., Jongman, R. H. G., Scholes, R. J., ... & Coops,
593 N. C. (2013). Essential biodiversity variables. *Science*, 339(6117), 277-278.
- 594 Popescu, V. D., Artelle, K. A., Pop, M. I., Manolache, S., & Rozyłowicz, L. (2016). Assessing
595 biological realism of wildlife population estimates in data-poor systems. *Journal of Applied*
596 *Ecology*, 53(4), 1248–1259. <https://doi.org/10.1111/1365-2664.12660>
- 597 Pöysä, H., & Väänänen, V. M. (2018). Changes in the proportion of young birds in the hunting
598 bag of Eurasian wigeon: long-term decline, but no association with climate. *European*
599 *Journal of Wildlife Research*, 64(2). <https://doi.org/10.1007/s10344-018-1179-9>
- 600 Ranta, E., Lindström, J., Lindén, H., & Helle, P. (2008). How reliable are harvesting data for
601 analyses of spatio-temporal population dynamics?. *Oikos*, 117(10), 1461-1468.
- 602 Reimoser, S., Smidt, S., Reimoser, F., & Wildauer, L. (2014). Changes of hunting bag and habitat
603 in the southern Vienna-Woods since 1891. *Allgemeine Forst-Und Jagdzeitung*, 185(1/2),
604 16-27.
- 605 Reljic, S., Jerina, K., Nilsen, E. B., Huber, D., Kusak, J., Jonozovic, M., & Linnell, J. D. C. (2018).
606 Challenges for transboundary management of a European brown bear population. *Global*
607 *Ecology and Conservation*, 16, e00488. <https://doi.org/10.1016/j.gecco.2018.e00488>
- 608 Rutten, A., Casaer, J., Swinnen, K. R., Herremans, M., & Leirs, H. (2019). Future distribution of
609 wild boar in a highly anthropogenic landscape: Models combining hunting bag and citizen
610 science data. *Ecological Modelling*, 411, 108804.
- 611 Silvertown, J. (2009). A new dawn for citizen science. *Trends in Ecology and Evolution*, 24, 467–
612 471. <https://doi.org/10.1016/j.chemosphere.2018.03.203>
- 613 Skrbinšek, T., Luštrik, R., Majič-Skrbinšek, A., Potočnik, H., Kljun, F., Jelenčič, M., ... Trontelj, P.
614 (2019). From science to practice: genetic estimate of brown bear population size in
615 Slovenia and how it influenced bear management. *European Journal of Wildlife Research*,
616 65(2). <https://doi.org/10.1007/s10344-019-1265-7>
- 617 Smedshaug, C. A. (2018). The effect of a natural reduction of red fox *Vulpes vulpes* on small
618 game hunting bags in Norway. *Wildlife Biology*, 5(1), 157.
619 <https://doi.org/10.2981/wlb.1999.020>
- 620 Solberg, E. J., Saether, B. E., Strand, O., & Loison, A. (1999). Dynamics of a harvested moose
621 population in a variable environment. *Journal of Animal Ecology*, 68(1), 186–204.
622 <https://doi.org/10.1046/j.1365-2656.1999.00275.x>
- 623 Solberg, J., & Saether, B. (1999). Hunter observations of moose *Alces alces* as a management
624 tool. *Wildlife Biology*, 5(1), 107. <https://doi.org/10.2981/wlb.1999.014>
- 625 Stewart, G. B., Coles, C. F., & Pullin, A. S. (2005). Applying evidence-based practice in
626 conservation management: Lessons from the first systematic review and dissemination
627 projects. *Biological Conservation*, 126(2), 270–278.
628 <https://doi.org/10.1016/j.biocon.2005.06.003>

629 Swenson, J. E., Sandegren, F., Bjärvall, A., Söderberg, A., Wabakken, P., & Franzén, R. (1994).
630 Size, trend, distribution and conservation of the brown bear *Ursus arctos* population in
631 Sweden. *Biological Conservation*, Vol. 70, pp. 9–17. [https://doi.org/10.1016/0006-](https://doi.org/10.1016/0006-3207(94)90293-3)
632 [3207\(94\)90293-3](https://doi.org/10.1016/0006-3207(94)90293-3)

633 Swenson, J. E., Sandegren, F., & Söderberg, A. (1998). Geographic expansion of an increasing
634 brown bear population: Evidence for presaturation dispersal. *Journal of Animal Ecology*,
635 *67*(5), 819–826. <https://doi.org/10.1046/j.1365-2656.1998.00248.x>

636 Tallmon, D. A., Luikart, G., & Waples, R. S. (2004). The alluring simplicity and complex reality of
637 genetic rescue. *Trends in Ecology and Evolution*, *19*(9), 489–496.
638 <https://doi.org/10.1016/j.tree.2004.07.003>

639 Tiago, P., Ceia-Hasse, A., Marques, T. A., Capinha, C., & Pereira, H. M. (2017). Spatial
640 distribution of citizen science casuistic observations for different taxonomic groups.
641 *Scientific Reports*, *7*(1), 1–9. <https://doi.org/10.1038/s41598-017-13130-8>

642 Ueno, M., Solberg, E. J., Iijima, H., Rolandsen, C. M., & Gangsei, L. E. (2014). Performance of
643 hunting statistics as spatiotemporal density indices of moose (*Alces alces*) in Norway.
644 *Ecosphere*, *5*(2). <https://doi.org/10.1890/ES13-00083.1>

645 UN General Assembly. (1992). *The Convention on Biological Diversity of 5 June 1992*.

646 UN General Assembly. (2015). *Transforming our world : the 2030 Agenda for Sustainable*
647 *Development, 21 October 2015, A/RES/70/1, available at:*
648 <https://www.refworld.org/docid/57b6e3e44.html> [accessed 1 December 2019]itle.

649 van der Jeugd, H. P., & Kwak, A. (2017). Management of a Dutch resident barnacle goose *Branta*
650 *leucopsis* population: How can results from counts, ringing and hunting bag statistics be
651 reconciled? *Ambio*, *46*(s2), 251–261. <https://doi.org/10.1007/s13280-017-0900-3>

652 Vincent, J.-P., Gaillard, J.-M., & Bideau, E. (1991). Kilometric index as biological indicator for
653 monitoring forest roe deer populations. *Acta Theriologica*, *36*, 315–328.
654 <https://doi.org/10.4098/at.arch.91-33>

655 Walters, M., Pereira, H., Ferrier, S., Jongman, R., & Geller, G. (2011). *Essential Biodiversity*
656 *Variables (EBVs)*. *2011*(December), 277–279.

657 Westgate, M. J. (2019). revtools: An R package to support article screening for evidence
658 synthesis. *Research Synthesis Methods*, (June), 606–614.
659 <https://doi.org/10.1002/jrsm.1374>

660

661

662

663

664
665
666
667
668
669
670

671 **Table 1:** List of criteria to classify hunters’ data collection methods.

Method	Criteria
Bag	If the EBV was calculated from data taken from official harvest numbers
Camera trap	If the EBV was calculated from data collected through camera traps operated by hunters
Carcass	If the EBV was calculated from the carcass of the shot animal including any invasive samples
Direct Observation	If the EBV was calculated from hunters' direct observations of a species
Indirect Observation	If the EBV was calculated from hunters' indirect observations of signs such as snow tracking, faeces sampling, observations of dens
Questionnaire	If the EBV was calculated from data coming from any sort of questionnaires, including written or digital questionnaires, distributed to hunters
Ringing	If the EBV was calculated as a result of the ringing of animals by hunters, this include birds rings or animal tags
Other	If the EBV was calculated from data collected through any sort hunters’ cooperation such as direct interview with hunters, non-invasive sampling or bringing shot ringed or tagged animals

672
673
674
675
676
677

678


679

680

681

682

683 **Figure 1:** Geographic extent of hunter-based monitoring per species group. A) Large carnivores,
684 B) ungulates, C) small game, D) other game birds and E) waterfowl.



685

686

687

688

689


690

691

692

693

694 **Figure 2:** Diversity of EBV monitored through hunter-based monitoring. A) Large carnivores, B)
 695 ungulates, C) small game, D) other game birds and E) waterfowl. A tick indicates if the Essential
 696 Biodiversity Variable has been found in our review. Colors represent the Essential Biodiversity
 697 Class



698 **Table 2:** Use of monitoring methods to infer different Essential Biodiversity Variables. (X)
 699 representing the finding of a source documenting the use of the method to infer the EBV, (-) if
 700 no sources have been found.

EBV class	EBV candidate	Bag	Cam	Carcass	D	I	Help	Questionnaire	Ringling	Other
Genetic composition	Allelic diversity	-	-	X	-	X	-	-	-	X
	Co-ancestry	-	-	X	-	-	-	-	-	-
Species population	Population abundance	X	X	X	X	X	-	X	-	X
	Species distribution	X	X	X	X	X	-	X	-	X
Species traits	Population structure	X	-	X	X	X	-	X	-	X
	Morphology	-	-	X	-	X	-	X	-	-
	Movement	-	-	X	-	-	X	-	X	X
	Phenology	X	-	X	X	X	-	-	-	-
	Physiology	-	-	X	-	X	-	X	-	X
Community composition	Reproduction	X	X	X	X	X	-	-	-	X
	Species interaction	X	-	X	X	X	-	-	-	X
	Taxonomic diversity	-	-	X	-	-	-	X	-	-

701
 702
 703
 704
 705
 706
 707
 708
 709
 710
 711
 712
 713
 714

715 **Figure 3:** Proportion of unique information yielded by the systematic search, unstructured
 716 search and both sources with regards to A) EBVs and B) Countries.

717

718

719

720

721

722

723

724

725

726

