

# Honey Bee Diversity is Swayed by Migratory Beekeeping and Trade Despite Conservation Practices: Genetic Evidences for the Impact of Anthropogenic Factors on Population Structure

1 **Mert Kükürer** \*<sup>1,2</sup>, **Meral Kence**<sup>1</sup>, **Aykut Kence**<sup>1,3</sup>

2 <sup>1</sup>Biology Department, Middle East Technical University, Ankara, Turkey

3 <sup>2</sup>Molecular Biology and Genetics Department, Kilis 7 Aralik University, Kilis, Turkey

4 <sup>3</sup>Deceased

5 **\* Correspondence:**

6 Mert Kükürer

7 mertkukrer@gmail.com

8 **Keywords:** migratory beekeeping, queen and colony trade, isolated regions, gene flow,  
9 population structure, biodiversity conservation, microsatellite markers, *Apis mellifera*  
10 subspecies

11 **Abstract**

12 Intense admixture of honey bee (*Apis mellifera* L.) populations is mostly attributed to migratory  
13 beekeeping practices and replacement of queens and colonies with non-native races or hybrids of  
14 different subspecies. These two practices are also heavily carried out in Anatolia and Thrace where 5  
15 subspecies reside naturally.

16 Here, we carried out an analysis of population structure of honey bees sampled from six different  
17 regions (n = 250) in order to test the genetic impacts of migratory beekeeping, queen and colony  
18 trade and conservation efficacy of isolated regions. A total of 30 microsatellite markers were used in  
19 four multiplex reactions.

20 Direct genetic impact of migratory beekeeping was demonstrated first time based on a comparison of  
21 assignment of individuals to their geographically native populations where migratory colonies  
22 showed less fidelity. We found genetic evidence for them acting as a hybrid zone mobile in space and  
23 time, becoming vectors of otherwise local gene combinations.

24 The effects of honey bee trade were revealed by the presence of very high introgression levels from  
25 the highly commercial Caucasian bees naturally limited to a narrow range. We also measured the  
26 direction and magnitude of this gene flow connected with bee trade.

27 Comparison between regions that are either open to migratory beekeeping or not let us evaluate the  
28 status of isolated regions as centers of limited gene flow and showed the importance of establishing  
29 such regions.

30 Despite signs of gene flow, our findings confirm high levels of geographically structured genetic  
31 diversity of four subspecies of honey bees in Turkey and emphasize the need to develop policies to  
32 maintain this diversity.

33 Our overall results might potentially bear a wider interest to the community since they constitute an  
34 important attempt to quantify the effects of anthropogenic impacts on established patterns of honey  
35 bee diversity. Our measurable and justified findings on migratory beekeeping, queen and colony  
36 replacements as well as conservation implications will hopefully be of use for the decision makers  
37 and other stakeholders.

### 38 **Introduction**

39 The Western honey bee, *Apis mellifera* L., is a species which plays role together with other  
40 pollinators in pollination of wild and cultivated plants while the species also have significant  
41 economic importance in terms of honey and other bee products output (Morse 1991; Breeze et al.  
42 2011). In addition to its ecological and economic importance, it is a model study organism both for  
43 evolution of eusociality and sophisticated cognitive abilities (Weinstock et al. 2006).

44 Natural distribution of *A. mellifera* includes Central and Southwest Asia, Europe and Africa but the  
45 species was also introduced to East and Southeast Asia, Australia and the Americas mainly on  
46 purpose for its economic benefits (Ruttner 1988). Morphological and molecular studies point to four  
47 major lineages of numerous –more than 20- subspecies (Ruttner, 1988; Whitfield et al. 2006). The  
48 four widely recognized lineages are A (Africa), M (western and northern Europe), O (Near East and  
49 Central Asia) and C (Eastern Europe) lineages.

50 Although bearing controversies, studies with Single Nucleotide Polymorphisms (SNPs) in the past  
51 decade supported the hypothesis that *A. mellifera* have originated in the tropics or subtropics in  
52 Africa and colonized its natural range by two main routes: one through Gibraltar and one through  
53 Suez and then Bosphorus, ending up with a secondary contact between the highly divergent A and C  
54 lineages around Alps (Whitfield et al. 2006; Han et al. 2012; Walberg et al. 2014; Harpur et al. 2014;  
55 Cridland et al. 2017).

56 Both the honey bees and wild pollinators are thought to be on decline (locally and/or globally  
57 depending on the species and region of concern) due to factors some of them relating closely to  
58 human activities. Among them, destruction and fragmentation of natural habitats, toxicity caused by  
59 pollution and pesticides –as such widely used neonicotinoids-, diseases and their spread getting  
60 easier, invasive species are leading the way (Meffe 1998; Brown & Paxton 2009; Van Engelsdorp &  
61 Meixner 2010; Blacquiere et al. 2012). Honey bees also, especially wild populations that are not  
62 managed by beekeepers (including the feral populations), take their share from the situation like the  
63 other species in the genus *Apis* –namely *Apis cerana*, *Apis florea*, *Apis dorsata* and other native bees  
64 of Asia (Oldroyd 2007; Dietemann et al. 2009; Van Engelsdorp et al. 2009; Genersch 2010; Evans &  
65 Schwarz 2011).

66 Besides such negative consequences created by human activities; the genetic admixture of honey bee  
67 populations due to bee trade, including complete replacement of local bees with non-natives and  
68 beekeeping practices involving movement of colonies from one region to the other impose another  
69 kind of pressure on the species: the loss and/or swamping of locally adapted gene combinations and  
70 local or global extinctions of native honey bees (De la Rua et al. 2009).

71 All those factors and their interactions, including genetic and environmental ones, when combined,  
72 may have an increased adverse effect on honey bees and may be the reasons behind continuous or  
73 discrete events of sudden colony losses with rapid depletion of worker bees while the queen  
74 continues laying eggs accompanied by lack of dead bees in and around the hive; the syndrome called

75 as Colony Collapse Disorder (CCD) or Colony Depopulation Syndrome (CDS) (Van Engelsdorp et  
76 al. 2009; Neumann & Carreck 2010).

77 Resilience of the honey bees may be lying in the adaptations they accumulated over thousands of  
78 years, and new potentials reside in their genetic diversity. It is highly probable that a combination of  
79 many above mentioned factors/threats are taking their places in the recent declines by weakening the  
80 colonies step by step. Due to altered rankings of performance of subspecies in varying environments,  
81 it is generally accepted that honey bees' resistance or tolerance to these factors differ greatly and  
82 locally adapted variants may be encountering less stress, thus remain standing strong (Büchler et al.  
83 2015). Hence, research on honey bee diversity in the global context and at various levels (genetic,  
84 individuals, colonies, populations, ecotypes and subspecies) is of great importance for maintaining  
85 the species' and ecosystem services they provide as well as their economic usefulness.

86 In recent years' research conducted on honey bee population structure in European countries, it was  
87 shown that the past structure was lost or strongly disturbed (Dall'Olio et al. 2007; Canovas et al.  
88 2011; Bouga et al. 2011). Introgression of non-native DNA was monitored in wild populations of  
89 Sudan (El-Niweiri & Moritz 2010). Among the anthropogenic effects, mainly queen and colony trade  
90 and replacement of native honey bees with non-natives as well as migratory beekeeping were the  
91 usual suspects.

92 Despite grounded suspicions there are very few studies that investigate and test the direct genetic  
93 consequences of human practices on honey bee diversity. Therefore, the aims of this series of  
94 experiments were testing different hypotheses about recent heavy/any admixture of honey bee  
95 populations across four subspecies by making use of microsatellite markers as well as i) evaluating  
96 the status of isolated regions as a conservation implication where migratory beekeeping is prohibited,  
97 restricted or very scarce due to lack of preference of migratory beekeepers or attitude of local  
98 beekeepers ii) acquiring and demonstrating the direct genetic outcomes of migratory beekeeping by a  
99 series of comparisons between migratory and stationary colonies iii) seeking for the effects of  
100 unregulated queen and colony trade by figuring out the origin, extent and direction of introgression  
101 between populations.

102 With five subspecies dwelling within its borders and with a variety of beekeeping strategies, Turkey  
103 makes a good stage for chasing genetic evidences for the impact of anthropogenic factors on one of  
104 the most important crop and wild plant pollinators. Beekeeping is an old tradition in those lands  
105 which dates back to 6600 BC and Hittites civilization (Akkaya & Serhat 2007), while still intensively  
106 practiced in Turkey where there are more than 8 million hives distributed all over the country. This is  
107 the third highest number in the world, alone tripling those of the USA and reaching the half of the EU  
108 total (USDA NASS 2019, European Parliament 2017).

109 Corresponding to one-fourth to one-fifth of all recognized subspecies of *A. mellifera*; *A. m. meda*, *A.*  
110 *m. syriaca*, *A. m. caucasica*, *A. m. anatoliaca* from the O-lineage and an ecotype from C subspecies  
111 group exist in Turkey (Kandemir et al. 2005). Even A-lineage genetic material was also characterized  
112 in native bees from the Levantine coast of Turkey (Kandemir et al. 2006) bringing together genetic  
113 elements from three continents. Major subspecies found in and around Anatolia are shown in Fig. 1a.

114 Anatolia and Thrace, when taken together, harbor a vast diversity: honey bees belonging to three  
115 different lineages meet, exchange genes and adapt to local conditions determined by diverse climatic,  
116 topographical and floristic variations available (Bouga et al. 2011). Refugial status of Anatolia during  
117 the ice ages contributed present enhanced levels of biodiversity (Hewitt 1999). Studies concerning

118 honey bee populations of Turkey (Bodur et al. 2007; Kence et al. 2009) demonstrated high genetic  
119 structuring among them and confirmed the presence of divergent populations pointing to different  
120 subspecies. They, all together, drew attention to this rich diversity hotspot present and particularly  
121 under threat in Anatolia and Thrace as well as importance to its conservation.

122 Despite that, still prevail the arguments in popular opinion -with significant effect on decision makers  
123 and other stakeholders- that honey bee ecotypes are inevitably lost due to gene flow facilitated by  
124 anthropogenic factors, so the relevance of investing in a strategy involving conservation of locally  
125 adapted variants are unremittingly questioned. This study aimed to quantify and weigh the impacts of  
126 anthropogenic factors and conservation efforts on the present condition of honey bee genetic  
127 diversity.

## 128 **Materials and Methods**

### 129 Sampling

130 We sampled a total of 250 honey bees each from different colonies from 18 provinces during the  
131 period of March 2010 and August 2012. Of those 250 honey bees, 174 were from apiaries that were  
132 stationary and 76 were from migratory ones. Beekeepers declared that they used original honey bees  
133 from stocks native to the area and that they did not purchase non-native queens or colonies in the last  
134 ten years.

135 We grouped samples from provinces with small sample sizes together with nearby provinces to form  
136 10 major localities: Kırklareli, Edirne+ (Edirne and Tekirdağ), Muğla, Eskişehir+ (Eskişehir,  
137 Kütahya and Bilecik), Düzce+ (Düzce, Zonguldak and Bolu), Ankara, Hatay, Bitlis+ (Bitlis, Elazığ,  
138 Erzurum and Ordu), Ardahan, and Artvin. Those localities correspond to natural distribution range of  
139 four subspecies. Those subspecies are *A. m. syriaca* in Hatay, *A. m. caucasica* in Ardahan and Artvin,  
140 *A. m. anatoliaca* in Düzce, Eskişehir+, Muğla and Ankara from the O lineage and an ecotype from C  
141 subspecies group in Kırklareli and Edirne+ by excluding the fifth subspecies *A. m. meda*. We carried  
142 out combinations of locations according to geographical proximity; similarity in terms of climatic,  
143 topographic and floral variables; results of previous studies as well as results of preliminary analysis  
144 of this study. Sampling sites and sample sizes can be seen in Fig. 1b.

145 The samples were kept in -80 °C until genetic analysis.

### 146 Genotyping

147 We isolated DNA from bee heads by QIAGEN DNeasy Blood and Tissue Kit following the  
148 procedure of the producer for insect samples with slight modifications. We grouped a set of 30  
149 microsatellite loci into four clusters for two 7-plex (set 1: AP218, A113, AB024, AP249, A088,  
150 AP001, AP043 and set 2: AP049, AP238, AC006, AP243, AP288, HBC1602, A107) and two 8-plex  
151 (set 3: A079, AC306, AP226, A007, HBC1601, AP068, A014, AP223 and set 4: AP019, AB124,  
152 A043, A076, AP273, AP289, HBC1605, A028) polymerase chain reactions (Estoup et al. 1995;  
153 Solignac et al. 2003; Bodur et al. 2007; Shaibi et al. 2008; Tunca et al. 2009). A software program,  
154 Multiplex Manager 1.2 (Holleley & Geerts 2009), was used for constructing the multiplex groups.  
155 Information on primer pairs, fluorescent dyes and PCR conditions are provided in the supplementary  
156 file.

157 Detection of microsatellite allele sizes was achieved by capillary electrophoresis with ABI 3730XL  
158 sequencing machines. We were not able to amplify locus A076 consistently across the samples thus  
159 we definitely excluded it from the data set and the downstream analysis.

### 160 Population structure

161 We calculated pairwise  $F_{ST}$  values by Arlequin 3.5 (Excoffier et al. 2005), Mantel test was applied to  
162 account for isolation by distance procedure. Pairwise population distances were calculated (Reynolds  
163 et al. 1983) by Populations 1.2.32 software (Langella 2011) and visualized by the online tool  
164 Interactive Tree of Life v4 (Letunic and Bork 2019). We used PAST4 and PCAGEN software to plot  
165 populations on a two-dimensional space by a PCA based on correlation matrix between groups  
166 (Goudet 1999; Hammer 2001).

167 Population structure was estimated by Structure 2.3.3 (Pritchard et al. 2000), K values of distinct  
168 populations were analyzed by Structure Harvester software (Earl & von Holdt 2012), and we used  
169 Clumpp software (Jakobsson & Rosenberg 2007) to permute the membership coefficients of  
170 individuals determined by Structure 2.3.3 and Distruct software to (Rosenberg 2004) visualize the  
171 results obtained by Clumpp.

172 Other population genetic parameters and diversity indicators were also estimated and they are  
173 provided as supplementary file. These parameters and indicators contain frequency of null alleles,  
174 allelic richness and diversities, inbreeding and prevalence of close relatives, number effective alleles,  
175 levels of heterozygosity, deviations from Hardy-Weinberg and linkage disequilibrium, bottlenecks,  
176 effective population sizes and microsatellite information index.

### 177 Statistical analyses

178 We then used membership coefficients obtained, to test hypotheses about beekeeping practices,  
179 isolated regions and queen/colony trade. For the analysis, we first arcsine root square transformed the  
180 coefficients since the data was composed of proportions and non-normally distributed. Then we  
181 carried out Shapiro, Mann-Whitney  $U$ , Kruskal-Wallis, Dunn's,  $F$ , ANOVA, Tukey's and  $t$  tests  
182 wherever necessary and applicable to compare mean membership coefficients and estimated Cohen's  
183  $d$  to determine effect sizes. Those were carried out in R statistical software using packages pwr,  
184 effsize, dunn.test and dabestr (R Core Team 2013, Torchiano 2016; Dinno 2017; Champely et al.  
185 2018; Ho et al. 2019). R code is provided as a supplementary material.

186 We made use of estimation plots to visualize untransformed data for membership coefficients and  
187 impact of various factors on them. This is a less conventional method when compared to bar or  
188 boxplots and reporting of significance tests but much more convenient and powerful method to  
189 summarize the whole data in an unbiased way by displaying all measurements and effect sizes as  
190 well as precision of estimates and distribution of mean differences (Ho et al. 2019).

### 191 Beekeeping practice: migratory vs stationary

192 For the first hypothesis to be tested, we compared membership coefficients of migratory and  
193 stationary colonies in Ankara, Muğla and Hatay separately, for the three provinces combined and for  
194 the total data set. If the migratory colonies acted as a potential vector of foreign alleles then they  
195 would have much lower probabilities of being assigned to their own clusters.

### 196 Isolated regions as a conservation practice

197 The second hypothesis was about isolated regions. If the isolated regions were efficient in preserving  
 198 genetic diversity by preventing gene flow between different clusters then one would expect to see  
 199 higher membership coefficients for stationary individuals belonging to these regions and lower for  
 200 stationary individuals that belong to regions open to migratory beekeeping.

201 Kırklareli is a province that is declared officially as an isolated region where migratory beekeepers  
 202 could not visit for years at first thanks to local beekeepers' negative attitude towards them. The  
 203 region is home to a Carniolan ecotype carefully maintained by local beekeepers. Ardahan is legally  
 204 declared a conservation and breeding area for *A. m. caucasica* so migratory beekeepers cannot enter  
 205 the province and queen import from other subspecies is forbidden. Parts of Artvin province are also  
 206 officially declared as isolated regions for conservation of *A. m. caucasica* as a pure race. The  
 207 province in general is rarely visited by migratory beekeepers for geographical reasons and beekeepers  
 208 there, dealing with mass queen breeding, do not use non-native queens. We compared these three  
 209 provinces with the other six regions (Edirne+, Muğla, Düzce+, Eskişehir+, Ankara and Hatay) where  
 210 migratory beekeeping and bee trade are freely exercised.

### 211 Effect of queen and colony trade

212 Third set of tests were about the impacts of honey bee trade. We compared the estimated proportion  
 213 of genomes assigned to a different cluster than the native cluster among individuals of the total data  
 214 set to find out which cluster contributed most to other clusters' gene pools.

215 Ardahan and Artvin provinces host the *A. m. caucasica* subspecies which is also widely used for  
 216 commercial purposes and the *caucasica* queens and their hybrids are sold all over the country. But  
 217 these provinces are also limited to a very narrow range in the Northeast of the country and are  
 218 declared isolated regions. So, a possible high introgression of their alleles would mostly, if not  
 219 completely, be due to replacement of queens and colonies.

220 We also investigated further patterns across populations to understand the magnitude and direction of  
 221 the gene flow by tracing the signs of those misassigned proportions within localities.

### 222 **Results**

223 We calculated  $F_{ST}$  values by using both the frequencies obtained in the study and by using the null  
 224 allele corrected frequencies. We calculated for the stationary ( $n = 174$ ) colonies an overall  $F_{ST}$  of  
 225 0.065 and an  $F_{ST}$  of 0.067 after null allele corrections. For migratory colonies the values were 0.011  
 226 and 0.015 respectively and for all the 250 samples the values were 0.046 and 0.047.

227 Phylogenetic tree we constructed by using pairwise population distances based on stationary colonies  
 228 only resolves four distinct branches corresponding to four subspecies (Fig. 2b). Thracian samples  
 229 constitute the extreme end of the unrooted tree. The other end is divided to three almost equidistant  
 230 branches of Caucasian, Levantine and Anatolian samples.

231 We plotted stationary colonies, migratory colonies and the overall data for the regions of sampling on  
 232 2D spaces by carrying out Principle Component Analysis (Fig. 2a) which showed a similar pattern  
 233 with the UPGMA tree. First axis designating the first principle component differentiated samples  
 234 those in Thrace whereas the second one corresponding to the second component differentiated  
 235 subspecies in Anatolia (*syriaca*, *anatoliaca* and *caucasica*). The x and y axes explained 41.8% and  
 236 32.1% of the variance within the samples.

237 Genetic distances in stationary colonies showed significant correlation ( $p < 0.001$ ) with geographic  
238 distance but those of migratory colonies were not correlated with geographic distances. Results of  
239 Mantel test point to an isolation by distance pattern in stationary colonies that is lost in migratory  
240 ones.

241 Concerning the Structure results, the best K values were selected by the Structure Harvester program  
242 as 2 and 4 with similar outcomes, K=2 being slightly likelier than K=4 which hint for lineage level  
243 diversification of C and O ancestries. We calculated membership coefficients of individuals to the  
244 observed clusters in K=4 since it can be biologically attributed to relevant subspecies under  
245 investigation and we used them for further hypothesis testing. Clustering analyses showed no  
246 population structuring for migratory colonies (Fig. 3a) in contrast to stationary colonies and the  
247 overall data (Fig. 3b and 3c). Concerning the overall data, however, distortion in the population  
248 structure caused by migratory colonies is evident in higher admixture levels observed.

249 We compared individuals from stationary and migratory colonies according to their membership  
250 coefficients belonging to their native clusters (or it can be called their expected natural populations  
251 alike). The mean values and effect sizes as well as the significance level of the differences were  
252 summarized in Table 1. Boxplots contrasting the arcsine root square transformed membership  
253 coefficients for migratory and stationary colonies are shown in Fig. 4a and scatter plots are very  
254 much similar but visualizing raw membership coefficients for each sample are shown in Fig. 4b.

255 Estimation plots not only fairly visualize the real distribution of the data but also let us compare the  
256 effect sizes and their precision. Stationary colonies are annotated as <Group name> 0 and migratory  
257 colonies are as <Group name> 1 (Fig. 4). Bars right to the data points refer to the 25% and 75%  
258 quartiles and the gap between them is the median value for the sample. The zero line below  
259 correspond to the mean membership coefficients of stationary colonies in each pairwise comparison.  
260 The Euclidean distances from those means for the migratory colonies are shown as dots with a 95%  
261 confidence interval bar around. Also, distributions of the estimation statistics are included. So that we  
262 can comprehensively compare the strength of the drift for different populations and subsets of the  
263 data.

264 Stationary colonies from Muğla and Hatay were quite more likely to be assigned to their own clusters  
265 than the migratory colonies from these provinces, the same held when we compared the combined  
266 data from the three provinces or all the migratory and stationary colonies. However, the situation was  
267 the reverse in Ankara possibly due to factors we discuss below. Stationary colonies from that  
268 province reflected patterns of high admixture. The difference between stationary colonies and  
269 migratory in Muğla are much less when compared against the ones in Hatay, signaling for a possible  
270 higher level of admixture in Muğla.

271 For that first comparison we used the complete ( $n = 250$ ) data set to be able to quantify the  
272 differences in membership coefficients for migratory and stationary colonies. But for the rest of the  
273 analysis we used the subset of data which is only composed of stationary colonies ( $n = 174$ ) since this  
274 would better reflect the population structure.

275 In the first scatter and the corresponding boxplots (respectively Fig. 5b for raw membership  
276 coefficients and Fig. 5a for transformed values) one can observe that within each locality samples are  
277 assigned with high proportions to their native clusters despite some admixed individuals. Also, one  
278 can see through observation of unpaired mean differences that Kırklareli, Ardahan, Artvin, Hatay and



279 to a lesser extent Düzce play role as centers of genuine subspecies diversity with exceptionally high  
280 levels and few individuals of admixed origin.

281 But when we compared isolated regions (Kırklareli, Ardahan, Artvin) and regions open to migratory  
282 beekeeping (consisting of Edirne+, Muğla, Ankara, Düzce+, Eskişehir+ and Hatay provinces in our  
283 sample) in terms of their arcsine transformed membership coefficients (Table 1 for means, effect  
284 sizes and significance of the difference and Fig. 5a second boxplot) we witnessed that -as expected-  
285 stationary colonies within isolated regions showed significantly higher fidelity to the original  
286 clusters. This is also obvious in the estimation plot in Fig. 5c where the mean membership  
287 coefficients of samples that are from regions open to migratory beekeeping (green colored group  
288 designated as <0>) fall beyond the 95% confidence interval of the estimated mean of the samples  
289 from the isolated regions (orchid colored group designated as <1>).

290 Even if the individuals are assigned with high probability to their own clusters, let's say with a 90%  
291 of probability, this means that 10% of their genome still belongs to other clusters. Given that there  
292 are four clusters, we investigated if any of these misassigned genome parts were enriched for any of  
293 them. Mean transformed values for Thracian cluster misassignments among individuals of the other  
294 populations were 0.16 and 0.25 for Anatolian cluster, 0.26 for the Caucasian and 0.20 for the  
295 Levantine (Fig. 6a).

296 A significant Kruskal-Wallis test ( $p < 0.001$ ) and a *post hoc* Dunn's test, accompanied by a  
297 significant ANOVA result ( $p < 0.001$ ) followed by a Tukey's test, showed that misassignments to *A.*  
298 *m. caucasica* and *A. m. anatoliaca* clusters were significantly more frequent than the others ( $p < 0.001$   
299 for both subspecies against C-lineage Thracian bees and  $p < 0.05$  against *syriaca* group). The effect  
300 sizes according to Cohen's *d* varied from 0.34 to 0.54 with estimation plots verifying the precision of  
301 the difference observed (Fig. 6b). Despite observation of the highest values in *A. m. caucasica*  
302 misassignments, the results were not significant between *A. m. caucasica* and *A. m. anatoliaca*  
303 clusters.

304 We checked if those differences result from many individuals with high admixture levels but such  
305 data only constituted the 7.5% of all the observations. This is with a threshold level of 0.5 for  
306 transformed values which corresponds to a second hybrid with a 25% contribution of non-native  
307 origin. So, we concluded that rather the main effect is due to consistent mid to low subspecies'  
308 contributions to other populations.

309 We also investigated if these small drifts in admixture proportions were more prominent in some  
310 localities and if populations differed in the subspecies they are receiving gene flow. This led us to  
311 comprehend the extent, magnitude and direction of the patterns of gene flow among the subspecies  
312 with a particular sensitivity to the populations. The results are summarized in Fig 7. We applied  
313 Dunn's test for each pairwise comparison between the populations. Significance of results will be  
314 mentioned but details of the test results can be found in the supplementary file.

315 Contributions from the Thracian cluster seem to be high in Düzce+ and Eskişehir+ which reside in  
316 the southeast of Marmara Sea across the Bosphorus and also there is some non-significant surplus in  
317 Muğla province in the Aegean coast. This is consistent with the Structure result for  $K = 2$ . While  
318 Thracian populations of Kırklareli and Edirne+ receive most gene flow from the Anatolian cluster  
319 these are not significantly different than Anatolian contributions to other regions which points to a  
320 balanced, uniform contribution of this subspecies to each group. Although non-significant, Edirne+  
321 receives much gene flow from that cluster. Caucasian cluster on the other hand contributes most to



322 Ankara, Muğla, Eskişehir+ and Düzce+ populations. Only significant differences are observed  
 323 between Ankara and Muğla receiver populations and Kırklareli in terms of admixture with Caucasian  
 324 populations. The same populations also significantly differed from Kırklareli in their admixture  
 325 levels with the Levantine cluster. In both cases Kırklareli population in Thrace had lower gene flow.  
 326 This is interesting since Muğla in the southwest and Caucasus region at the northeast lie at the  
 327 different extremes of the country.

## 328 Discussion

329  $F_{ST}$  values obtained were highly significant but they were lower than what Bodur et al. (2007)  
 330 estimated -a total  $F_{ST}$  of 0.077 together with higher values for pairwise comparisons among  
 331 populations- by samples collected ten years before ours. This may indicate a recent increased gene  
 332 flow and can be an alarm signal for a trend. Constant monitoring studies are needed in the future to  
 333 see if it is a persistent trend really. The high degree of structuring in stationary colonies according to  
 334  $F_{ST}$  results was lost in migratory ones, meaning they are less differentiated from each other due to  
 335 high degree of gene flow.

336 Phylogenetic tree clearly showed that Thracian samples were completely distinct from others  
 337 pointing to an early division of populations and limited gene flow. This supports the hypothesis for a  
 338 Carniolan (C-lineage) descent of Thracian bees in Turkey. Directly including samples from the major  
 339 C-lineage subspecies would confirm the subspecies of these bees highly differentiated from  
 340 Anatolian samples. In our initial observations of another research, Thracian samples grouped with C-  
 341 lineage European breeds rather than the samples throughout Anatolia (Kükreler, unpublished data).  
 342 This is a challenging point to Ruttner's claim (Ruttner 1988) that Thracian bees belong to *anatoliaca*  
 343 subspecies and needs further investigation.

344 West Anatolian, Levant and Caucasian populations did also form separate clusters in the tree. PCA  
 345 results confirmed those 4 different clusters inferred from tree topology. Bitlis+ samples resided with  
 346 Central and West Anatolian populations in both phylogenetic tree and PCA results (supplementary  
 347 file) but it should be kept in mind that all samples from that locality belonged to migratory colonies  
 348 so resampling with inclusion of stationary colonies from East Anatolia would be beneficial to  
 349 understand the real phylogenetic relations.

350 The two most possible K values in structure analysis for the whole sample and the stationary colonies  
 351 were  $K = 2$  and  $K = 4$ , both results supporting the hypotheses of populations belonging to 2 separate  
 352 lineages (C and O) and 4 distinct subspecies (a Carniolan ecotype in Thrace, *A. m. caucasica* in  
 353 Artvin and Ardahan, *A. m. syriaca* in Hatay and *A. m. anatoliaca* widely distributed covering the rest  
 354 of the country). In contrast to the expectations of migratory beekeepers of making use of native  
 355 stocks, results involving migratory beekeepers' samples lacked any population structuring in the  
 356 cluster analysis further clarifying the highly hybridized status of migratory apiaries.

357 Stationary apiaries, as expected, yielded highly structured groups where all the subspecies could be  
 358 detected. When K was 2, the structure analysis of two distinct clusters showed that there was a  
 359 transition zone between Thracian and Anatolian samples around Marmara Sea and Aegean. This may  
 360 be a hybrid zone between the C and O lineages like the ones identified before between M and C  
 361 lineages in Alps and Apennine Peninsula and between A and M lineages at the Iberian Peninsula and  
 362 Mediterranean islands (De la Rua et al. 2009). An analysis of ecological niches under species  
 363 distribution models suggest an intersection of habitat suitability of both subspecies within the  
 364 aforementioned geographic area (Kükreler, unpublished data).

365 When K was considered as 4, all four subspecies were easily differentiated from each other, in  
 366 accordance with the expectances. The significance of two distinct clusters ( $K = 2$ ) was higher than  
 367 four ( $K = 4$ ) which means that the differences between the populations belonging to C (in Thrace)  
 368 and O (in Anatolia) lineages are more clear-cut than differences between the populations of four  
 369 different subspecies.

370 *A. m. anatoliaca* samples fell in the middle of the other subspecies in ordinations, being similar to all  
 371 other populations according to FST values despite being a distinct cluster in structure analysis which  
 372 may point to a significant historical contribution to *A. m. anatoliaca* populations from the  
 373 neighboring regions. Another explanation can be that *anatoliaca* subspecies' putative basal position  
 374 for O-lineage honey bees places it as a center of genetic diversity. With *anatoliaca* bees exhibiting a  
 375 distinct identity, the situation was quite different than what was observed in all-migratory Bitlis+  
 376 samples where a mixture of different clusters surpassed instead of a separate identity.

377 A better understanding in terms of phylogenetic relationships between the populations in Turkey can  
 378 be developed if populations neighboring Anatolia and Thrace in Balkans, Iran, Caucasus and the  
 379 Middle East are also sampled. This can be a direction for future research, for shedding light on the  
 380 complicated taxonomic status within and between the C and O lineages and for drawing edges and  
 381 transition zones of the subspecies present across the whole region.

382 Results from different analyses conducted here confirmed the presence of clusters but also, they all  
 383 together pointed to the status of migratory colonies: they might be acting as a hybrid zone mobile in  
 384 space and time, being at one region in spring and at others in summer and fall, becoming vectors of  
 385 otherwise local gene combinations. Statistical results concerning a comparison between migratory  
 386 and stationary colonies confirmed the significant gene flow towards the migrants from local bees.

387 A significant gene flow towards local bees was also observed by a comparison between isolated  
 388 regions and those are not. This result, derived from directly contrasting two settings in an  
 389 experimental framework, is pointing to the vitality of establishing areas away and free from  
 390 migratory beekeeping for preservation of honey bee genetic diversity in conclusion with other studies  
 391 on conservation practices (Pinto et al. 2015; Oleksa et al. 2015).

392 One interesting point in the results was that the trend of the stationary colonies in Ankara. They had a  
 393 significantly lower probability of being assigned to their own clusters than the migratory colonies of  
 394 their province. This may be related with the regions migratory beekeepers of Ankara prefer to visit  
 395 during their migratory cycle or due to the insistent preference of using native queen bees by  
 396 migratory beekeepers. The low assignment degree of stationary colonies in Ankara may also be  
 397 related with Kazan apiary of TKV (Development Foundation of Turkey) placed there where hundreds  
 398 of colonies of Caucasian bees are raised and sold around for more than 30 years. The same practice is  
 399 also carried out by many queen bee breeders in Kazan region. Gene flow through these apiaries and  
 400 queen bees distributed locally by trade may contribute quietly to such an admixture observed in  
 401 stationary colonies in Ankara. The high misassignment probability of colonies in Ankara to the  
 402 Caucasian cluster also revealed such a process as probable.

403 It's hard to directly quantify the effect of queen and colony trade but unique features of Anatolia and  
 404 Thrace by availability of a number of naturally occurring subspecies renders possible the  
 405 understanding of their relative roles. Honeybees from stationary colonies were assigned more often to  
 406 their native clusters but they were also assigned to other clusters with lower probabilities. Samples in

407 the whole range of the study misassigned to Caucasian cluster more often than they were misassigned  
408 to others.

409 This is most probably due to wide distribution of Caucasian queen bees by trade. Migratory  
410 beekeeping is not practiced in Ardahan and Artvin where highly commercial Caucasian bees are  
411 native. Hence no bees go in or leave out the region as migratory colonies. So, the observed  
412 introgression of Caucasian alleles to the stationary colonies elsewhere whose beekeepers let them  
413 change their queens on their own rather than purchasing queens of different origins, could mainly be  
414 attributed to frequent queen bee and colony replacements in neighboring apiaries within those  
415 regions.

416 It is shown here that practicing of honey bee replacements increase the level of admixture within the  
417 gene pool. As previously discussed, a very high level of Caucasian introgression was observed in  
418 Ankara. *A. m. anatoliaca* alleles also showed high introgression especially in Edirne+ of Thrace  
419 region but also at average levels in other regions. These high levels may be related to this subspecies'  
420 geographical proximity to other populations which might have led to historical and recent gene  
421 exchange. By another explanation it can be related to the widespread practicing of migratory  
422 beekeeping by Western and Central Anatolian beekeepers throughout Turkey, rather than queen or  
423 colony replacements since there are very few commercial queen breeders within the distribution  
424 range of *A. m. anatoliaca*.

425 Results of the various statistical tests carried out and analysis applied in this study clearly showed  
426 that the genetic structure of honeybee populations in Turkey were highly conserved and still  
427 maintained. But it doesn't mean that the structure and diversity observed is secure. Rather it should  
428 be considered under threat since the anthropogenic factors leading to gene flows are still underway  
429 and keep admixing the populations.

430 A quiet interesting point was that, the preservation of population structure was achieved despite a  
431 very high number of colonies moved from one location to the other by migratory beekeeping practice  
432 and despite unregulated and frequent queen and colony sales. Future research may also need to focus  
433 on how this biodiversity and its structuring were preserved and its relation to natural selection.  
434 Further hypothesis can be formulated to distinguish the relative effects of natural selection and gene  
435 flow, the former could be so significant that it could potentially counterbalance the latter.

436 Genetic variation eventually leading to local adaptations with such significant outweighing effect can  
437 be considered as a valuable resource for honey bee populations in the global context at this time of  
438 unusual bee losses as well as global climate change. So, a better understanding of both present  
439 adaptation to local climate and geographic conditions as well as adaptive capacity to future changes  
440 would better be developed for the sake of the bees and their beneficiaries. A fair amount of effort  
441 should be invested on more studies focusing on candidate functional variants at the genome level that  
442 play role in due process in different parts of the world. Novel and innovative ways of coping with  
443 environmental and climatic stressors developed by honey bee populations or exploration of  
444 interesting patterns of convergent evolution are waiting ahead to be yet discovered.

445 Importance of establishing isolated regions was highlighted with genetic data. The results of the  
446 statistical tests showed a significant difference between the conservation of identity in and out of  
447 isolated regions with isolated regions staying purer in terms of subspecies composition. Such regions  
448 were proven to be effective in conservation of unique diversity present within.

449 In the light of this study we propose a renewed effort to address the need for massive establishing of  
450 such regions for conserving locally adapted native bees throughout the whole natural distribution of  
451 the species. This especially holds for underrepresented regions in terms of local diversity hotspots. A  
452 gap analysis aiming for complementarity in the planning of systematic conservation efforts are  
453 urgently needed globally.

454 In such isolated regions, naturally, migratory beekeeping as well as replacement of queen bees with  
455 non-native ones must be strictly prohibited and checked by relevant molecular monitoring  
456 techniques. However, these isolated regions should also be wide enough involving additional buffer  
457 zones where further restrictions on migratory beekeeping and bee trade are applied for efficient  
458 isolation and for fulfilling sufficient effective population sizes.

459 Thanks to increasing awareness in the last decade within the industry, now there are at least 11  
460 isolated regions in service or being established in Turkey through significant efforts of scientists and  
461 their collaboration with Turkish Beekeepers Association. There is an ever-growing need for  
462 establishing closer links with decision makers and stakeholders and necessity of investing more effort  
463 in communicating results of scientific studies in order to make the most out of them.

464 Queen bee trade is not currently subjected to any restrictions or regulations in Turkey and there are  
465 still very few pioneering measures within the natural distribution range, obviously not enough to  
466 guarantee the realized preservation in the next decades. Such measures should be applied from a  
467 conservation perspective to avoid extinction of native races, ecotypes and diversity present in these  
468 populations. Genetic similarity of donor and recipient populations should be considered while  
469 determining migration routes for migratory beekeepers and determining permissions for bee sales.

470 Central and western Anatolian populations suffer from significant gene flow from Caucasian  
471 populations as demonstrated by our results. Muğla and Ankara especially showed alarming levels of  
472 significant gene flow from other subspecies. This is not unexpected since the former receives  
473 millions of migratory colonies during the honeydew season.

474 Despite its wide range of distribution spanning Anatolia from one side to other, special consideration  
475 should be taken for preserving *A. m. anatoliaca* subspecies. The large and heterogenous native range  
476 of this subspecies permitted the evolution of numerous ecotypes including those in coastal, inner step  
477 or rainy forest ecosystems with noteworthy adaptations linked to their local environments.

478 The case with Hatay's *syriaca* populations too, can get worse and worse since the migratory  
479 beekeeping practice is heavily carried out in the region and queen bee replacement with non-native  
480 races was frequent throughout the last decade. This is mainly due to aggressive behavior, high  
481 swarming tendency and an infame for low levels of honey piling but the subspecies is also shown to  
482 exhibit some natural forms of varroa resistance (Kence et al. 2013). In the future this may end up in  
483 *A. m. syriaca* colonies getting limited to a few localities and apiaries since the range of the subspecies  
484 in Turkey is very narrow. A long-term conservation program considering improvement of traits that  
485 result in beekeepers staying away from that subspecies should be actualized immediately in this  
486 region too.

487 Thracian populations show a significant differentiation from the rest of the bees in Anatolia but the  
488 subspecies which they belong to is not characterized on a strong basis yet and this unique population  
489 is not registered officially like the case with *A. m. syriaca* of Hatay. Only subspecies officially  
490 recognized in Turkey is *A. m. caucasica* so identification and registration procedures for the others  
491 should be put into practice as soon as possible.

492 An improvement based on molecular genetic techniques can be applied to the ongoing conservation  
493 programs for the *A. m. caucasica* subspecies. It is interesting to note that we even detected hybrid  
494 individuals within the range of largest, oldest and heavily invested conservation area. This proposal  
495 for application of molecular monitoring techniques holds for other subspecies too.

496 Recently a registration procedure for Muğla bees as an Aegean coastal ecotype of *anatoliaca*  
497 subspecies with specific adaptations to resource phenology in the form of availability of honeydew  
498 obtained from scale insect *M. hellenica* of Turkish red pine *P. brutia* is under process. During the  
499 conservation and breeding efforts, an adequate level of use of molecular markers was achieved  
500 (Kükrer, unpublished data). More attention should be paid to genetically characterize *A. m. meda*  
501 subspecies that was out of the reach of this study and which can be threatened by anthropogenic  
502 factors listed and studied here.

503 Rather than queen bee replacement, it should be encouraged to use native bees improved for desired  
504 characters which are also locally adapted by definition. Such improved breeds would be used locally  
505 and not distributed in a country-wide manner so that local adaptations would still be preserved while  
506 bees are selected for resistance to pests and pathogens, hygienic behavior, reduced aggressiveness,  
507 reduced tendency for swarming, higher winter survival, higher productivity or for increased  
508 pollination. For obtaining better results in that, research concerning the smoothing, development and  
509 extension of breeding locally adapted native bees and artificial insemination techniques should be  
510 given higher priority and be adopted globally throughout the natural distribution range of local  
511 subspecies.

512 Our overall results answer arguments about the present situation of honey bee subspecies in Turkey  
513 but they also bear a wider interest to the community since they constitute an important pioneer  
514 attempt to quantify the effects of human impact. Our measurable and justified scientific findings on  
515 migratory beekeeping, queen and colony trade as well as conservation implications will hopefully be  
516 of some use for the decision makers and other stakeholders.

### 517 **Conflict of Interest**

518 The authors declare that the research was conducted in the absence of any commercial or financial  
519 relationships that could be construed as a potential conflict of interest.

### 520 **Author Contributions**

521 All authors conceived and planned the experiments and contributed to the field work. Mert Kükrer  
522 carried out the experiments and statistical analyses. All authors contributed to the interpretation of the  
523 results. Mert Kükrer took the lead in writing the manuscript. All authors provided critical feedback  
524 and helped shape the research, analysis and manuscript. Aykut Kence was in charge of overall  
525 direction and planning.

### 526 **Funding**

527 This study was funded by The Scientific and Technological Research Council of Turkey (project no:  
528 109T547), Republic of Turkey Ministry of Agriculture and Forestry (project no: TAGEM /11/AR-  
529 GE/13) and Middle East Technical University Revolving Funds (project no: 2007-16-12-00-3008).

### 530 **Acknowledgments**

531 We dedicate this paper to the memory of late Professor Aykut Kence who spent His last 20 years  
532 studying honey bee diversity in Turkey trying strongly to draw attention of researchers, decision  
533 makers and beekeepers to the conservation of locally adapted native bees as a precious legacy of our  
534 kind. He will be remembered also for defending theory of evolution to be taught in science  
535 curriculum and for training many valuable young evolutionary biologists under very harsh  
536 conditions. He passed away on Feb. 1. 2014 after the completion of the study and at the very  
537 beginning of manuscript preparation.

538 We'd like to thank numerous beekeepers who provided samples to the study and The Turkish  
539 Beekeepers Association. Special thanks to Dr. Devrim Oskay for his contribution in the form of  
540 consumables. We'd like to thank Mustafa Nail Cırık, Mehmet Ali Döke, Okan Can Arslan, Cansu  
541 Özge Tozkar, Mehmet Kayım and Eda Gazel Karakaş for the field work and Esin Öztürk and Ezgi  
542 Ersin as well as Batuhan Çağrı Yapan, Ayshin Ghalici, Gizem Kars and Batuhan Elçin for  
543 contributing lab sessions. Special thanks to Çiğdem Akın Pekşen and Cemal Can Bilgin for their  
544 advice on the early versions of the manuscript.

545 This manuscript has been released as a pre-print at  
546 <https://www.biorxiv.org/content/10.1101/154195v1>, (Kükrer et al. 2017) and includes elements from  
547 the corresponding authors thesis (Kükrer 2013).

### 548 **References**

549 Akkaya H, Alkan S. 2007. Beekeeping in Anatolia from the Hittites to the present days. *Journal of*  
550 *Apicultural Research* 46:120-124.

551 Blacquiere T, Smaghe G, Van Gestel CA, Mommaerts V. 2012. Neonicotinoids in bees: a review on  
552 concentrations, side-effects and risk assessment. *Ecotoxicology* 21:973-992.

553 Bodur C, Kence M, Kence A. 2007. Genetic structure of honey bee, *Apis mellifera* L.  
554 (Hymenoptera:Apidae) populations of Turkey inferred from microsatellite analysis. *Journal of*  
555 *Apicultural Research* 46:50-56.

556 Bouga M et al. 2011. A review of methods for discrimination of honey bee populations as applied to  
557 European beekeeping. *Journal of Apicultural Research* 50:51-84.

558 Breeze TD, Bailey AP, Balcombe KG, Potts SG. 2011. Pollination services in the UK: how important  
559 are honey bees? *Agriculture, Ecosystems & Environment* 142:137–143.

560 Brown MJF, Paxton RJ. 2009. The conservation of bees: a global perspective. *Apidologie* 40:410–416.

561 Büchler R, Cecilia Costa, Fani Hatjina, Sreten Andonov, Marina D Meixner, Yves Le Conte,  
562 Aleksandar Uzunov, Stefan Berg, Malgorzata Bienkowska, Maria Bouga, Maja Drazic, Winfried  
563 Dyrba, Per Kryger, Beata Panasiuk, Hermann Pechhacker, Plamen Petrov, Nikola Kezić, Seppo  
564 Korpela & Wilde J. 2014. The influence of genetic origin and its interaction with environmental effects  
565 on the survival of *Apis mellifera* L. colonies in Europe, *Journal of Apicultural Research*, 53:2, 205-  
566 214, DOI: 10.3896/IBRA.1.53.2.03

567 Cánovas F, de la Rúa P, Serrano J, Galián J. 2011. Microsatellite variability reveals beekeeping  
568 influences on Iberian honeybee populations. *Apidologie* 42:235-251.

- 569 Champely, S., Ekstrom, C., Dalgaard, P., Gill, J., Weibelzahl, S., Anandkumar, A., ... & De Rosario,  
570 M. H. 2018. Package 'pwr'. R package version, 1-2.
- 571 Cridland, J. M., Tsutsui, N. D., & Ramírez, S. R. (2017). The Complex Demographic History and  
572 Evolutionary Origin of the Western Honey Bee, *Apis Mellifera*. *Genome biology and evolution*, 9(2),  
573 457-472.
- 574 Dall'Olio R, Marino A, Lodesani M, Moritz RF. 2007. Genetic characterization of Italian honeybees,  
575 *Apis mellifera ligustica*, based on microsatellite DNA polymorphisms. *Apidologie* 38:207-217.
- 576 De la Rúa P, Jaffé R, Dall'Olio R, Muñoz I, Serrano J. 2009. Biodiversity, conservation and current  
577 threats to European honey bees. *Apidologie* 40:263-284.
- 578 Dietemann V, Pirk CWW, Crewe R. 2009. Is there a need for conservation of honey bees in Africa?  
579 *Apidologie* 40:285-295.
- 580 Dinno, A. 2017. Package 'dunn.test'. CRAN Repos.
- 581 Earl DA, vonHoldt BM. 2012. STRUCTURE HARVESTER: a website and program for visualizing  
582 STRUCTURE output and implementing the Evanno method. *Conservation Genetics Resources* 4:359-  
583 361.
- 584 El-Niweiri MA, Moritz RF. 2010. The impact of apiculture on the genetic structure of wild honeybee  
585 populations (*Apis mellifera*) in Sudan. *Journal of Insect Conservation* 14:115-124.
- 586 Estoup, A., Garnery, L., Solignac, M., & Cornuet, J. M. (1995). Microsatellite variation in honey bee  
587 (*Apis mellifera* L.) populations: hierarchical genetic structure and test of the infinite allele and stepwise  
588 mutation models. *Genetics*, 140(2), 679-695.
- 589 European Parliament. (2017). The EU's beekeeping sector. At a Glance.  
590 [https://www.europarl.europa.eu/RegData/etudes/ATAG/2017/608786/EPRS\\_ATA%282017%29608](https://www.europarl.europa.eu/RegData/etudes/ATAG/2017/608786/EPRS_ATA%282017%29608786_EN.pdf)  
591 [786\\_EN.pdf](https://www.europarl.europa.eu/RegData/etudes/ATAG/2017/608786/EPRS_ATA%282017%29608786_EN.pdf) (Accessed 28.04.2020)
- 592 Evans JD, Schwarz RS. 2011. Bees brought to their knees: microbes affecting honey bee health. *Trends*  
593 *in Microbiology* 19:614-620.
- 594 Excoffier L, Lischer HEL. 2010. Arlequin suite ver 3.5: a new series of programs to perform population  
595 genetics analyses under Linux and Windows. *Molecular Ecology Resources* 10:564-567.
- 596 Genersch E. 2010. Honey bee pathology: current threats to honey bees and beekeeping. *Applied*  
597 *Microbiology and Biotechnology* 87:87-97.
- 598 Goudet J. 1999. PCAGEN, a computer package which performs principal component analysis (PCA)  
599 on gene frequency data. Université de Lausanne, Lausanne, Switzerland. Available from  
600 <http://www2.unil.ch/popgen/softwares/pcagen.htm> (accessed July 2015).
- 601 Hammer, Ø., Harper, D.A.T., Ryan, P.D. 2001. PAST: Paleontological statistics software package for  
602 education and data analysis. *Palaeontologia Electronica* 4(1): 9pp. [http://palaeo-](http://palaeo-electronica.org/2001_1/past/issue1_01.htm)  
603 [electronica.org/2001\\_1/past/issue1\\_01.htm](http://palaeo-electronica.org/2001_1/past/issue1_01.htm)



- 604 Han, F., Wallberg, A., & Webster, M. T. (2012). From where did the Western honeybee (*Apis*  
605 *mellifera*) originate? *Ecology and evolution*, 2(8), 1949-1957.
- 606 Harpur, B. A., Kent, C. F., Molodtsova, D., Lebon, J. M., Alqarni, A. S., Owayss, A. A., & Zayed, A.  
607 (2014). Population genomics of the honey bee reveals strong signatures of positive selection on worker  
608 traits. *Proceedings of the National Academy of Sciences*, 111(7), 2614-2619.
- 609 Hewitt, G. M. 1999. Post-glacial re-colonization of European biota. *Biological Journal of the Linnean*  
610 *Society*, 68, 87–112.
- 611 Ho, J., Tumkaya, T., Aryal, S. et al. 2019. Moving beyond P values: data analysis with estimation  
612 graphics. *Nat Methods* 16, 565–566 <https://doi.org/10.1038/s41592-019-0470-3>
- 613 Holleley C.E. and Geerts P.G. (2009). Multiplex Manager 1.0: a crossplatform computer program that  
614 plans and optimizes multiplex PCR. *BioTechniques*, 46(7), 511-517.
- 615 Jakobsson M, Rosenberg NA. 2007. CLUMPP: a cluster matching and permutation program for  
616 dealing with label switching and multimodality in analysis of population structure. *Bioinformatics*  
617 23:1801-1806.
- 618 Kandemir I, Kence M, Kence A. 2005. Morphometric and electrophoretic variation in different honey  
619 bee (*Apis mellifera* L.) populations. *Turkish Journal of Veterinary and Animal Science* 29:885-890.
- 620 Kandemir, I., Kence, M., Sheppard, W. S., & Kence, A. (2006). Mitochondrial DNA variation in honey  
621 bee (*Apis mellifera* L.) populations from Turkey. *Journal of apicultural research*, 45(1), 33-38.
- 622 Kence M, Farhoud HJ, Tunca RI. 2009. Morphometric and genetic variability of honey bee (*Apis*  
623 *mellifera* L.) populations from northern Iran. *Journal of Apicultural Research* 48:247-255.
- 624 Kence M, Oskay D, Giray T et al (2013) Honeybee colonies from different races show variation in  
625 defenses against the varroa mite in a ‘common garden’. *Entomol Exp Appl* 149(1):36–43
- 626 Kükreler M, Kence M and Kence A. 2017. Genetic evidences for the impact of anthropogenic factors on  
627 honey bee diversity. *bioRxiv* <https://www.biorxiv.org/content/10.1101/154195v1> (Accessed  
628 28.04.2020)
- 629 Kükreler M 2013. Genetic diversity of honey bee populations in turkey based on microsatellite markers:  
630 a comparison between migratory versus stationary apiaries and isolated regions versus regions open to  
631 migratory beekeeping. (Master’s thesis) Ankara Turkey: Middle East Technical University.  
632 <http://etd.lib.metu.edu.tr/upload/12615972/index.pdf> (Accessed 28.04.2020)
- 633 Langella O. 2011. Population genetic software: individuals or populations distances based on allelic  
634 frequencies, phylogenetic trees, file conversions. *Bioinformatics Organization, Inc.* Hudson,  
635 Massachusetts, USA. Available from [http://www.bioinformatics.org/project/?group\\_id=84](http://www.bioinformatics.org/project/?group_id=84) (accessed  
636 July 2015).
- 637 Letunic I, Peer Bork, Interactive Tree Of Life (iTOL) v4: recent updates and new developments,  
638 *Nucleic Acids Research*, Volume 47, Issue W1, 02 July 2019, Pages W256–W259,  
639 <https://doi.org/10.1093/nar/gkz239>

- 640 Meffe GK. 1998. The potential consequences of pollinator declines on the conservation of biodiversity  
641 and stability of food crop yields. *Conservation Biology* 12:8-17.
- 642 Morse RA. 1991. Honey bees forever. *Trends in Ecology & Evolution* 6:337–338.
- 643 Neumann P, Carreck NL. 2010. Honey bee colony losses. *Journal of Apicultural Research* 49:1-6.
- 644 Oldroyd BP. 2007. What's killing American honey bees? *PLoS Biology* (e168) DOI:  
645 10.1371/journal.pbio.0050168.
- 646 Oleksa A, Igor Chybicki, Adam Tofilski & Burczyk J. 2011. Nuclear and mitochondrial patterns of  
647 introgression into native dark bees (*Apis mellifera mellifera*) in Poland, *Journal of Apicultural*  
648 *Research*, 50:2, 116-129, DOI: 10.3896/IBRA.1.50.2.03
- 649 Pinto MA, Dora Henriques, Julio Chávez-Galarza, Per Kryger, Lionel Garnery, Romée van der Zee,  
650 Bjørn Dahle, Gabriele Soland-Reckeweg, Pilar de la Rúa, Raffaele Dall' Olio, Norman L Carreck &  
651 Johnston JS. 2014. Genetic integrity of the Dark European honey bee (*Apis mellifera mellifera*) from  
652 protected populations: a genome-wide assessment using SNPs and mtDNA sequence data, *Journal of*  
653 *Apicultural Research*, 53:2, 269-278, DOI: 10.3896/IBRA.1.53.2.08
- 654 Pritchard JK, Stephens M, Donnelly P. 2000. Inference of population structure using multilocus  
655 genotype data. *Genetics* 155:945-959.
- 656 Reynolds J, B. S. Weir and C. Clark Cockerham 1983. Estimation of the coancestry coefficient: basis  
657 for a short-term genetic distance. *Genetics* 105:767-779.
- 658 Rosenberg NA. 2004. DISTRUCT: a program for the graphical display of population structure.  
659 *Molecular Ecology Notes* 4:137-138.
- 660 R Core Team. (2013). R: A language and environment for statistical computing.
- 661 Ruttner F. 1988. Biogeography and taxonomy of honey bees. In Springer V editor. New York.
- 662 Shaibi, T., Lattorff, H. M. G., & Moritz, R. F. A. (2008). A microsatellite DNA toolkit for studying  
663 population structure in *Apis mellifera*. *Molecular Ecology Resources*, 8(5), 1034-1036.
- 664 Solignac, M., Vautrin, D., Loiseau, A., Mougel, F., Baudry, E., Estoup, A. ... & Cornuet, J. M. (2003).  
665 Five hundred and fifty microsatellite markers for the study of the honeybee (*Apis mellifera* L.) genome.  
666 *Molecular Ecology Notes*, 3(2), 307-311.
- 667 Torchiano, Marco. (2016, November 13). Effsize - a package for efficient effect size computation.  
668 Zenodo. <http://doi.org/10.5281/zenodo.196082>
- 669 Tunca R. I. (2009). Determination and comparison of genetic variation in honey bee (*Apis mellifera*  
670 L.) populations of Turkey by random amplified polymorphic DNA and microsatellite analyses, Ph. D.  
671 Thesis, Middle East Technical University. Van Engelsdorp D et al. 2009. Colony collapse disorder: a  
672 descriptive study. *PloS One* (e6481) DOI: 10.1371/journal.pone.0006481.
- 673 USDA NASS. (2019). Statistical Summary: Honey Bees. NASS Highlights  
674 [https://www.nass.usda.gov/Publications/Highlights/2019/2019\\_Honey\\_Bees\\_StatisticalSummary.pdf](https://www.nass.usda.gov/Publications/Highlights/2019/2019_Honey_Bees_StatisticalSummary.pdf)  
675 (Accessed 28.04.2020)

676 Van Engelsdorp D, Meixner MD. 2010. A historical review of managed honey bee populations in  
 677 Europe and the United States and the factors that may affect them. *Journal of Invertebrate Pathology*  
 678 103:S80-S95.

679 Wallberg, A., Han, F., Wellhagen, G., Dahle, B., Kawata, M., Haddad, N., ... Webster, M. T. (2014).  
 680 A worldwide survey of genome sequence variation provides insight into the evolutionary history of the  
 681 honeybee *Apis mellifera*. *Nature Genetics*, 46(August), 1081–1088. <http://doi.org/10.1038/ng.3077>

682 Weinstock GM et al. 2006. Insights into social insects from the genome of the honey bee *Apis*  
 683 *mellifera*. *Nature* 443:931-949.

684 Whitfield CW, Behura SK, Berlocher SH, Clark AG, Johnston JS, Sheppard WS, Smith DR, Suarez  
 685 AV, Weaver D, Tsutsui ND. 2006. Thrice out of Africa: ancient and recent expansions of the honey  
 686 bee, *Apis mellifera*. *Science* 314:642-645.

687 **Tables**

688 **Table 1.** Genetic impact of beekeeping and conservation practices on (arcsine root square  
 689 transformed) membership coefficients to native clusters (\*\*  $p < 0.05$ , \*\*  $p < 0.01$  and \*\*\*  $p < 0.001$ ).

690

	n	n				
	Migratories	Stationaries	Stationaries	Migratories	Cohen's <i>d</i>	<i>U</i> and <i>t</i> tests significance
<b>Beekeeping Practice</b>						
<b>Ankara</b>	9	18	0,82	1,11	-1,00	*
<b>Muğla</b>	15	21	0,93	0,70	0,89	*
<b>Hatay</b>	13	23	1,20	0,66	2,01	***
<b>Combined</b>	37	62	1,00	0,79	0,66	**
<b>Overall</b>	76	174	1,06	0,72	1,22	***
<b>Conservation Practice</b>						
<b>Isolated</b>	NA	79	NA	1,21	0,49	***
<b>Not</b>						
<b>Isolated</b>	NA	95	NA	1,08		

691

692 **Supplementary material**

693 R codes and Supplementary File.

694 **Figure legends**

695 Figure 1. Geographic distribution of (a) major honey bee (*A. mellifera*) subspecies in and around  
 696 Anatolia (b) sampling sites and sample sizes.

697 Figure 2. (a) PCA of stationary colonies, Component 1: 41.8%, Component 2: 32.1% (b) UPGMA  
 698 tree of honey bee populations based on Reynolds' 1983 genetic distances (orange: Thracian, yellow:  
 699 Anatolian, blue: Caucasian, violet: Levantine clusters).

700 Figure 3. Estimated population structure and clustering of honeybees in Anatolia and Thrace for (a)  
701 migratory colonies (b) stationary colonies (c) the whole sample (orange: Thracian, yellow: Anatolian,  
702 blue: Caucasian, violet: Levantine clusters).

703 Figure 4. Comparison between stationary (Sta\_) and migratory (Mig\_) colonies in Ankara, Muğla  
704 and Hatay, as well as these three provinces combined and the whole data set,  $n = 250$  (a) boxplot  
705 display of arcsine root square transformed membership coefficients (b) scatter plot with estimations  
706 of mean differences based on raw individual membership coefficients (yellow: Ankara and Muğla  
707 belonging to Anatolian cluster, violet: Levantine cluster, coral: for a combination of three provinces,  
708 firebrick: whole data ).

709 Figure 5. Comparison between isolated regions and regions that are open to migratory beekeeping (a)  
710 first boxplot display the arcsine root square transformed membership coefficients for 9 populations  
711 whereas the second one presents a comparison of samples within isolated regions and those are not  
712 (b) scatter plot with estimations of mean differences based on raw individual membership  
713 coefficients to the native clusters (c) scatter plot contrasting individual raw membership coefficients  
714 with an estimation of mean differences (orange: Thracian, yellow: Anatolian, blue: Caucasian, violet:  
715 Levantine clusters, orchid and “1”: isolated regions, green and “0”: regions open to migratory  
716 beekeeping).

717 Figure 6. Comparison of misassignment proportions between the major clusters (a) boxplot display of  
718 arcsine root square transformed membership coefficients (b) scatter plot with estimations of mean  
719 differences based on raw individual membership coefficients (orange: Thracian, yellow: Anatolian,  
720 blue: Caucasian, violet: Levantine clusters).

721 Figure 7. Patterns of gene flow between populations (a) boxplot display of arcsine root square  
722 transformed membership coefficients (b) scatter plot with estimations of mean differences based on  
723 raw individual membership coefficients (orange: Thracian, yellow: Anatolian, blue: Caucasian,  
724 violet: Levantine clusters).

725

726

727

728

729


730

731



732

733




734



735









737



(a)




(b)




739


(a)



(b)




(c)






(a)

Misassignments to Clusters



(b)





741