

# **Moving on from the insect apocalypse narrative: engaging with evidence-based insect conservation**

Manu E. Saunders<sup>1,2</sup>, Jasmine Janes<sup>1,3</sup>, James O'Hanlon<sup>1</sup>

<sup>1</sup>School of Environmental and Rural Science, University of New England Armidale NSW  
Australia

<sup>2</sup>UNE Business School, University of New England Armidale NSW Australia

<sup>3</sup>Biology Department, Vancouver Island University, Nanaimo, BC, Canada

This is the author's version of a manuscript published in BioScience. Please cite as:

Saunders ME, Janes J, O'Hanlon J (2019) Moving on from the insect apocalypse narrative: engaging with evidence-based insect conservation. *BioScience*  
<https://doi.org/10.1093/biosci/biz143>

## **Abstract**

Recent studies showing temporal changes in local and regional insect populations received exaggerated global media coverage. Confusing and inaccurate science communication on this important issue could have counter-productive effects on public support for insect conservation. The ‘insect apocalypse’ narrative is fuelled by a limited number of studies that are restricted geographically (predominantly UK, Europe, USA) and taxonomically (predominantly bees, macrolepidoptera, and ground beetles). Biases in sampling and analytical methods (e.g. categorical vs. continuous time series, different diversity metrics) limit the relevance of these studies as evidence of generalised global insect decline. Rather, the value of this research lies in highlighting important areas for priority investment. We summarise research, communication and policy priorities for evidence-based insect conservation, including key areas of knowledge to increase understanding of insect population dynamics. Importantly, we advocate for a balanced perspective in science communication to better serve both public and scientific interests.

## **Introduction**

Insects are the most diverse and abundant group of animals on Earth and are critical drivers of ecosystem function in terrestrial and aquatic systems; yet the majority of insect taxa are understudied, publicly misunderstood and face numerous environmental threats (Samways 2007; Cardoso et al. 2011). Public support is essential to address knowledge gaps and build global support for insect conservation. We need immediate solutions to human behaviours and management strategies that have detrimental impacts on insect biodiversity (e.g. habitat destruction, pesticide overuse), and we need broad support for publicly-funded research that fills critical knowledge gaps. Science communication efforts can help build this support, but maintaining accuracy in communication narratives, whether presented via popular media platforms, scientific literature or research institutions, is essential (Dahlstrom and Ho 2012). Sensationalising geographically and taxonomically limited studies as evidence of global patterns may grab attention, but can also have unwanted side effects. In particular, doom and gloom messaging rarely works to galvanise public support (McAfee et al. 2019) and strong negative messaging (e.g. apocalypse narratives) can undermine the credibility of science, especially as more facts become available (Horeis 2009).

## **The Insect Apocalypse narrative**

“The Insect Apocalypse is here” - *New York Times*, 27 November 2018


“Plummeting insect numbers ‘threaten collapse of nature’” - *The Guardian*, 11 February 2019

“Global insect decline may see ‘plague of pests’” - *BBC News*, 11 February 2019

Recent popular media coverage claimed that we are facing apocalyptic global insect declines and extinction of all insects within 100 years (e.g. Carrington 2017; Jarvis 2018; McGrath 2019). The only exceptions, according to some news reports, are pest insects, which are apparently on track to reproduce out of control (e.g. McGrath 2019). This exaggerated and unlikely narrative, which spread across multiple media platforms over the course of many months, grew from coverage of two empirical studies that show temporal changes in local insect populations in Germany (Hallmann et al. 2017) and Puerto Rico (Lister and Garcia 2018), and a review paper that claimed to show evidence of “worldwide decline” of insect fauna (Sánchez-Bayo and Wyckhuys 2019). While Hallmann et al. and Lister and Garcia did not specifically connote global insect apocalypse in their peer-reviewed articles, the associated media releases from the lead authors’ respective institutions extrapolated the limited results to suggest broader scale declines; in the case of Hallmann et al. (2017), a Radboud University (2017) release implied the study demonstrated global declines, while a Rensselaer Polytechnic Institute release suggested Lister and Garcia’s (2018) results represented declines across all of Puerto Rico’s insects, with implications for tropical rainforests generally (Martialay 2018). In their peer-reviewed paper, Sánchez-Bayo and Wyckhuys (2019) extrapolated beyond the limitations of their review to suggest evidence of global decline across all insect taxa, and the associated The University of Sydney (2019) release corroborated this. The ‘insect apocalypse’ narrative spread unchecked via popular and social media platforms and subsequent peer-reviewed literature have cited all of these studies uncritically as evidence of global-scale insect decline or ecosystem collapse (e.g. Pascal et al. 2019; Vilcinskas 2019; Gillespie et al. 2019).

## **Limitations of the apocalypse narrative**

The aforementioned studies certainly deserve discussion, and contribute to understanding the extent of our knowledge of insect communities. But they do not provide evidence of global-scale insect declines and impending extinction of all insect life on Earth. The reality is far more complex (Saunders 2019; Wagner 2019; Thomas et al. 2019; Simmons et al. 2019). The popular ‘insect apocalypse’ narrative is largely based on a body of literature comprised of Hallmann et al. (2017), Lister and Garcia (2018) and the 78 studies presented as evidence of global decline by Sánchez-Bayo and Wyckhuys (2019). The narrative is presented as fact, but despite the strong negative messaging, the evidence of global insect apocalypse is ambiguous (see supplemental materials). Other researchers have published general critiques of these papers and their results (Wagner 2019; Willig et al. 2019; Thomas et al. 2019; Simmons et al. 2019), but there has been little critical assessment of the variation in methodological details and population trends behind the insect apocalypse narrative (see supplemental materials). Most of these studies are from Europe and the UK (Figure 1), and many are highly localised, collecting data from one specific location (e.g. one nature reserve) or from one region within a country. All studies found increases or no changes for some of the focal taxa across the analysed time period. Indeed, some studies could not confirm declines for any of the focal taxa e.g. (Petanidou et al. 2011; Gardner and Spivak 2014) and, in most studies, more taxa showed increases or no change relative to the number of taxa showing declines (see supplemental materials).



**Figure 1:** Geographic and taxonomic distribution of the body of literature behind the popular insect apocalypse narrative. This figure does not show trends, only the presence of an empirical study measuring changes in insect populations. See supplemental materials for full details of studies depicted in the figure. Insect taxa are grouped based on the groupings used within studies. ‘Unspecified’ indicates studies that sampled insects generally without specifying particular taxa. Each cell indicates the number of studies focused on a particular taxonomic group in a particular country i.e. (country-taxa pairs). White cells indicate no studies, and cell colour shade increases in darkness with a greater number of studies focused on that country-taxa pair.

The limitations of the insect apocalypse narrative become clear when attempting to scale these studies up to global long-term patterns. Most studies treated time as a categorical variable, analysing differences between time periods rather than change over time. This approach can be misleading, as arbitrary time periods may be a poor proxy for environmental change or ecological interactions. In addition, analysis methods based on comparing occupancy across spatial grid cells in different time periods (Williams 1982; Desender and Turin 1989) can overlook correlations within the dataset (e.g. co-dependant or interacting species) or other influential covariates that vary over time (e.g. land use change). More recently, trait-based analyses have been used to assess some of these confounding factors. For example, Paukkunen et al. (2018) investigated cuckoo wasp (Hymenoptera: Chrysididae) abundance and distribution over time in Finland. They found that trends of cuckoo wasps were positively correlated with those of their host species, and declining trends were more apparent in scarce, small-bodied species that used above-ground nesting hosts. Considering feeding habit and life cycle traits can also reveal pattern variation between species. Warren et al. (2001) show that specialist butterfly species in the UK have declined in distribution, while mobile generalist species have increased, and Ball-Damerow et al. (2014) showed contrasting patterns for migratory and habitat specialist Odonata in arid western parts of the United States.

Confirmed declines for a taxonomic group in one location do not always translate to other regions or similar species. Biesmeijer et al. (2006) found contrasting patterns for bees and hoverflies when comparing trends in Britain and the Netherlands, while Brooks et al. (2012) found that carabid (Coleoptera: Carabidae) abundance across the UK declined significantly in

some regions and increased in others. Bumble bees (Hymenoptera: *Bombus* spp.) were one of the most commonly studied groups, with some species showing variable trends in different studies. For example, *B. pensylvanicus* has been identified as declining in southern Ontario Colla and Packer (2008), data deficient in New Hampshire (Jacobson et al. 2018), and the most common and stable species in Oklahoma (Figueroa and Bergey 2015). Similarly, the diversity of community-level and population-level metrics used across studies make it difficult to identify patterns at larger scales, because different components of biodiversity can tell vastly different stories. Schuch et al. (2012) studied Auchenorrhyncha communities in central European grasslands, and found declines in abundance but no change in species richness. Jacobson et al. (2018) found that abundance of some bumble bee species in New Hampshire have declined, but there was no trend apparent for plant-bumble bee interactions. The relevance of different metrics to identifying overall global population trends was also largely overlooked or misrepresented in popular media coverage; for example, The Guardian coverage of the Hallmann et al. (2017) study claimed that insect ‘abundance’ had fallen, when the study only measured biomass (Carrington 2017).

The insect apocalypse narrative is built from exaggerated interpretations of a limited number of studies, and ignores other recent empirical literature on insect populations showing long-term increases for some taxa (Herrera 2018; Gibb et al. 2019), no discernible trends in other taxa (Bell et al. 2015), and new species or records in understudied locations (Leijs et al. 2018; Meiners et al. 2019). Very few popular and social media reports on the ‘insect apocalypse’ made an attempt to reconcile the complexities inherent in insect population dynamics for non-specialist audiences. We are still not certain how many insect species exist on Earth, nor do we know the distribution, life history, or ecology of most described species. Therefore, our



current knowledge of global insect biodiversity is too limited to support strong negative messaging about global patterns in insect population dynamics.

Coordinated monitoring systems that collect standardised observations on multiple taxonomic orders across consecutive years (e.g. UK's Rothamsted Insect Survey) are extremely valuable for identifying temporal patterns in insect populations, but are not the only priority for understanding global insect diversity (Cardoso and Leather 2019). We believe that the insect apocalypse narrative detracts from the key issues of insect conservation, and potentially damages public understanding of insect ecology. Here, we highlight a more constructive path for evidence-based discussion of insect conservation involving: (i) priorities for research, communication and policy to support evidence-based insect conservation (Box 1); and (ii) a more balanced presentation of insect ecology in science communication to better serve both public and scientific interests.

**Box 1: Priorities for building support for evidence-based insect conservation**

*Fundamental and applied research*

Develop:

- cost-effective non-destructive insect monitoring techniques.
- robust statistical methods for analysing unverified citizen science data.
- trait-based frameworks for understanding human impacts on insect communities.
- guidelines to facilitate standardised protocols for the use and analysis of genetic-based monitoring (e.g. eDNA, metabarcoding, metagenomics) that complement field-based efforts.

Identify:

- how species turnover influences population trends within communities and meta-communities.
- relationships between community-level processes and species population trends.
- how population fluctuations vary in time and space.
- how different components of insect biodiversity (including trait variability) contribute to ecosystem function and associated ecosystem services.

Increase:

- availability of unpublished long-term datasets.
- use of multidisciplinary approaches to better understand species biology, interactions and overall community resilience, e.g. field survey + eDNA identification; pollinator network + metabarcoding; trophic assessments + metagenomics/metabarcoding of gut/faeces; ecological + genetic distance/resistance layers to identify landscape drivers.

*Communication and education*

Develop:

- robust citizen science programs that combine community education and training with long-term monitoring data collection
- new technologies for user-friendly insect identification

Increase:

- school and community education programs focused on the value of natural history observation.
- funding and support for local and regional natural history journals, including digitisation programs.
- taxonomy and entomology training in universities.
- media ethics training for scientists.
- science communication focused on scientific methods and processes.

### *Policy and management*

#### Identify:

- how policies and management strategies influence changes in insect communities.
- better regulatory frameworks for sale and use of insecticides and sustainable land management to protect insect biodiversity.

#### Increase:

- incentives and consumer support for non-conventional food production systems (e.g. organic, regenerative agriculture).
- incorporation of genetic/genomic data with ecological data to better inform policy and management.

## **Priority areas for research and funding**

### *Population dynamics of insect species*

Animal populations are never truly stable (Solomon 1949; Wallner 1987). They constantly fluctuate in response to multiple factors including resource availability, climatic conditions, and biotic interactions (Wallner 1987; Gaston and Lawton 1988). Insects generally have short, seasonal life cycles, large generation sizes, and can exhibit significant niche separation between life stages; hence, insect population fluxes are incredibly dynamic (Wallner 1987; Gaston and Lawton 1988). Cyclic outbreaks are common for many species, often influenced by trophic interactions and resource pulses (Ostfeld and Keesing 2000). Indeed, mass insect outbreaks occur regularly with significant impacts on ecosystems, industry and agriculture (Esper et al. 2007; Büntgen et al. 2009) and some systems, such as temperate forests and agroecosystems, are predicted to experience increasing insect outbreaks because of global climate change (Porter et al. 1991; Lovett et al. 2006). Yet factors that contribute to outbreaks are numerous and often difficult to identify (Wallner 1987; Buma 2015).

Managed systems (e.g. crop fields or tree plantations) tend to be more susceptible to pest insect outbreaks because of frequent disturbance, low diversity and spatial uniformity (Coyle et al. 2005; Dalin et al. 2009). Identifying variation in outbreaks between different groups of species (e.g. herbivores and predators) in managed and natural systems is important to understand how to manage systems sustainably to reduce pest impacts (Knops et al. 1999; Tschardtke et al. 2012). Such fluctuations are a fundamental component of ecosystems (Buma 2015), but it is unclear how important these sporadic outbreaks are for the maintenance of insect populations in the long term. Community-level patterns and processes that influence local abundance also fluctuate as the size and distribution of local populations varies over time (Elton 1924). In addition, sex ratios for many insect species are often overdispersed (Godfray and Werren 1996) and can be readily adjusted by environmental drivers, such as resource limitation or mate competition (Charnov et al. 1981; Chapuisat et al.

1997). Invasive species can also alter population structure and long-term trends; for example, the arrival of an exotic gall wasp, *Andricus quercuscalicis*, in the UK has caused a significant male bias in native parasitoid populations that emerge from the galls (Schönrogge et al. 2000), but it is unclear what long-term effect this will have on parasitoid populations in the long term. These are some of the reasons why relying on few sampling seasons, or comparing surveys conducted many years apart, can be misleading when attempting to pinpoint population declines.

Long-term monitoring of insect communities at particular sites can help elucidate these dynamics, but can be costly and counter-productive if not designed effectively (Tepedino et al. 2015). In addition, any changes observed at individual sites can only be interpreted as variation in characteristics of that site, not true variations in the insect population (McArdle and Gaston 1993). For many species, local extinction is often impossible to confirm (McArdle 1990). To rigorously confirm a decline in populations requires ongoing censuses of those populations, which for insects is rarely feasible given their fast generation time, elusiveness, cryptic life cycles, and the inherent difficulties of defining a population in the first place. A more immediate solution is to invest in building knowledge of global insect diversity, insect species ecology and life histories, and the effect of human impacts and environmental change on taxonomic groups (Box 1).

### *Accessible taxonomy*

Estimates of numbers of insect species appear regularly in the literature, ranging from 3 million (Blaxter 2003) to over 30 million (Stork 1988). Whatever the correct number, we have described only a fraction of the hypothetical global diversity; approximately 1 million

insect species are currently known (Stork 2018). Although all taxonomic levels and functional groupings are valuable units of enquiry, 'species' remain the basic unit of biology and conservation. To understand the ecology of an organism, it is essential to first accurately identify it (Mallet and Willmott 2003).

Our current identification system relies on people with highly-specialised training and knowledge (i.e. taxonomists) using discipline-specific rules and hypotheses (Lipscomb et al. 2003). The process of describing a new species can take decades, but in brief, a formal description is published and a type specimen is designated and then stored as a reference for all other material (Godfray 2002). The problems with this system are numerous and well-known: (i) type specimens can be lost or damaged; (ii) access to specialist knowledge for general users is limited (Tautz et al. 2003); (iii) complex descriptive terminology make existing keys inaccessible; and (iv) reference specimens are typically held in restricted access museum departments (Alberch 1993).

A more concerning issue is the decline in funding and succession for taxonomic specialists. Much information is lost when a specialist taxonomist retires (Tautz et al. 2003). The number of universities offering dedicated upper-level taxonomic and systematics courses is dwindling (Mallet and Willmott 2003). Reduced popularity and employability for taxonomic skills have resulted in fewer enthusiasts. Resources for museum collections are frequently cut back (Alberch 1993) and, for an individual, investing time in publishing user-friendly identification keys has limited reward for career progression or recognition. A number of solutions can reduce the burden on specialists, promote information exchange, and contribute to the next generation of skilled specialists, including greater support for online collections,

user-friendly identification initiatives, and more training opportunities (e.g. public workshops, formal classes, placement programmes).

### *Cost-efficient genetic tools*

An alternative to morphological taxonomy is genetic identification. DNA holds much promise in that it can be useful in delimiting species, populations and individuals (Pons et al. 2006). Yet genetic methods also present many challenges: the choice of molecular marker can profoundly affect the taxonomic resolution of a study (Dupuis et al. 2012); the cost of genetic methods and the amount of data generated can be prohibitive (Hunter et al. 2018); and specialists are still required to develop methods and training protocols. Environmental DNA (eDNA) and similar metabarcoding methods can survey large numbers of samples with relatively little investment (Shaw et al. 2017), thereby facilitating rapid, low impact biodiversity surveys with comparable, and sometimes superior, detection rates to traditional survey methods (Hoffmann et al. 2016). Further, more affordable next-generation sequencing (NGS), improved protocols and commercial sample collection kits, and portable third generation sequencers, provide excellent opportunities to engage scientists in widespread, long-term monitoring.

However, such methods are a complementary approach, not an absolute replacement for on-ground field assessments. The efficacy of eDNA is currently limited by the number of ‘known’ (i.e. previously described and sequenced) species available for building comparative genetic databases, and it is also difficult to statistically derive relative abundance information from eDNA samples (Deiner et al. 2017). Thus, it is easier to answer the question of ‘what is present?’ rather than ‘how many are present?’.

Despite limitations, genetic methods continue to offer valuable insight into global biodiversity and its broader interactions. For example, it is often difficult to directly study species- and population-level dynamics of rare, cryptic or under-studied organisms. However, genetic approaches can provide an indirect means of identifying: species (Deiner et al. 2017); behaviour (e.g. mating systems (Janes et al. 2016), trophic interactions (de Vere et al. 2017), migration (Chapuis et al. 2009)); demography (e.g. effective population sizes (Janes et al. 2018)); and subsequent population-level changes (Gloss et al. 2016). Where possible, better integration of genetic approaches with traditional surveys could greatly advance our knowledge of insect biodiversity and biology (e.g. (Cullingham et al. 2018)). Integration of these methods is becoming increasingly time- and cost-effective because: (i) material for DNA-based work can often be collected during planned surveys or opportunistic site visits by non-specialists; (ii) improved specimen preservation methods are increasing the longevity of samples for research (Matos-Maraví et al. 2019); (iii) commercial DNA extraction kits are decreasing the amount of starting material (e.g. a single leg) required to generate sufficient template (Richards and Murali 2015), meaning that a single specimen might be used multiple times; (iv) the yield of DNA is generally increasing while the amount of DNA required for sequencing is typically decreasing, thereby facilitating numerous applications (Matos-Maraví et al. 2019); and (v) sequencing costs continue to decline making genetic/genomic approaches more accessible. Lastly, the increasing need and pressure to catalogue biodiversity and combat biodiversity declines continues to inspire novel and broad collaborations (e.g. academic, government, non-government organizations, field naturalists, citizen scientists); these collaborations are essential to better combine observational and genetic-based methods and to better integrate these methods with policy and decision-making.



### *Insect Behaviour*

The sheer diversity of insects, and the huge variation in life histories and behavioural traits, explains why there is such a variety of trapping methods in the entomologist's toolkit (e.g. observation, pitfall traps, pan traps, malaise traps, aerial traps, netting, light traps, Lindgren traps, beat trays etc.). The most appropriate sampling method will depend almost entirely on the focal taxa and the system and season being sampled. This means there is no single trapping method that can identify population declines for all insect taxa at the global scale. Importantly, it means that any public discussion around studies showing declines of insect taxa need to clearly identify which taxa were targeted in the study's sampling methods.

Within species, insects can exhibit incredible plasticity in behaviour and a range of personality syndromes (Kralj-Fišer and Schuett 2014; Modlmeier et al. 2015). The speed and flexibility with which behavioural responses occur give an insect greater capacity to deal with environmental change, particularly in the face of rapid or catastrophic events. Insects can respond to environmental changes behaviourally; for example changes in temperature can lead insects to modify their dispersal, diapause, thermoregulation, foraging, and reproductive behaviours (Parmesan 2006; Schilthuizen and Kellermann 2014). Furthermore, environmental change can exert differing selective pressures on behavioural traits. Invasive insects can exhibit drastic shifts in foraging behaviour, activity and boldness in their new environments, and some species have shown rapid behavioural changes in response to environmental variation (Singer et al. 1993; Lombaert et al. 2014).

The effect of evolutionary changes in behaviour on insect population dynamics or measures of insect occurrence (e.g. trapping biases) remains to be seen. Currently data on the extent of behavioural plasticity in insects is scarce, and further research is needed to understand whether this capacity, in conjunction with genetic responses, will buffer insects against future environmental changes.

### *Natural history observation*

Natural history observations are valuable information sources to help understand historical insect declines, and identify potential drivers that can be tested as research hypotheses. A great example of this are the annual natural history notes from Inverness-shire, Scotland, published by G. W. Harper in ‘The Entomologist’s Record and journal of variation’. Harper followed Lepidoptera populations in the highlands from 1952 until his death in 1973. In his 1970 record, he notes that “distinct climatic change” and “human destruction of our small local entomological habitats” have driven the decline in Lepidoptera he had witnessed over the previous 14 years (Harper 1971). Declines of some lepidopteran taxa in Great Britain have been confirmed more recently (Fox 2013), and it is uncertain whether these populations will recover.

Comprehensive records such as these are now rare in most countries, mostly because of declining support for traditional natural history among funding bodies and the broader research community (Tewksbury et al. 2014). It is never too late to start recording natural patterns. Observing nature can lead to discovery of new species, expand known ranges of species, build knowledge of a species’ role in its community, and inform novel research hypotheses (Wheeler 2013; Saunders 2016). We must support and promote natural history

observation as an essential component of scientific research, and as a rewarding opportunity to leave a legacy of knowledge for future generations (Box 1).

### *Links between insect communities and ecosystem function*

Insects contribute to ecosystem function and associated ecosystem services in multiple ways throughout their life cycles (Losey and Vaughan 2006; Saunders 2018). Understanding the importance of insects in maintaining ecosystem function and human well-being is an excellent engagement tool to build public support for insect conservation. Yet reductionist approaches to classifying an insect's role in the ecosystem and quantifying relevant ecosystem services have resulted in many complex interactions and functional roles being overlooked (Yang and Gratton 2014; Saunders et al. 2016). In addition, taxonomic biases in community ecology and ecosystem function research mean we still have limited understanding of how insects mediate ecosystem function at different scales in many systems (Basset et al. 2019). Greater investment in cross-taxon (e.g. plant-insect, insect-vertebrate interactions) and cross-system (e.g. land-water, social-ecological) research is essential to fill these knowledge gaps. Network and trait-based analyses are promising approaches for understanding how complex interactions influence ecosystem function across different scales, including trait linkage frameworks (Hevia et al. 2017), species-habitat bipartite networks (Saunders and Rader 2019), and multilayer network frameworks (Bohan and et al. 2016).

### *Human impacts on insect communities*

Human activity impacts insect taxa in many ways. Climate change is already causing phenological shifts in insect life cycles, resulting in potentially damaging effects on trophic interactions that influence ecosystem function, e.g. pollination and herbivory (Hegland et al.

2009; Bell et al. 2015). Human-mediated introduction of invasive alien insects can affect populations of native species in many ways. For example, the biological control agent *Cotesia glomerata* was introduced to the USA to control *Pieris rapae*, subsequently contributing to local declines of a native pierid butterfly in northeast USA (Benson et al. 2003). Effects of these complex interactions on the resilience of insect populations are still uncertain.

Overuse of synthetic chemicals (pesticides, herbicides, fungicides) has lethal and sub-lethal effects on many insect taxa, particularly beneficial species (Desneux et al. 2007). The physiological and biological effects that pesticides have on particular insect taxa is known but, more broadly, we still know very little about how environmental concentrations of these chemicals interfere with biological and ecological processes at the scale of populations, communities and ecosystems (Köhler and Triebkorn 2013; Wood and Goulson 2017). Synergistic interactions between pesticides, herbicides and fungicides can enhance detrimental effects beyond what is observed in isolation, even for individual chemicals that are perceived as apparently ‘harmless’, yet we know very little about the true extent of these interactions in environmental contexts (Desneux et al. 2007).

Habitat removal and landscape modification are also key drivers of change in local insect populations. In particular, the increasing amount of land converted to homogeneous crop production can provide ideal environments for more pest insect outbreaks and have detrimental effects on populations of beneficial insects (Haddad et al. 2011; Nicholls and Altieri 2013). Broad-scale land clearing and agricultural intensification reduce the diversity of habitats and resources available to insects at the landscape scale (Tschardt et al. 2012).

Most insect taxa have complex life cycles and require access to multiple different habitats and resources in different life stages. For example, adult dragonflies (Odonata) are winged predators in terrestrial systems, while larval stages live in water; some hoverfly species (Syrphidae) have predatory larvae that hunt herbivorous insects on plants, while the adult feeds on pollen and nectar. We still have very limited knowledge of how multiple drivers impact insects across the whole of their life cycle, or of how these drivers affect community structure and functional diversity.

### **Communicating science for insect conservation**

Communicating research results is an essential part of the research process and is critical to engage public audiences with insect conservation. This includes publishing results in scholarly literature and disseminating summaries through popular and social media. Using generalised framing or amplified language can be useful to grab audience attention, especially for critical issues like biodiversity declines. However, this should never be at the expense of accuracy. Evidence of geographically-restricted declines should not be framed as global in scale, and evidence of declines for particular species or taxonomic groups is not evidence that all insects are disappearing. Such misinformation can be counter-productive and affect public trust in scientific knowledge.

Hyping science is not a new phenomenon (Rinaldi 2012). However, in recent decades, it has arguably become more common, and potentially more damaging, due to enhanced competition among scientists and the rapid increase in an individual's potential global reach via online media (Bubela 2006; Caulfield and Condit 2012). The responsibility to support accuracy in science communication lies with all stakeholders, including scientists,

institutions, communicators, journalists and public audiences. We need public and political support for the informative and transformative power of scientific evidence, but to achieve this, we need to ensure that any text published under the auspices of scientific evidence is rigorous and accurate.

Achieving accuracy in ecological science communication also means discussing uncertainty. The role of science is not to produce absolute truths, but to document and understand variation in patterns and processes. Yet the deficit model of science communication (i.e. the belief that the public are ignorant of scientific truths and communication should focus on fact-based enlightenment) is common among scientists and science reporters (Frewer et al. 2003; Nisbet and Scheufele 2009). In reality, effective science communication should be a mutual relationship that focuses on engagement and dialogue, and discusses uncertainty relative to the needs and beliefs of the audience (Rabinovich and Morton 2012; Fischhoff and Davis 2014). A critical point often missing in popular science communication is that uncertainty in scientific knowledge is relative to the breadth of literature on a specific topic. In the case of insect declines, there are not enough independent studies to reach the level of consensus needed to prove global declines of all insect species. However, we also stress that, while discussing uncertainty is essential to science communication, uncertainty is not a valid argument for doing nothing to reduce a potential harm (Oreskes 2004).

We may never know if global declines are truly happening, and we don't need to wait for proof. Uncertainty can frame a more constructive and hopeful message. To gain global support for insect conservation, we need people to believe the truth: we have not yet lost all insects, and there is still time to protect global insect diversity if we act now. Building

knowledge of the mechanisms that may drive future insect declines, and how those declines would impact ecosystems and human wellbeing, presents valuable opportunities for conservation investment, engagement and action.

### **Author contributions**

MES conceived the idea and led the study; all authors contributed equally to content and manuscript development.

### **Author Biographical**

Manu Saunders, Jasmine Janes and James O’Hanlon are affiliated with the University of New England in Armidale, Australia; Jasmine Janes is also at Vancouver Island University, Canada.

## References

- Alberch P. 1993. Museums, collections and biodiversity inventories. *Trends in Ecology & Evolution*. 8(10):372–375. doi:10.1016/0169-5347(93)90222-B.
- Ball-Damerow JE, M’Gonigle LK, Resh VH. 2014. Changes in occurrence, richness, and biological traits of dragonflies and damselflies (Odonata) in California and Nevada over the past century. *Biodiversity Conservation*. 23(8):2107–2126. doi:10.1007/s10531-014-0707-5.
- Basset Y, Miller SE, Gripenberg S, Ctvrticka R, Dahl C, Leather SR, Didham RK. 2019. An entomocentric view of the Janzen–Connell hypothesis. *Insect Conservation & Diversity*. 12(1):1–8. doi:10.1111/icad.12337.
- Bell JR, Alderson L, Izera D, Kruger T, Parker S, Pickup J, Shortall CR, Taylor MS, Verrier P, Harrington R. 2015. Long-term phenological trends, species accumulation rates, aphid traits and climate: five decades of change in migrating aphids. *Journal of Animal Ecology*. 84(1):21–34. doi:10.1111/1365-2656.12282.
- Benson J, Van Driesche RG, Pasquale A, Elkinton J. 2003. Introduced braconid parasitoids and range reduction of a native butterfly in New England. *Biological Control*. 28(2):197–213. doi:10.1016/S1049-9644(03)00058-6.
- Biesmeijer JC, Roberts SPM, Reemer M, Ohlemüller R, Edwards M, Peeters T, Schaffers AP, Potts SG, Kleukers R, Thomas CD, et al. 2006. Parallel declines in pollinators and insect-pollinated plants in Britain and the Netherlands. *Science*. 313(5785):351–354. doi:10.1126/science.1127863.
- Blaxter M. 2003. Molecular systematics: Counting angels with DNA. *Nature*. 421(6919):122–124. doi:10.1038/421122a.
- Bohan D, et al. 2016. Networking our way to better ecosystem service provision. *Trends in Ecology & Evolution*. 31(2):105–115.
- Brooks DR, Bater JE, Clark SJ, Monteith DT, Andrews C, Corbett SJ, Beaumont DA, Chapman JW. 2012. Large carabid beetle declines in a United Kingdom monitoring network increases evidence for a widespread loss in insect biodiversity. *Journal of Applied Ecology*. 49(5):1009–1019. doi:10.1111/j.1365-2664.2012.02194.x.
- Bubela T. 2006. Science communication in transition: genomics hype, public engagement, education and commercialization pressures. *Clinical Genetics*. 70(5):445–450. doi:10.1111/j.1399-0004.2006.00693.x.
- Buma B. 2015. Disturbance interactions: characterization, prediction, and the potential for cascading effects. *Ecosphere*. 6(4):art70. doi:10.1890/ES15-00058.1.
- Büntgen U, Frank D, Liebhold A, Johnson D, Carrer M, Urbinati C, Grabner M, Nicolussi K, Levanić T, Esper J. 2009. Three centuries of insect outbreaks across the European Alps. *New Phytologist*. 182(4):929–941. doi:10.1111/j.1469-8137.2009.02825.x.



- Cardoso P, Erwin TL, Borges PAV, New TR. 2011. The seven impediments in invertebrate conservation and how to overcome them. *Biological Conservation*. 144(11):2647–2655. doi:10.1016/j.biocon.2011.07.024.
- Cardoso P, Leather SR. 2019. Predicting a global insect apocalypse. *Insect Conservation & Diversity*. 12(4):263–267. doi:10.1111/icad.12367.
- Carrington D. 2017 Oct 18. Warning of ‘ecological Armageddon’ after dramatic plunge in insect numbers. *The Guardian*. [accessed 2019 Apr 25]. <https://www.theguardian.com/environment/2017/oct/18/warning-of-ecological-armageddon-after-dramatic-plunge-in-insect-numbers>.
- Caulfield T, Condit C. 2012. Science and the Sources of Hype. *Public Health Genomics*. 15(3–4):209–217. doi:10.1159/000336533.
- Chapuis M-P, Loiseau A, Michalakis Y, Lecoq M, Franc A, Estoup A. 2009. Outbreaks, gene flow and effective population size in the migratory locust, *Locusta migratoria*: a regional-scale comparative survey. *Molecular Ecology*. 18(5):792–800. doi:10.1111/j.1365-294X.2008.04072.x.
- Chapuisat M, Sundström L, Keller L. 1997. Sex–ratio regulation: the economics of fratricide in ants. *Proceedings of the Royal Society of London B Biological Sciences*. 264(1385):1255–1260. doi:10.1098/rspb.1997.0173.
- Charnov EL, Los-den Hartogh RL, Jones WT, van den Assem J. 1981. Sex ratio evolution in a variable environment. *Nature*. 289(5793):27–33. doi:10.1038/289027a0.
- Colla SR, Packer L. 2008. Evidence for decline in eastern North American bumblebees (Hymenoptera: Apidae), with special focus on *Bombus affinis* Cresson. *Biodiversity & Conservation*. 17(6):1379. doi:10.1007/s10531-008-9340-5.
- Coyle DR, Nebeker TE, Hart ER, Mattson WJ. 2005. Biology and Management of Insect Pests in North American Intensively Managed Hardwood Forest Systems. *Annual Reviews of Entomology*. 50(1):1–29. doi:10.1146/annurev.ento.50.071803.130431.
- Cullingham CI, Janes JK, Hamelin RC, James PMA, Murray BW, Sperling FAH. 2018. The contribution of genetics and genomics to understanding the ecology of the mountain pine beetle system. *Canadian Journal of Forest Research*. doi:10.1139/cjfr-2018-0303.
- Dahlstrom MF, Ho SS. 2012. Ethical Considerations of Using Narrative to Communicate Science. *Science Communication*. 34(5):592–617. doi:10.1177/1075547012454597.
- Dalin P, Kindvall O, Björkman C. 2009. Reduced Population Control of an Insect Pest in Managed Willow Monocultures. *PLOS ONE*. 4(5):e5487. doi:10.1371/journal.pone.0005487.
- Deiner K, Bik HM, Mächler E, Seymour M, Lacoursière-Roussel A, Altermatt F, Creer S, Bista I, Lodge DM, Vere N de, et al. 2017. Environmental DNA metabarcoding: Transforming how we survey animal and plant communities. *Molecular Ecology*. 26(21):5872–5895. doi:10.1111/mec.14350.

- Desender K, Turin H. 1989. Loss of habitats and changes in the composition of the ground and tiger beetle fauna in four West European countries since 1950 (Coleoptera: Carabidae, cicindelidae). *Biological Conservation*. 48(4):277–294. doi:10.1016/0006-3207(89)90103-1.
- Desneux N, Decourtye A, Delpuech J-M. 2007. The Sublethal Effects of Pesticides on Beneficial Arthropods. *Annual Reviews of Entomology*. 52(1):81–106. doi:10.1146/annurev.ento.52.110405.091440.
- Dupuis JR, Roe AD, Sperling F a. H. 2012. Multi-locus species delimitation in closely related animals and fungi: one marker is not enough. *Molecular Ecology*. 21(18):4422–4436. doi:10.1111/j.1365-294X.2012.05642.x.
- Elton CS. 1924. Periodic Fluctuations in the Numbers of Animals: Their Causes and Effects. *Journal of Experimental Biology*. 2(1):119–163.
- Esper J, Büntgen U, Frank DC, Nievergelt D, Liebhold A. 2007. 1200 years of regular outbreaks in alpine insects. *Proceedings of the Royal Society B Biological Sciences*. 274(1610):671–679. doi:10.1098/rspb.2006.0191.
- Figuerola LL, Berguey EA. 2015. Bumble Bees (Hymenoptera: Apidae) of Oklahoma: Past and Present Biodiversity. *Journal of the Kansas Entomological Society*. 88(4):418–429. doi:10.2317/0022-8567-88.4.418.
- Fischhoff B, Davis AL. 2014. Communicating scientific uncertainty. *Proceedings of the National Academy of Sciences*. 111(Supplement 4):13664–13671. doi:10.1073/pnas.1317504111.
- Fox R. 2013. The decline of moths in Great Britain: a review of possible causes. *Insect Conservatio & Diversity*. 6(1):5–19. doi:10.1111/j.1752-4598.2012.00186.x.
- Frewer L, Hunt S, Brennan M, Kuznesof S, Ness M, Ritson C. 2003. The views of scientific experts on how the public conceptualize uncertainty. *Journal of Risk Research*. 6(1):75–85. doi:10.1080/1366987032000047815.
- Gardner JD, Spivak M. 2014. A Survey and Historical Comparison of the Megachilidae (Hymenoptera: Apoidea) of Itasca State Park, Minnesota. *Annals of the Entomological Society of America*. 107(5):983–993. doi:10.1603/AN14023.
- Gaston KJ, Lawton JH. 1988. Patterns in the distribution and abundance of insect populations. *Nature*. 331(6158):709–712. doi:10.1038/331709a0.
- Gibb H, Grossman BF, Dickman CR, Decker O, Wardle GM. 2019. Long-term responses of desert ant assemblages to climate. *Journal of Animal Ecology*. doi:10.1111/1365-2656.13052.
- Gillespie MAK, Alfredsson M, Barrio IC, Bowden JJ, Convey P, Culler LE, Coulson SJ, Krogh PH, Koltz AM, Koponen S, et al. 2019 Mar 16. Status and trends of terrestrial arthropod abundance and diversity in the North Atlantic region of the Arctic. *Ambio*. doi:10.1007/s13280-019-01162-5.
- Gloss AD, Groen SC, Whiteman NK. 2016. A Genomic Perspective on the Generation and Maintenance of Genetic Diversity in Herbivorous Insects. *Annual Reviews of Ecology Evolution & Systematics*. 47(1):165–187. doi:10.1146/annurev-ecolsys-121415-032220.

- Godfray HCJ. 2002. Challenges for taxonomy. *Nature*. 417:17–19. doi:10.1038/417017a.
- Godfray HCJ, Werren JH. 1996. Recent developments in sex ratio studies. *Trends in Ecology & Evolution*. 11(2):59–63. doi:10.1016/0169-5347(96)81043-3.
- Haddad NM, Crutsinger GM, Gross K, Haarstad J, Tilman D. 2011. Plant diversity and the stability of foodwebs. *Ecology Letters*. 14(1):42–46. doi:10.1111/j.1461-0248.2010.01548.x.
- Hallmann CA, Sorg M, Jongejans E, Siepel H, Hofland N, Schwan H, Stenmans W, Müller A, Sumser H, Hörrén T, et al. 2017. More than 75 percent decline over 27 years in total flying insect biomass in protected areas. *Plos One*. 12(10):e0185809. doi:10.1371/journal.pone.0185809.
- Harper GW. 1971. Inverness-shire in 1970. *Entomological Record*. 83:94–97.
- Hegland SJ, Nielsen A, Lázaro A, Bjerknes A-L, Totland Ø. 2009. How does climate warming affect plant-pollinator interactions? *Ecology Letters*. 12(2):184–195. doi:10.1111/j.1461-0248.2008.01269.x.
- Herrera CM. 2018. Complex long-term dynamics of pollinator abundance in undisturbed Mediterranean montane habitats over two decades. *Ecological Monographs*. 89(1): e01338. doi:10.1002/ecm.1338.
- Hevia V, Martín-López B, Palomo S, García-Llorente M, Bello F de, González JA. 2017. Trait-based approaches to analyze links between the drivers of change and ecosystem services: Synthesizing existing evidence and future challenges. *Ecology & Evolution*. 7(3):831–844. doi:10.1002/ece3.2692.
- Hoffmann C, Schubert G, Calvignac-Spencer S. 2016. Aquatic biodiversity assessment for the lazy. *Molecular Ecology*. 25(4):846–848. doi:10.1111/mec.13535.
- Horeis H. 2009. Once upon a time, there was a dying forest. *Technology in Society*. 31(1):111–116. doi:10.1016/j.techsoc.2008.10.012.
- Hunter ME, Hoban SM, Bruford MW, Segelbacher G, Bernatchez L. 2018. Next-generation conservation genetics and biodiversity monitoring. *Evolutionary Applications*. 11(7):1029–1034. doi:10.1111/eva.12661.
- Jacobson MM, Tucker EM, Mathiasson ME, Rehan SM. 2018. Decline of bumble bees in northeastern North America, with special focus on *Bombus terricola*. *Biological Conservation*. 217:437–445. doi:10.1016/j.biocon.2017.11.026.
- Janes JK, Roe AD, Rice AV, Gorrell JC, Coltman DW, Langor DW, Sperling F a. H. 2016. Polygamy and an absence of fine-scale structure in *Dendroctonus ponderosae* (Hopk.) (Coleoptera: Curculionidae) confirmed using molecular markers. *Heredity*. 116(1):68–74. doi:10.1038/hdy.2015.71.
- Janes JK, Worth JRP, Batista PD, Sperling FAH. 2018. Inferring Ancestry and Divergence Events in a Forest Pest Using Low-Density Single-Nucleotide Polymorphisms. *Insect Systematics and Diversity*. 2(6). doi:10.1093/isd/ixy019.

- Jarvis B. 2018 Nov 27. The Insect Apocalypse Is Here. N Y Times. [accessed 2019 Sep 30]. <https://www.nytimes.com/2018/11/27/magazine/insect-apocalypse.html>.
- Knops JMH, Tilman D, Haddad NM, Naeem S, Mitchell CE, Haarstad J, Ritchie ME, Howe KM, Reich PB, Siemann E, et al. 1999. Effects of plant species richness on invasion dynamics, disease outbreaks, insect abundances and diversity. *Ecology Letters*. 2(5):286–293. doi:10.1046/j.1461-0248.1999.00083.x.
- Köhler H-R, Triebkorn R. 2013. Wildlife Ecotoxicology of Pesticides: Can We Track Effects to the Population Level and Beyond? *Science*. 341(6147):759–765. doi:10.1126/science.1237591.
- Kralj-Fišer S, Schuett W. 2014. Studying personality variation in invertebrates: why bother? *Animal Behaviour*. 91:41–52. doi:10.1016/j.anbehav.2014.02.016.
- Leijs R, Dorey J, Hogendoorn K. 2018. Twenty six new species of *Leioproctus* (Colletellus): Australian Neopasiphaeinae, all but one with two submarginal cells (Hymenoptera, Colletidae, Leioproctus). *ZooKeys* (811):109–168. doi:10.3897/zookeys.811.28924.
- Lipscomb D, Platnick N, Wheeler Q. 2003. The intellectual content of taxonomy: a comment on DNA taxonomy. *Trends in Ecology & Evolution*. 18(2):65–66. doi:10.1016/S0169-5347(02)00060-5.
- Lister BC, Garcia A. 2018. Climate-driven declines in arthropod abundance restructure a rainforest food web. *Proceedings of the National Academy of Sciences U S A*. 115(44):E10397–E10406. doi:10.1073/pnas.1722477115.
- Lombaert E, Estoup A, Facon B, Joubard B, Grégoire J-C, Jannin A, Blin A, Guillemaud T. 2014. Rapid increase in dispersal during range expansion in the invasive ladybird *Harmonia axyridis*. *Journal of Evolutionary Biology*. 27(3):508–517. doi:10.1111/jeb.12316.
- Losey J, Vaughan M. 2006. The Economic Value of Ecological Services Provided by Insects. *Bioscience*. 56(4):311. doi:10.1641/0006-3568(2006)56[311:TEVOES]2.0.CO;2.
- Lovett GM, Canham CD, Arthur MA, Weathers KC, Fitzhugh RD. 2006. Forest Ecosystem Responses to Exotic Pests and Pathogens in Eastern North America. *BioScience*. 56(5):395–405. doi:10.1641/0006-3568(2006)056[0395:FERTEP]2.0.CO;2.
- Mallet J, Willmott K. 2003. Taxonomy: renaissance or Tower of Babel? *Trends in Ecology & Evolution*. 18(2):57–59. doi:10.1016/S0169-5347(02)00061-7.
- Martialay M. 2018. Two Degrees Decimated Puerto Rico’s Insect Populations. *Rensselaer Polytech*. [accessed 2019 Sep 30]. /content/2018/10/15/two-degrees-decimated-puerto-rico%E2%80%99s-insect-populations.
- Matos-Maraví P, Ritter CD, Barnes CJ, Nielsen M, Olsson U, Wahlberg N, Marquina D, Sääksjärvi I, Antonelli A. 2019. Biodiversity seen through the perspective of insects: 10 simple rules on methodological choices and experimental design for genomic studies. *PeerJ*. 7:e6727. doi:10.7717/peerj.6727.

- McAfee D, Doubleday ZA, Geiger N, Connell SD. 2019. Everyone Loves a Success Story: Optimism Inspires Conservation Engagement. *BioScience*. 69(4):274–281. doi:10.1093/biosci/biz019.
- McArdle B, Gaston K. 1993. The Temporal Variability of Populations. *Oikos*. 67(1):187–191. doi:10.2307/3545110.
- McArdle BH. 1990. When Are Rare Species Not There? *Oikos*. 57(2):276–277. doi:10.2307/3565950.
- McGrath M. 2019 Feb 11. Insect decline may see ‘plague of pests’. BBC News. [accessed 2019 Sep 30]. <https://www.bbc.com/news/science-environment-47198576>.
- Meiners JM, Griswold TL, Carril OM. 2019. Decades of native bee biodiversity surveys at Pinnacles National Park highlight the importance of monitoring natural areas over time. *PLOS ONE*. 14(1):e0207566. doi:10.1371/journal.pone.0207566.
- Modlmeier AP, Keiser CN, Wright CM, Lichtenstein JL, Pruitt JN. 2015. Integrating animal personality into insect population and community ecology. *Current Opinion in Insect Science*. 9:77–85. doi:10.1016/j.cois.2015.03.008.
- Myers JH. 1993. Population Outbreaks in Forest Lepidoptera. *American Scientist*. 81(3):240–251.
- Nicholls CI, Altieri MA. 2013. Plant biodiversity enhances bees and other insect pollinators in agroecosystems. A review. *Agronomy for Sustainable Development*. 33(2):257–274. doi:10.1007/s13593-012-0092-y.
- Nisbet MC, Scheufele DA. 2009. What’s next for science communication? Promising directions and lingering distractions. *American Journal of Botany*. 96(10):1767–1778. doi:10.3732/ajb.0900041.
- Oreskes N. 2004. Science and public policy: what’s proof got to do with it? *Environmental Science & Policy*. 7(5):369–383. doi:10.1016/j.envsci.2004.06.002.
- Ostfeld RS, Keesing F. 2000. Pulsed resources and community dynamics of consumers in terrestrial ecosystems. *Trends in Ecology & Evolution*. 15(6):232–237. doi:10.1016/S0169-5347(00)01862-0.
- Parmesan C. 2006. Ecological and Evolutionary Responses to Recent Climate Change. *Annual Review of Ecology, Evolution & Systematics*. 37(1):637–669. doi:10.1146/annurev.ecolsys.37.091305.110100.
- Pascal M, Beaudeau P, Medina S, Hamilton NC. 2019. Global Change: a Public Health Researcher’s Ethical Responsibility. *Current Environmental Health Reports*. 6(3):160–166. doi:10.1007/s40572-019-00238-4.
- Paukkunen J, Pöyry J, Kuussaari M. 2018. Species traits explain long-term population trends of Finnish cuckoo wasps (Hymenoptera: Chrysididae). *Insect Conservation & Diversity*. 11(1):58–71. doi:10.1111/icad.12241.

Petanidou T, Vujić A, Ellis WN. 2011. Hoverfly diversity (Diptera: Syrphidae) in a Mediterranean scrub community near Athens, Greece. *Annales Société Entomologique France NS*. 47(1–2):168–175. doi:10.1080/00379271.2011.10697709.

Pons J, Barraclough TG, Gomez-Zurita J, Cardoso A, Duran DP, Hazell S, Kamoun S, Sumlin WD, Vogler AP. 2006. Sequence-Based Species Delimitation for the DNA Taxonomy of Undescribed Insects. *Systematic Biology*. 55(4):595–609. doi:10.1080/10635150600852011.

Porter JH, Parry ML, Carter TR. 1991. The potential effects of climatic change on agricultural insect pests. *Agricultural and Forest Meteorology*. 57(1):221–240. doi:10.1016/0168-1923(91)90088-8.

Rabinovich A, Morton TA. 2012. Unquestioned Answers or Unanswered Questions: Beliefs About Science Guide Responses to Uncertainty in Climate Change Risk Communication. *Risk Analysis*. 32(6):992–1002. doi:10.1111/j.1539-6924.2012.01771.x.

Radboud University. 2017. Three-quarters of the total insect population lost in protected nature reserves. Radboud Univ. [accessed 2019 Sep 30]. <https://www.ru.nl/english/news-agenda/news/vm/iwwr/2017/three-quarters-total-insect-population-lost/>.

Richards S, Murali SC. 2015. Best practices in insect genome sequencing: what works and what doesn't. *Current Opinion in Insect Science*. 7:1–7. doi:10.1016/j.cois.2015.02.013.

Rinaldi A. 2012. To hype, or not to(o) hype: Communication of science is often tarnished by sensationalization, for which both scientists and the media are responsible. *EMBO Reports*. 13(4):303–307. doi:10.1038/embor.2012.39.

Samways MJ. 2007. Insect Conservation: A Synthetic Management Approach. *Annual Review of Entomology*. 52(1):465–487. doi:10.1146/annurev.ento.52.110405.091317.

Sánchez-Bayo F, Wyckhuys KAG. 2019. Worldwide decline of the entomofauna: A review of its drivers. *Biological Conservation*. 232:8–27. doi:10.1016/j.biocon.2019.01.020.

Saunders ME. 2016. Species that are 'known unknowns' to science. Ecology is not a Dirty Word. [accessed 2019 Jan 11]. <https://ecologyisnotadirtyword.com/2016/08/30/species-that-are-known-unknowns-to-science/>.

Saunders ME. 2019. Insectageddon is a great story. But what are the facts? Ecology is not a Dirty Word. [accessed 2019 Apr 6]. <https://ecologyisnotadirtyword.com/2019/02/16/insectageddon-is-a-great-story-but-what-are-the-facts/>.

Saunders ME. 2018. Ecosystem services in agriculture: understanding the multifunctional role of invertebrates. *Agricultural and Forest Entomology*. 20(2):298–300. doi:10.1111/afe.12248.

Saunders ME, Peisley RK, Rader R, Luck GW. 2016. Pollinators, pests, and predators: Recognizing ecological trade-offs in agroecosystems. *Ambio*. 45(1):4–14. doi:10.1007/s13280-015-0696-y.

Saunders ME, Rader R. 2019. Network modularity influences plant reproduction in a mosaic tropical agroecosystem. *Proceedings of the Royal Society B Biological Sciences*. 286(1899):20190296. doi:10.1098/rspb.2019.0296.

Schilthuizen M, Kellermann V. 2014. Contemporary climate change and terrestrial invertebrates: evolutionary versus plastic changes. *Evolutionary Applications*. 7(1):56–67. doi:10.1111/eva.12116.

Schönrogge K, Walker P, Crawley MJ. 2000. Parasitoid and inquiline attack in the galls of four alien, cynipid gall wasps: host switches and the effect on parasitoid sex ratios. *Ecological Entomology*. 25(2):208–219. doi:10.1046/j.1365-2311.2000.00244.x.

Schuch S, Wesche K, Schaefer M. 2012. Long-term decline in the abundance of leafhoppers and planthoppers (Auchenorrhyncha) in Central European protected dry grasslands. *Biological Conservation*. 149(1):75–83. doi:10.1016/j.biocon.2012.02.006.

Shaw JLA, Weyrich L, Cooper A. 2017. Using environmental (e)DNA sequencing for aquatic biodiversity surveys: a beginner's guide. *Marine and Freshwater Research*. 68(1):20–33. doi:10.1071/MF15361.

Simmons BI, Balmford A, Bladon AJ, Christie AP, Palma AD, Dicks LV, Gallego-Zamorano J, Johnston A, Martin PA, Purvis A, et al. 2019. Worldwide insect declines: An important message, but interpret with caution. *Ecology & Evolution*. doi:10.1002/ece3.5153.

Singer MC, Thomas CD, Parmesan C. 1993. Rapid human-induced evolution of insect–host associations. *Nature*. 366(6456):681–683. doi:10.1038/366681a0.

Solomon ME. 1949. The Natural Control of Animal Populations. *Journal of Animal Ecology*. 18(1):1–35. doi:10.2307/1578.

Stork NE. 1988. Insect diversity: facts, fiction and speculation. *Biological Journal of the Linnean Society*. 35(4):321–337. doi:10.1111/j.1095-8312.1988.tb00474.x.

Stork NE. 2018. How Many Species of Insects and Other Terrestrial Arthropods Are There on Earth? *Annual Review of Entomology*. 63(1):31–45. doi:10.1146/annurev-ento-020117-043348.

Tautz D, Arctander P, Minelli A, Thomas RH, Vogler AP. 2003. A plea for DNA taxonomy. *Trends in Ecology & Evolution*. 18(2):70–74. doi:10.1016/S0169-5347(02)00041-1.

Tepedino VJ, Durham S, Cameron SA, Goodell K. 2015. Documenting bee decline or squandering scarce resources. *Conserv Biol*. 29(1):280–282. doi:10.1111/cobi.12439.

Tewksbury JJ, Anderson JGT, Bakker JD, Billo TJ, Dunwiddie PW, Groom MJ, Hampton SE, Herman SG, Levey DJ, Machnicki NJ, et al. 2014. Natural History's Place in Science and Society. *BioScience*. 64(4):300–310. doi:10.1093/biosci/biu032.

The University of Sydney. 2019. Insect population faces 'catastrophic' collapse: Sydney research. *Univ Syd*. [accessed 2019 Oct 2]. <https://sydney.edu.au/news-opinion/news/2019/02/12/insect-population-faces--catastrophic--collapse--sydney-research.html>.

- Thomas CD, Jones TH, Hartley SE. 2019. “Insectageddon”: a call for more robust data and rigorous analyses. *Global Change Biology*. 25(6): 1891-1892.
- Tscharntke T, Tylianakis JM, Rand TA, Didham RK, Fahrig L, Batáry P, Bengtsson J, Clough Y, Crist TO, Dormann CF, et al. 2012. Landscape moderation of biodiversity patterns and processes - eight hypotheses. *Biological Reviews*. 87(3):661–685. doi:10.1111/j.1469-185X.2011.00216.x.
- de Vere N, Jones LE, Gilmore T, Moscrop J, Lowe A, Smith D, Hegarty MJ, Creer S, Ford CR. 2017. Using DNA metabarcoding to investigate honey bee foraging reveals limited flower use despite high floral availability. *Scientific Reports*. 7:42838. doi:10.1038/srep42838.
- Vilcinskas A. 2019. Pathogens associated with invasive or introduced insects threaten the health and diversity of native species. *Current Opinion in Insect Science*. 33:43–48. doi:10.1016/j.cois.2019.03.004.
- Wagner DL. 2019. Global insect decline: Comments on Sánchez-Bayo and Wyckhuys (2019). *Biological Conservation*. 233:332–333. doi:10.1016/j.biocon.2019.03.005.
- Wallner WE. 1987. Factors Affecting Insect Population Dynamics: Differences Between Outbreak and Non-Outbreak Species. *Annual Review of Entomology*. 32(1):317–340. doi:10.1146/annurev.en.32.010187.001533.
- Warren MS, Hill JK, Thomas JA, Asher J, Fox R, Huntley B, Roy DB, Telfer MG, Jeffcoate S, Harding P, et al. 2001. Rapid responses of British butterflies to opposing forces of climate and habitat change. *Nature*. 414(6859):65. doi:10.1038/35102054.
- Wheeler T. 2013. Why we do science: the paradox of natural history. *Lyman Entomol Mus*. [accessed 2019 Jan 11]. <https://lymanmuseum.wordpress.com/2013/11/03/why-we-do-science-the-paradox-of-natural-history/>.
- Williams PH. 1982. The Distribution and Decline of British Bumble Bees (*Bombus* Latr.). *Journal of Apicultural Research*. 21(4):236–245. doi:10.1080/00218839.1982.11100549.
- Willig MR, Woolbright L, Presley SJ, Schowalter TD, Waide RB, Scalley TH, Zimmerman JK, González G, Lugo AE. 2019. Populations are not declining and food webs are not collapsing at the Luquillo Experimental Forest. *Proceedings of the National Academy of Science*. 116(25):12143–12144. doi:10.1073/pnas.1820456116.
- Wood TJ, Goulson D. 2017. The environmental risks of neonicotinoid pesticides: a review of the evidence post 2013. *Environmental Science and Pollution Research*. 24(21):17285–17325. doi:10.1007/s11356-017-9240-x.
- Yang LH, Gratton C. 2014. Insects as drivers of ecosystem processes. *Current Opinion in Insect Science*. 2:26–32. doi:10.1016/j.cois.2014.06.004.