

1 **Into the wild: microbiome transplant** 2 **studies need broader ecological reality**

3 4 **Authors**

5 Christopher J. Greyson-Gaito^{*1,†}, Timothy J. Bartley^{1,2,†}, Karl Cottenie^{1,†}, Will M.C.
6 Jarvis^{3,†}, Amy E.M. Newman^{1,†}, Mason R. Stothart^{4,†}

7
8 *Corresponding Author - christopher@greyson-gaito.com

9 10 **Affiliations**

- 11 1. University of Guelph, Department of Integrative Biology, Guelph, ON, Canada
- 12 2. University of Toronto Mississauga, Mississauga, ON, Canada
- 13 3. University of Ottawa, Department of Biology, Ottawa, ON, Canada
- 14 4. University of Calgary, Calgary, Department of Ecosystem and Public Health,
15 Calgary, AB, Canada

16
17 † All authors contributed equally

18 19 **ORCID**

20 CJGG - 0000-0001-8716-0290

21 TJB - 0000-0002-5898-0588

22 KC - 0000-0001-9498-8483

23 WMCJ - 0000-0001-8525-6707

24 AEMN - 0000-0003-1005-8380

25 MRS - 0000-0002-2863-908X

26
27 **Note: This version is a preprint and has not been accepted by a peer review.**

28 **The peer-reviewed publication is:**

29 Christopher J. Greyson-Gaito, Timothy J. Bartley, Karl Cottenie, Will M. C. Jarvis, Amy E. M.
30 Newman, and Mason R. Stothart Into the wild: microbiome transplant studies need broader
31 ecological reality. **287**. Proc. R. Soc. B. <https://doi.org/10.1098/rspb.2019.2834>.

32 **Keywords**

33 conservation, ecological adaptation, ecophysiology, holobiont, metacommunity,
34 ecosystem on a leash

36 **Abstract**

37

38 Gut microbial communities (microbiomes) profoundly shape the ecology and
39 evolution of multicellular life. Interactions between host and microbiome appear to be
40 reciprocal, and ecological theory is now being applied to better understand how hosts
41 and their microbiome influence each other. However, some ecological processes that
42 underlie reciprocal host-microbiome interactions may be obscured by the current
43 convention of highly-controlled transplantation experiments. Although these approaches
44 have yielded invaluable insights, there is a need for a broader array of approaches to
45 fully understand host-microbiome reciprocity. Using a directed review, we surveyed the
46 breadth of ecological reality in the current literature on gut microbiome transplants with
47 non-human recipients. For 55 studies, we categorized 9 key experimental conditions
48 that impact the ecological reality (EcoReality) of the transplant, including host taxon
49 match and donor environment. Using these categories, we rated the EcoReality of each
50 transplant. Encouragingly, the breadth of EcoReality has increased over time, but some
51 components of EcoReality are still relatively unexplored, including recipient host
52 environment and microbiome state. The conceptual framework we develop here maps
53 the landscape of possible EcoReality to highlight where fundamental ecological
54 processes can be considered in future transplant experiments.

55

56 **A Quest for Ecological Reality**

57

58

We shall not cease from exploration

59

And the end of all our exploring

60

Will be to arrive where we started

61

And know the place for the first time.

62

T.S. Eliot - Little Gidding (1)

63

64 Far from passive passengers, resident microbial communities (microbiomes) are
65 integral to the basic biological functioning of multicellular life. This revelation, ushered in
66 by advances in sequencing and computing technology, is grounded in a growing
67 understanding that microbiomes profoundly shape their host's biology, influencing
68 factors such as immunity (2), adiposity (3), thermogenesis (4), hormonal regulation (5),
69 physiological development (6), memory (7), and behaviour (8). To date, highly-
70 controlled experiments with laboratory rodent microbiomes have provided foundational
71 and indispensable knowledge on host-microbiome interactions. Furthermore, these
72 initial experiments have set the stage for integrative contributions by comparative
73 animal physiologists, ecologists, and evolutionary biologists to fill knowledge gaps in our
74 understanding of host-microbiome evolution and the interactions which underlay these
75 partnerships (9).

76

77 Recently, researchers have started to appreciate the intertwining nature of host-
78 microbiome interactions. Evidence is mounting that hosts can shape the composition of
79 their microbiome community (10), and that microbiomes can influence their host's
80 behaviour (8) and physiology (5). Based on differing cases of how host and microbiome
81 might interact, Foster et al. (11) proposed four distinct models: 1) 'host control', in which
82 the host unilaterally governs the composition of its microbiome; 2) 'symbiont control', in
83 which the microbiome shapes the host phenotype; 3) 'open ecosystem', in which the
84 host and microbiome do not interact; and 4) 'ecosystem on a leash', in which the host
85 influences the microbiome by selecting upon microbial function rather than for specific
86 microbial taxa. These connections can be so intimate that some researchers (12,13)
87 proposed that a host and its associated microorganisms are a single biological entity—
88 termed the 'holobiont'—on which selection acts, challenging notions of organismal
89 individuality. Using this holobiont perspective, Alberdi et al. (14) posited that the
90 microbial component of the holobiont, with its greater mutability compared to the host
91 genome, may be an important mechanism facilitating host adaptation to rapid
92 environmental change. Therefore, understanding the interplay between the host and the
93 microbiome is crucial for addressing both fundamental and applied questions about the
94 microbiome.

95

96 Host-microbiome interactions are shaped by ecological and evolutionary
97 processes (15,16). Because host-microbiome interactions are potentially reciprocal,
98 these processes act on three levels: the assembly and dynamics of the microbiome, the
99 influence of the host on the microbiome, and the influence of the microbiome on the
100 host. Microbiome assembly is governed by a variety of factors including environmental
101 filtering, priority effects, random sampling, and dispersal limitation (16,17). The within-
102 microbiome community dynamics are influenced by new invasions, competition,
103 mutualisms, and other interactions (15). A host's actions can also shape their
104 associated microbiomes. For example, the host's social behaviour can impact microbial
105 dispersal (18). Conversely, the dynamics of the microbiome can impact the host; the
106 change in microbiome community composition leading to *Clostridium difficile*
107 colonization and pathogenicity is a classic example in humans (19). Evolutionary
108 processes also occur in tandem with all the ecological processes mentioned previously
109 because of the short timescales associated with microbial turnover relative to microbial
110 evolutionary rates (17). Consequently, considering the ecological processes that
111 underlie host-microbiome interactions is critical for making sense of the reciprocity
112 between the host and its microbiome.

113


114 The most convincing evidence for host-microbiome interactions has been
115 gleaned through microbiome transplantation studies. In these studies, researchers
116 experimentally translocate microbial species or communities from donor hosts or

117 external substrates to recipient hosts. Highly-controlled transplantation studies have
118 been and will continue to be invaluable to experimentally probe the host-microbiome
119 relationship. However, there is a trade-off: highly-controlled experiments isolate
120 mechanisms of interest, but they cannot simultaneously capture the full suite of
121 ecological processes (drift, dispersal, competition, etc.) that influence reciprocal host-
122 microbiome interactions in nature. For example, the use of germ-free recipients may
123 preclude competition between introduced and resident microbes (20), and isolated
124 laboratory conditions may limit the potential for microbial dispersal from influencing the
125 composition of the resulting microbiome (21). How researchers weigh this trade-off
126 depends on the research question of interest. If a researcher's goal is to understand the
127 effect of specific microbes on host physiology, or to develop applications for human
128 health and domestic animal production, controlled experimental conditions are
129 preferred. In contrast, when examining the role and consequences of reciprocal host-
130 microbiome interactions in ecological and evolutionary contexts (e.g., fitness effects,
131 intergenerational microbial transmission, speciation, species persistence, etc.),
132 ecological complexity needs to be considered (22). Therefore, a comparison of highly-
133 controlled transplants and ecologically realistic (which we term EcoReal, see Box 1 for a
134 full definition) transplants that match what the host plus its microbiome would
135 experience in a wild ecosystem is required. The trade-offs of laboratory approaches and
136 the need for comparison to studies that use ecologically realistic conditions have long
137 been recognized by comparative animal physiologists (23), though to date there does
138 not seem to have been a similar recognition in microbiome research. Specifically, the
139 breadth of EcoReality in microbiome transplant studies has not been examined,
140 meaning such an evaluation remains an exciting potential avenue for future work.

141

142 Here, we probe the current EcoReality of microbiome transplantation studies.
143 Our work here is not unlike Hanage's (24) questioning of the reality and applicability of
144 biomedical microbiome studies. By taking advantage of the recent explosion of studies
145 conducting microbiome transplants, we evaluated whether the current microbiome
146 transplant literature limits opportunities for ecological processes to influence study
147 outcomes. We investigated two key questions: 1) how EcoReal are the experimental
148 conditions in the current microbiome transplant literature? and 2) does the literature
149 currently cover the full potential range of EcoReality? Using long-established ecological
150 concepts, we categorized microbiome transplantations into different experimental
151 conditions which can impact the EcoReality of the transplant (Figure 1 and Box 2).
152 Using this framework, we scored the EcoReality of microbiome transplant studies that
153 used non-human recipients. We show that, overall, the breadth of EcoReality of the
154 present microbiome transplant literature has increased over time. However, EcoReality
155 has been constrained by hosts bred and kept in lab conditions, and with transplants into
156 germ-free recipient hosts. Importantly, we provide a conceptual framework, illustrated in
157 Figure 1, to help broaden the range of EcoReality in transplant experiments and to

158 facilitate comparisons between transplants of varying EcoReality.
 159



161 Figure 1: Conceptual framework of all the experimental conditions in a microbial
 162 transplant where EcoReality can vary. See Box 2 for explanations for each experimental
 163 condition.

164

165 **Box 1.** Key terms and definitions

166

Term	Definition
Transplant Instance	A transplant of a microbial strain or community from its native host or substrate to a different host population. A given study can involve multiple transplant instances, which are delineated based on non-substitutability of host populations or of transplant parameters.
Experimental Conditions	A decision or step in a transplant instance where there is the potential for variation in ecological reality.
Level of EcoReality	The degree to which an experimental condition matches the conditions that a host-microbiome interaction would experience in a wild ecosystem. Each experimental condition possesses its own intrinsic EcoReality. Each transplant instance can also be assigned an EcoReality score.

167

168 **Box 2.** Ecological reasoning for each experimental conditions within a transplant

Experimental condition	Reasoning Ecological theory + application to microbiome
Taxon Match	Organisms can become locally adapted (25). Local adaptation of a microbial species to a given host may mean it is not adapted to hetero-specific hosts and will perform poorly after transplantation (26).
Donor & Recipient Environment	During community assembly, the local environment acts as a filter, incorporating species from a wider species pool (21). From the microbiome’s perspective, the host’s physiology and the external environment are one intertwined environment. Therefore, the external environment can affect microbiome dynamics in two ways: indirectly through impacting the host physiology (27), and directly through the wider microbial species pool that the host and its microbiome has access to.
Donor & Recipient Physiology	The local environment acts as a filter in community assembly (21). For this experimental condition, we define physiology as physiological states that would occur regardless of the external environmental context (e.g. gene knock-out, disease-state). We differentiate intrinsic physiology from mutable host physiological responses to the external environmental context. Although these indirect environmental effects acting through host physiology are relevant (27), they are captured by the ‘Environment’ experimental condition. A host’s physiology is the <i>de facto</i> environment of inhabitant microbes, and changes or dysregulation in the host may disrupt associations between host and the microbes that persist under homeostatic physiological conditions.
Transplanted Microbiome	The interactions within an invading community, including predation or mutualism, can impact whether colonisation is successful or not (20). Thus, a full community microbiome transplantation may differ significantly from the transplantation of a single microbe monoculture at artificially high densities.
Transplant Method	Species have different dispersal abilities (27) and local environments filter species from the wider species pool (28). Active transplantations may circumvent differing dispersal abilities of microbial species and may undermine host filtering of

	the microbial community. Furthermore, active transplant methods can stress the host thereby changing host physiology and disrupting endogenous microbial communities (29).
Recipient Pre-transplant Microbiome	High species diversity in a community is predicted to reduce niche opportunities and to increase invasion resistance (20). Germ-free or antibiotic perturbed recipients are likely to have lower invasion resistance than recipients with intact microbiomes.
Housing conditions	Dispersal between patches is an integral ecological process which can maintain stable populations or can rescue extirpated populations (21,30). Recipient host cohabitation allows for further transmissions of the microbiome.

169

170 **Lay of the land**

171

172 *Literature Search*

173 We conducted a directed review of the existing literature on gut microbiome
174 transplants, finishing on October 26th 2018. We conducted our literature search in three
175 stages. First, to gauge the extent of the current literature, we did a preliminary search of
176 gut microbiome transplant studies using both Google Scholar and Web of Science
177 (University of Guelph subscription). Based on this preliminary search, we conducted a
178 more methodical search using both Google Scholar and Web of Science. Search terms
179 can be found in the Supporting Information (SI). We then sought additional publications
180 through searching the citations of papers already collected using the Web of Science
181 citations tool. We retained only those studies that conducted at least one gut
182 microbiome transplant into a non-human recipient organism. To ensure our findings
183 were generalizable to ecological and evolutionary frameworks across a broad range of
184 taxa and ecosystems, we excluded studies focused on a single human disease, *C.*
185 *difficile*.

186

187 *Literature Evaluation*

188 For each study that met our criteria, we determined the number of transplant
189 instances, which we defined as the transfer of a microbial strain or community from its
190 native host or substrate to a different host population (see Box 1). We used transplant
191 instances as our unit of focus because many studies contained multiple transplant
192 instances which sometimes differed substantially in EcoReality (e.g. Seedorf et al. (31)).
193 For studies that had sequential transplants (i.e., transplant from donor to a first
194 recipient, which then was the donor for a second recipient, e.g. Seedorf et al. (31)), we
195 used only the first phase of the transplant experiment.

196 We identified nine key experimental conditions in a transplant where variation in
197 EcoReality might substantially affect the outcome of the experiment: host taxon match,
198 donor environment, donor physiology, transplanted microbiome, transplant method,
199 recipient pre-transplant microbiome, recipient environment, recipient physiology, and
200 recipient housing conditions (see Figure 1 & Box 2). Each experimental condition was
201 given an ordinal data scale (see SI Table S1) based on the range of observed and
202 possible levels for that condition, with one always representing the lowest level of
203 EcoReality. Our goal was to maintain similar resolution for each highly dimensional
204 experimental condition within our framework. For example, Taxon Match could have
205 included more levels to capture phylogenetic distance, geographic distance, and
206 feeding relationships between host and recipient (32). However given the variation in
207 scale, generalization across host taxa would have then been difficult. The levels in each
208 experimental condition were based on likely conditions found in the wild. For example,
209 with respect to the Transplanted Microbiome experimental condition, a single bacterial
210 strain at high densities entering a host in the wild is less likely than invasion by mixed
211 communities. For each transplant instance, we characterized the level of EcoReality in
212 each of the 9 experimental conditions. To ensure consistent evaluation methods,
213 EcoReality scores for each transplant instance were determined independently by two
214 co-authors (separate pairs of co-authors randomized per paper). The co-author pairs
215 then compared their scores and agreed upon the final transplant EcoReality scores.


216 To determine the overall standardized EcoReality score of a transplant instance,
217 we divided each score by its corresponding maximum potential EcoReality score and
218 then added the scaled scores for each experimental condition. Thus all experimental
219 conditions were equally weighted in the overall calculation of standardized EcoReality.

220 We separated lab rodents from other animals in our results for each experimental
221 condition because the ecology, physiology and genetics of lab-reared, inbred rodent
222 models are heavily modified from wild-type rodents and other wild animals in ways that
223 may affect our understanding of reciprocal host-microbiome interactions (for example
224 Newman et al. (23) and Walter et al. (33)).

225 226 *Literature EcoReality patterns*


227 Our literature search returned 55 articles that met our criteria for inclusion. These
228 articles ranged from having one to 13 transplant instances with an average of 2.91
229 transplant instances per article and a total of 160 from all articles. There was a clear
230 shift over time in the number of articles using microbiome transplants. Notably, there
231 were 20 articles in the first 10 years of our search period in comparison to almost 40
232 articles during 2015-2018 (SI Figure 1). This increase coincided with a switch from
233 mainly lab rodent studies to a more diverse group of donor hosts (Figure 2A, around
234 2013), and later also to more diverse recipient hosts (Figure 2B, around 2016).

235




236
 237 Figure 2: Number of transplant instances over time where the donor or recipient animal
 238 was either a lab rodent (mouse or rat) or another animal.
 239

240 The transplant conditions Donor and Recipient Physiology had the highest
 241 EcoReality with average scores of 1.8 out of 2 (Figure 3C & H). Taxon match (score 1.6
 242 out of 2, Figure 3A), transplanted microbiome (score 2.5 out of 3, Figure 3D), transplant
 243 method (score 1.7 out of 2, Figure 3E), and housing condition (score 1.5 out of 2, Figure
 244 3I) were moderately EcoReal. Donor environment (score 2.4 out of 5, Figure 3B),
 245 recipient environment (score 1.6 out of 5, Figure 3G), and recipient microbiome (score
 246 1.8 out of 3, Figure 3F) had the lowest EcoReality. Breaking EcoReality into recipient
 247 lab rodents and other animals, we see that active transplant methods (score of 1) were
 248 used more for lab rodents, and passive transplant methods (score of 2) were used more
 249 for other animals (Figure 3E). Interestingly, there were fewer transplants with germ-free
 250 recipient lab rodents than germ-free recipient other animals (score of 1) (Figure 3F).
 251 This pattern was driven by bees (19 out of 85 transplant instances from five articles)
 252 and zebrafish (14 out of 85 transplant instances from two articles). Overall, most
 253 transplants were performed with matching (score of 2, Figure 3A) wild-type, non-
 254 diseased donor and recipient hosts (score of 2, Figures 3C & H) using passive
 255 transplant methods (score of 2, Figure 3E) of whole microbial communities (score of 3,
 256 Figure 3D) and with a mixture of individual (score of 1, Figure 3I) and cohousing (score
 257 of 2, Figure 3I) of recipient hosts. However, transplants were mostly in sterile or normal
 258 lab conditions (score of 1 & 2, Figures 3B & G) with germ-free recipient hosts (score of
 259 1, Figure 3F).



260
 261 Figure 3: Number of transplant instances in each experimental condition, separated into
 262 whether the recipient animal was a lab rodent or another animal. The X-axis is the level
 263 of EcoReality, with 1 always the lowest EcoReality. The levels are explained in Table S1
 264 of our Supporting Information.

265
 266 Although the maximal EcoReality score increased, increasing the breadth of
 267 EcoReality studied, the maximal EcoReality score was still below the theoretical
 268 maximum standardized EcoReality score of 9 possible outlined in our framework (Figure
 269 4).



270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283 Figure 4: Standardized EcoReality score for each transplant instance. The grey area
 284 identifies the zone of EcoReality that has been studied in the literature, and the “Here
 285 be Dragons” area is the unexplored zone of EcoReality that is bound at the top by the
 286 theoretical maximum standardized EcoReality score of 9.

287 **Here Be Dragons!**

288

289 The burgeoning field of microbiome research is integrating the traditionally
290 disparate disciplines of ecology, evolution, and physiology, which examine distinct but
291 interrelated processes at different scales. Yet, these interrelated processes across
292 scales are inherent in host-microbiome relationships (e.g. Stothart et al. (34)), and thus
293 further integration of ecology, evolution, and physiology with microbiology will be crucial
294 for unlocking important insights about the interactions between hosts and their
295 microbiome. As microbiome research expands further to include ecological processes
296 that are well established in traditional ecosystems, studies that can capture these
297 processes will be necessary. Here, we expand on the insights from foundational highly-
298 controlled experiments that identified key mechanisms in host-microbiome interactions.
299 We surveyed the state of the microbiome transplant literature and identified gaps in how
300 well ecological processes are captured in transplants, what we term as ecological
301 reality, (i.e., EcoReality, see Box 1). Our results are promising; the breadth of
302 EcoReality is increasing over time in transplant experiments (Figure 4), but there are
303 still some key gaps in the types of studies conducted on host-microbiome interactions
304 (Figure 3). We suggest that a critical step in understanding reciprocal host-microbiome
305 interactions includes explicitly designing a broader array of studies that can evaluate the
306 role of various ecological processes that are known to shape traditional ecological
307 systems.

308

309 Our evaluation of the microbiome transplantation literature revealed broadening
310 EcoReality in experimental procedures. Lately, there has been a sharp increase in
311 taxonomic diversity of both donor and recipient hosts (Figure 2). Transplants often used
312 passive transplantation methods with wild-type non-diseased donors as well as a
313 mixture of individual and cohabitation housing conditions (Figure 3). Finally, the
314 maximal EcoReality score of microbiome transplant studies has increased over time
315 (Figure 4). These results are encouraging because they suggest that researchers are
316 building on the initial flurry of highly-controlled transplant experiments and designing
317 diverse studies that differ in their degree of EcoReality in several of the categories we
318 examined. Continuing to broaden EcoReality will be essential for understanding the
319 ecological and evolutionary processes at work in reciprocal host-microbiome
320 interactions.

321

322 However, our results show that the current literature lacks EcoReality in two key
323 areas: host environment and the state of the recipient microbiome (Figure 3). Although
324 the environment of the donor hosts was on average more EcoReal than the
325 environment of the recipient hosts, the EcoReality of the donor and recipient host's
326 environments was generally low. Most studies that we evaluated used laboratory
327 settings that exclude the chance for hosts to encounter the broader microbial species

328 pool in the environment (18,21). Laboratory conditions can also either increase or
329 decrease conspecific interactions relative to what would be observed in nature, thus
330 affecting the dispersal of microbes between hosts (18). Furthermore, laboratory
331 conditions may be obscuring feedbacks between the host and its microbiome that can
332 impact diet and habitat choice (18). The second key area lacking EcoReality is the state
333 of the recipient microbiome where most recipient hosts were germ-free. Although some
334 animals naturally start out with germ-free gastrointestinal tracts (e.g., newly eclosed
335 worker bees (35)) or do not have a resident microbiome (36), most animal species host
336 substantial microbial communities (37). Germ-free gastrointestinal tracts may lack key
337 biotic processes such as predation, competition, and facilitation, which are important
338 filters in classic ecological communities that act to mediate incoming species (20,38).
339 Overall, neglecting natural environments and intact recipient microbiomes risks
340 constraining the fundamental processes that impact reciprocal host-microbiome
341 interactions.

342
343 Consequently, we advocate for more breadth in EcoReality in microbiome
344 transplant experiments. This breadth includes highly-controlled laboratory transplants,
345 which offer critical points of comparison, and provide a focused understanding of
346 particular mechanisms. The wider breadth of EcoReality for which we are advocating for
347 requires that we venture into the largely untested realm of highly EcoReal experimental
348 conditions (Figure 4), despite the logistical challenges likely associated with wild
349 conditions. There are many ways in which we might venture beyond our present
350 frontier. For example, using wild-caught animals that are either allowed to roam freely or
351 are housed in outdoor enclosures, and to use recipient animals with intact microbiomes,
352 rather than germ-free microbiomes (Figure 3). We also suggest identifying and
353 addressing the major phylogenetic gaps in the tree of life for the donor and recipient
354 host taxa. Overall, we call for a balance of studies dealing with all permutations of
355 EcoReality in each experimental condition. We hope researchers will use and adapt our
356 conceptual framework (Figure 1) in their own systems to incorporate EcoReality and,
357 where appropriate, consider how constrained EcoReality may impact their conclusions.
358 Likewise, we encourage researchers to report the methodological details pertaining to
359 each experimental condition we have identified. We hope that our literature evaluation
360 and conceptual framework will stimulate new avenues of collaborative research that will
361 evaluate the role of ecological processes in host-microbiome interactions.

362
363 Our literature evaluation suggests that we may understand only a small subset of
364 possible reciprocal host-microbiome interactions impacting our ability to assess the
365 conservation potential of the microbiome. Because we are presently likely constraining
366 fundamental ecological and evolutionary processes, host-microbiome studies may be
367 biased towards results that indicate a strong role of the microbiome on the host. Yet
368 researchers have already made strong and general assertions about the role of the

369 microbiome in the biology of the host. Due to the large effects of the microbiome on its
370 host and its mutability, Alberdi et al. (14) argued that the microbiome could act as an
371 additional axis of ecological adaptation for hosts. If the microbiome does act as an
372 additional axis, conserving microbial diversity and using bioaugmentation tools
373 (probiotic therapy and transplantation of microbiomes) would then be critical tools for
374 animal conservation (39,40). We caution that experimental protocols that lack
375 EcoReality might lead us to overestimate the capacity for microbiome variation to shape
376 host phenotypes in nature by biasing our understanding of the host-microbiome
377 relationship towards models of symbiont control (11). We suspect that a full reckoning of
378 the spectrum of EcoReality in microbiome transplant studies will uncover more
379 examples of the ‘ecosystem on a leash’ model (11), which posits an important but more
380 limited reciprocity between the host and the ‘ecosystem’ of the microbiome. These sorts
381 of nuanced interactions may or may not include the large microbiome effects which
382 underpin the ecological adaptation and conservation arguments above. Thus, we may
383 not yet have the level of understanding about reciprocal host-microbiome interactions
384 that is required to know the role of the microbiome in host adaptation or to confidently
385 inform conservation efforts. Moving forward, we assert that a consideration of
386 EcoReality is required in the design and interpretation of every study that explores how
387 the host-microbiome relationship impacts ecological adaptation.

388

389 Microbiome research has undoubtedly fascinated biologists across disciplines,
390 prompting advances in both pure and applied research and raising questions about
391 some of the most fundamental ideas in biology (13). Yet, the lay of the land in terms of
392 ecological reality of this rapidly growing research area was unexplored. Our objective
393 here—to survey the breadth of EcoReality in the microbiome transplant literature and
394 identify key areas lacking EcoReality—was not unlike a fact-finding mission expanding
395 the map of our understanding of reciprocal host-microbiome interactions. We
396 recommend a full, extended journey into the wilds to round out the literature’s coverage
397 of the landscape of possible EcoReality. Charting all territories, from highly controlled
398 lab studies to free-ranging organisms, is necessary to fully comprehend the interplay
399 between microbiomes and their hosts.

400

401 **Acknowledgements**

402 We thank the authors, animals, and microbes of the many studies that provided the
403 peaks and valleys that we traversed in our exploration. CJGG thanks his PhD
404 supervisor, Kevin McCann, for his support. We thank the attendees of the Evelyn Pielou
405 Discussion Group for inspiring this voyage.

406

407 **Funding**

408 CJGG was supported by a Natural Sciences and Engineering Research Council of
409 Canada (NSERC) CGS-D. KC acknowledges the support of NSERC, RGPIN-2018-

410 04399. MRS was supported by a NSERC Vanier CGS.

411

412 **Author contributions**

413 All authors conceived of and produced the directed review. CJGG wrote the first draft
414 and all authors contributed to editing the manuscript.

415

416 **Data accessibility**

417 The supporting information is included with this paper. The supporting information plus
418 the data, and R script for this manuscript can be found on Zenodo/Github (41). The full
419 list of transplant studies used in this article can be found in the dataset on
420 Zenodo/GitHub (41), and (10,26,31,35,42–92) in the bibliography below.

421

422 **Bibliography**

- 423 1. Eliot T, Eliot T, Hodgson A, Mairet P. Little Gidding. London, United Kingdom: Faber and
424 Faber; 1942.
- 425 2. Lee YK, Mazmanian SK. Has the microbiota played a critical role in the evolution of the
426 adaptive immune system? *Science*. 2010 Dec 24;330(6012):1768–73.
- 427 3. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An obesity-
428 associated gut microbiome with increased capacity for energy harvest. *Nature*. 2006
429 Dec;444(7122):1027–31.
- 430 4. Chevalier C, Stojanović O, Colin DJ, Suarez-Zamorano N, Tarallo V, Veyrat-Durebex C, et
431 al. Gut microbiota orchestrates energy homeostasis during cold. *Cell*. 2015
432 Dec;163(6):1360–74.
- 433 5. Sudo N, Chida Y, Aiba Y, Sonoda J, Oyama N, Yu X-N, et al. Postnatal microbial
434 colonization programs the hypothalamic-pituitary-adrenal system for stress response in
435 mice: Commensal microbiota and stress response. *J Physiol*. 2004 Jul;558(1):263–75.
- 436 6. Sommer F, Bäckhed F. The gut microbiota — masters of host development and
437 physiology. *Nat Rev Microbiol*. 2013 Apr;11(4):227–38.
- 438 7. Gareau MG, Wine E, Rodrigues DM, Cho JH, Whary MT, Philpott DJ, et al. Bacterial
439 infection causes stress-induced memory dysfunction in mice. *Gut*. 2011 Mar 1;60(3):307–
440 17.
- 441 8. Desbonnet L, Clarke G, Shanahan F, Dinan TG, Cryan JF. Microbiota is essential for
442 social development in the mouse. *Mol Psychiatry*. 2014 Feb;19(2):146–8.
- 443 9. Kohl KD, Carey HV. A place for host–microbe symbiosis in the comparative physiologist’s
444 toolbox. *J Exp Biol*. 2016 Nov 15;219(22):3496–504.
- 445 10. Zhang X-Y, Sukhchuluun G, Bo T-B, Chi Q-S, Yang J-J, Chen B, et al. Huddling remodels
446 gut microbiota to reduce energy requirements in a small mammal species during cold
447 exposure. *Microbiome*. 2018 Dec;6(1):103.
- 448 11. Foster KR, Schluter J, Coyte KZ, Rakoff-Nahoum S. The evolution of the host microbiome
449 as an ecosystem on a leash. *Nature*. 2017 Aug 2;548(7665):43–51.
- 450 12. Rosenberg E, Koren O, Reshef L, Efrony R, Zilber-Rosenberg I. The role of
451 microorganisms in coral health, disease and evolution. *Nat Rev Microbiol*. 2007
452 May;5(5):355–62.
- 453 13. Zilber-Rosenberg I, Rosenberg E. Role of microorganisms in the evolution of animals and
454 plants: the hologenome theory of evolution. *FEMS Microbiol Rev*. 2008 Aug;32(5):723–35.
- 455 14. Alberdi A, Aizpurua O, Bohmann K, Zepeda-Mendoza ML, Gilbert MTP. Do vertebrate gut
456 metagenomes confer rapid ecological adaptation? *Trends Ecol Evol*. 2016 Sep;31(9):689–

- 457 99.
- 458 15. Koskella B, Hall LJ, Metcalf CJE. The microbiome beyond the horizon of ecological and
459 evolutionary theory. *Nat Ecol Evol*. 2017 Nov;1(11):1606–15.
- 460 16. Costello EK, Stagaman K, Dethlefsen L, Bohannan BJM, Relman DA. The application of
461 ecological theory toward an understanding of the human microbiome. *Science*.
462 2012;336:1255–62.
- 463 17. Ley RE, Lozupone CA, Hamady M, Knight R, Gordon JI. Worlds within worlds: evolution of
464 the vertebrate gut microbiota. *Nat Rev Microbiol*. 2008 Oct;6(10):776–88.
- 465 18. Miller ET, Svanbäck R, Bohannan BJM. Microbiomes as metacommunities: Understanding
466 host-associated microbes through metacommunity ecology. *Trends Ecol Evol*. 2018
467 Sep;33(12):926–35.
- 468 19. Samarkos M, Mastrogianni E, Kampouropoulou O. The role of gut microbiota in
469 *Clostridium difficile* infection. *Eur J Intern Med*. 2018;50:28–32.
- 470 20. Shea K, Chesson P. Community ecology theory as a framework for biological invasions.
471 *Trends Ecol Evol*. 2002 Apr 1;17(4):170–6.
- 472 21. Leibold MA, Holyoak M, Mouquet N, Amarasekare P, Chase JM, Hoopes MF, et al. The
473 metacommunity concept: a framework for multi-scale community ecology. *Ecol Lett*.
474 2004;7(7):601–13.
- 475 22. Carrier TJ, Reitzel AM. The Hologenome Across Environments and the Implications of a
476 Host-Associated Microbial Repertoire. *Front Microbiol*. 2017 May 11;8:802.
- 477 23. Newman AEM, Edmunds NB, Ferraro S, Heffell Q, Merritt GM, Pakkala JJ, et al. Using
478 ecology to inform physiology studies: implications of high population density in the
479 laboratory. *Am J Physiol-Regul Integr Comp Physiol*. 2015 Mar 15;308(6):R449–54.
- 480 24. Hanage WP. Microbiome science needs a healthy dose of scepticism: to guard against
481 hype, those interpreting research on the body’s microscopic communities should ask five
482 questions. *Nature*. 2014;512(7514):247–8.
- 483 25. Kawecki TJ, Ebert D. Conceptual issues in local adaptation. *Ecol Lett*. 2004
484 Dec;7(12):1225–41.
- 485 26. Brooks AW, Kohl KD, Brucker RM, van Opstal EJ, Bordenstein SR. Phyllosymbiosis:
486 relationships and functional effects of microbial communities across host evolutionary
487 history. Relman D, editor. *PLOS Biol*. 2016 Nov 18;14(11):e2000225.
- 488 27. Bie T, Meester L, Brendonck L, Martens K, Goddeeris B, Ercken D, et al. Body size and
489 dispersal mode as key traits determining metacommunity structure of aquatic organisms.
490 *Ecol Lett*. 2012 Jul;15(7):740–7.
- 491 28. Chase JM. Community assembly: when should history matter? *Oecologia*.
492 2003;136(4):489–498.
- 493 29. Allen-Blevins CR, You X, Hinde K, Sela DA. Handling stress may confound murine gut
494 microbiota studies. *PeerJ*. 2017 Jan 11;5:e2876.
- 495 30. Zelnik YR, Arnoldi J, Loreau M. The three regimes of spatial recovery. *Ecology*. 2019
496 Feb;100(2):e02586.
- 497 31. Seedorf H, Griffin NW, Ridaura VK, Reyes A, Cheng J, Rey FE, et al. Bacteria from
498 diverse habitats colonize and compete in the mouse gut. *Cell*. 2014 Oct;159(2):253–66.
- 499 32. Moeller AH, Suzuki TA, Lin D, Lacey EA, Wasser SK, Nachman MW. Dispersal limitation
500 promotes the diversification of the mammalian gut microbiota. *Proc Natl Acad Sci*. 2017
501 Dec 26;114(52):13768–73.
- 502 33. Walter J, Armet AM, Finlay BB, Shanahan F. Establishing or Exaggerating Causality for
503 the Gut Microbiome: Lessons from Human Microbiota-Associated Rodents. *Cell*. 2020
504 Jan;180(2):221–32.
- 505 34. Stothart MR, Palme R, Newman AEM. It’s what’s on the inside that counts: stress
506 physiology and the bacterial microbiome of a wild urban mammal. *Proc R Soc B Biol Sci*.
507 2019 Oct 23;286(1913):20192111.

- 508 35. Näpflin K, Schmid-Hempel P. Immune response and gut microbial community structure in
509 bumblebees after microbiota transplants. *Proc R Soc B Biol Sci.* 2016 May
510 25;283(1831):20160312.
- 511 36. Hammer TJ, Janzen DH, Hallwachs W, Jaffe SP, Fierer N. Caterpillars lack a resident gut
512 microbiome. *Proc Natl Acad Sci.* 2017 Sep 5;114(36):9641–6.
- 513 37. Russell JA, Dubilier N, Rudgers JA. Nature's microbiome: introduction. *Mol Ecol.* 2014
514 Mar;23(6):1225–37.
- 515 38. Jackson DA, Peres-Neto PR, Olden JD. What controls who is where in freshwater fish
516 communities – the roles of biotic, abiotic, and spatial factors. *Can J Fish Aquat Sci.*
517 2001;58(1):157–70.
- 518 39. Trevelline BK, Fontaine SS, Hartup BK, Kohl KD. Conservation biology needs a microbial
519 renaissance: a call for the consideration of host-associated microbiota in wildlife
520 management practices. *Proc R Soc B Biol Sci.* 2019 Jan 30;286(1895):20182448.
- 521 40. Banerjee A, Cornejo J, Bandopadhyay R. Emergent climate change impact throughout the
522 world: call for “Microbiome Conservation” before it’s too late. *Biodivers Conserv.* 2019
523 Oct;1–4.
- 524 41. Greyson-Gaito CJ, Bartley TJ, Cottenie K, Jarvis W, Newman AEM, Stothart MR. Data, SI
525 & R Script for: Into the wild: microbiome transplant studies need broader ecological reality.
526 (Version v2.1). Zenodo/Github. <https://doi.org/10.5281/zenodo.2652255>. 2020.
- 527 42. Rawls JF, Mahowald MA, Ley RE, Gordon JI. Reciprocal gut microbiota transplants from
528 zebrafish and mice to germ-free recipients reveal host habitat selection. *Cell.* 2006
529 Oct;127(2):423–33.
- 530 43. Pang X, Hua X, Yang Q, Ding D, Che C, Cui L, et al. Inter-species transplantation of gut
531 microbiota from human to pigs. *ISME J.* 2007 Jun;1(2):156–62.
- 532 44. Manichanh C, Reeder J, Gibert P, Varela E, Llopis M, Antolin M, et al. Reshaping the gut
533 microbiome with bacterial transplantation and antibiotic intake. *Genome Res.* 2010 Oct
534 1;20(10):1411–9.
- 535 45. Bindels LB, Beck R, Schakman O, Martin JC, De Backer F, Sohet FM, et al. Restoring
536 specific Lactobacilli levels decreases inflammation and muscle atrophy markers in an
537 acute leukemia mouse model. *PLoS ONE.* 2012;7(6):e37971.
- 538 46. Bindels LB, Neyrinck AM, Claus SP, Le Roy CI, Grangette C, Pot B, et al. Synbiotic
539 approach restores intestinal homeostasis and prolongs survival in leukaemic mice with
540 cachexia. *ISME J.* 2016 Jun;10(6):1456–70.
- 541 47. Koch H, Schmid-Hempel P. Gut microbiota instead of host genotype drive the specificity in
542 the interaction of a natural host-parasite system. *Ecol Lett.* 2012 Oct;15(10):1095–103.
- 543 48. Upadhyay V, Poroyko V, Kim T, Devkota S, Fu S, Liu D, et al. Lymphotoxin regulates
544 commensal responses to enable diet-induced obesity. *Nat Immunol.* 2012 Aug
545 26;13(10):947–53.
- 546 49. Toh MC, Goodyear M, Daigneault M, Allen-Vercoe E, Van Raay TJ. Colonizing the
547 embryonic zebrafish gut with anaerobic bacteria derived from the human gastrointestinal
548 tract. *Zebrafish.* 2013 Jun;10(2):194–8.
- 549 50. Ridaura VK, Faith JJ, Rey FE, Cheng J, Duncan AE, Kau AL, et al. Gut microbiota from
550 twins discordant for obesity modulate metabolism in mice. *Science.*
551 2013;341(6150):1241214–1241214.
- 552 51. Kohl KD, Weiss RB, Cox J, Dale C, Denise Dearing M. Gut microbes of mammalian
553 herbivores facilitate intake of plant toxins. *Ecol Lett.* 2014 Oct;17(10):1238–46.
- 554 52. Li M, Liang P, Li Z, Wang Y, Zhang G, Gao H, et al. Fecal microbiota transplantation and
555 bacterial consortium transplantation have comparable effects on the re-establishment of
556 mucosal barrier function in mice with intestinal dysbiosis. *Front Microbiol.* 2015 Jul 7;6:692.
- 557 53. Bruce-Keller AJ, Salbaum JM, Luo M, Blanchard E, Taylor CM, Welsh DA, et al. Obese-
558 type gut microbiota induce neurobehavioral changes in the absence of obesity. *Biol*

- 559 Psychiatry. 2015;77(7):607–15.
- 560 54. Di Luccia B, Crescenzo R, Mazzoli A, Cigliano L, Venditti P, Walser J-C, et al. Rescue of
561 fructose-induced metabolic syndrome by antibiotics or faecal transplantation in a rat model
562 of obesity. PLOS ONE. 2015 Aug 5;10(8):e0134893.
- 563 55. Hensley-McBain T, Zevin AS, Manuzak J, Smith E, Gile J, Miller C, et al. Effects of fecal
564 microbial transplantation on microbiome and immunity in simian immunodeficiency virus-
565 infected macaques. J Virol. 2016 May 15;90(10):4981–9.
- 566 56. Twitchell EL, Tin C, Wen K, Zhang H, Becker-Dreps S, Azcarate-Peril MA, et al. Modeling
567 human enteric dysbiosis and rotavirus immunity in gnotobiotic pigs. Gut Pathog. 2016
568 Dec;8(1).
- 569 57. Yoon MY, Min KB, Lee K-M, Yoon Y, Kim Y, Oh YT, et al. A single gene of a commensal
570 microbe affects host susceptibility to enteric infection. Nat Commun. 2016 May 13;7:11606.
- 571 58. Kelly JR, Borre Y, O' Brien C, Patterson E, El Aidy S, Deane J, et al. Transferring the
572 blues: Depression-associated gut microbiota induces neurobehavioural changes in the rat.
573 J Psychiatr Res. 2016 Nov;82:109–18.
- 574 59. Tian Z, Liu J, Liao M, Li W, Zou J, Han X, et al. Beneficial effects of fecal microbiota
575 transplantation on ulcerative colitis in mice. Dig Dis Sci. 2016 Aug;61(8):2262–71.
- 576 60. Wang Y, Xie J, Li Y, Dong S, Liu H, Chen J, et al. Probiotic *Lactobacillus casei* Zhang
577 reduces pro-inflammatory cytokine production and hepatic inflammation in a rat model of
578 acute liver failure. Eur J Nutr. 2016;55(2):821–831.
- 579 61. Yan H, Diao H, Xiao Y, Li W, Yu B, He J, et al. Gut microbiota can transfer fiber
580 characteristics and lipid metabolic profiles of skeletal muscle from pigs to germ-free mice.
581 Sci Rep. 2016 Oct;6(1):31786.
- 582 62. Mikaelyan A, Thompson CL, Hofer MJ, Brune A. Deterministic assembly of complex
583 bacterial communities in guts of germ-free cockroaches. Appl Environ Microbiol.
584 2016;82(4):1256–63.
- 585 63. Schwarz RS, Moran NA, Evans JD. Early gut colonizers shape parasite susceptibility and
586 microbiota composition in honey bee workers. Proc Natl Acad Sci. 2016 Aug
587 16;113(33):9345–50.
- 588 64. Kohl KD, Stengel A, Dearing MD. Inoculation of tannin-degrading bacteria into novel hosts
589 increases performance on tannin-rich diets: Tannin-degrading bacteria aid hosts
590 consuming tannins. Environ Microbiol. 2016 Jun;18(6):1720–9.
- 591 65. Sonowal R, Swimm A, Sahoo A, Luo L, Matsunaga Y, Wu Z, et al. Indoles from
592 commensal bacteria extend healthspan. Proc Natl Acad Sci. 2017 Sep 5;114(36):E7506–
593 15.
- 594 66. Miller AW, Dale C, Dearing MD. The induction of oxalate metabolism *In Vivo* is more
595 effective with functional microbial communities than with functional microbial species.
596 mSystems. 2017;2(5):e00088-17.
- 597 67. Schmidt V, Gomez-Chiarri M, Roy C, Smith K, Amaral-Zettler L. Subtle microbiome
598 manipulation using probiotics reduces antibiotic-associated mortality in fish. mSystems.
599 2017;2(6):e00133-17.
- 600 68. Warne RW, Kirschman L, Zeglin L. Manipulation of gut microbiota reveals shifting
601 community structure shaped by host developmental windows in Amphibian larvae. Integ
602 Comp Biol. 2017;57(4):786–94.
- 603 69. Ericsson AC, Personett AR, Turner G, Dorfmeier RA, Franklin CL. Variable colonization
604 after reciprocal fecal microbiota transfer between mice with low and high richness
605 microbiota. Front Microbiol. 2017 Feb 23;8:196.
- 606 70. Ekmekciu I, von Klitzing E, Fiebiger U, Escher U, Neumann C, Bacher P, et al. Immune
607 responses to broad-spectrum antibiotic treatment and fecal microbiota transplantation in
608 mice. Front Immunol. 2017;8:397.
- 609 71. Smith P, Willemsen D, Popkes M, Metge F, Gandiwa E, Reichard M, et al. Regulation of

- 610 life span by the gut microbiota in the short-lived African turquoise killifish. *Elife*.
611 2017;6:e27014.
- 612 72. von Klitzing E, Ekmekci I, Kühl AA, Bereswill S, Heimesaat MM. Intestinal, extra-intestinal
613 and systemic sequelae of *Toxoplasma gondii* induced acute ileitis in mice harboring a
614 human gut microbiota. *PLOS ONE*. 2017;12(4):e0176144.
- 615 73. Macke E, Callens M, De Meester L, Decaestecker E. Host-genotype dependent gut
616 microbiota drives zooplankton tolerance to toxic cyanobacteria. *Nat Commun*.
617 2017;8(1):1608.
- 618 74. Berasategui A, Salem H, Paetz C, Santoro M, Gershenzon J, Kaltenpoth M, et al. Gut
619 microbiota of the pine weevil degrades conifer diterpenes and increases insect fitness. *Mol*
620 *Ecol*. 2017 Aug;26(15):4099–110.
- 621 75. Kwong WK, Mancenido AL, Moran NA. Immune system stimulation by the native gut
622 microbiota of honey bees. *R Soc Open Sci*. 2017 Feb;4(2):170003.
- 623 76. Alberoni D, Baffoni L, Gaggia F, Ryan PM, Murphy K, Ross PR, et al. Impact of beneficial
624 bacteria supplementation on the gut microbiota, colony development and productivity of
625 *Apis mellifera* L. *Benef Microbes*. 2018;9(2):269–78.
- 626 77. Billiet A, Meeus I, Cnockaert M, Vandamme P, Van Oystaeyen A, Wäckers F, et al. Effect
627 of oral administration of lactic acid bacteria on colony performance and gut microbiota in
628 indoor-reared bumblebees (*Bombus terrestris*). *Apidologie*. 2017;48(1):41–50.
- 629 78. Erkosar B, Kolly S, van der Meer JR, Kawecki TJ. Adaptation to chronic nutritional stress
630 leads to reduced dependence on microbiota in *Drosophila melanogaster*. *mBio*.
631 2017;8:e01496-17.
- 632 79. Moghadam NN, Thorshauge PM, Kristensen TN, de Jonge N, Bahrndorff S, Kjeldal H, et
633 al. Strong responses of *Drosophila melanogaster* microbiota to developmental
634 temperature. *Fly (Austin)*. 2018;12(1):1–12.
- 635 80. Shi J, Wang Y, He J, Li P, Jin R, Wang K, et al. Intestinal microbiota contributes to colonic
636 epithelial changes in simulated microgravity mouse model. *FASEB J*. 2017;31(8):3695–
637 3709.
- 638 81. Ji S, Hui Y, Jiang T, Guo C, Liu J, Su H, et al. Ecological restoration of antibiotic-disturbed
639 gastrointestinal microbiota in foregut and hindgut of cows. *Front Cell Infect Microbiol*.
640 2018;8:79.
- 641 82. Schroeder BO, Birchenough GMH, Ståhlman M, Arike L, Johansson MEV, Hansson GC, et
642 al. Bifidobacteria or fiber protects against diet-induced microbiota-mediated colonic mucus
643 deterioration. *Cell Host Microbe*. 2018;23(1):27-40.e7.
- 644 83. Näpflin K, Schmid-Hempel P. Host effects on microbiota community assembly. *J Anim*
645 *Ecol*. 2018;87(2):331–40.
- 646 84. Mockler BK, Kwong WK, Moran NA, Koch H. Microbiome structure influences infection by
647 the parasite *Crithidia bombi* in bumble bees. *Appl Environ Microbiol*. 2018;84(7):e02335-
648 17.
- 649 85. Hu L, Geng S, Li Y, Cheng S, Fu X, Yue X, et al. Exogenous fecal microbiota
650 transplantation from local adult pigs to crossbred newborn piglets. *Front Microbiol*. 2018
651 Jan 9;8:2663.
- 652 86. Siegerstetter S-C, Petri RM, Magowan E, Lawlor PG, Zebeli Q, O'Connell NE, et al. Fecal
653 microbiota transplant from highly feed-efficient donors shows little effect on age-related
654 changes in feed-efficiency-associated fecal microbiota from chickens. *Appl Environ*
655 *Microbiol*. 2018;84(2):e02330-17.
- 656 87. Bidu C, Escoula Q, Bellenger S, Spor A, Galan M, Geissler A, et al. The transplantation of
657 ω 3 PUFA–altered gut microbiota of fat-1 mice to wild-type littermates prevents obesity and
658 associated metabolic disorders. *Diabetes*. 2018;67(8):1512–23.
- 659 88. Wrzosek L, Ciocan D, Borentain P, Spatz M, Puchois V, Hugot C, et al. Transplantation of
660 human microbiota into conventional mice durably reshapes the gut microbiota. *Sci Rep*.

- 661 2018 Dec;8(1):6854.
- 662 89. Lee H, Lee Y, Kim J, An J, Lee S, Kong H, et al. Modulation of the gut microbiota by
663 metformin improves metabolic profiles in aged obese mice. *Gut Microbes*. 2018;9(2):155–
664 65.
- 665 90. Moeller AH, Suzuki TA, Phifer-Rixey M, Nachman MW. Transmission modes of the
666 mammalian gut microbiota. 2018;5.
- 667 91. Sommer F, Ståhlman M, Ilkayeva O, Arnemo JM, Kindberg J, Josefsson J, et al. The gut
668 microbiota modulates energy metabolism in the hibernating brown bear *Ursus arctos*. *Cell*
669 *Rep*. 2016 Feb;14(7):1655–61.
- 670 92. Miller AW, Oakeson KF, Dale C, Dearing MD. Microbial community transplant results in
671 increased and long-term oxalate degradation. *Microb Ecol*. 2016 Aug;72(2):470–8.
672

Supporting Information

Into the wild: microbiome transplant studies need broader ecological reality

Authors

Christopher J. Greyson-Gaito*^{1†}, Timothy J. Bartley^{1,2†}, Karl Cottenie^{1†}, William Jarvis^{3†}, Amy E.M. Newman^{1†}, Mason R. Stothart^{4†}

Affiliations

*Corresponding author - christopher@greyson-gaito.com

1. University of Guelph, Department of Integrative Biology, Guelph, ON, Canada
2. University of Toronto Mississauga, Mississauga, ON, Canada
3. University of Ottawa, Department of Biology, Ottawa, ON, Canada
4. University of Calgary, Calgary, AB, Canada

†All authors contributed equally

Search terms used for methodological literature search

- Google scholar
 - (transplant* AND microbio) (gut OR fecal OR feces OR gastrointestinal OR gastro-intestin* OR faecal OR faeces OR forces OR faecal) -human -patient -"homo sapiens" -man -woman -child*
- Web of Science
 - ((TOPIC:(transplant) AND TOPIC: (microbio))AND (TOPIC: (((((((gut OR fecal) OR feces) OR gastrointestinal) OR gastro-intestin) OR faecal) OR faeces) OR foeces) OR faecal) NOT TOPIC:((((human OR patient) OR homo sapiens) OR man) OR woman) OR child*)))

Table S1 Ordinal data scale (EcoReality score) for each experimental condition

Experimental Condition	Ordinal Data Scale
Taxon Match	1 = Mismatch (different species) 2 = Match (same species)
Donor Environment	1 = Lab animal host in sterile lab 2 = Lab animal host in non-sterile lab 3 = Captive bred wildlife (multiple generations bred in captivity) 4 = Wildlife brought into captivity (no generations bred in captivity) 5 = Free-ranging wildlife (capture and release)
Donor Physiology	1 = Gene knockout or disease harbouring (non-microbial) 2 = Wildtype non-diseased
Transplanted Microbiome	1 = Single strain 2 = Consortium, mixture of select strains

	3 = Whole community (no sorting or altering of community sampled for transplantation)
Transplant Method	1 = Active (microbiome sample forcefully added to recipient gut e.g., by a suppository or oral gavage) 2 = Passive (microbiome sample passively given to recipient e.g., mixed into food)
Recipient Microbiome	1 = Germ-free 2 = Antibiotic perturbed/pathologic 3 = Whole community (no experimental alteration of community)
Recipient Environment	1 = Lab animal host in sterile lab 2 = Lab animal host in non-sterile lab 3 = Captive bred wildlife (multiple generations bred in captivity) 4 = Wildlife brought into captivity (no generations bred in captivity) 5 = Free-ranging wildlife (capture and release)
Recipient Physiology	1 = Gene knockout or disease harbouring (non-microbial) 2 = Wildtype or non-diseased
Housing Conditions	1 = Housed singly (after microbiome transplantation) 2 = Co-housed (after microbiome transplantation with either other replicates in the experiment or with individuals of the same species that were not replicates. Co-housing could also have been used as the method of transplantation)




Figure S1 Cumulative sum of articles from our directed review between 2006 and 2018.

Data accessibility

The data, the above supporting information, and the R script for this manuscript are in a repository on GitHub. This repository can be cloned or downloaded straight from Github (https://github.com/cgreysongaito/Intothewild_Microbiome) or from Zenodo (<https://doi.org/10.5281/zenodo.2652255>).

Folder and file structure of Github repository (Intothewild_Microbiome)

- data
 - EcoRealTable_2019-10-09_Data.csv
- figs
 - 2019-10-09 CountAnimals.pdf – Figure 2 in manuscript
 - 2019-10-09 Eco-realityComparisons.pdf – Figure 3 in manuscript
 - 2019-10-09 Eco-realityAverageStandardOverTime.pdf – Figure 4 in manuscript
 - 2019-10-09 CumulativeSumArticles.pdf – Supporting Information Figure 1
- .gitignore – File containing files or folders that git should ignore
- IntotheWild_Microbiome_Greyson-Gaito_etal_2019.R – R script for analysis and figure creation
- SupportingInformation_Intothewild_GreysonGaitoetal.pdf – Supporting information (search terms, ordinal data scales, figure)
- LICENSE – Mozilla Public License 2.0
- README.md – Important information
- meta_transplant_microbiome.Rproj – R Project to increase ease of use

Instructions for use

- Download the whole repository (either by forking and cloning or by downloading a ZIP folder)
- In RStudio, open the project called meta_transplant_microbiome.Rproj and open the file Intothewild_Microbiome_Greyson-Gaito_etal_2019.R
 - If not using RStudio, open the file Intothewild_Microbiome_Greyson-Gaito_etal_2019.R and edit the path to the data file called EcoRealTable_2019-10-09_Data.csv to whatever path is required on your computer.
- Run the script in RStudio or however you normally run R scripts