

1 **This pre-print has been accepted for publication in the journal *Estuaries and Coasts***

2 Alice Stocco^{a*}, Laura Basconi^a, Silvia Rova^a, Fabio Pranovi^a

3 **Like little lagoons: the contribution of *valli da pesca* to the**
4 **Ecosystem Services supply of the Venice Lagoon**

5
6 Original Report of Research Results Article

7 ^a Department of Environmental Sciences, Informatics and Statistics,
8 Ca' Foscari University of Venice,
9 Via Torino, 155
10 30170 Mestre – Venezia
11 Italy

12
13 *Corresponding author:
14 Alice Stocco
15 Ca' Foscari University of Venice,
16 Via Torino, 155
17 30170 Mestre – Venezia
18 Italy
19 alice.stocco@unive.it
20 + 39 333 847 9659
21 +39 041 234 7713
22

Like little lagoons: the contribution of *valli da pesca* to the Ecosystem Services supply of the Venice Lagoon

Alice Stocco^{a*}, Laura Basconi^a, Silvia Rova^a, Fabio Pranovi^a

^a Department of Environmental Sciences, Informatics and Statistics, Università Ca' Foscari, Venezia, Italy

*Corresponding author: alice.stocco@unive.it

Abstract

The Venice lagoon social-ecological system is characterized by a strong relationship between the natural environment and human activities. This is especially noticeable in the aquaculture and hunting reserves of the lagoon, locally known as *valli da pesca*. Previous works about Ecosystem Services (ESs) in the Venice lagoon focused on the so-called “open lagoon”, overlooking the role of the *valli da pesca*. Nonetheless, despite being completely managed ecosystems, the *valli da pesca* have conserved typical elements of transitional water environments that the other parts of the lagoon have lost. By evaluating nine ESs using a spatially explicit approach, we found that the *valli da pesca*, despite covering 17% of the surface, are contributing for 38% of the ESs total capacity, and for 24% of the ESs total flow, in comparison to the open part of the lagoon. Moreover, the management that aims to maximize in a perspective of sustainability some provisioning ESs, such as extensive aquaculture, can positively influence the presence of factors on which other ESs capacity is also based. As a result, the open lagoon benefits from a sort of spill-over effect for lifecycle support, hunting, and cultural ESs such as tourism, information for cognitive development, and birdwatching. Such significant contributions could be endangered in the context of a lagoon subjected to increasing pressures from anthropic activities, where adaptations to impacts, as well as to climate change and sea-level rise effects, in the long run will modify the lagoon hydrodynamics and the sea-lagoon connectivity, threatening the *valli da pesca* and so their ESs supply.

Keywords: Ecosystem Services, ‘Side Effects’, Valli da pesca, Extensive aquaculture, Waterfowls Hunting, Venice lagoon

1. Introduction

In the last decades, assessment and mapping of Ecosystem Services (ESs) have become effective methods to highlight all the contributions that humankind receives from Nature (Costanza et al. 1997; Burkhard and Maes 2017; Baró et al. 2016; MEA 2005), making evident also services that would otherwise go unnoticed by the mere fact that they flow effortlessly to human beings (Liu et al. 2007).

Nature and humans always interact, creating complex social-ecological systems, where society receives positive contributions to their well-being through the ecological processes.

In Italy, the Venice lagoon ecosystem is an emblematic example of such a complex social-ecological system. Since remote times, interactions between natural factors, social dynamics, and economic activities have shaped and affected its morphology and functioning (Solidoro et al. 2010). Given the complexity of these interactions,

58 understanding spatial and temporal patterns of ESs capacity and flow (sensu Villamagna et al., 2013) plays a
59 key role in environmental decision-making regarding the Venice lagoon, as suggested by Rova et al. (2015,
60 2019).

61 Previous works, however, focused on the so-called “open lagoon”, composed of the principal islands and
62 water bodies (Newton et al. 2018; la Notte et al. 2017; Rova et al. 2015; D’alpaos and D’alpaos 2021; Rova et
63 al. 2019; Rova et al. 2022). Until now, no data have been gathered about the possible contribution to ESs by
64 some confined, man-managed areas along the lagoon edges, called in Italian “*valli da pesca*”. These areas,
65 considered as Heavily Modified Water Bodies (HMWB) under the Water Framework Directive (European
66 Commission 2000), are located at the interface between the land and the lagoon water and are almost
67 completely closed, covering a total surface of approximately 97 km².

68 Like other similar environments in other coastal lagoons of the Northern Adriatic, the *valli da pesca* were
69 established during the XIV Century as aquaculture facilities, where temporary boundaries were conceived to
70 entrap fish without affecting water flows. Over time, the *valli da pesca* have been progressively isolated from
71 the lagoon with permanent embankments. Nowadays they are used especially for fish farming and waterfowl
72 hunting, and depend almost completely on human intervention for functioning, in terms of freshwater and
73 brackish water supply, as well as for the maintenance of morphological features, resulting in an ecosystem that
74 can be considered like an “artificial ecosystem”. Quite paradoxically, the *valli da pesca*, due to their private
75 management regime, have conserved typical elements of transitional ecosystems that the lagoon itself has
76 instead progressively lost. Consequently, it is legit to ask whether these environments, while including and
77 preserving brackish water basins, freshwater lakes, and saltmarshes, could act as important conservation
78 areas, while providing ESs for the lagoon social-ecological system.

79 This work presents the first quantitative, GIS-based assessment of ESs in the *valli da pesca* of the Venice
80 lagoon. On the one hand, it widens the knowledge about today’s status of these ancient but poorly investigated
81 environments, and on the other hand, it suggests some reflections about the effects of private land
82 conservation. Furthermore, it explores the multiple ESs capacity and flow relationships emerging within these
83 areas, and sheds light on their contribution to the whole lagoon ‘ESs budget’.

84


85 2 Materials and methods

86 2.1 Study area

87 The study area is included in the Venice lagoon, the widest Italian transitional ecosystem (Fig.1). Each of the
88 27 *valli da pesca* that are currently still operative consists of a series of basins at different water salinity,
89 separated from each other, and from the lagoon, by means of artificial embankments, imitating the typical
90 transitional water gradients.

91 Indeed, a typical *valle da pesca* receives the lagoon water that enters through a barrage, flows in the
92 brackish lakes, and then glides towards the land. On the opposite side, some basins store freshwater inputs
93 coming from inland rivers, creating a freshwater wetland area from which the water flows out to reach the
94 major brackish basin. The mixed water then circulates in different sectors of the *valle* and finally, with the low
95 tide, flows out into the lagoon (more information in Supplementary Materials I, Fig. S.M. 1.1). This creates a
96 heterogeneous landscape with strong ecological gradients and a multitude of habitats, as much as a natural
97 lagoon.

98 Among the *valli da pesca*, different types of management can be distinguished, ranging from aquaculture
99 and hunting reserves to tourist estates, but all of them show the common trait of having a restricted access
100 regime, making it difficult to collect data.
101



102

103 **Figure 1** Study area; 1 =Valle Dogà, 2 = Valle Grassabò, 3 = Vallesina, 4 = Valle Fosse, 5 = Valle Lio Maggiore, 6 =
104 Valle Bianca, 7 = Valle Dragojesolo, 8 = Valle Cavallino, 9 = Valle Falconera, 10 = Valle Liona, 11 = Valle Olivara, 12 =
105 Saline-Manciane-Sparasera, 13 = Valle Paleazza, 14 = Valle Sacchettina, 15 = Valle Sacchetta, 16 = Valle Ca' Zane, 17 =
106 Santa Cristina island, 18 = Valle Perini, 19 = Valle Miana-Serraglia, 20 = Valle Avertò, 21 = Valle A.M.A., 22 = Valle
107 Contarina, 23 = Valle Cornio Alto e Cornio Basso, 24 = Valle Zappa, 25 = Valle Figheri, 26 = Valle Pierimpiè, 27 = Valle
108 Morosina-Ghebo Storto.

109 2.2 ESs' data collection, analysis, and mapping

110 The nine Ecosystem Services and related indicators for both capacity and flow are reported in Table 1. The ES
111 category refers to the nomenclature of the Common International Classification of Ecosystem Services (CICES
112 framework).

113

114

CICES section	Ecosystem Service	Capacity indicator	Flow indicator
Regulating & Maintenance services	Climate regulation	Carbon sequestration rate by saltmarshes and seagrasses [$\text{gC m}^{-2} \text{y}^{-1}$] *	
	Water purification	Percentage of Nitrogen load removed by denitrification process in brackish water [%] *	
	*According to Schröter et al. (2014) and Hein et al. (2006) capacity and flow indicators of climate regulation and water purification ESs have been considered coincident		
	Lifecycle support for fish and of avian migratory species	Attractiveness for migratory waterbirds and potentially hostable juveniles fish biomass normalized to a 0-1 scale	Number of wintering migratory waterbirds and sown juveniles fish biomass normalized to a 0-1 scale
Provisioning services	Aquaculture production	Fish biomass [$\text{kg ha}^{-1}\text{y}^{-1}$]	Harvested fish biomass [$\text{kg ha}^{-1}\text{y}^{-1}$]
	Waterbirds' hunting	Number of huntable waterbirds [$\text{n ha}^{-1}\text{y}^{-1}$]	Number of catches [$\text{n ha}^{-1}\text{y}^{-1}$]
	Wild edible plants and honey production	<i>Salicornia sp.</i> biomass [$\text{kg ha}^{-1}\text{y}^{-1}$] Honey [$\text{kg ha}^{-1}\text{y}^{-1}$]	Harvested <i>Salicornia sp.</i> biomass [$\text{kg ha}^{-1}\text{y}^{-1}$] Harvested honey [$\text{kg ha}^{-1}\text{y}^{-1}$]
Cultural services	Tourism	Tourism attractiveness [0-1 scale]	Number of tourists [n y^{-1}]
	Information for cognitive development	Environmental education attractiveness [0-1 scale]	Number of one-day guided excursionists and students [n y^{-1}]
	Birdwatching	Birdwatching attractiveness [0-1 scale]	Mean number of active birdwatchers [n y^{-1}]

115 **Table 1** ESs selected for the present assessment and the adopted indicator.

116 2.2.1 Regulating and maintenance ESs

117 The climate regulation ES has been assessed as carbon sequestration process by saltmarshes accretion (Day et
118 al. 1998; Roner et al. 2015) and seagrasses meadows (Sfriso and Francesco Ghetti, 1998; Sfriso and Marcomini,
119 1999; Sfriso et al., 2007) that are present in the studied reserves.

120 The water purification ES is expressed as the nitrogen removal capacity of brackish lakes, in proportion to
121 the water volume and its turnover time in each *valle da pesca*. Because of the unavailability of local data, the
122 estimation was based on denitrification data by Ravagnan (1982), who measured the difference between the
123 TIN in the inflowing water and outlet water in *valli da pesca* which are in all aspects managed similarly to the
124 *valli da pesca* considered in this case study.

125 According to the literature, the capacity and flow indicators of the aforementioned regulating ESs are
126 considered to be coincident (Schröter et al. 2014; Hein et al. 2006a).

127 The lifecycle support was assessed by focusing on the migration patterns of both the fish and waterbirds.
128 Fish lifecycle support capacity was estimated by extrapolating the distribution of mugilids fingerlings in the
129 most confined part of the lagoon, retrieved from a spatialized food-web model of the Venice lagoon based on
130 functional groups (Anelli Monti et al., 2021). For migratory birds, we mapped for each one of the *valli da pesca*
131 the favorable factors, namely saltmarshes, freshwater presence, shrubs, and herbaceous vegetation, that
132 enhance the attractiveness for resting and molt changing (Korschgen et al., 1985; Havera et al., 1992; Arzel et
133 al., 2006; Hatziordanou et al., 2019). Fish and waterbird lifecycle support capacities were normalized on a 0-
134 1 scale. To assess the flow for fish lifecycle support we referred to the actual mullets' biomass sown per hectare

135 (kg ha⁻¹ y⁻¹), referring to the species *Mugil cephalus*, *Chelon labrosus*, *Chelon aurata*, *Chelon saliens*, and
136 *Chelon ramada*, as officially declared by the *valli da pesca* managers. We focused on mullets because they are
137 the only taxonomic group for which data regarding the sowing of juveniles were available, and because the
138 origin of their fingerlings was local. In contrast, for other farmed species (for example *Sparus aurata*,
139 *Dicentrarchus labrax*), it is more frequent that fry often come from intensive hatcheries, not necessarily
140 located in Northern Italy; thus, we decided to not take them into account for assessing the lagoon fish lifecycle
141 support ES. For birds lifecycle support flow, we referred to the average number of migratory waterbirds that
142 winter within the *valli da pesca*, from the last ten years' waterbirds annual censuses (AFV 2020). The resulting
143 indicators were combined and then normalized on a 0-1 scale.

144

145 2.2.2 Provisioning ESs

146 The aquaculture practiced in the *valli da pesca* is mainly extensive aquaculture. In this type of aquaculture,
147 fish are free to move into a wide brackish basin and rely only on natural food (Costa Pierce 2002). Therefore,
148 aquaculture capacity is represented by the potential biomass hosted by the brackish basins of each *valle da*
149 *pesca* (kg ha⁻¹ y⁻¹), estimated from a spatialized food-web model (Anelli Monti et al. 2021). The flow is expressed
150 as the average fish catches per hectare of brackish water surfaces per year (kg ha⁻¹ y⁻¹), according to the 2010-
151 2019 official data (Regione Veneto).

152 The waterfowl hunting was evaluated in terms of capacity by using the time series of waterbird censuses
153 from 2010 to 2019 (Associazione Faunisti Veneti). Since the census data were associated with point features,
154 we interpolated the average number of huntable birds censused in the last ten year by Dirichlet-Voronoi
155 tessellation for implicit surfaces to obtain the most likely spatial distribution of huntable waterbirds per
156 hectare per year. Hunting flow was obtained from waterbird catch historical series, collected from 2010 to
157 2020 (Ente Produttori Selvaggina Veneto).

158 The food production ES focused on wild edible plants of the genus *Salicornia* growing in saltmarshes and
159 on honey obtained from flowers of sea lavender (plants of the genus *Limonium*).

160 Perennial saltmarsh vegetation dominated by halophytic dwarf shrubs has been identified by visual census
161 method in 12 patches of two different *valli da pesca* (Valle Dogà and Valle Cavallino, Figure 1 - 1, 8), and then
162 through the identification of their predictive range in R, G, B bands reflectance values, filtered through a cut-
163 off NDVI value calculated from aerial and satellite images (listed in Supplementary Materials II Tabs. S.M. II
164 - 1, 2). The identified vegetation patches allowed for the evaluation of the capacity for edible plants, in terms
165 of kilograms of *Salicornia* biomass that could be harvested in the *valli da pesca* per year. Considering the
166 *Limonium* inflorescence cover in the patches (Fantinato and Buffa 2019) and the ratio between the number of
167 sea lavender flowers and grams of honey potentially produced per unit area, the honey capacity was assessed
168 as kilograms of honey that can be produced in the *valli da pesca*. The flow indicator refers to the kilograms of
169 harvested plants and honey. Edible plants harvesting data were reported by 2020 market data (Veneto
170 Agricoltura), by interviewed local people and restaurant chefs, while the amount of sea lavender honey was
171 witnessed by 5 beekeepers.

172 2.2.3 Cultural ESs

173 Tourism attractiveness was assessed by a survey carried out in the second half of 2019, addressing tourists who
174 recreate and travel in the Venice lagoon.

175 To assess the tourism ES capacity, we mapped for each one of the *valli da pesca* the attractiveness factors
176 evaluated by tourists in the questionnaires, namely saltmarshes presence, the possibility to observe birds and
177 fauna, good water quality, and the chance to contemplate natural terrestrial habitats. Each element was
178 weighted to depict the interest declared by the people and normalized to a 0-1 scale index.

179 Tourism flow was represented by the number of persons who, during a year, had passed at least one night
180 in one of the accommodation facilities within the *valli da pesca*.

181 Regarding the ES information for cognitive development, the natural factors that enhance touristic
182 attractiveness have been mapped along with the inclusivity toward the necessities of people with disabilities.
183 The resulting map was normalized to a 0-1 scale. The flow indicator summarized data concerning the number
184 of persons who annually attend outdoor educational activities or guided one-day trips, as reported by touristic
185 guides and associations (ATN Laguna Sud, Cooperativa Limosa, Ente di promozione turistica di Cavallino
186 Treporti, Oasi WWF Valle Averno).

187 Birdwatching ES capacity was expressed with the birdwatching attractiveness normalized map, based on
188 the factors reported as important by 30 interviewed birdwatchers: the presence of pedestrian paths,
189 saltmarshes presence, nesting areas in the visual field, and a high probability of observing birds. All these
190 factors were summed and scaled to obtain a normalized attractiveness map. Birdwatching ES flow indicator
191 was considered the mean number of active birdwatchers derived from the observers' activity trend, recorded
192 from 2010 to 2020 in the Italian birdwatchers' database (<https://www.ornitho.it/>).

193 *2.2.4 Aggregated ESs indicators*

194 Each ES indicator was spatially assessed in the *valli da pesca* and in the open lagoon, that is all the lagoon
195 surface which is not privately managed. The open lagoon ESs assessment was conducted accordingly with the
196 one presented by Rova, Stocco and Pranovi (2022), and detailed in Supplementary Materials III. Once we
197 obtained the raster layer for each indicator, for both the *valli da pesca* and the open lagoon, the results were
198 normalized to a 0-1 scale and finally aggregated to have a normalized sum of all the ESs capacities and a
199 normalized sum of all the ESs flows, allowing quantitative spatial comparisons. In particular, we ran the
200 algebraic sum of the normalized raster values through the core zonal statistics plugin in QGIS, by overlapping
201 the polygons of the areas of interest on each ES normalized raster. The results were compared between the
202 *valli da pesca* and the non-managed part of the lagoon and expressed as percentage of the overall capacity and
203 flow, as well as the percentage of each ES capacity and flow of the *valli da pesca* with respect to the related
204 values found in the open lagoon. All the analyses were performed using the software R 4.1.2 (R Core team,
205 2021) within the RStudio 2021.09.2 environment (RStudio Team, 2021).

206 **2.3 ESs spatialization**

207 To assess all the considered ESs with a spatially explicit approach, remote sensing imageries of the study area
208 were collected and analyzed to obtain the land cover/land use map (LULC) on which ES mapping was based.

209 To do so, a machine-learning, scalable gradient-boosting decision trees XgBoost algorithm (Chen and
210 Guestrin 2016) was used to classify land cover classes from multispectral and hyperspectral remote sensing
211 data. Very High-Resolution multi-spectral satellite scenes (VHR), collected by Worldview-02, Worldview-03,
212 and GeoEye-01 constellations, were granted by the European Space Agency Services via the "ESA On-Demand
213 Restricted Data Access program" (see Supplementary Materials II for details).

214 For some uncovered areas, satellite tiles were pan-sharpened and mosaicked in a custom composition of
 215 RGB + NIR raster stack, along with AGEA aerial photograms at 20 cm resolution, granted by Regione Veneto.

216 The classification was performed on the multi-band images through the tuning of the XgBoost algorithm
 217 and was assessed with an accuracy test, resulting in a mean of 96% correct class prediction.

218 The classification and the following spatialization tasks have been performed with the open source software
 219 QGIS 3.16 (QGIS Association: QGIS Geographic Information System, 2022), R 4.1.2 (R Core, 2022), and
 220 RStudio 2021.09.2 (RStudio Team, 2022).

221 3 Results

222 The assessment results are summarized in Table 2. The spatial distribution of each ES capacity and flow are
 223 illustrated in Supplementary Materials IV.

Ecosystem service	Measure unit	Capacity		Flow		
		Mean ± s.d.	Values range	Measure unit	Mean ± s.d.	Values range
Climate regulation	gC m ⁻² y ⁻¹	60.86 ± 54.60	10.00 ÷ 245.00	gC m ⁻² y ⁻¹	60.86 ± 54.60	10.00 ÷ 245.00
Water purification	% removed nitrogen	9.37 ± 11.67	0.00 ÷ 34.61	% removed nitrogen	9.37 ± 11.67	0.00 ÷ 34.61
Lifecycle support	0-1 scale	0.58 ± 0.23	0.30 ÷ 0.73	0-1 scale	0.31 ± 0.24	0.00 ÷ 1.00
Aquaculture	kg ha ⁻¹ y ⁻¹	60.79 ± 59.99	0.00 ÷ 119.92	kg ha ⁻¹ y ⁻¹	29.83 ± 34.77	0.00 ÷ 159.13
Hunting	n ha ⁻¹ y ⁻¹	29.56 ± 36.56	0.02 ÷ 376.67	n ha ⁻¹ y ⁻¹	6.12 ± 4.77	0.00 ÷ 20.94
Wild hedible herbs & honey production	edible plants kg ha ⁻¹ y ⁻¹	0.20 ± 0.37	0.00 ÷ 2.03	edible plants kg ha ⁻¹ y ⁻¹	0.00 ± 0.00	0.00 ÷ 0.00
	honey kg ha ⁻¹ y ⁻¹	0.11 ± 0.01	0.00 ÷ 0.11	honey kg ha ⁻¹ y ⁻¹	0.03 ± 0.10	0.00 ÷ 0.45
Tourism	attractiveness, 0-1 scale	0.49 ± 0.14	0.06 ÷ 1.00	tourists n y ⁻¹	7.86 ± 34.07	0.00 ÷ 20.00
Information for cognitive development	attractiveness, 0-1 scale	0.30 ± 0.12	0.04 ÷ 0.89	students n y ⁻¹	102 ± 470	0.00 ÷ 3640.0
Birdwatching capacity	attractiveness, 0-1 scale	0.33 ± 0.23	0.00 ÷ 1.00	birdwatchers n y ⁻¹	1.47 ± 5.99	0.00 ÷ 38.67

224 **Table 2** Capacity and flow results in the valli da pesca.

225 Overall, in the *valli da pesca* the highest values for carbon sequestration were detected in the areas where
 226 both saltmarshes and seagrasses are present, reaching values up to 245 gCm⁻²y⁻¹. The mean carbon
 227 sequestration has been estimated at 60.86±54.60 gCm⁻²y⁻¹, for a total amount of 2448 tonC y⁻¹, or 8982
 228 tons of CO₂y⁻¹ (Supplementary Materials IV, a). The water purification ES is represented by an average removal
 229 of 9.37% (±11.67%) of the nitrogen loadings; the greater the extent of the brackish lakes in the *valli*, the higher
 230 results the removal capacity. However, it must be noticed that this average estimate may likely be influenced
 231 by the water quality of the river water entering the *valli da pesca* (Salveti et al. 2008; Bettiol et al. 2005) and
 232 must be considered as a first estimate of the average value for this ES.

233 As regards the lifecycle support, the capacity indicator is homogenously distributed with a mean of 0.58
 234 (±0.23), on a 0-1 scale, while the flow indicator shows a mean of 0.31±0.24 (Supplementary Materials IV, c-
 235 d).

236 The aquaculture capacity of the study area results in a mean of 60.79 ± 59.99 kg per hectare per year, but
237 the flow of this ES varies greatly between different *valli da pesca* (Supplementary Materials IV, f) because it is
238 influenced by several factors, both natural and social. Fish production is related to fish sowing, which in turn
239 depends on wild fry availability; moreover, fish sowing is dependent on both fish survival to the previous years
240 and precedent production. Based on the 2014-2019 period, the average total quantity of juveniles sown per
241 year was 6'867'213, with a value per *valle da pesca* ranging between 26'000 and 2'793'033 individuals. On this
242 basis, the mean sowing value was $1'227 (\pm 958)$ juveniles per hectare per year. Thus, when faced with an average
243 production of the whole area of $29.83 (\pm 34.77)$ kg ha⁻¹ y⁻¹, we must distinguish that the *valli da pesca* carrying
244 out semi-intensive aquaculture show a value of 70.55 ± 23.78 kg ha⁻¹ y⁻¹ (e.g., *valli da pesca* n. 2, 22, 24) while
245 those practicing only extensive aquaculture drop to a mean of 15.45 ± 16.35 kg ha⁻¹ y⁻¹.

246 Regarding waterfowl hunting, during the period 2010-2019, about 280'000 huntable waterfowls have been
247 wintering inside the *valli da pesca*, where a yearly average of 29.56 ± 36.56 huntable waterfowls per hectare
248 were hosted. The highest number of waterfowls was consistently recorded during 2010-2019 in the *valli da*
249 *pesca* n. 1, 7, 8, 19, 25, 26, and 27, while in the *valli da pesca* from n. 10 to n. 15, most exposed to nautical and
250 tourist traffic, the number of censused waterfowls was lower (Supplementary Materials IV, g).

251 Regarding the hunting flow, in 2010-2019 the average catch was estimated at 38'404 (± 7299) waterfowls per
252 year, considering the total surface of the *valli da pesca*. On average, 6.12 ± 4.77 waterbirds ha⁻¹ y⁻¹ have been
253 caught in the considered period. These catches were homogeneously distributed, with a maximum value of 20
254 catches ha⁻¹ y⁻¹ in the most confined *valli da pesca* 3, 4, 5, 6, 26, 27 (Supplementary Materials IV, h).

255 Although a homogeneous potential for provisioning edible wild plants and honey has been estimated in the
256 *valli* area, harvesting occurs almost exclusively in the saltmarshes located in the open part of the lagoon, except
257 from three *valli da pesca* – n. 5, 6, 8 – which host beehives (Supplementary Materials IV, j). Harvesting of
258 *Salicornia* and other edible herbs has not been reported in the *valli da pesca* saltmarshes, apart from tiny
259 quantities in occasional occurrences.

260 Tourism capacity results in all the areas with a mean attractiveness of 0.49 ± 0.14 (Supplementary Materials
261 IV, k); nevertheless, very few of them are open for touristic trips, and not even regularly, which restricts the
262 flow for tourism as well as for other cultural ESs (Supplementary Materials IV, l). The flow of tourists is limited
263 to few *valli da pesca* in the Northern lagoon, involving Valle Lio Maggiore (5), Valle Cavallino (8), Valle
264 Falconera (9), Valle Sacchetta (15), and Santa Cristina (17).

265 The capacity and flow for the ES information for cognitive development cover an even narrower extent:
266 only the WWF Oasis Valle Averno (20), Valle Olivara (11), and Valle Liona (12) have a capacity higher than 0.5
267 (Supplementary Materials IV, m), but it resulted that the 3'154 students per year, who visit the *valli da pesca*
268 for educational purposes, take part into guided tours in *valli da pesca* n. 12, 20, 23, 24 (Supplementary
269 Materials IV, l).

270 Finally, the overall attractiveness for birdwatching is evenly distributed (Supplementary Materials IV, o)
271 but the ES flow is not homogeneous and confirms also for this cultural ES that the Northern part of the lagoon
272 is the most frequented for such outdoor activities, with 72 active birdwatchers – who spend a mean of $84.41 \pm$
273 21.36 hours per year birdwatching – converging in the nearing of the area n. 5, 11, 12 (Supplementary Materials
274 IV, p).

275 Table 3 shows how much the total ESs capacity and flow assessed in all the *valli da pesca* is comparable to
276 that observed in the open, not privately managed part of the lagoon. The most important contributions from

277 the *valli da pesca* to the ESs supply of the open lagoon are noticeable for lifecycle support, aquaculture,
 278 hunting, information for cognitive development, and birdwatching.
 279

<i>Ecosystem service</i>	<i>% vs. lagoon capacity</i>	<i>% vs. lagoon flow</i>
<i>Climate regulation</i>	12.79	12.79
<i>Water purification</i>	12.50	12.50
<i>Lifecycle support</i>	23.93	25.78
<i>Aquaculture</i>	12.79	31.68
<i>Hunting</i>	99.8	65.76
<i>Wild hedible herbs</i>	6.46	0.00
<i>Honey production</i>	83.48	1.82
<i>Tourism</i>	27.22	0.01
<i>Information for cognitive development</i>	25.97	23.88
<i>Birdwatching</i>	21.57	13.66

Table 3 Proportion of capacity and flow assessed in the *valli da pesca* compared to each ES capacity and flow in the open lagoon.

280 3.4 Aggregated ESs indexes analyses

281 Figure 2 and figure 3 show the normalized maps for the aggregated indicators of all the ESs capacity and the
 282 flow intensities, respectively. The sum of all the capacities shows that the highest values have been registered
 283 inside the *valli da pesca* which maximize both aquaculture and hunting (Fig. 2 n. 1, 2, 4, 26, 27).

284 As regards the sum of all the flows, the highest values have been recorded especially where provisioning
 285 and cultural ESs flows are higher, e.g., Valle Grassabò, Valle Lio Maggiore, and Valle Dragojesolo (Fig. 3 n. 2,
 286 5, 7).





Figure 2 Spatialized total capacity indicator, as the sum of all the ESs' capacities

287
 288
 289



290

291


Figure 3 Spatialized total flow indicator, as the sum of all the ESs' flows

292

293

294

In general, the extension of the area where the sum of all the ESs' capacity is higher than the sum of all the ESs' flow is 71.1 km²; on the contrary, the area where the sum of all the assessed ESs' flow results greater than the sum of all the ESs' capacity is 23.4 km² (Fig. 4)



295

296

Figure 4 Spatialized capacity – flow difference (C-F). Green = $C > F$; red = $C < F$.

297

298

Discussion

299

300

301

302

303

304

305

306

The lagoon of Venice is a complex social-ecological system with a historical link between natural habitats and human activities, which is particularly strong for the *valli da pesca*. Despite being “artificially maintained”, the *valli da pesca* conserves the typical features and the related ecological processes of transitional environments that the open lagoon has progressively lost, especially due to erosion and relative sea level rise (Day et al. 1998; Madricardo and Donnici 2014). For these reasons, it would be interesting to analyze the possible role of the *valli da pesca* in terms of ESs.

The ESs approach has already proven to be effective in visualizing and assessing the contribution to people of similar man-managed ecosystems, and in suggesting sustainable management strategies as well (Gaglio et

307 al. 2019; Aschonitis et al. 2016; Weitzman 2019; Walton et al. 2015). In this study, we proposed for the first
308 time the assessment of multiple ESs in the *valli da pesca* of the Venice lagoon to better understand their
309 possible ecological role within the context of the entire lagoon environment, highlighting a non-negligible
310 contribution of the *valli da pesca* to the entire supply of ESs of the Venice lagoon.

311 With reference to carbon sequestration, for instance, about 12% of the CO₂ sequestered every year by the
312 lagoon is due to saltmarshes and seagrasses located inside the *valli da pesca*, where these habitats are
313 preserved by different impacts, erosion, and trampling. Also, the mean water purification through
314 denitrification is higher inside the boundaries of the *valli da pesca* than in the outer part of the lagoon. Indeed,
315 it has been estimated that the open lagoon can remove about 10.08% of nitrogen loadings (Rova et al. 2022),
316 but no area reaches the maximum values recorded in some lakes located inside the *valli da pesca*. These
317 findings are in accordance with those of other studies that have focused on water management for extensive
318 aquaculture (Walton, Vilas, Coccia, et al. 2015; Gamito 1997). This result, however, should inspire further
319 research because the denitrification process is influenced by several factors, including the water quality of the
320 river entering the *valli da pesca*, and the organic matter loading that can increase in response to eventual
321 higher densities of fish and waterbirds inside the lakes. Today, extensive aquaculture is prevalent in *the valli*
322 *da pesca*. Therefore, fish feed on the available food they naturally find in the *valle da pesca*, without the need
323 to artificially add feed, and the density of fish is such that the brackish lake ecological processes can process
324 naturally dead organic matter and nutrient loadings, avoiding the risk of eutrophication (Anras et al. 2010;
325 Costa Pierce 2002). Thus, the denitrification process is worthy of further experimental field studies. New data
326 and models about this aspect, along with an exploration of the most likely future trends, may prove extremely
327 useful to better depict this regulating ES not only at the local scale but also for other similar Heavily Modified
328 Water Bodies, such as in the Po River Delta, in the Marano lagoon or in other areas along the Adriatic coasts.

329 The most confined parts of the lagoon were identified centuries ago as the most suitable places for fish
330 farming and hunting purposes, as confirmed more recently for fish (Cavraro et al. 2017; Brigolin et al. 2014)
331 and waterfowl (Arzel et al., 2006; Scarton and Bon, 2009; Guillemain et al., 2013). Within this context, the ES
332 lifecycle support provided by the *valli da pesca* turned out to be quite significant compared to the open lagoon,
333 confirming the important ecological role still played by these areas despite the presence of levees and
334 embankments. However, the naturally recruited fish is not sufficient for meeting the *valli* requirements, to the
335 point that today the aquaculture ES in the *valli da pesca* of the Venice lagoon depends on personal, social, and
336 economic factors, which lead the owner to decide whether and how much fish to sow. As a consequence, the
337 practice of fishing wild fry in the lagoon and then sowing them inside the *valli da pesca* is the only method that
338 grants the fraction of the biomass actually hosted by each area. From such a point of view, this tradition,
339 besides enhancing the aquaculture productivity, could be seen as a good practice, capable to maintain in the
340 long run the presence of some species. On one hand, fry fishery is strictly regulated and controlled, being
341 entrusted to only a very few licensed operators who annually update the quota that can be caught in order not
342 to affect lagoon fish population dynamics (Fortibuoni et al. 2014; Granzotto et al. 2001).

343 On the other hand, while the fish migration and the recruitment processes taking place outside in the lagoon
344 plays a fundamental role for sustaining fish lifecycle (Lanzoni et al. 2021; Scapin et al. 2022; Cavraro et al.
345 2017) and the aquaculture ES in the *valli da pesca*, we could in turn affirm that the aquaculture ES in the *valli*
346 *da pesca* could positively affect the regulating ES of lifecycle support in the entire lagoon. As happens for
347 waterbirds, indeed, the persistency of suitable habitats in the *valli da pesca* depends on human maintenance,
348 which in turn depend on economic resources that come from the provisioning ESs incomes. This generates

349 positive feedback that rises considerations about how a provisioning ES can support also other ESs belonging
350 to a different category, in agreement with the findings of other authors (Liquete et al. 2016; Grizzetti et al.
351 2019)

352 The extensive aquaculture practiced today in the *valli da pesca*, moreover, can be considered quite sustainable,
353 considering also that the production value of 60 kg ha⁻¹y⁻¹ falls in the range of 45 - 110 kg ha⁻¹y⁻¹ reported by
354 (Koutrakis et al. 2007) for extensive aquaculture in Mediterranean lagoon, but is quite lower than 150 kg ha⁻¹
355 y⁻¹ expected for a desirable production in a typical Italian *valle da pesca*, according to Ravagnan (1982) and
356 Shang (1982).

357 Since the aquaculture ES in the *valli da pesca* is getting less worthwhile in the last decade, as shown by the
358 data, other activities have started to play a key role. Among them, the most important is waterfowls hunting,
359 which also depends both on natural and anthropogenic factors. To say to what extent the great capacity of bird
360 attraction is granted by the *valli da pesca* environment itself, or whether it is due to management choices, is
361 difficult. To attract waterfowl and increase hunting opportunities, indeed, the managers do regulate water
362 regimes, construct mild sloping islands, take care of windbreak hedges made of reeds and tamarisks, and often
363 spread millet or grains.

364 The management which aims to maximize hunting ES is possibly influencing the capacity and the flow of
365 this ES not only inside, but also outside the boundaries of the *valli da pesca*. Our assessments found that most
366 huntable waterfowl censused in the lagoon are reaching for the *valli da pesca* during their migration, where
367 they rest, feed, and breed as well (at least some species). Such evidence is in accordance with other authors,
368 who reported about the high importance of the *valli da pesca* for waterbird populations, whereas the lagoon is
369 losing suitable habitats (Scarton 2017; AFV 2020). Also, the waterfowls hunting flow in the lagoon results
370 strongly related, for timing and catches as well, to the flow happening inside the *valli*. As a comparison, the
371 hunting flow in the lagoon is estimated to be in the range from 12'635 to 12'770 catches per year and takes
372 place just in proximity of the allowed hunting blinds, where the catches have a mean of 4.44 ± 1.56 per hectare
373 per year. Even if the distribution of this flow is homogeneous in the whole lagoon, the highest values have been
374 detected in the blinds between contiguous *valli da pesca*, in the North-Eastern as well as in the Southern part
375 of the lagoon. Indeed, waterfowl hosted inside the *valli da pesca*, while moving across different *valli da pesca*
376 or from a *valle da pesca* to another place just outside, represent the "stock" which not only the *valli da pesca*
377 hunters, but also the "open lagoon" hunters rely on.

378 Another example of driving relationships between different ESs is the substantial contributions of the *valli da*
379 *pesca* to the ESs that depend on the consistency of the birds' presence. For example, the management which
380 aims to attract the highest possible number of huntable waterfowls is possibly also dragging the attractiveness
381 for cultural ESs, especially for birdwatching. Accordingly, many aquatic birds are attracted into the *valli da*
382 *pesca* managed environment and this also can increase the overall capacity, as bird movements between the
383 *valli da pesca* and the open lagoon create a positive side-effect in increasing attractiveness to birdwatchers,
384 tourists, and hikers even outside the *valli da pesca* boundaries.

385 However, despite the high overall capacity for all cultural ESs in the *valli da pesca*, the flow is strongly shaped
386 by accessibility and inclusiveness, ending up with flow occurring just on a few areas. Anyway, this restricted
387 flow should not be read as totally negative: the *valli da pesca* are seen as extremely attractive especially for the
388 preservation of lagoon landscapes and fauna in a quiet and uncrowded space. This consideration should make
389 one think carefully about whether, how, and how often the *valli da pesca* could be open for touristic and
390 educational visits. Similar considerations have been aroused also on different environments by other authors

391 (Villamagna et al., 2015; Drescher et al., 2017; Drescher and Brenner, 2018), who proposed to give more
392 support to private natural areas for enhancing conservation and educational purposes, provided that
393 excursionists' entrances must be controlled.

394 Management initiatives that also take these perspectives into account should seek to harmonize with the
395 current situation of the *valli da pesca*, seen that, to date, the difference between capacity and flow aggregated
396 indexes shows that the majority of the *valli da pesca* display a total capacity higher than the total flow (as it
397 can be seen in figs. 2, 3, 4). Just a few zones with the highest flows of mediated ESs (*sensu* Rova and Pranovi
398 2017) stand out amongst other areas: in fact, some *valli da pesca* where hunting ES' flow is on average more
399 important than the aquaculture production display an aggregated flow index slightly higher than the ESs'
400 capacity. This may be because waterfowls hunting relies on a moving resource so that the catches in one *valle*
401 *da pesca* can affect the waterfowls which have been censused in another *valle* and, consequently, represent the
402 capacity assessed on a different place, but not strictly linked to it.

403 The analysis of the contribution of the *valli da pesca* sub-system to the whole transitional ecosystem confirms
404 that the *valli da pesca* are highly significant to the whole Venice lagoon. The comparison of the ESs' aggregated
405 index between the open part of the lagoon and the *valli* areas brings out that the *valli da pesca*, despite covering
406 just 17% of the total lagoon, play an important role in terms of ESs contributions: 38% of the capacity and 24%
407 of the flow in comparison to the open lagoon capacity and flow, respectively.

408 This non-negligible contribution to ESs, habitats, and landscape features conservation, makes the *valli da*
409 *pesca* a good example of effective ecosystem-based management of Heavily Modified Water Bodies, capable to
410 represent a good example to set up a sustainable management for similar productive areas in the
411 Mediterranean.

412 Like little lagoons that are artificially managed, the *valli da pesca* are regulated by a private regime that, on
413 one hand, denies the access and minimizes cultural ESs flow, but on the other hand, conserves the landscape,
414 the ecological processes and the ESs provided by them, to the point that the open lagoon receives the benefits
415 of a spill-over effect especially for lifecycles support, aquaculture, hunting, and birdwatching ESs. For these
416 ESs, the *valli da pesca* provide in fact ESs that give benefits to a wider community, spread into a larger spatial
417 scale than that the ES are generated at. In addition, the *valli da pesca* act as a buffer area between the land and
418 the open lagoon, in particular for climate regulation and water purification, as the results about carbon
419 sequestration and nitrogen removal show.

420 The role of *valli da pesca* as buffers and as conservation opportunities leads also to consider how important
421 it might be to restore a buffer strip between the land and the lagoon, which has been lost in some areas. As the
422 *valli da pesca* prove to be effective in maintaining saltmarshes, reed meadows and ecological gradients,
423 similarly it might be useful to consider ecological restoration efforts in areas that are most affected by the
424 transfiguration of the original lagoon landscape (Feola et al., 2022).

425 Since the connection between lagoon and *valli da pesca* persists, albeit in different ways and intensities
426 depending on the processes being considered, we must take into account the possible evolution of such
427 interconnected systems as a whole. The lagoon, indeed, is now subjected to manifold pressures from anthropic
428 activities (Fortibuoni et al. 2015; Zucchetta et al. 2021; Anelli Monti et al. 2021; Solidoro et al. 2010) that are
429 likely going to increase (Jennerjahn and Mitchell 2013). The adaptations to such impacts, as well as to climate
430 change and sea-level rise effects, in the long run might modify the lagoon hydrodynamics and the sea-lagoon
431 connectivity, threatening the *valli da pesca* and so their ESs supply (Solidoro et al. 2010; Cristiano and Gonella
432 2020).

433 Eventually, we suggest that decisions about the MOSE barrier functioning must be well pondered in order
434 not to put the *valli da pesca* at risk of being endangered. Considering that the *valli da pesca* are definitely
435 continuing to exchange energy and matter with the lagoon, such equilibria can be at risk in the context of a
436 “regulated lagoon”, where the MOSE system (with its mobile barriers at the inlet) controls the marine water
437 flow to mitigating the effect of climate change and sea-level rise on cultural heritage, but could conversely
438 threaten the daily exchange of water in the most confined part of the lagoon, and the *valli* waterfronts as a
439 consequence.

440 Indeed, it is very important to preserve the cultural heritage of the historic center of Venice and the islands so
441 as not to lose cultural ESs and economic activities flowing from them. Nevertheless, we are facing the
442 possibility that the lagoon will remain closed for longer periods of time. According to Umgiesser (2020) and
443 Lionello et al. (2021), the period of closure of the mobile barriers is very likely to grow up to 3 weeks per year
444 by the end of 2050, and up to 2 months per year by 2080. This would mean affecting the sea-lagoon
445 connectivity, the lagoon hydrodynamics, and landscape evolution patterns, as several studies have already
446 argued (Ghezzi et al. 2010; 2011; Pérez-Ruzafa et al. 2019). Such a perspective could put the *valli da pesca* at
447 risk, firstly because a lower connectivity with the sea can affect the fish fry availability for the aquaculture ES
448 even more than today, and because a lower frequency of exchanges through the inlets is expected to worsen
449 the water quality, nutrient concentration (Solidoro et al. 2005), and microcirculation (Ghezzi et al. 2010)
450 involved in the processes on which the regulating ESs are based.

451 In light of our results, it must be carefully considered that losing the *valli da pesca* might represent an issue
452 for the entire community because of their contribution to conserving habitats and providing regulating,
453 provisioning and cultural ESs. This trade-off perspective takes great importance, especially in the light of future
454 scenarios including climate change, subsidence, and management of health problems, which claim a
455 responsible attendance of the lagoon.

456 Conclusions

457 This work has shown, by using quantitative indicators, the spatial distribution of ESs capacity and flow that
458 take place within the managed *valli da pesca* of the lagoon of Venice. The multiple ESs approach enhanced the
459 areas where capacity and flow have the highest values but has also revealed that, due to different accessibility
460 and availability of some places, the capacity is not always and everywhere fully granted to translate itself in a
461 flow towards the local population. The comparison between different areas of the same wide ecosystem
462 highlights that, although the capacity of the provided ESs tends to be overall high, the relative flows are
463 influenced by a multiplicity of factors, both social and behavioral.

464 In this context, confirmation of the usefulness of assessing the value of ESs in ecosystem management is
465 given (Costanza 2006), especially when a trade-off must be accomplished between the moral duty to preserve
466 and emphasize regulation and support services, and the need to build on the economic revenues that come
467 from provisioning ESs. This dynamical network stimulates to continue the research with a modeling approach,
468 which could consider behavioral, social, and economic factors that act as fulcrums for decision-making.

469 Increasing the knowledge about the *valli da pesca* and other similar socio-ecological systems can help to
470 enlighten not only the relationships between ESs, landscape, and management choices, but also to rethink the
471 interaction between the environment, the society, and the institutions involved in the Venice lagoon ecosystem.

472 The assessment of ESs, such as lifecycle support, hunting, and birdwatching, showed that the open side of
473 the lagoon is often strictly related to the management of the most confined areas, corroborating the hypothesis
474 that the *valli da pesca* play an important role in defining the ESs pattern of the whole lagoon of Venice. Indeed,
475 the *valli da pesca* represent a buffer between the aquatic ecosystem of the lagoon and its land boundaries,
476 mitigating anthropogenic pressures; in addition, their provisioning and cultural ESs also show a series of
477 positive side-effects in the open lagoon.

478 Thus, it is of great importance to take them into account to have a full comprehension of the Venice lagoon
479 ecosystem, especially when investigating the effects under different health, social and economic scenarios, as
480 well as when facing climate change and sea-level rise.

481 Acknowledgments

482 This scientific work has been developed in the context of the “Venezia 2021” project, with the contribution of
483 the Provveditorato Interregionale Opere Pubbliche per il Veneto, Trentino Alto Adige e Friuli Venezia Giulia
484 through the Consorzio Venezia Nuova and coordinated by CORILA.

485 We acknowledge the owners and the operators of the *valli da pesca* for providing data and explanations, as
486 well as the Regione Veneto, the Associazione Faunisti Veneti, the EPS, and the local hunting associations.

487 We thank the European Space Agency for granting the VHR satellite imageries for this research work.

488 We thank Marco Anelli Monti for providing the food web model data, and Matteo Zucchetta and Alberto Caccin
489 for their helpful suggestions.

490 We are grateful to the reviewers for their kind comments, which helped to improve the manuscript.

- 492 AFV, Associazione Faunisti Veneti. 2020. 'CENSIMENTO DEGLI UCCELLI ACQUATICI SVERNANTI IN
493 PROVINCIA DI VENEZIA'.
- 494 Anelli Monti, M., D. Brigolin, P. Franzoi, S. Libralato, R. Pastres, C. Solidoro, M. Zucchetta, and F. Pranovi. 2021a.
495 'Ecosystem Functioning and Ecological Status in the Venice Lagoon, Which Relationships?' *Ecological Indicators*
496 133 (December): 108461. <https://doi.org/10.1016/j.ecolind.2021.108461>.
- 497 Anras, L., Clara Boglione, Stefano Cataudella, M. T. Dinis, Pavlos Makridis, Giovanna Marino, Ana Ramalho Ribeiro,
498 and Manuel Yúfera. 2010. 'The Current Status of Extensive and Semi-Intensive Aquaculture Practices in Southern
499 Europe'. *Aquaculture Europe* 35 (2): 12–16. <https://digital.csic.es/handle/10261/50633>.
- 500 Arzel, Celine, J. Elmberg, and M. Guillemain. 2006. 'Ecology of Spring-Migrating Anatidae: A Review'. *Journal of*
501 *Ornithology*. Springer. <https://doi.org/10.1007/s10336-006-0054-8>.
- 502 Aschonitis, Vassilis G., Mattias Gaglio, Elisa A. Fano, Elena Gissi, Giuseppe Castaldelli, Vassilis G. Aschonitis, Elena
503 Gissi, Giuseppe Castaldelli, and Elisa A. Fano. 2016. 'Land Use Change Effects on Ecosystem Services of River
504 Deltas and Coastal Wetlands: Case Study in Volano–Mesola–Goro in Po River Delta (Italy)'. *Wetlands Ecology*
505 *and Management*. <https://doi.org/10.1007/s11273-016-9503-1>.
- 506 Baró, Francesc, Ignacio Palomo, Grazia Zulian, Pilar Vizcaino, Dagmar Haase, and Erik Gómez-Baggethun. 2016.
507 'Mapping Ecosystem Service Capacity, Flow and Demand for Landscape and Urban Planning: A Case Study in the
508 Barcelona Metropolitan Region'. *Land Use Policy* 57 (November): 405–17.
509 <https://doi.org/10.1016/j.landusepol.2016.06.006>.
- 510 Bettiol, Cinzia, Flaviano Collavini, Stefano Guerzoni, Emanuela Molinaroli, Paolo Rossini, Luca Zaggia, and Roberto
511 Zonta. 2005. 'Atmospheric and Riverine Inputs of Metals, Nutrients and Persistent Organic Pollutants into the
512 Lagoon of Venice'. *Hydrobiologia* 2005 550:1 550 (1): 151–65. <https://doi.org/10.1007/S10750-005-4372-2>.
- 513 Brigolin, Daniele, Chiara Facca, Anita Franco, Piero Franzoi, Roberto Pastres, Adriano Sfriso, Marco Sigovini, et al.
514 2014. 'Linking Food Web Functioning and Habitat Diversity for an Ecosystem Based Management: A
515 Mediterranean Lagoon Case-Study'. *Marine Environmental Research* 97: 58–66.
516 <https://doi.org/10.1016/j.marenvres.2014.02.006>.
- 517 Burkhard, Benjamin, and Joachim Maes. 2017. 'Mapping Ecosystem Services'. *Advanced Books* 1 (March): e12837.
518 <https://doi.org/10.3897/AB.E12837>.
- 519 Cavraro, Francesco, Matteo Zucchetta, Stefano Malavasi, and Piero Franzoi. 2017. 'Small Creeks in a Big Lagoon: The
520 Importance of Marginal Habitats for Fish Populations'. *Ecological Engineering* 99 (February): 228–37.
521 <https://doi.org/10.1016/j.ecoleng.2016.11.045>.
- 522 Chen, Tianqi, and Carlos Guestrin. 2016. 'XGBoost'. In *Proceedings of the 22nd ACM SIGKDD International*
523 *Conference on Knowledge Discovery and Data Mining*, 13-17-Aug:785–94. New York, NY, USA: ACM.
524 <https://doi.org/10.1145/2939672.2939785>.
- 525 Costa Pierce, Barry A. 2002. 'Ecological Aquaculture'. *Ecological Aquaculture*, April.
526 <https://doi.org/10.1002/9780470995051>.
- 527 Costanza, Robert. 2006. 'Nature: Ecosystems without Commodifying Them'. *Nature* 2006 443:7113 443 (7113): 749–
528 749. <https://doi.org/10.1038/443749b>.
- 529 Costanza, Robert, Ralph D'Arge, Rudolf de Groot, Stephen Farber, Monica Grasso, Bruce Hannon, Karin Limburg, et al.
530 1997. 'The Value of the World's Ecosystem Services and Natural Capital'. *Nature* 387 (6630): 253–60.
531 <https://doi.org/10.1038/387253a0>.
- 532 Cristiano, Silvio, and Francesco Gonella. 2020. "'Kill Venice": A Systems Thinking Conceptualisation of Urban Life,
533 Economy, and Resilience in Tourist Cities'. *Humanities and Social Sciences Communications* 2020 7:1 7 (1): 1–
534 13. <https://doi.org/10.1057/s41599-020-00640-6>.
- 535 D'alpaos, Chiara, and Andrea D'alpaos. 2021. 'The Valuation of Ecosystem Services in the Venice Lagoon: A
536 Multicriteria Approach'. *Sustainability* 2021, Vol. 13, Page 9485 13 (17): 9485.
537 <https://doi.org/10.3390/SU13179485>.

538 Day, J. W., A. Rismondo, F. Scarton, D. Are, and G. Cecconi. 1998. 'Relative Sea Level Rise and Venice Lagoon
539 Wetlands'. *Journal of Coastal Conservation* 4 (1): 27–34. <https://doi.org/10.1007/BF02806486>.

540 European Commission. 2000. 'Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000
541 Establishing a Framework for Community Action in the Field of Water Policy'. *Official Journal of the European
542 Communities* L327.

543 Fantinato, Edy, and Gabriella Buffa. 2019. 'Animal-Mediated Interactions for Pollination in Saltmarsh Communities'.
544 *Plant Sociology* 56 (2): 35–42. <https://doi.org/10.7338/pls2019562/02>.

545 Feola, Alessandra, Emanuele Ponis, Michele Cornello, Rossella Boscolo Brusà, Federica Cacciatore, Federica Oselladore,
546 Bruno Matticchio, et al. 2022. 'An Integrated Approach for Evaluating the Restoration of the Salinity Gradient in
547 Transitional Waters: Monitoring and Numerical Modeling in the Life Lagoon Refresh Case Study'. *Environments*
548 *2022, Vol. 9, Page 31* 9 (3): 31. <https://doi.org/10.3390/ENVIRONMENTS9030031>.

549 Fortibuoni, T., R. Gertwagen, O. Giovanardi, and S. Raicevich. 2014. 'The Progressive Deregulation of Fishery
550 Management in the Venetian Lagoon after the Fall of the Repubblica Serenissima: Food for Thought on
551 Sustainability'. *Global Bioethics* 25 (1): 42–55. <https://doi.org/10.1080/11287462.2014.894707>.

552 Fortibuoni, Tomaso, Federica Aldighieri, Otello Giovanardi, Fabio Pranovi, and Matteo Zucchetta. 2015. 'Climate Impact
553 on Italian Fisheries (Mediterranean Sea)'. *Regional Environmental Change* 15 (5): 931–37.

554 Gaglio, Mattias, Mattia Lanzoni, Giovanni Nobili, Diego Viviani, Giuseppe Castaldelli, and Elisa Anna Fano. 2019.
555 'Ecosystem Services Approach for Sustainable Governance in a Brackish Water Lagoon Used for Aquaculture'.
556 *Journal of Environmental Planning and Management* 62 (9): 1501–24.
557 <https://doi.org/10.1080/09640568.2019.1581602>.

558 Gamito, S. 1997. 'Sustainable Management of a Coastal Lagoonal System (Ria Formosa, Portugal): An Ecological Model
559 for Extensive Aquaculture'. *International Journal of Salt Lake Research* 1997 6:2 6 (2): 145–73.
560 <https://doi.org/10.1007/BF02441891>.

561 Ghezzi, Michol, Stefano Guerzoni, Andrea Cucco, and Georg Umgiesser. 2010. 'Changes in Venice Lagoon Dynamics
562 Due to Construction of Mobile Barriers'. *Coastal Engineering* 57 (7): 694–708.
563 <https://doi.org/10.1016/J.COASTALENG.2010.02.009>.

564 Ghezzi, Michol, Alessandro Sarretta, Marco Sigovini, Stefano Guerzoni, Davide Tagliapietra, and Georg Umgiesser.
565 2011. 'Modeling the Inter-Annual Variability of Salinity in the Lagoon of Venice in Relation to the Water
566 Framework Directive Typologies'. *Ocean and Coastal Management* 54 (9): 706–19.
567 <https://doi.org/10.1016/j.ocecoaman.2011.06.007>.

568 Granzotto, A, P Franzoi, A Longo, and F Pranovi P Torricelli. 2001. 'La Pesca Nella Laguna Di Venezia : Un Percorso
569 Di Sostenibilità Nel Recupero Delle Tradizioni Lo Stato Dell ' Arte'.

570 Grizzetti, B., C. Liqueste, A. Pistocchi, O. Vigiak, G. Zulian, F. Bouraoui, A. de Roo, and A. C. Cardoso. 2019.
571 'Relationship between Ecological Condition and Ecosystem Services in European Rivers, Lakes and Coastal
572 Waters'. *Science of the Total Environment* 671 (June): 452–65.
573 <https://doi.org/10.1016/J.SCITOTENV.2019.03.155>.

574 Guillemain, Matthieu, Hannu Pöysä, Anthony D. Fox, Céline Arzel, Lisa Dessborn, Johan Ekroos, Gunnar Gunnarsson,
575 et al. 2013. 'Effects of Climate Change on European Ducks: What Do We Know and What Do We Need to Know?'
576 *Wildlife Biology* 19 (4): 404–19. <https://doi.org/10.2981/12-118>.

577 Hatziiordanou, Lena, Eleni Fitoka, Elena Hadjicharalampous, Nefta Eleftheria Votsi, Dimitris Palaskas, and Dania Abdul
578 Malak. 2019. 'Indicators for Mapping and Assessment of Ecosystem Condition and of the Ecosystem Service
579 Habitat Maintenance in Support of the EU Biodiversity Strategy to 2020'. *One Ecosystem* 4.
580 <https://doi.org/10.3897/oneeco.4.e32704>.

581 Havera, S. P., Lonie R. Boens, M. M. Georgi, and R. Shealy. 1992. 'HUMAN DISTURBANCE OF WATERFOWL ON
582 KEOKUK POOL'. *Undefined*.

583 Hein, Lars, Kris van Koppen, Rudolf S. de Groot, and Ekko C. van Ierland. 2006a. 'Spatial Scales, Stakeholders and the
584 Valuation of Ecosystem Services'. *Ecological Economics* 57 (2): 209–28.
585 <https://doi.org/10.1016/J.ECOLECON.2005.04.005>.

586 Jennerjahn, Tim C., and Steve B. Mitchell. 2013. 'Pressures, Stresses, Shocks and Trends in Estuarine Ecosystems – An
587 Introduction and Synthesis'. *Estuarine, Coastal and Shelf Science* 130 (September): 1–8.
588 <https://doi.org/10.1016/J.ECSS.2013.07.008>.

589 Korschgen, C., L. George, and W. L. Green. 1985. 'Disturbance of Diving Ducks by Boaters on a Migrational Staging
590 Area'. *Undefined*.

591 Koutrakis, E.T., A. Conides, A.C. Parpoura, E.H. van Ham, G. Katselis, and & C. Koutsikopoulos. 2007. 'Lagoon
592 Fisheries Resources in Hellas Alexis Conides Hellenic Centre for Marine Research'.

593 Lanzoni, Mattia, Mattias Gaglio, Anna Gavioli, Elisa Anna Fano, and Giuseppe Castaldelli. 2021. 'Seasonal Variation of
594 Functional Traits in the Fish Community in a Brackish Lagoon of the Po River Delta (Northern Italy)'.
595 <https://doi.org/10.3390/w13050679>.

596 Lionello, Piero, Robert J. Nicholls, Georg Umgiesser, and Davide Zanchettin. 2021. 'Venice Flooding and Sea Level:
597 Past Evolution, Present Issues, and Future Projections (Introduction to the Special Issue)'. *Natural Hazards and
598 Earth System Sciences* 21 (8): 2633–41. <https://doi.org/10.5194/NHESS-21-2633-2021>.

599 Liqueste, Camino, Núria Cid, Denis Lanzanova, Bruna Grizzetti, and Arnaud Reynaud. 2016. 'Perspectives on the Link
600 between Ecosystem Services and Biodiversity: The Assessment of the Nursery Function'. *Ecological Indicators* 63
601 (April): 249–57. <https://doi.org/10.1016/j.ecolind.2015.11.058>.

602 Liu, Jianguo, Thomas Dietz, Stephen R. Carpenter, Marina Alberti, Carl Folke, Emilio Moran, Alice N. Pell, et al. 2007.
603 'Complexity of Coupled Human and Natural Systems'. *Science*. American Association for the Advancement of
604 Science. <https://doi.org/10.1126/science.1144004>.

605 Madricardo, Fantina, and S. Donnici. 2014. 'Mapping Past and Recent Landscape Modifications in the Lagoon of Venice
606 through Geophysical Surveys and Historical Maps'. *Anthropocene* 6 (June): 86–96.
607 <https://doi.org/10.1016/J.ANCENE.2014.11.001>.

608 MEA. 2005. 'MILLENNIUM ECOSYSTEM ASSESSMENT (PROGRAM). (2005). Ecosystems and Human Well-
609 Being.' Washington, D.C.

610 Newton, Alice, Ana C. Brito, John D. Icely, Valérie Derolez, Inês Clara, Stewart Angus, Gerald Schernewski, et al. 2018.
611 'Assessing, Quantifying and Valuing the Ecosystem Services of Coastal Lagoons'. *Journal for Nature
612 Conservation*, 1 July 2018. <https://doi.org/10.1016/j.jnc.2018.02.009>.

613 Notte, Alessandra la, Margherita Turvani, and Sergio Giaccaria. 2017. 'Economic Valuation of Ecosystem Services at
614 Local Level for Policy Makers and Planners. The Case of the Island of St. Erasmo in the Lagoon of Venice'.
615 *Environmental Economics* 2 (3). [https://www.businessperspectives.org/index.php/journals/environmental-
616 economics/issue-212/economic-valuation-of-ecosystem-services-at-local-level-for-policy-makers-and-planners-
617 the-case-of-the-island-of-st-erasmo-in-the-lagoon-of-venice](https://www.businessperspectives.org/index.php/journals/environmental-economics/issue-212/economic-valuation-of-ecosystem-services-at-local-level-for-policy-makers-and-planners-the-case-of-the-island-of-st-erasmo-in-the-lagoon-of-venice).

618 Pérez-Ruzafa, Angel, Francesca de Pascalis, Michol Ghezzeo, Jhoni Ismael Quispe-Becerra, Raquel Hernández-García,
619 Irene Muñoz, Carlos Vergara, Isabel María Pérez-Ruzafa, Georg Umgiesser, and Concepción Marcos. 2019.
620 'Connectivity between Coastal Lagoons and Sea: Asymmetrical Effects on Assemblages' and Populations'
621 Structure'. *Estuarine, Coastal and Shelf Science* 216 (January): 171–86.
622 <https://doi.org/10.1016/J.ECSS.2018.02.031>.

623 QGIS Association: QGIS Geographic Information System. 2022. 'QGIS'. 2022. <http://www.qgis.org>.

624 R Core team. 2022. 'R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing,
625 Vienna, Austria.' 2022. <https://www.r-project.org/>.

626 Ravagnan, Gino. 1982. 'Vallicoltura Moderna. Proposte Operative per La Ristrutturazione e Lo Sviluppo Della Itticoltura
627 Salmastra Italiana'. Edagricole. 1982. [https://www.libreriauniversitaria.it/vallicoltura-moderna-proposte-
628 operative-ristrutturazione/libro/9788820617769](https://www.libreriauniversitaria.it/vallicoltura-moderna-proposte-operative-ristrutturazione/libro/9788820617769).

629 Roner, M., A. D'Alpaos, M. Ghinassi, M. Marani, S. Silvestri, E. Franceschinis, and N. Realdon. 2015. 'Spatial Variation
630 of Salt-Marsh Organic and Inorganic Deposition and Organic Carbon Accumulation: Inferences from the Venice
631 Lagoon, Italy'. *Advances in Water Resources* 000: 1–12. <https://doi.org/10.1016/j.advwatres.2015.11.011>.

632 Rova, Silvia, Felix Müller, Patrick Meire, and Fabio Pranovi. 2019. 'Sustainability Perspectives and Spatial Patterns of
633 Multiple Ecosystem Services in the Venice Lagoon: Possible Roles in the Implementation of the EU Water
634 Framework Directive'. *Ecological Indicators* 98 (March): 556–67. <https://doi.org/10.1016/j.ecolind.2018.11.045>.

- 635 Rova, Silvia, and Fabio Pranovi. 2017. 'Analysis and Management of Multiple Ecosystem Services within a Social-
636 Ecological Context'. *Ecological Indicators* 72: 436–43. <https://doi.org/10.1016/j.ecolind.2016.07.050>.
- 637 Rova, Silvia, Fabio Pranovi, and Felix Müller. 2015. 'Provision of Ecosystem Services in the Lagoon of Venice (Italy):
638 An Initial Spatial Assessment'. *Ecohydrology and Hydrobiology* 15 (1): 13–25.
639 <https://doi.org/10.1016/j.ecohyd.2014.12.001>.
- 640 Rova, Silvia, Alice Stocco, and Fabio Pranovi. 2022. 'Ecosystem Services' Capacity and Flow in the Venice Lagoon and
641 the Relationship with Ecological Status'. *One Ecosystem* 7 (September): e79715-
642 <https://doi.org/10.3897/oneeco.7.e79715>.
- 643 RStudio Team. n.d. 'RStudio'.
- 644 Salvetti, Roberta, Marco Acutis, Arianna Azzellino, Marta Carpani, Carlo Giupponi, Paolo Parati, Manfredi Vale, and
645 Renato Vismara. 2008. 'Modelling the Point and Non-Point Nitrogen Loads to the Venice Lagoon (Italy): The
646 Application of Water Quality Models to the Dese-Zero Basin'. *Desalination* 226 (1–3): 81–88.
647 <https://doi.org/10.1016/J.DESAL.2007.01.236>.
- 648 Scapin, Luca, Matteo Zucchetta, Fabio Pranovi, and Piero Franzoi. 2022. 'Influence of Seascape on Coastal Lagoon
649 Fisheries: The Role of Habitat Mosaic in the Venice Lagoon'. *Estuaries and Coasts* 45 (3): 793–811.
650 <https://doi.org/10.1007/S12237-021-00986-3/TABLES/3>.
- 651 Scarton, Francesco. 2017. 'Long-Term Trend of the Waterbird Community Breeding in a Heavily Man-Modified Coastal
652 Lagoon: The Case of the Important Bird Area "Lagoon of Venice"'. *Journal of Coastal Conservation* 21 (1): 35–
653 45. <https://doi.org/10.1007/S11852-016-0470-8/FIGURES/5>.
- 654 Scarton, Francesco, and Mauro Bon. 2009. 'Gli Uccelli Acquatici Svernanti Nella Laguna Di Venezia Nel Periodo 1993-
655 2007: Analisi Delle Dinamiche Temporali e Spaziali'. *Avocetta*.
- 656 Schröter, Matthias, David N. Barton, Roy P. Remme, and Lars Hein. 2014. 'Accounting for Capacity and Flow of
657 Ecosystem Services: A Conceptual Model and a Case Study for Telemark, Norway'. *Ecological Indicators* 36: 539–
658 51. <https://doi.org/10.1016/j.ecolind.2013.09.018>.
- 659 Seitzinger, S., J.A. Harrison, A.F. Bohlke, A.F. Bouwman, R. Lowrance, B. Peterson, C. Tobias, and G. van Drecht. 2006.
660 'Denitrification Across Landscapes and Waterscapes : A Synthesis'. *Ecological Applications* 16 (6): 2064–90.
661 [https://doi.org/10.1890/1051-0761\(2006\)016\[2064:DALAWA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[2064:DALAWA]2.0.CO;2).
- 662 Sfriso, Adriano, and Chiara Facca. 2007a. 'Distribution and Production of Macrophytes and Phytoplankton in the Lagoon
663 of Venice: Comparison of Actual and Past Situation'. *Hydrobiologia*. <https://doi.org/10.1007/s10750-006-0418-3>.
- 664 Sfriso, Adriano, Chiara Facca, and Sonia Ceoldo. 2004. 'Growth and Production of *Cymodocea Nodosa* (Ucria)
665 *Ascherson* in the Venice Lagoon'. *Scientific Research and Safeguarding of Venice* 96: 229–36.
- 666 Sfriso, Adriano, and Pier Francesco Ghetti. 1998a. 'Seasonal Variation in Biomass, Morphometric Parameters and
667 Production of Seagrasses in the Lagoon of Venice'. *Aquatic Botany*. [https://doi.org/10.1016/S0304-3770\(98\)00064-](https://doi.org/10.1016/S0304-3770(98)00064-3)
668 3.
- 669 Sfriso, Adriano, and Antonio Marcomini. 1999. 'Macrophyte Production in a Shallow Coastal Lagoon. Part II: Coupling
670 with Sediment, SPM and Tissue Carbon, Nitrogen and Phosphorus Concentrations'. *Marine Environmental*
671 *Research*. [https://doi.org/10.1016/S0141-1136\(98\)00122-6](https://doi.org/10.1016/S0141-1136(98)00122-6).
- 672 Shang, Yung Cheng. 1982. *Aquaculture Economics: Basic Concepts and Methods of Analysis*. *Journal of the Marine*
673 *Biological Association of the United Kingdom*. Vol. 62. Cambridge University Press.
674 <https://doi.org/10.1017/S0025315400057519>.
- 675 Solidoro, Cosimo, Vinko Bandelj, Fabrizio Aubry Bernardi, Elisa Camatti, Stefano Ciavatta, Gianpiero Cossarini, Chiara
676 Facca, et al. 2010. 'Response of the Venice Lagoon Ecosystem to Natural and Anthropogenic Pressures over the
677 Last 50 Years'. *Coastal Lagoons: Critical Habitats of Environmental Change*, no. January: 483–511.
678 <https://doi.org/10.1201/EBK1420088304>.
- 679 Solidoro, Cosimo, Roberto Pastres, and Gianpiero Cossarini. 2005. 'Nitrogen and Plankton Dynamics in the Lagoon of
680 Venice'. *Ecological Modelling* 184 (1): 103–23. <https://doi.org/10.1016/j.ecolmodel.2004.11.009>.
- 681 Umgiesser, Georg. 2020. 'The Impact of Operating the Mobile Barriers in Venice (MOSE) under Climate Change'.
682 *Journal for Nature Conservation* 54 (April). <https://doi.org/10.1016/j.jnc.2019.125783>.

683 Umgiesser, Georg, Donata Melaku Canu, Andrea Cucco, and Cosimo Solidoro. 2004. 'A Finite Element Model for the
684 Venice Lagoon. Development, Set up, Calibration and Validation'. *Journal of Marine Systems* 51 (1-4 SPEC. ISS.):
685 123–45. <https://doi.org/10.1016/j.jmarsys.2004.05.009>.

686 Villamagna, Amy M., Paul L. Angermeier, and Elena M. Bennett. 2013. 'Capacity, Pressure, Demand, and Flow: A
687 Conceptual Framework for Analyzing Ecosystem Service Provision and Delivery'. *Ecological Complexity* 15: 114–
688 21. <https://doi.org/10.1016/j.ecocom.2013.07.004>.

689 Walton, M. E.M., C. Vilas, J. P. Cañavate, E. Gonzalez-Ortegon, A. Prieto, S. A. van Bergeijk, A. J. Green, M. Librero,
690 N. Mazuelos, and L. le Vay. 2015. 'A Model for the Future: Ecosystem Services Provided by the Aquaculture
691 Activities of Veta La Palma, Southern Spain'. *Aquaculture* 448 (November): 382–90.
692 <https://doi.org/10.1016/j.aquaculture.2015.06.017>.

693 Walton, M. E.M., C. Vilas, C. Coccia, A. J. Green, J. P. Cañavate, A. Prieto, S. A. van Bergeijk, et al. 2015. 'The Effect
694 of Water Management on Extensive Aquaculture Food Webs in the Reconstructed Wetlands of the Doñana Natural
695 Park, Southern Spain'. *Aquaculture* 448 (November): 451–63.
696 <https://doi.org/10.1016/J.AQUACULTURE.2015.06.011>.

697 Weitzman, Jenny. 2019. 'Applying the Ecosystem Services Concept to Aquaculture: A Review of Approaches,
698 Definitions, and Uses'. *Ecosystem Services* 35 (February): 194–206. <https://doi.org/10.1016/j.ecoser.2018.12.009>.

699 Zucchetta, Matteo, Fabrizio Capoccioni, Piero Franzoi, Eleonora Ciccotti, and Chiara Leone. 2021. 'Fish Response to
700 Multiple Anthropogenic Stressors in Mediterranean Coastal Lagoons: A Comparative Study of the Role of Different
701 Management Strategies'. *Water* 13 (2): 130. <https://doi.org/10.3390/w13020130>.

702
703

704
705

Supplementary materials I

Scheme of a valle da pesca




Figure S.M. I - 1 Sketch map of a valle da pesca structure. a) brackish water inlet; b) brackish lake; c) winter fishponds; d) freshwater lakes; e) freshwater intake.

706

Supplementary materials II

707

Remote sensing images

708

709

710

711

712

The following tables list the remote sensing images analyzed in this work. Very High-Resolution (VHR) satellite imageries have been granted by the European Space Agency (ESA) via the ESA On-Demand Restricted Data Access program (Tab. S. M. II - 1). Aerial photographs has been granted by Regione Veneto, “Direzione Pianificazione Territoriale, U.O. Pianificazione territoriale strategica e cartografia” (Tab. S. M. II - 2).

Satellite scenes					
Collection data	Satellite/sensor	bands	resolution	clouds coverage	off-nadir:
2019/07/01	WV 02	8-bands	40 cm	0.0%	13.0°
2019/01/10	WV 02	8-bands	40 cm	1.0%	12.6°
2019/11/07	WV 03	8-bands	32 cm	0.0%	8.1°
2018/11/29	GE 01	4-bands	51 cm	0.0%	26.7°

Table S. M. II - 1 VHR satellite imageries characteristics

Collection data	Flight name	bands	resolution
2018	REVEN 2018	3-bands (RGB)	20 cm
2019	AGEA REVEN	3-bands (RGB)	20 cm

Table S.M. II - 2 Aerial photographs characteristics

713 **Supplementary materials III**714 **ESs assessment methodology**

ES category	ES	Assessment methodology	
		Valli da pesca	Lagoon
Regulating & maintenance	Climate regulation Carbon sequestration rate by saltmarshes and seagrasses [gC m ⁻² y ⁻¹]	<p><i>Capacity/flow methodology</i></p> <p>The spatial distribution of saltmarshes in the valli da pesca has been achieved with a machine learning classification of very high-resolution satellite and aerial imageries (listed in table S. M. II - 1), with calibration points collected during field work.</p> <p>Average saltmarshes' carbon sequestration rate based on literature data (Day et al. 1998; Roner et al. 2015)</p> <p>The spatial distribution of seagrasses meadows has been achieved by photointerpretation of very high-resolution satellite imageries (WV-2A, GeoEye), with the support of UAV aerial imageries and visual census field surveys data collected during periodical visits within the valli da pesca.</p> <p>Seagrasses' C sequestration rate based on species-specific belowground production and organic C content (Sfriso, Facca, and Ceoldo 2004; Sfriso and Francesco Ghetti 1998b; Sfriso, Facca, and Ceoldo 2007; Sfriso and Facca 2007; 2007)</p> <p>According to the literature, capacity and flow indicators of regulating ESs have been considered coincident (Schröter et al. 2014; Hein et al. 2006b).</p>	<p><i>Capacity/flow methodology</i></p> <p>The saltmarshes spatial distribution in the open, not privately managed part of the lagoon has been retrieved from the saltmarshes map achieved in 2003 by Magistrato alle Acque.</p> <p>Average saltmarshes' carbon sequestration rate based on literature data (Day et al. 1998; Roner et al. 2015)</p> <p>The seagrasses distribution refers to the seagrasses map in 2017 (Provveditorato OO. PP. del Triveneto; SELC, 2018).</p> <p>Seagrasses' C sequestration rate based on species-specific belowground production and organic C content (Sfriso and Facca 2007; Sfriso, Facca, and Ceoldo 2007; 2004; Sfriso and Francesco Ghetti 1998b)</p> <p>According to the literature, capacity and flow indicators of regulating ESs have been considered coincident (Schröter et al. 2014; Hein et al. 2006b).</p>
Regulating & maintenance	Water purification Percentage of Nitrogen load removed by denitrification process in brackish water [%]	<p><i>Capacity/flow methodology</i></p> <p>Nitrogen removal capacity of the brackish lakes estimated in proportion to the lake extension and the turnover time of the volume of water in the saltwater basin, based on experimental data about denitrification measured by Ravagnan (1982) in other valli da pesca managed in the same way as the valli da pesca considered in this study.</p> <p>According to the literature, capacity and flow indicators of regulating ESs have been considered coincident (Schröter et al. 2014; Hein et al. 2006a).</p>	<p><i>Capacity/flow methodology</i></p> <p>N load removed through denitrification estimated based on residence time, according to the equation proposed by Seitzinger et al., (2006) for estuarine systems. Residence time calculated with SHYFEM model (Umgiesser et al. 2004) referred to the year 2014 (courtesy of G. Umgiesser, ISMAR-CNR).</p> <p>According to the literature, capacity and flow indicators of regulating ESs have been considered coincident (Schröter et al. 2014; Hein et al. 2006a).</p>


ES category	ES	Assessment methodology	
		Valli da pesca	Lagoon
Regulating & maintenance	Lifecycle support 0-1 scale	<p>Capacity methodology Attractiveness for migratory waterbirds related to the presence of waterbirds lifecycle support factors (saltmarshes, freshwater presence, shrubs, and herbaceous vegetation) that enhance the suitability for resting and molt changing (Korschgen et al., 1985; Havera et al., 1992; Arzel et al., 2006; Hatziiordanou et al., 2019), retrieved from aerial imageries photo interpretation and geospatial layers data. Estimated distribution in the most confined part of the lagoon of juveniles fish biomass [kg ha⁻¹y⁻¹] of the food-web group "Mugilidae", retrieved from a spatialized foodweb model of the Venice lagoon (Anelli Monti et al. 2021a). Fish and waterbirds lifecycle support capacities were normalized to a 0-1 scale, then aggregated.</p> <p>Flow methodology Waterbirds: Average number of migratory waterbirds [n ha⁻¹y⁻¹] wintering in the valli da pesca, according to the last ten years' waterbirds annual censuses (AFV 2020). Fish: Fry biomass [kg ha⁻¹y⁻¹] sown in the valli da pesca for the species Mugil cephalus, Chelon labrosus, Chelon aurata, Chelon saliens and Chelon ramada according to the official management data.</p> <p>Waterbirds and fish lifecycle support capacities were normalized to a 0-1 scale, then aggregated.</p>	<p>Capacity methodology Attractiveness for migratory waterbirds related to the presence of waterbirds lifecycle support factors (saltmarshes, freshwater presence, shrubs, and herbaceous vegetation) that enhance the suitability for resting and molt changing (Korschgen et al., 1985; Havera et al., 1992; Arzel et al., 2006; Hatziiordanou et al., 2019), retrieved from aerial imageries photo interpretation and geospatial layers data. Estimated distribution of juveniles fish biomass [kg ha⁻¹y⁻¹] of the food-web group "Mugilidae", retrieved from a spatialized foodweb model of the Venice lagoon (Anelli Monti et al. 2021a). Fish and waterbirds lifecycle support capacities were normalized to a 0-1 scale, then aggregated.</p> <p>Flow methodology Waterbirds: Average number of migratory waterbirds [n ha⁻¹y⁻¹] wintering in the open and not privately managed part of the lagoon, according to the last ten years' waterbirds annual censuses (AFV 2020). Fish: Fry biomass [kg ha⁻¹y⁻¹] distribution estimated in the lagoon, retrieved from a spatialized foodweb model of the Venice lagoon (Anelli Monti et al., 2021).</p> <p>Waterbirds and fish lifecycle support capacities were normalized to a 0-1 scale, then aggregated.</p>
Provisioning	Fish production Fish biomass [kg ha ⁻¹ y ⁻¹]	<p>Capacity methodology Potential fish biomass hosted by each <i>valle da pesca</i> (kg ha⁻¹ y⁻¹), estimated basing on an Ecopath-Ecosim with Ecospace spatialized foodweb model (Anelli Monti et al. 2021a).</p> <p>Flow methodology Average fish catches per hectare of brackish water surfaces per year (kg ha⁻¹ y⁻¹) in the valli da pesca, according to the 2010-2019 official data (Regione Veneto).</p>	<p>Capacity methodology Sum of the biomass of fish functional groups targeted by artisanal fishing, calculated by an Ecopath-Ecosim with Ecospace spatialized foodweb model (Anelli Monti et al. 2021).</p> <p>Flow methodology Average fish catches from artisanal fishing per hectare of water per year (kg ha⁻¹ y⁻¹), calculated by an Ecopath-Ecosim with Ecospace spatialized foodweb model (Anelli Monti et al. 2021).</p>
Provisioning	Waterfowl hunting Number of huntable waterbirds [n ha ⁻¹ y ⁻¹]	<p>Capacity methodology Mean number of huntable waterbirds per hectare per year wintering in the valli da pesca, according to the time series of waterbirds censuses from 2010 to 2019 (Associazione Faunisti Veneti, 2020), spatially interpolated by Dirichlet tessellation.</p> <p>Flow methodology Average number of waterbirds catches calculated on 2010 - 2020 hunting registries data of the <i>valli da pesca</i>, spatialized considering the effective hunting reserve area.</p>	<p>Capacity methodology Mean number of huntable waterbirds per hectare per year wintering in the open part of the lagoon, according to the time series of waterbirds censuses from 2010 to 2019 (Associazione Faunisti Veneti, 2020), spatially interpolated by Dirichlet tessellation</p> <p>Flow methodology Estimated considering the catches per capita (n. birds/person/hunting trip), the hunting effort (n. of hunting trips/person/year) and the proportion of active hunters in the lagoon outside the hunting farms. Data were gathered through 84 interviews to hunters and by asking the total number of hunters active in the lagoon to the "Ambito Territoriale di caccia VE5" in 2020. Spatial distribution based on the location of the hunting blinds in the lagoon (Regione Veneto, 2019), weighted by the average effective shotgun range.</p>
Provisioning	Wild herbs & honey	<p>Capacity methodology Kg of <i>Salicornia spp.</i> biomass that could be harvested in the <i>valli da pesca</i>, per year, estimated from the vegetational patches distribution in the valli da pesca retrieved with both visual census method and through the identification of their predictive range in R, G, B bands</p>	<p>Capacity methodology Kg of <i>Salicornia</i> biomass that could be harvested in the lagoon, per year, estimated from the vegetational patches distribution in the open lagoon retrieved with both visual census method and through the identification of their predictive range in R, G, B bands reflectance values,</p>

ES category	ES	Assessment methodology	
		Valli da pesca	Lagoon
		<p>reflectance values, filtered through a cut-off NDVI value calculated from aerial and satellite imageries (listed in Supplementary Materials II).</p> <p>Kg of honey potentially produced per unit area in the valli da pesca, considering the <i>Limonium sp.</i> inflorescence cover in the patches (Fantinato and Buffa 2019) and the ratio between the number of sea lavender flowers and the grams of honey that can be obtained from them.</p> <p><i>Flow methodology</i> Kg of harvested edible plants reported by 2020 market data (courtesy of Veneto Agricoltura), by interviewed local people and restaurant chefs. Kg of produced sea-lavender honey witnessed by 5 interviewed beekeepers.</p>	<p>filtered through a cut-off NDVI value calculated from aerial and satellite imageries (listed in Supplementary Materials II).</p> <p>Kg of honey potentially produced per unit area in the lagoon, considering the <i>Limonium</i> inflorescence cover in the patches (Fantinato and Buffa 2019) and the ratio between the number of sea lavender flowers and the grams of honey that can be obtained from them.</p> <p><i>Flow methodology</i> Kg of harvested edible plants reported by 2020 market data (courtesy of Veneto Agricoltura), by interviewed local people and restaurant chefs. Kg of produced sea-lavender honey witnessed by 5 interviewed beekeepers.</p>
Cultural	Tourism	<p><i>Capacity methodology</i> Tourism attractiveness map, assessed by a survey carried out in the second half of 2019. Spatialization achieved by considering the attractiveness factors evaluated by tourists in the questionnaires, namely saltmarshes presence, the possibility to observe birds and fauna, good water quality, and the chance to contemplate natural terrestrial habitats in the valli da pesca. Each element was weighted to depict the interest declared by people and normalized to a 0-1 scale index.</p> <p><i>Flow methodology</i> Number of people who, during a year, have passed at least one night in one of the accommodation facilities within the valli da pesca.</p>	<p><i>Capacity methodology</i> Tourism attractiveness map, assessed by a survey carried out in the second half of 2019. Spatialization achieved by considering the attractiveness factors evaluated by tourists in the questionnaires, namely saltmarshes presence, the possibility to observe birds and fauna, good water quality, and the chance to contemplate natural terrestrial habitats in the lagoon. Each element was weighted to depict the interest declared by people and normalized to a 0-1 scale index.</p> <p><i>Flow methodology</i> Number of people visiting the lagoon in the year 2019, excluding the historical center of Venice, obtained from tourism and transportation operators (public transport company AVM-ACTV S.p.a., 17 private navigation companies, 9 ecotourism associations).</p>
Cultural	Information for cognitive development	<p><i>Capacity methodology</i> The natural factors that enhance touristic attractiveness assessed by a survey carried out in the second half of 2019 have been mapped along with the inclusivity for the necessities of people with disabilities. The resulting map was normalized to a 0-1 scale.</p> <p><i>Flow methodology</i> Number of persons that yearly attend outdoor educational activities or guided one-day trip in the valli da pesca, as reported by the valli da pesca managers, 6 touristic guides and 6 major ecotourism associations.</p>	<p><i>Capacity methodology</i> The natural factors that enhance touristic attractiveness assessed by a survey carried out in the second half of 2019 have been mapped along with the inclusivity for the necessities of people with disabilities. The resulting map was normalized to a 0-1 scale.</p> <p><i>Flow methodology</i> Number of persons that yearly attend outdoor educational activities or guided one-day trip in the lagoon, as reported by 6 touristic guides and 6 major ecotourism associations.</p>
Cultural	Birdwatching	<p><i>Capacity methodology</i> Birdwatching attractiveness normalized map, scaled 0-1, based on the factors reported as important by 30 interviewed birdwatchers, namely the presence of pedestrian paths, saltmarshes presence, birds nesting areas in the visual field, and high probability for observing birds. All these factors were summed up to obtain a normalized attractiveness map.</p> <p><i>Flow methodology</i> Average number of active birdwatchers [$n y^{-1}$] calculated from the observers' activity trend, recorded from 2010 to 2020 in the Italian birdwatchers' database (https://www.ornitho.it/).</p>	<p><i>Capacity methodology</i> Birdwatching attractiveness normalized map, scaled 0-1, based on the factors reported as important by 30 interviewed birdwatchers, namely the presence of pedestrian paths, saltmarshes presence, birds nesting areas in the visual field, and high probability for observing birds. All these factors were summed up to obtain a normalized attractiveness map.</p> <p><i>Flow methodology</i> Average number of active birdwatchers [$n y^{-1}$] calculated from the observers' activity trend, recorded from 2010 to 2020 in the Italian birdwatchers' database (https://www.ornitho.it/).</p>


- 717 AFV, A.F.V. (2020) CENSIMENTO DEGLI UCCELLI ACQUATICI SVERNANTI IN PROVINCIA DI VENEZIA.
- 718 Anelli Monti, M. et al. (2021) 'Ecosystem functioning and ecological status in the Venice lagoon, which relationships?',
719 *Ecological Indicators*, 133, p. 108461. Available at: <https://doi.org/10.1016/j.ecolind.2021.108461>.
- 720 Arzel, C., Elmberg, J. and Guillemain, M. (2006) 'Ecology of spring-migrating Anatidae: A review', *Journal of*
721 *Ornithology*. Springer, pp. 167–184. Available at: <https://doi.org/10.1007/s10336-006-0054-8>.
- 722 Day, J.W. et al. (1998) 'Relative sea level rise and Venice lagoon wetlands', *Journal of Coastal Conservation*, 4(1), pp.
723 27–34. Available at: <https://doi.org/10.1007/BF02806486>.
- 724 Fantinato, E. and Buffa, G. (2019) 'Animal-mediated interactions for pollination in saltmarsh communities', *Plant*
725 *Sociology*, 56(2), pp. 35–42. Available at: <https://doi.org/10.7338/pls2019562/02>.
- 726 Hatziordanou, L. et al. (2019) 'Indicators for mapping and assessment of ecosystem condition and of the ecosystem
727 service habitat maintenance in support of the EU biodiversity strategy to 2020', *One Ecosystem*, 4. Available at:
728 <https://doi.org/10.3897/oneeco.4.e32704>.
- 729 Havera, S.P. et al. (1992) 'HUMAN DISTURBANCE OF WATERFOWL ON KEOKUK POOL', undefined [Preprint].
- 730 Hein, L. et al. (2006a) 'Spatial scales, stakeholders and the valuation of ecosystem services', *Ecological Economics*,
731 57(2), pp. 209–228. Available at: <https://doi.org/10.1016/J.ECOLECON.2005.04.005>.
- 732 Hein, L. et al. (2006b) 'Spatial scales, stakeholders and the valuation of ecosystem services', *Ecological Economics*,
733 57(2), pp. 209–228. Available at: <https://doi.org/10.1016/J.ECOLECON.2005.04.005>.
- 734 Korschgen, C., George, L. and Green, W.L. (1985) 'Disturbance of diving ducks by boaters on a migrational staging
735 area', undefined [Preprint].
- 736 Ravagnan, G. (1982) *Vallicoltura moderna. Proposte operative per la ristrutturazione e lo sviluppo della itticoltura*
737 *salmastra italiana, Edagricole*. Available at: [https://www.libreriauniversitaria.it/vallicoltura-moderna-proposte-](https://www.libreriauniversitaria.it/vallicoltura-moderna-proposte-operative-ristrutturazione/libro/9788820617769)
738 [operative-ristrutturazione/libro/9788820617769](https://www.libreriauniversitaria.it/vallicoltura-moderna-proposte-operative-ristrutturazione/libro/9788820617769) (Accessed: 27 March 2021).
- 739 Roner, M. et al. (2015) 'Spatial variation of salt-marsh organic and inorganic deposition and organic carbon
740 accumulation: Inferences from the Venice lagoon, Italy', *Advances in Water Resources*, 000, pp. 1–12. Available at:
741 <https://doi.org/10.1016/j.advwatres.2015.11.011>.
- 742 Schröter, M. et al. (2014) 'Accounting for capacity and flow of ecosystem services: A conceptual model and a case
743 study for Telemark, Norway', *Ecological Indicators*, 36, pp. 539–551. Available at:
744 <https://doi.org/10.1016/j.ecolind.2013.09.018>.
- 745 Seitzinger, S. et al. (2006) 'Denitrification Across Landscapes and Waterscapes : a Synthesis', *Ecological Applications*,
746 16(6), pp. 2064–2090. Available at: [https://doi.org/10.1890/1051-0761\(2006\)016\[2064:DALAWA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[2064:DALAWA]2.0.CO;2).
- 747 Sfriso, A. and Facca, C. (2007) 'Nanozostera noltii growth and production in the lagoon of Venice', *Biologia Marina*
748 *Mediterranea*, 14, pp. 326–327.
- 749 Sfriso, A., Facca, C. and Ceoldo, S. (2004) 'Growth and production of *Cymodocea nodosa* (Ucria) ascherson in the
750 Venice Lagoon', *Scientific Research and Safeguarding of Venice*, 96, pp. 229–236.
- 751 Sfriso, A., Facca, C. and Ceoldo, S. (2007) *Scientific Research and Safeguarding of Venice*.
- 752 Sfriso, A. and Francesco Ghetti, P. (1998) 'Seasonal variation in biomass, morphometric parameters and production of
753 seagrasses in the lagoon of Venice', *Aquatic Botany*, 61(3), pp. 207–223. Available at: [https://doi.org/10.1016/S0304-](https://doi.org/10.1016/S0304-3770(98)00064-3)
754 [3770\(98\)00064-3](https://doi.org/10.1016/S0304-3770(98)00064-3).
- 755 Umgiesser, G. et al. (2004) 'A finite element model for the Venice Lagoon. Development, set up, calibration and
756 validation', *Journal of Marine Systems*, 51(1-4 SPEC. ISS.), pp. 123–145. Available at:
757 <https://doi.org/10.1016/j.jmarsys.2004.05.009>
758

Supplementary materials IV


a) Climate regulation




b) Denitrification




c) Lifecycles support - capacity




d) Lifecycles support - flow




e) Aquaculture - capacity




f) Aquaculture - flow




g) Waterfowls hunting - capacity




h) Waterfowls hunting - flow




i) Edible halophytes & honey production - cap




j) Edible halophytes & honey production - flow




k) Tourism - capacity




l) Tourism - flow




m) Information for c.d. - capacity



n) Information for c.d. - flow



o) Birdwatching - capacity



p) Birdwatching - flow



760

761

762

763

Fig. S.M. IV Results of the ESs assessment. a), b), c), d) refer to regulating and maintenance ESs. e), f), g), h), i), j) refer to provisioning ESs. k), l), m), n), o), p) refer to cultural ESs.