

1 Alice Stocco^{a*}, Laura Basconi^a, Silvia Rova^a, Fabio Pranovi^a

2 Like little lagoons: the contribution of *valli da pesca* to the
3 Ecosystem Services supply of the Venice Lagoon
4

5 Original Report of Research Results Article

6 ^a Department of Environmental Sciences, Informatics and Statistics,
7 Ca' Foscari University of Venice,
8 Via Torino, 155
9 30170 Mestre – Venezia
10 Italy

11
12 *Corresponding author:
13 Alice Stocco
14 Ca' Foscari University of Venice,
15 Via Torino, 155
16 30170 Mestre – Venezia
17 Italy
18 alice.stocco@unive.it
19 + 39 333 847 9659
20 +39 041 234 7713
21

22 Like little lagoons: the contribution of *valli da pesca* to the 23 Ecosystem Services supply of the Venice Lagoon 24

25 Alice Stocco^{a*}, Laura Basconi^a, Silvia Rova^a, Fabio Pranovi^a

26 ^a Department of Environmental Sciences, Informatics and Statistics, Università Ca' Foscari, Venezia, Italy

27 *Corresponding author: alice.stocco@unive.it

28 Abstract

29 The Venice lagoon social-ecological system is characterized by a strong relationship between the natural
30 environment and human activities. This is especially noticeable in the aquaculture and hunting reserves of the
31 lagoon, locally known as *valli da pesca*. Previous works about Ecosystem Services (ESs) in the Venice lagoon
32 focused on the so-called “open lagoon”, overlooking the role of the *valli da pesca*. Despite being completely
33 managed ecosystems, the *valli da pesca* have conserved typical elements of transitional water environments
34 that the other parts of the lagoon have lost. By evaluating nine ESs using a spatially explicit approach, we found
35 that the *valli da pesca*, despite covering 17% of the surface, are contributing for 38% of the ESs total capacity,
36 and for 24% of the ESs total flow, in comparison to the open part of the lagoon. Moreover, the management
37 that aims to maximize in a perspective of sustainability some provisioning ESs, such as extensive aquaculture,
38 can positively influence the presence of factors on which other ESs capacity is also based. As a result, the open
39 lagoon benefits from a sort of spill-over effect for lifecycle support, hunting, and cultural ESs such as tourism,
40 information for cognitive development, and birdwatching. Such significant contributions could be endangered
41 in the context of a lagoon subjected to increasing pressures from anthropic activities. Adaptations to impacts,
42 as well as to climate change and sea-level rise effects, will modify the lagoon hydrodynamics and the sea-lagoon
43 connectivity and threaten the *valli da pesca* and so their ESs supply.

44 **Keywords:** Ecosystem Services, ‘Side Effects’, Valli da pesca, Extensive aquaculture, Waterfowls Hunting, Venice lagoon
45

46 1. Introduction

47 In the last decades, assessment and mapping of Ecosystem Services (ESs) have become effective methods to
48 highlight all the contributions that humankind receives from Nature (Costanza et al. 1997; Burkhard and Maes
49 2017; Baró et al. 2016; MEA 2005). The ESs approach also makes evident services that would otherwise go
50 unnoticed because they flow effortlessly to human beings (Liu et al. 2007).

51 Nature and humans always interact, creating complex social-ecological systems where society receives
52 positive contributions to their well-being through the ecological processes.

53 In Italy, the Venice lagoon ecosystem is an emblematic example of such a complex social-ecological system.

54 Since remote times, interactions between natural factors, social dynamics, and economic activities have
55 shaped and affected its morphology and functioning (Solidoro et al. 2010). Given the complexity of these
56 interactions, understanding spatial and temporal patterns of ESs capacity and flow (*sensu* Villamagna et al.,

57 2013) plays a key role in environmental decision-making regarding the Venice lagoon, as suggested by Rova et
58 al. (2015, 2019).

59 Previous works, however, focused on the so-called “open lagoon”, composed of the principal islands and
60 water bodies (Newton et al. 2018; la Notte et al. 2017; Rova et al. 2015; D’alpaos and D’alpaos 2021; Rova et
61 al. 2019; Rova et al. 2022). Until now, no data have been gathered about the possible contribution to ESs by
62 some confined, man-managed areas along the lagoon edges, called in Italian “*valli da pesca*”. These areas,
63 being considered Heavily Modified Water Bodies (HMWB) under the Water Framework Directive (European
64 Commission 2000), are located at the interface between the land and the lagoon water and are almost
65 completely closed, covering a total surface of approximately 97 km².

66 Like other similar environments in other coastal lagoons of the Northern Adriatic, the *valli da pesca* were
67 established during the XIV Century as aquaculture facilities, where temporary boundaries were conceived to
68 entrap fish without affecting water flows. Over time, the *valli da pesca* have been progressively isolated from
69 the lagoon with permanent embankments. Nowadays, they are used especially for fish farming and waterfowl
70 hunting and depend almost completely on human intervention for functioning, in terms of freshwater and
71 brackish water supply, as well as for the maintenance of morphological features, resulting in an ecosystem that
72 can be considered like an “artificial ecosystem”. Quite paradoxically, the *valli da pesca*, due to their private
73 management regime, have conserved typical elements of transitional ecosystems that the lagoon itself has
74 instead progressively lost. Consequently, it is legit to ask whether these environments, while including and
75 preserving brackish water basins, freshwater lakes, and saltmarshes, could act as important conservation
76 areas, while providing ESs for the lagoon social-ecological system.

77 This work presents the first quantitative, GIS-based assessment of ESs in the *valli da pesca* of the Venice
78 lagoon. On the one hand, it widens the knowledge about today’s status of these ancient but poorly investigated
79 environments, and on the other hand, it suggests some reflections about the effects of private land
80 conservation. Furthermore, it explores the multiple ESs capacity and flow relationships emerging within these
81 areas, and sheds light on their contribution to the whole lagoon ‘ESs budget’.

82

83 2 Materials and methods


84 2.1 Study area

85 The study area is included in the Venice lagoon, the widest Italian transitional ecosystem (Fig.1). Each of the
86 27 *valli da pesca* that are currently still operative consists of a series of basins at different water salinity,
87 separated from each other, and the lagoon, by means of artificial embankments, imitating the typical
88 transitional water gradients.

89 Indeed, a typical *valle da pesca* receives the lagoon water that enters through a barrage, flows in the
90 brackish lakes, and then glides towards the land. On the opposite side, some basins store freshwater inputs
91 from inland rivers, creating a freshwater wetland area from which the water flows out to reach the major
92 brackish basin. The mixed water then circulates in different sectors of the *valle* and finally, with the low tide,
93 flows out into the lagoon (more information in Supplementary Materials I, Fig. S.M. 1.1). This creates a
94 heterogeneous landscape with strong ecological gradients and a multitude of habitats, as much as a natural
95 lagoon.

96 Among the *valli da pesca*, different types of management can be distinguished, ranging from aquaculture
97 and hunting reserves to tourist estates. However, they all show the common trait of having a restricted access
98 regime, making it difficult to collect data.

99



100

101 **Figure 1** Study area; 1 = Valle Dogà, 2 = Valle Grassabò, 3 = Vallesina, 4 = Valle Fosse, 5 = Valle Lio Maggiore, 6 =
102 Valle Bianca, 7 = Valle Dragojesolo, 8 = Valle Cavallino, 9 = Valle Falconera, 10 = Valle Liona, 11 = Valle Olivara, 12 =
103 Saline-Manciane-Sparasera, 13 = Valle Paleazza, 14 = Valle Sacchettina, 15 = Valle Sacchetta, 16 = Valle Ca' Zane, 17 =
104 Santa Cristina island, 18 = Valle Perini, 19 = Valle Miana-Serraglia, 20 = Valle Averno, 21 = Valle A.M.A., 22 = Valle
105 Contarina, 23 = Valle Cornio Alto e Cornio Basso, 24 = Valle Zappa, 25 = Valle Figheri, 26 = Valle Pierimpiè, 27 = Valle
106 Morosina-Ghebo Storto.

107

2.2 ESS' data collection, analysis, and mapping

108 The nine Ecosystem Services and related indicators for both capacity and flow are reported in Table 1. The ES
109 category refers to the nomenclature of the Common International Classification of Ecosystem Services (CICES
110 framework).

111

112

CICES section	Ecosystem Service	Capacity indicator	Flow indicator
Regulating & Maintenance services	Climate regulation	Carbon sequestration rate by saltmarshes and seagrasses [gC m ⁻² y ⁻¹]*	
	Water purification	Percentage of Nitrogen load removed by denitrification process in brackish water [%]*	
	*According to Schröter et al. (2014) and Hein et al. (2006) capacity and flow indicators of climate regulation and water purification ESs have been considered coincident		
Provisioning services	Lifecycle support for fish and of avian migratory species	Attractiveness for migratory waterbirds and potentially hostable juveniles fish biomass normalized to a 0-1 scale	Number of wintering migratory waterbirds and sown juveniles fish biomass normalized to a 0-1 scale
	Aquaculture production	Fish biomass [kg ha ⁻¹ y ⁻¹]	Harvested fish biomass [kg ha ⁻¹ y ⁻¹]
	Waterbirds' hunting	Number of huntable waterbirds [n ha ⁻¹ y ⁻¹]	Number of catches [n ha ⁻¹ y ⁻¹]
Cultural services	Wild edible plants and honey production	<i>Salicornia sp.</i> biomass [kg ha ⁻¹ y ⁻¹] Honey [kg ha ⁻¹ y ⁻¹]	Harvested <i>Salicornia sp.</i> biomass [kg ha ⁻¹ y ⁻¹] Harvested honey [kg ha ⁻¹ y ⁻¹]
	Tourism	Tourism attractiveness [0-1 scale]	Number of tourists [n y ⁻¹]
	Information for cognitive development	Environmental education attractiveness [0-1 scale]	Number of one-day guided excursionists and students [n y ⁻¹]
	Birdwatching	Birdwatching attractiveness [0-1 scale]	Mean number of active birdwatchers [n y ⁻¹]

113 **Table 1** ESs selected for the present assessment and the adopted indicator.

114 2.2.1 Regulating and maintenance ESs

115 The climate regulation ES has been assessed as carbon sequestration process by saltmarshes accretion (Day et al. 1998; Roner et al. 2015) and seagrasses meadows (Sfriso and Francesco Ghetti, 1998; Sfriso and Marcomini, 116 1999; Sfriso et al., 2007) that are present in the studied reserves.

118 The water purification ES is expressed as the nitrogen removal capacity of brackish lakes, in proportion to the water volume and its turnover time in each *valle da pesca*. Because of the unavailability of local data, the 119 estimation was based on denitrification data by Ravagnan (1982), who measured the difference between the 120 TIN in the inflowing water and outlet water in *valli da pesca* which are in all aspects managed similarly to the 121 those considered in this case study.

123 According to the literature, the capacity and flow indicators of the aforementioned regulating ESs are 124 considered coincident (Schröter et al. 2014; Hein et al. 2006a).

125 The lifecycle support was assessed by focusing on the migration patterns of both the fish and waterbirds. 126 Fish lifecycle support capacity was estimated by extrapolating the distribution of mugilids fingerlings in the 127 most confined part of the lagoon, retrieved from a spatialized foodweb model of the Venice lagoon based on 128 functional groups (Anelli Monti et al., 2021). For migratory birds, we mapped for each one of the *valli da pesca* 129 the favorable factors, namely saltmarshes, freshwater presence, shrubs, and herbaceous vegetation, that 130 enhance the attractiveness for resting and molt changing (Korschgen et al., 1985; Havera et al., 1992; Arzel et 131 al., 2006; Hatziiordanou et al., 2019). Fish and waterbird lifecycle support capacities were normalized on a 0- 132 1 scale. To assess the flow for fish lifecycle support, we referred to the actual mullets' biomass sown per hectare

133 (kg ha⁻¹ y⁻¹), referring to the species *Mugil cephalus*, *Chelon labrosus*, *Chelon aurata*, *Chelon saliens*, and
134 *Chelon ramada*, as officially declared by the *valli da pesca* managers. We focused on mullets because they are
135 the only taxonomic group for which data regarding the sowing of juveniles were available and because the
136 origin of their fingerlings was local. In contrast, for other farmed species (for example *Sparus aurata*,
137 *Dicentrarchus labrax*), it is more frequent that fry often come from intensive hatcheries, not necessarily
138 located in Northern Italy. Subsequently, we decided to not take them into account for assessing the lagoon fish
139 lifecycle support ES. For birds lifecycle support flow, we referred to the average number of migratory
140 waterbirds that winter within the *valli da pesca*, from the last ten years' waterbirds annual censuses (AFV
141 2020) The resulting indicators were combined and then normalized on a 0-1 scale.

142

143 2.2.2 Provisioning ESs

144 The aquaculture practiced in the *valli da pesca* is mainly extensive aquaculture. In this type of aquaculture,
145 fish are free to move into a wide brackish basin and rely only on natural food (Costa Pierce 2002). Therefore,
146 aquaculture capacity is represented by the potential biomass hosted by the brackish basins of each *valle da*
147 *pesca* (kg ha⁻¹ y⁻¹), estimated from a spatialized foodweb model (Anelli Monti et al. 2021). The flow is expressed
148 as the average fish catches per hectare of brackish water surfaces per year (kg ha⁻¹ y⁻¹), according to the 2010-
149 2019 official data (Regione Veneto).

150 Waterfowl hunting was evaluated in terms of capacity using the time series of waterbird censuses from 2010
151 to 2019 (Associazione Faunisti Veneti). Since the census data were associated with point features, we
152 interpolated the average number of huntable birds censused in the last ten year by Dirichlet-Voronoi
153 tessellation for implicit surfaces to obtain the most likely spatial distribution of huntable waterbirds per
154 hectare per year. Hunting flow was obtained from waterbird catch historical series, collected from 2010 to
155 2020 (Ente Produttori Selvaggina Veneto).

156 The food production ES focused on wild edible plants of the genus *Salicornia* growing in saltmarshes and
157 on honey obtained from flowers of sea lavender (plants of the genus *Limonium*).

158 Perennial saltmarsh vegetation dominated by halophytic dwarf shrubs has been identified by visual census
159 method in 12 patches of two different *valli da pesca* (Valle Dogà and Valle Cavallino, Figure 1 - 1, 8), and then
160 through the identification of their predictive range in R, G, B bands reflectance values, filtered through a cut-
161 off NDVI value calculated from aerial and satellite images (listed in Supplementary Materials II Tabs. S.M. II
162 - 1, 2). The identified vegetation patches allowed for evaluating the capacity for edible plants, in terms of
163 kilograms of *Salicornia* biomass that could be harvested in the *valli da pesca* per year. Considering the
164 *Limonium* inflorescence cover in the patches (Fantinato and Buffa 2019) and the ratio between the number of
165 sea lavender flowers and grams of honey potentially produced per unit area, the honey capacity was assessed
166 as kilograms of honey that can be produced in the *valli da pesca*. The flow indicator refers to the kilograms of
167 harvested plants and honey. Edible plant harvesting data were retrieved from 2020 market data (Veneto
168 Agricoltura) and interview to local people and restaurant chefs; the amount of sea lavender honey was
169 witnessed by five beekeepers.

170 2.2.3 Cultural ESs

171 Tourism attractiveness was assessed by a survey carried out in the second half of 2019, addressing tourists who
172 recreate and travel in the Venice lagoon.

173 To assess the tourism ES capacity, we mapped for each one of the *valli da pesca* the attractiveness factors
174 evaluated by tourists in the questionnaires, namely saltmarshes presence, the possibility to observe birds and
175 fauna, good water quality, and the chance to contemplate natural terrestrial habitats. Each element was
176 weighted to depict the interest declared by the people and normalized to a 0-1 scale index.

177 Tourism flow was represented by the number of persons who, during a year, had passed at least one night
178 in one of the accommodation facilities within the *valli da pesca*.

179 Regarding the ES information for cognitive development, the natural factors that enhance touristic
180 attractiveness have been mapped along with the inclusivity toward the necessities of people with disabilities.
181 The resulting map was normalized to a 0-1 scale. The flow indicator summarized data concerning the number
182 of persons who annually attend outdoor educational activities or guided one-day trips, as reported by touristic
183 guides and associations (ATN Laguna Sud, Cooperativa Limosa, Ente di promozione turistica di Cavallino
184 Treporti, Oasi WWF Valle Averno).

185 Birdwatching ES capacity was expressed with the birdwatching attractiveness normalized map, based on
186 the factors reported as important by 30 interviewed birdwatchers: the presence of pedestrian paths,
187 saltmarshes presence, nesting areas in the visual field, and a high probability of observing birds. All these
188 factors were summed and scaled to obtain a normalized attractiveness map. Birdwatching ES flow indicator
189 was considered the mean number of active birdwatchers derived from the observers' activity trend, recorded
190 from 2010 to 2020 in the Italian birdwatchers' database (<https://www.ornitho.it/>).

191 *2.2.4 Aggregated ESs indicators*

192 Each ES indicator was spatially assessed in the *valli da pesca* and in the open lagoon, that is all the lagoon
193 surface which is not privately managed. The open lagoon ESs assessment was conducted accordingly with the
194 one presented by Rova, Stocco and Pranovi (2022), and detailed in Supplementary Materials III. Once we
195 obtained the raster layer for each indicator, for both the *valli da pesca* and the open lagoon, the results were
196 normalized to a 0-1 scale. Finally, the indicators were aggregated to have a normalized sum of all the ESs
197 capacities and a normalized sum of all the ESs flows, allowing quantitative spatial comparisons. In particular,
198 we ran the algebraic sum of the normalized raster values through the core zonal statistics plugin in QGIS, by
199 overlapping the polygons of the areas of interest on each ES normalized raster. The results were compared
200 between the *valli da pesca* and the non-managed part of the lagoon and expressed as a percentage of the overall
201 capacity and flow, as well as the percentage of each ES capacity and flow of the *valli da pesca* with respect to
202 the related values found in the open lagoon. All the analyses were performed using the software R 4.1.2 (R Core
203 team, 2021) within the RStudio 2021.09.2 environment (RStudio Team, 2021).

204 **2.3 ESs spatialization**

205 In order to assess all the considered ESs with a spatially explicit approach, remote sensing imageries of the
206 study area were collected and analyzed to obtain the land cover/land use map (LULC) on which ES mapping
207 was based.

208 A machine-learning, scalable gradient-boosting decision trees XgBoost algorithm (Chen and Guestrin
209 2016) was used to classify land cover classes from multispectral and hyperspectral remote sensing data. Very
210 High-Resolution multispectral satellite scenes (VHR), collected by Worldview-02, Worldview-03, and GeoEye-
211 01 constellations, were granted by the European Space Agency Services via the "ESA On-Demand Restricted
212 Data Access program" (see Supplementary Materials II for details).

213 For some uncovered areas, satellite tiles were pan-sharpened and mosaicked in a custom composition of
 214 RGB + NIR raster stack, along with AGEA aerial photograms at 20 cm resolution, granted by Regione Veneto.

215 The classification was performed on the multi-band images through the tuning of the XgBoost algorithm
 216 and was assessed with an accuracy test, resulting in a mean of 96% correct class prediction.

217 The classification and the following spatialization tasks have been performed with the open-source software
 218 QGIS 3.16 (QGIS Association: QGIS Geographic Information System, 2022), R 4.1.2 (R Core, 2022), and
 219 RStudio 2021.09.2 (RStudio Team, 2022).

220 3 Results

221 The assessment results are summarized in Table 2. The spatial distribution of each ES capacity and flow are
 222 illustrated in Supplementary Materials IV.

Ecosystem service	Capacity			Flow		
	Measure unit	Mean ± s.d.	Values range	Measure unit	Mean ± s.d.	Values range
Climate regulation	gC m ⁻² y ⁻¹	60.86 ± 54.60	10.00 ÷ 245.00	gC m ⁻² y ⁻¹	60.86 ± 54.60	10.00 ÷ 245.00
Water purification	% removed nitrogen	9.37 ± 11.67	0.00 ÷ 34.61	% removed nitrogen	9.37 ± 11.67	0.00 ÷ 34.61
Lifecycle support	0-1 scale	0.58 ± 0.23	0.30 ÷ 0.73	0-1 scale	0.31 ± 0.24	0.00 ÷ 1.00
Aquaculture	kg ha ⁻¹ y ⁻¹	60.79 ± 59.99	0.00 ÷ 119.92	kg ha ⁻¹ y ⁻¹	29.83 ± 34.77	0.00 ÷ 159.13
Hunting	n ha ⁻¹ y ⁻¹	29.56 ± 36.56	0.02 ÷ 376.67	n ha ⁻¹ y ⁻¹	6.12 ± 4.77	0.00 ÷ 20.94
Wild edible herbs & honey production	edible plants kg ha ⁻¹ y ⁻¹	0.20 ± 0.37	0.00 ÷ 2.03	edible plants kg ha ⁻¹ y ⁻¹	0.00 ± 0.00	0.00 ÷ 0.00
	honey kg ha ⁻¹ y ⁻¹	0.11 ± 0.01	0.00 ÷ 0.11	honey kg ha ⁻¹ y ⁻¹	0.03 ± 0.10	0.00 ÷ 0.45
Tourism	attractiveness, 0-1 scale	0.49 ± 0.14	0.06 ÷ 1.00	tourists n y ⁻¹	7.86 ± 34.07	0.00 ÷ 20.00
Information for cognitive development	attractiveness, 0-1 scale	0.30 ± 0.12	0.04 ÷ 0.89	students n y ⁻¹	102 ± 470	0.00 ÷ 3640.0
Birdwatching capacity	attractiveness, 0-1 scale	0.33 ± 0.23	0.00 ÷ 1.00	birdwatchers n y ⁻¹	1.47 ± 5.99	0.00 ÷ 38.67

223 **Table 2** Capacity and flow results in the valli da pesca.

224 Overall, in the *valli da pesca* the highest values for carbon sequestration were detected in the areas where
 225 both saltmarshes and seagrasses are present, reaching values up to 245 gCm⁻²y⁻¹. The mean carbon
 226 sequestration has been estimated at 60.86±54.60 gCm⁻²y⁻¹, for a total amount of 2448 tonC y⁻¹, or 8982
 227 tons of CO₂y⁻¹ (Supplementary Materials IV, a). The water purification ES is represented by an average removal
 228 of 9.37% (±11.67%) of the nitrogen loadings; the greater the extent of the brackish lakes in the *valli*, the higher
 229 results the removal capacity. However, it must be noticed that this average estimate may likely be influenced
 230 by the water quality of the river water entering the *valli da pesca* (Salveti et al. 2008; Bettiol et al. 2005) and
 231 must be considered as a first estimate of the average value for this ES.

232 Regarding lifecycle support, the capacity indicator is homogeneously distributed with a mean of 0.58
 233 (±0.23), while the flow indicator shows a mean of 0.31±0.24 (Supplementary Materials IV, c-d).

234 The aquaculture capacity of the study area results in a mean of 60.79±59.99 kg per hectare per year, but
 235 the flow of this ES varies greatly between different *valli da pesca* (Supplementary Materials IV, f) because it is

236 influenced by several factors, both natural and social. Fish production is related to fish sowing, which in turn
237 depends on wild fry availability; moreover, fish sowing is dependent on both fish survival to the previous years
238 and precedent production. Based on the 2014-2019 period, the average total quantity of juveniles sown per
239 year was 6'867'213, with a value per *valle da pesca* ranging between 26'000 and 2'793'033 individuals. On this
240 basis, the mean sowing value was 1'227 (± 958) juveniles per hectare per year. Thus, when faced with an average
241 production of the whole area of 29.83 (± 34.77) kg ha⁻¹ y⁻¹, we must distinguish that the *valli da pesca* carrying
242 out semi-intensive aquaculture show a value of 70.55 \pm 23.78 kg ha⁻¹ y⁻¹ (e.g., *valli da pesca* n. 2, 22, 24), while
243 those practicing only extensive aquaculture drop to a mean of 15.45 \pm 16.35 kg ha⁻¹ y⁻¹.

244 Regarding waterfowl hunting, during the period 2010-2019, about 280'000 huntable waterfowls have been
245 wintering inside the *valli da pesca*, where a yearly average of 29.56 \pm 36.56 huntable waterfowls per hectare
246 were hosted. The highest number of waterfowls was consistently recorded during 2010-2019 in the *valli da*
247 *pesca* n. 1, 7, 8, 19, 25, 26, and 27, while in the *valli da pesca* from n. 10 to n. 15, most exposed to nautical and
248 tourist traffic, the number of censused waterfowls was lower (Supplementary Materials IV, g).

249 Regarding the hunting flow, in 2010-2019 the average catch was estimated at 38'404 (± 7299) waterfowls per
250 year, considering the total surface of the *valli da pesca*. On average, 6.12 \pm 4.77 waterbirds ha⁻¹ y⁻¹ have been
251 caught in the considered period. These catches were homogeneously distributed, with a maximum value of 20
252 catches ha⁻¹ y⁻¹ in the most confined *valli da pesca* 3, 4, 5, 6, 26, 27 (Supplementary Materials IV, h).

253 Although a homogeneous potential for provisioning edible wild plants and honey has been estimated in the
254 *valli* area, harvesting occurs almost exclusively in the saltmarshes located in the open part of the lagoon, except
255 from three *valli da pesca* – n. 5, 6, 8 – which host beehives (Supplementary Materials IV, j). Harvesting of
256 *Salicornia* and other edible herbs has not been reported in the *valli da pesca* saltmarshes, apart from tiny
257 quantities in occasional occurrences.

258 Tourism capacity results in all the areas with a mean attractiveness of 0.49 \pm 0.14 (Supplementary Materials
259 IV, k); nevertheless, very few of them are open for touristic trips, and not even regularly, which restricts the
260 flow for tourism as well as for other cultural ESs (Supplementary Materials IV, l). The flow of tourists is limited
261 to few *valli da pesca* in the Northern lagoon, involving Valle Lio Maggiore (5), Valle Cavallino (8), Valle
262 Falconera (9), Valle Sacchetta (15), and Santa Cristina (17).

263 The capacity and flow for the ES information for cognitive development cover an even narrower extent:
264 only the WWF Oasis Valle Averno (20), Valle Olivara (11), and Valle Liona (12) have a capacity higher than 0.5
265 (Supplementary Materials IV, m), but it resulted that the 3'154 students per year, who visit the *valli da pesca*
266 for educational purposes, take part into guided tours in *valli da pesca* n. 12, 20, 23, 24 (Supplementary
267 Materials IV, l).

268 Finally, the overall attractiveness for birdwatching is evenly distributed (Supplementary Materials IV, o)
269 but the ES flow is not homogeneous and confirms that the Northern part of the lagoon is the most frequented
270 for such outdoor activities, with 72 active birdwatchers – who spend a mean of 84.41 \pm 21.36 hours per year
271 birdwatching – converging in the nearing of the area n. 5, 11, 12 (Supplementary Materials IV, p).

272 Table 3 shows how much the total ESs capacity and flow assessed in all the *valli da pesca* is comparable to
273 that observed in the open, not privately managed part of the lagoon. The most important contributions from
274 the *valli da pesca* to the ESs supply of the open lagoon are noticeable for lifecycle support, aquaculture,
275 hunting, information for cognitive development, and birdwatching.

276


Ecosystem service	% vs. lagoon capacity	% vs. lagoon flow
Climate regulation	12.79	12.79
Water purification	12.50	12.50
Lifecycle support	23.93	25.78
Aquaculture	12.79	31.68
Hunting	99.8	65.76
Wild hedible herbs	6.46	0.00
Honey production	83.48	1.82
Tourism	27.22	0.01
Information for cognitive development	25.97	23.88
Birdwatching	21.57	13.66

Table 3 Proportion of capacity and flow assessed in the valli da pesca compared to each ES capacity and flow in the open lagoon.

277 **3.4 Aggregated ESs indexes analyses**

278 Figure 2 and figure 3 show the normalized maps for the aggregated indicators of all the ESs capacity and the
 279 flow intensities, respectively. The sum of all the capacities shows that the highest values have been registered
 280 inside the valli da pesca which maximize both aquaculture and hunting (Fig. 2 n. 1, 2, 4, 26, 27).


281 As regards the sum of all the flows, the highest values have been recorded especially where provisioning
 282 and cultural ESs flows are the highest as well, e.g., Valle Grassabò, Valle Lio Maggiore, and Valle Dragojesolo
 283 (Fig. 3 n. 2, 5, 7).



284 **Figure 2** Spatialized total capacity indicator, as the sum of all the ESs' capacities

285

286



287

288


Figure 3 Spatialized total flow indicator, as the sum of all the ESs' flows

289

290

291

In general, the extension of the area where the sum of all the ESs' capacity is higher than the sum of all the ESs' flow is 71.1 km². In contrast, the area where the sum of all the assessed ESs' flow results greater than the sum of all the ESs' capacity is 23.4 km² (Fig. 4)



292

293

294

Figure 4 Spatialized capacity – flow difference (C-F). Green = C > F; red = C < F.

Discussion

296

297

298

299

300

301

302

303

The lagoon of Venice is a complex social-ecological system with a historical link between natural habitats and human activities, which is particularly strong for the *valli da pesca*. Despite being “artificially maintained”, the *valli da pesca* conserved the typical features and related ecological processes of transitional environments that the open lagoon has progressively lost, especially due to erosion and relative sea level rise (Day et al. 1998; Madricardo and Donnici 2014). For these reasons, it would be interesting to analyze the possible role of the *valli da pesca* in terms of ESs.

The ESs approach has already proven to be effective in visualizing and assessing the contribution to people of similar man-managed ecosystems, and in suggesting sustainable management strategies as well (Gaglio et

304 al. 2019; Aschonitis et al. 2016; Weitzman 2019; Walton et al. 2015). In this study, we proposed for the first
305 time the assessment of multiple ESs in the *valli da pesca* of the Venice lagoon to better understand their
306 possible ecological role within the context of the entire lagoon environment. Obtained results highlighted a
307 non-negligible contribution of the *valli da pesca* to the entire supply of ESs of the Venice lagoon.

308 Regarding carbon sequestration, for instance, about 12% of the CO₂ sequestered every year by the lagoon is
309 due to saltmarshes and seagrasses located inside the *valli da pesca*, where these habitats are preserved by
310 different impacts, erosion, and trampling. Also, the mean water purification through denitrification is higher
311 inside the boundaries of the *valli da pesca* than in the outer part of the lagoon. Indeed, it has been estimated
312 that the open lagoon can remove about 10.08% of nitrogen loadings (Rova et al. 2022) but no area reaches the
313 maximum values recorded in some lakes located inside the *valli da pesca*. These findings are in accordance
314 with those of other studies that have focused on water management for extensive aquaculture (Walton, Vilas,
315 Coccia, et al. 2015; Gamito 1997). This result, however, should inspire further research. Indeed, extensive
316 aquaculture is today prevalent in *the valli da pesca*, meaning that fish feed on the available food they naturally
317 find in the *valle da pesca* without the need to add feed artificially. In addition, the density of fish is such that
318 the brackish lake ecological processes can process naturally dead organic matter and nutrient loadings,
319 avoiding the risk of eutrophication (Anras et al. 2010; Costa Pierce 2002). If things are going to change, the
320 denitrification process would also be influenced by several factors, including the water quality of the rivers
321 entering the *valli da pesca*, and the organic matter loading that can increase in response to eventual higher
322 densities of fish and waterbirds inside the lakes. Thus, the denitrification process is worthy of further
323 experimental field studies. New data and models about this aspect, along with an exploration of the most likely
324 future trends, may prove extremely useful to better depict this regulating ES not only at the local scale but also
325 for other similar Heavily Modified Water Bodies, such as in the Po River Delta, in the Marano lagoon or other
326 areas along the Adriatic coasts.

327 The most confined parts of the lagoon were identified centuries ago as the most suitable places for fish
328 farming and hunting purposes, as confirmed more recently for fish (Cavraro et al. 2017; Brigolin et al. 2014)
329 and waterfowl (Arzel et al., 2006; Scarton and Bon, 2009; Guillemain et al., 2013). Within this context, the ES
330 lifecycle support provided by the *valli da pesca* turned out to be quite significant compared to the open lagoon,
331 confirming the important ecological role still played by these areas despite the presence of levees and
332 embankments. However, the naturally recruited fish is not sufficient for meeting the *valli* requirements, to the
333 point that today the aquaculture ES in the *valli da pesca* of the Venice lagoon depends on personal, social, and
334 economic factors, which lead the owner to decide whether and how much fish to sow. As a consequence, the
335 practice of fishing wild fry in the lagoon and then sowing them inside the *valli da pesca* is the only method that
336 grants the biomass actually hosted by each area. From such a point of view, besides enhancing aquaculture
337 productivity, this tradition could be seen as a good practice, capable of maintaining some species' presence in
338 the long run. On the one hand, fry fishery is strictly regulated and controlled, being entrusted to only a very
339 few licensed operators who annually update the quota that can be caught in order not to affect lagoon fish
340 population dynamics (Fortibuoni et al. 2014; Granzotto et al. 2001).

341 On the other hand, since fish migration and the recruitment processes taking place in the open lagoon play a
342 fundamental role in sustaining fish lifecycle (Lanzoni et al. 2021; Scapin et al. 2022; Cavraro et al. 2017) and
343 the aquaculture ES in the *valli da pesca*, we could also affirm that the aquaculture ES in the *valli da pesca*
344 could positively affect the regulating ES of lifecycle support in the entire lagoon. As happens for waterbirds,
345 the persistence of suitable habitats in the *valli da pesca* depends on human maintenance, which in turn

346 depends on economic resources from the provisioning ESs incomes. This generates positive feedback that
347 raises considerations about how a provisioning ES can also support other ESs belonging to a different category,
348 in agreement with the findings of other authors (Liquete et al. 2016; Grizzetti et al. 2019)

349 The extensive aquaculture practiced today in the *valli da pesca*, moreover, can be considered quite sustainable,
350 considering that the production value of 60 kg ha⁻¹ y⁻¹ falls in the range of 45 - 110 kg ha⁻¹ y⁻¹ reported by
351 (Koutrakis et al. 2007) for extensive aquaculture in Mediterranean lagoon, but is quite lower than 150 kg ha⁻¹
352 y⁻¹ expected for a desirable production in a typical Italian *valle da pesca*, according to Ravagnan (1982) and
353 Shang (1982).

354 Since the aquaculture ES in the *valli da pesca* is getting less worthwhile in the last decade, as shown by the
355 data, other activities have started to play a key role. Among them, the most important is waterfowls hunting,
356 which also depends on both natural and anthropogenic factors. To say to what extent the great capacity of bird
357 attraction is granted by the *valli da pesca* environment itself, or whether it is due to management choices, is
358 difficult. Indeed, to attract waterfowl and increase hunting opportunities the managers regulate water regimes,
359 construct mild sloping islands, take care of windbreak hedges made of reeds and tamarisks, and often spread
360 millet or grains.

361 The management which aims to maximize hunting ES is possibly influencing the capacity and the flow of
362 this ES not only inside, but also outside the boundaries of the *valli da pesca*. Our assessments found that most
363 huntable waterfowl censused in the lagoon are reaching for the *valli da pesca* during their migration, where
364 they rest, feed, and breed as well (at least some species). Such evidence is in accordance with other authors,
365 who reported about the high importance of the *valli da pesca* for waterbird populations, whereas the lagoon is
366 losing suitable habitats (Scarton 2017; AFV 2020) Also, the waterfowls hunting flow in the lagoon results
367 strongly related, for timing and catches as well, to the flow happening inside the *valli*. As a comparison, the
368 hunting flow in the lagoon is estimated from 12'635 to 12'770 catches per year and takes place just in proximity
369 of the allowed hunting blinds, where the catches have a mean of 4.44 ± 1.56 per hectare per year. Even if the
370 distribution of this flow is homogeneous in the whole lagoon, the highest values have been detected in the
371 blinds between contiguous *valli da pesca*, in the North-Eastern and Southern parts of the lagoon. Indeed,
372 waterfowl hosted inside the *valli da pesca*, while moving across different *valli da pesca* or from a *valle da pesca*
373 to another place just outside, represent the "stock" on which not only the *valli da pesca* hunters, but also the
374 "open lagoon" hunters rely.

375 Another example of driving relationships between different ESs is the substantial contributions of the *valli da*
376 *pesca* to the ESs that depend on the consistency of the birds presence. For example, the management which
377 aims to attract the highest possible number of huntable waterfowls is possibly also dragging the attractiveness
378 for cultural ESs, especially for birdwatching. Accordingly, many aquatic birds are attracted into the *valli da*
379 *pesca* managed environment and this also can increase the overall capacity, as bird movements between the
380 *valli da pesca* and the open lagoon create a positive side-effect in increasing attractiveness to birdwatchers,
381 tourists, and hikers even outside the *valli da pesca* boundaries.

382 However, despite the high overall capacity for all cultural ESs in the *valli da pesca*, the flow is strongly shaped
383 by accessibility and inclusiveness, ending up with flow occurring just on a few areas. Nevertheless, this
384 restricted flow should not be read as totally negative: the *valli da pesca* are seen as extremely attractive
385 especially for the preservation of lagoon landscapes and fauna in a quiet and uncrowded space. This
386 consideration should make one think carefully about whether, how, and how often the *valli da pesca* could be
387 open for touristic and educational visits. Similar considerations have also been aroused on different

388 environments by other authors (Villamagna et al., 2015; Drescher et al., 2017; Drescher and Brenner, 2018),
389 who proposed to give more support to private natural areas for enhancing conservation and educational
390 purposes, provided that excursionists' entrances must be controlled.

391 Management initiatives that also take these perspectives into account should seek to harmonize with the
392 current situation of the *valli da pesca*, seeing that the difference between capacity and flow aggregated indexes
393 shows that the majority of the *valli da pesca* display a total capacity higher than the total flow (figs. 2, 3, 4).
394 Just a few zones with the highest flows of mediated ESs (*sensu* Rova and Pranovi 2017) stand out amongst
395 other areas: in fact, some *valli da pesca* where hunting ES' flow is, on average, more important than the
396 aquaculture production display an aggregated flow index slightly higher than the ESs' capacity. This may be
397 because waterfowls hunting relies on a moving resource so that the catches in one *valle da pesca* can affect the
398 waterfowls which have been censused in another *valle* and, consequently, represent the capacity assessed on a
399 different place, but not strictly linked to it.

400 The analysis of the contribution of the *valli da pesca* sub-system to the whole transitional ecosystem confirms
401 that the *valli da pesca* are highly significant to the whole Venice lagoon. The comparison of the ESs' aggregated
402 index between the open part of the lagoon and the *valli da pesca* brings out that the latter, despite covering
403 just 17% of the total lagoon, play an important role in terms of ESs contributions: 38% of the capacity and 24%
404 of the flow in comparison to the open lagoon capacity and flow, respectively.

405 This non-negligible contribution to ESs, habitats, and landscape features conservation, makes the *valli da*
406 *pesca* a good example of effective ecosystem-based management of Heavily Modified Water Bodies,
407 representing a good example to set up sustainable management for similar productive areas in the
408 Mediterranean.

409 Like little lagoons that are artificially managed, the *valli da pesca* are regulated by a private regime that, on
410 the one hand, denies the access and minimizes cultural ESs flow, but on the other hand, conserves the
411 landscape, the ecological processes and ESs provided by them. The open lagoon receives from them the benefits
412 of a spill-over effect, especially for lifecycles support, aquaculture, hunting, and birdwatching ESs. For these
413 ESs, the *valli da pesca* provide ESs that benefit a wider community, spread into a larger spatial scale than that
414 the ES are generated at. In addition, the *valli da pesca* act as a buffer area between the land and the open
415 lagoon, in particular for climate regulation and water purification, as the results of carbon sequestration and
416 nitrogen removal show.

417 The role of *valli da pesca* as buffers and as conservation opportunities also leads us to consider how
418 important it might be to restore a buffer strip between the land and the lagoon, which has been lost in some
419 areas. As the *valli da pesca* proved to be effective in maintaining saltmarshes, reed meadows and ecological
420 gradients, similarly it might be useful to consider ecological restoration efforts in areas that are most affected
421 by the transfiguration of the original lagoon landscape (Feola et al., 2022).

422 Since the connection between the lagoon and *valli da pesca* persists, we must consider the possible
423 evolution of such interconnected systems as a whole. The lagoon now subjected to manifold pressures from
424 anthropic activities (Fortibuoni et al. 2015; Zucchetto et al. 2021; Anelli Monti et al. 2021; Solidoro et al. 2010)
425 that are likely going to increase (Jennerjahn and Mitchell 2013). The adaptations to such impacts, as well as to
426 climate change and sea-level rise effects, in the long run, might modify the lagoon hydrodynamics and the sea-
427 lagoon connectivity, threatening the *valli da pesca* and so their ESs supply (Solidoro et al. 2010; Cristiano and
428 Gonella 2020).

429 Eventually, we suggest that decisions about the MOSE barrier functioning must be well pondered to not put
430 the *valli da pesca* at risk of being endangered. Considering that the *valli da pesca* are continuing to exchange
431 energy and matter with the lagoon, such equilibria can be at risk in the context of a “regulated lagoon”. The
432 MOSE system (with its mobile barriers at the inlet) controls the marine water flow to mitigating the effect of
433 climate change and sea-level rise on cultural heritage, but could conversely threaten the daily exchange of water
434 in the most confined part of the lagoon, and the *valli* waterfronts as a consequence.
435 Indeed, it is very important to preserve the cultural heritage of the historic center of Venice and the islands so
436 as not to lose cultural ESs and economic activities flowing from them. Nevertheless, we are facing the
437 possibility that the lagoon will remain closed for longer periods of time. According to Umgiesser (2020) and
438 Lionello et al. (2021), the period of closure of the mobile barriers is very likely to grow up to 3 weeks per year
439 by the end of 2050, and up to 2 months per year by 2080. This would mean affecting the sea-lagoon
440 connectivity, the lagoon hydrodynamics, and landscape evolution patterns, as several studies have already
441 argued (Ghezzi et al. 2010; 2011; Pérez-Ruzafa et al. 2019). Such a perspective could put the *valli da pesca* at
442 risk: lower connectivity with the sea can affect the fish fry availability for the aquaculture ES even more than
443 today. In addition, a lower frequency of exchanges through the inlets is expected to worsen the water quality,
444 nutrient concentration (Solidoro et al. 2005), and microcirculation (Ghezzi et al. 2010) involved in the
445 processes on which the regulating ESs are based.

446 In light of our results, it must be carefully considered that losing the *valli da pesca* might represent an issue
447 for the entire community because of their contribution to conserving habitats and providing regulating,
448 provisioning, and cultural ESs. This trade-off perspective takes great importance, especially in the light of
449 future scenarios, including climate change, subsidence, and management of health problems, which claim a
450 responsible attendance of the lagoon.

451 Conclusions

452 This work has shown, by using quantitative indicators, the spatial distribution of ESs capacity and flow within
453 the managed *valli da pesca* of the lagoon of Venice. The multiple ESs approach enhanced the areas where
454 capacity and flow have the highest values. It also revealed that the capacity is not always and everywhere fully
455 granted to translate itself in a flow towards the local population due to different accessibility and availability
456 of some places. Therefore, the comparison between different areas of the same wide ecosystem highlights that,
457 although the capacity of the provided ESs tends to be overall high, a multiplicity of social and behavioral factors
458 influence the relative flows.

459 In this context, confirmation of the usefulness of assessing the value of ESs in ecosystem management is
460 given (Costanza 2006), especially when a trade-off must be accomplished between the moral duty to preserve
461 and emphasize regulation and support services, and the need to build on the economic revenues that come
462 from provisioning ESs. This dynamical network stimulates to continue the research with a modeling approach,
463 which could consider behavioral, social, and economic factors that act as fulcrums for decision-making.

464 Increasing the knowledge about the *valli da pesca* and other similar socio-ecological systems can help to
465 enlighten not only the relationships between ESs, landscape, and management choices, but also to rethink the
466 interaction between the environment, the society, and the institutions involved in the Venice lagoon ecosystem.

467 The assessment of ESs, such as lifecycle support, hunting, and birdwatching, showed that the open side of
468 the lagoon is often strictly related to the management of the most confined areas, corroborating the hypothesis

469 that the *valli da pesca* play an important role in defining the ESs pattern of the whole lagoon of Venice. Indeed,
470 the *valli da pesca* represent a buffer between the aquatic ecosystem of the lagoon and its land boundaries,
471 mitigating anthropogenic pressures; in addition, their provisioning and cultural ESs also show a series of
472 positive side-effects in the open lagoon.

473 Thus, it is of great importance to take them into account to fully comprehend the Venice lagoon ecosystem,
474 especially when investigating the effects under different health, social and economic scenarios, as well as when
475 facing climate change and sea-level rise.

476 Acknowledgments

477 This scientific work has been developed in the context of the “Venezia 2021” project, with the contribution of
478 Provveditorato Interregionale Opere Pubbliche per il Veneto, Trentino Alto Adige e Friuli Venezia Giulia
479 through Consorzio Venezia Nuova and coordinated by CORILA.

480 We acknowledge the owners and the operators of the *valli da pesca* for providing data and explanations, the
481 Regione Veneto, the Associazione Faunisti Veneti, the EPS, and the local hunting associations.

482 We thank the European Space Agency for granting the VHR satellite imageries for this research work.

483 We thank Marco Anelli Monti for providing the food web model outputs, as well as Matteo Zuchetta and
484 Alberto Caccin for their helpful suggestions.

485 We are grateful to the reviewers for their kind comments, which helped to improve the manuscript.

References

- 487 AFV, Associazione Faunisti Veneti. 2020. ‘CENSIMENTO DEGLI UCCELLI ACQUATICI SVERNANTI IN
488 PROVINCIA DI VENEZIA’.
- 489 Anelli Monti, M., D. Brigolin, P. Franzoi, S. Libralato, R. Pastres, C. Solidoro, M. Zucchetta, and F. Pranovi. 2021a.
490 ‘Ecosystem Functioning and Ecological Status in the Venice Lagoon, Which Relationships?’ *Ecological Indicators*
491 133 (December): 108461. <https://doi.org/10.1016/j.ecolind.2021.108461>.
- 492 Anras, L., Clara Boglione, Stefano Cataudella, M. T. Dinis, Pavlos Makridis, Giovanna Marino, Ana Ramalho Ribeiro,
493 and Manuel Yúfera. 2010. ‘The Current Status of Extensive and Semi-Intensive Aquaculture Practices in Southern
494 Europe’. *Aquaculture Europe* 35 (2): 12–16. <https://digital.csic.es/handle/10261/50633>.
- 495 Arzel, Celine, J. Elmerberg, and M. Guillemain. 2006. ‘Ecology of Spring-Migrating Anatidae: A Review’. *Journal of*
496 *Ornithology*. Springer. <https://doi.org/10.1007/s10336-006-0054-8>.
- 497 Aschonitis, Vassilis G., Mattias Gaglio, Elisa A. Fano, Elena Gissi, Giuseppe Castaldelli, Vassilis G. Aschonitis, Elena
498 Gissi, Giuseppe Castaldelli, and Elisa A. Fano. 2016. ‘Land Use Change Effects on Ecosystem Services of River
499 Deltas and Coastal Wetlands: Case Study in Volano–Mesola–Goro in Po River Delta (Italy)’. *Wetlands Ecology*
500 *and Management*. <https://doi.org/10.1007/s11273-016-9503-1>.
- 501 Baró, Francesc, Ignacio Palomo, Grazia Zulian, Pilar Vizcaino, Dagmar Haase, and Erik Gómez-Baggethun. 2016.
502 ‘Mapping Ecosystem Service Capacity, Flow and Demand for Landscape and Urban Planning: A Case Study in the
503 Barcelona Metropolitan Region’. *Land Use Policy* 57 (November): 405–17.
504 <https://doi.org/10.1016/j.landusepol.2016.06.006>.
- 505 Bettiol, Cinzia, Flaviano Collavini, Stefano Guerzoni, Emanuela Molinaroli, Paolo Rossini, Luca Zaggia, and Roberto
506 Zonta. 2005. ‘Atmospheric and Riverine Inputs of Metals, Nutrients and Persistent Organic Pollutants into the
507 Lagoon of Venice’. *Hydrobiologia* 2005 550:1 550 (1): 151–65. <https://doi.org/10.1007/S10750-005-4372-2>.
- 508 Brigolin, Daniele, Chiara Facca, Anita Franco, Piero Franzoi, Roberto Pastres, Adriano Sfriso, Marco Sigovini, et al.
509 2014. ‘Linking Food Web Functioning and Habitat Diversity for an Ecosystem Based Management: A
510 Mediterranean Lagoon Case-Study’. *Marine Environmental Research* 97: 58–66.
511 <https://doi.org/10.1016/j.marenvres.2014.02.006>.
- 512 Burkhard, Benjamin, and Joachim Maes. 2017. ‘Mapping Ecosystem Services’. *Advanced Books* 1 (March): e12837.
513 <https://doi.org/10.3897/AB.E12837>.
- 514 Cavarro, Francesco, Matteo Zucchetta, Stefano Malavasi, and Piero Franzoi. 2017. ‘Small Creeks in a Big Lagoon: The
515 Importance of Marginal Habitats for Fish Populations’. *Ecological Engineering* 99 (February): 228–37.
516 <https://doi.org/10.1016/j.ecoleng.2016.11.045>.
- 517 Chen, Tianqi, and Carlos Guestrin. 2016. ‘XGBoost’. In *Proceedings of the 22nd ACM SIGKDD International*
518 *Conference on Knowledge Discovery and Data Mining*, 13-17-Aug:785–94. New York, NY, USA: ACM.
519 <https://doi.org/10.1145/2939672.2939785>.
- 520 Costa Pierce, Barry A. 2002. ‘Ecological Aquaculture’. *Ecological Aquaculture*, April.
521 <https://doi.org/10.1002/9780470995051>.
- 522 Costanza, Robert. 2006. ‘Nature: Ecosystems without Commodifying Them’. *Nature* 2006 443:7113 443 (7113): 749–
523 749. <https://doi.org/10.1038/443749b>.
- 524 Costanza, Robert, Ralph D’Arge, Rudolf de Groot, Stephen Farber, Monica Grasso, Bruce Hannon, Karin Limburg, et al.
525 1997. ‘The Value of the World’s Ecosystem Services and Natural Capital’. *Nature* 387 (6630): 253–60.
526 <https://doi.org/10.1038/387253a0>.
- 527 Cristiano, Silvio, and Francesco Gonella. 2020. ‘“Kill Venice”: A Systems Thinking Conceptualisation of Urban Life,
528 Economy, and Resilience in Tourist Cities’. *Humanities and Social Sciences Communications* 2020 7:1 7 (1): 1–
529 13. <https://doi.org/10.1057/s41599-020-00640-6>.
- 530 D’alpaos, Chiara, and Andrea D’alpaos. 2021. ‘The Valuation of Ecosystem Services in the Venice Lagoon: A
531 Multicriteria Approach’. *Sustainability* 2021, Vol. 13, Page 9485 13 (17): 9485.
532 <https://doi.org/10.3390/SU13179485>.

- 533 Day, J. W., A. Rismondo, F. Scarton, D. Are, and G. Cecconi. 1998. 'Relative Sea Level Rise and Venice Lagoon
534 Wetlands'. *Journal of Coastal Conservation* 4 (1): 27–34. <https://doi.org/10.1007/BF02806486>.
- 535 European Commission. 2000. 'Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000
536 Establishing a Framework for Community Action in the Field of Water Policy'. *Official Journal of the European
537 Communities* L327.
- 538 Fantinato, Edy, and Gabriella Buffa. 2019. 'Animal-Mediated Interactions for Pollination in Saltmarsh Communities'.
539 *Plant Sociology* 56 (2): 35–42. <https://doi.org/10.7338/pls2019562/02>.
- 540 Feola, Alessandra, Emanuele Ponis, Michele Cornello, Rossella Boscolo Brusà, Federica Cacciatore, Federica Oselladore,
541 Bruno Matticchio, et al. 2022. 'An Integrated Approach for Evaluating the Restoration of the Salinity Gradient in
542 Transitional Waters: Monitoring and Numerical Modeling in the Life Lagoon Refresh Case Study'. *Environments*
543 2022, Vol. 9, Page 31 9 (3): 31. <https://doi.org/10.3390/ENVIRONMENTS9030031>.
- 544 Fortibuoni, T., R. Gertwagen, O. Giovanardi, and S. Raicevich. 2014. 'The Progressive Deregulation of Fishery
545 Management in the Venetian Lagoon after the Fall of the Repubblica Serenissima: Food for Thought on
546 Sustainability'. *Global Bioethics* 25 (1): 42–55. <https://doi.org/10.1080/11287462.2014.894707>.
- 547 Fortibuoni, Tomaso, Federica Aldighieri, Otello Giovanardi, Fabio Pranovi, and Matteo Zucchetta. 2015. 'Climate Impact
548 on Italian Fisheries (Mediterranean Sea)'. *Regional Environmental Change* 15 (5): 931–37.
- 549 Gaglio, Mattias, Mattia Lanzoni, Giovanni Nobili, Diego Viviani, Giuseppe Castaldelli, and Elisa Anna Fano. 2019.
550 'Ecosystem Services Approach for Sustainable Governance in a Brackish Water Lagoon Used for Aquaculture'.
551 *Journal of Environmental Planning and Management* 62 (9): 1501–24.
552 <https://doi.org/10.1080/09640568.2019.1581602>.
- 553 Gamito, S. 1997. 'Sustainable Management of a Coastal Lagoonal System (Ria Formosa, Portugal): An Ecological Model
554 for Extensive Aquaculture'. *International Journal of Salt Lake Research* 1997 6:2 6 (2): 145–73.
555 <https://doi.org/10.1007/BF02441891>.
- 556 Ghezzi, Michol, Stefano Guerzoni, Andrea Cucco, and Georg Umgiesser. 2010. 'Changes in Venice Lagoon Dynamics
557 Due to Construction of Mobile Barriers'. *Coastal Engineering* 57 (7): 694–708.
558 <https://doi.org/10.1016/J.COASTALENG.2010.02.009>.
- 559 Ghezzi, Michol, Alessandro Sarretta, Marco Sigovini, Stefano Guerzoni, Davide Tagliapietra, and Georg Umgiesser.
560 2011. 'Modeling the Inter-Annual Variability of Salinity in the Lagoon of Venice in Relation to the Water
561 Framework Directive Typologies'. *Ocean and Coastal Management* 54 (9): 706–19.
562 <https://doi.org/10.1016/j.ocecoaman.2011.06.007>.
- 563 Granzotto, A, P Franzoi, A Longo, and F Pranovi P Torricelli. 2001. 'La Pesca Nella Laguna Di Venezia : Un Percorso
564 Di Sostenibilità Nel Recupero Delle Tradizioni Lo Stato Dell ' Arte'.
- 565 Grizzetti, B., C. Liqueste, A. Pistocchi, O. Vigiak, G. Zulian, F. Bouraoui, A. de Roo, and A. C. Cardoso. 2019.
566 'Relationship between Ecological Condition and Ecosystem Services in European Rivers, Lakes and Coastal
567 Waters'. *Science of the Total Environment* 671 (June): 452–65.
568 <https://doi.org/10.1016/J.SCITOTENV.2019.03.155>.
- 569 Guillemain, Matthieu, Hannu Pöysä, Anthony D. Fox, Céline Arzel, Lisa Dessborn, Johan Ekroos, Gunnar Gunnarsson,
570 et al. 2013. 'Effects of Climate Change on European Ducks: What Do We Know and What Do We Need to Know?'
571 *Wildlife Biology* 19 (4): 404–19. <https://doi.org/10.2981/12-118>.
- 572 Hatziiordanou, Lena, Eleni Fitoka, Elena Hadjicharalampous, Nefta Eleftheria Votsi, Dimitris Palaskas, and Dania Abdul
573 Malak. 2019. 'Indicators for Mapping and Assessment of Ecosystem Condition and of the Ecosystem Service
574 Habitat Maintenance in Support of the EU Biodiversity Strategy to 2020'. *One Ecosystem* 4.
575 <https://doi.org/10.3897/oneeco.4.e32704>.
- 576 Havera, S. P., Lonie R. Boens, M. M. Georgi, and R. Shealy. 1992. 'HUMAN DISTURBANCE OF WATERFOWL ON
577 KEOKUK POOL'. *Undefined*.
- 578 Hein, Lars, Kris van Koppen, Rudolf S. de Groot, and Ekko C. van Ierland. 2006a. 'Spatial Scales, Stakeholders and the
579 Valuation of Ecosystem Services'. *Ecological Economics* 57 (2): 209–28.
580 <https://doi.org/10.1016/J.ECOLECON.2005.04.005>.

- 581 Jennerjahn, Tim C., and Steve B. Mitchell. 2013. 'Pressures, Stresses, Shocks and Trends in Estuarine Ecosystems – An
582 Introduction and Synthesis'. *Estuarine, Coastal and Shelf Science* 130 (September): 1–8.
583 <https://doi.org/10.1016/J.ECSS.2013.07.008>.
- 584 Korschgen, C., L. George, and W. L. Green. 1985. 'Disturbance of Diving Ducks by Boaters on a Migrational Staging
585 Area'. *Undefined*.
- 586 Koutrakis, E.T., A. Conides, A.C. Parpoura, E.H. van Ham, G. Katselis, and & C. Koutsikopoulos. 2007. 'Lagoon
587 Fisheries Resources in Hellas Alexis Conides Hellenic Centre for Marine Research'.
- 588 Lanzoni, Mattia, Mattias Gaglio, Anna Gavioli, Elisa Anna Fano, and Giuseppe Castaldelli. 2021. 'Seasonal Variation of
589 Functional Traits in the Fish Community in a Brackish Lagoon of the Po River Delta (Northern Italy)'.
590 <https://doi.org/10.3390/w13050679>.
- 591 Lionello, Piero, Robert J. Nicholls, Georg Umgiesser, and Davide Zanchettin. 2021. 'Venice Flooding and Sea Level:
592 Past Evolution, Present Issues, and Future Projections (Introduction to the Special Issue)'. *Natural Hazards and
593 Earth System Sciences* 21 (8): 2633–41. <https://doi.org/10.5194/NHESS-21-2633-2021>.
- 594 Liqueste, Camino, Núria Cid, Denis Lanzanova, Bruna Grizzetti, and Arnaud Reynaud. 2016. 'Perspectives on the Link
595 between Ecosystem Services and Biodiversity: The Assessment of the Nursery Function'. *Ecological Indicators* 63
596 (April): 249–57. <https://doi.org/10.1016/j.ecolind.2015.11.058>.
- 597 Liu, Jianguo, Thomas Dietz, Stephen R. Carpenter, Marina Alberti, Carl Folke, Emilio Moran, Alice N. Pell, et al. 2007.
598 'Complexity of Coupled Human and Natural Systems'. *Science*. American Association for the Advancement of
599 Science. <https://doi.org/10.1126/science.1144004>.
- 600 Madricardo, Fantina, and S. Donnici. 2014. 'Mapping Past and Recent Landscape Modifications in the Lagoon of Venice
601 through Geophysical Surveys and Historical Maps'. *Anthropocene* 6 (June): 86–96.
602 <https://doi.org/10.1016/J.ANCENE.2014.11.001>.
- 603 MEA. 2005. 'MILLENNIUM ECOSYSTEM ASSESSMENT (PROGRAM). (2005). Ecosystems and Human Well-
604 Being.' Washington, D.C.
- 605 Newton, Alice, Ana C. Brito, John D. Icelly, Valérie Derolez, Inês Clara, Stewart Angus, Gerald Schernewski, et al. 2018.
606 'Assessing, Quantifying and Valuing the Ecosystem Services of Coastal Lagoons'. *Journal for Nature
607 Conservation*, 1 July 2018. <https://doi.org/10.1016/j.jnc.2018.02.009>.
- 608 Notte, Alessandra la, Margherita Turvani, and Sergio Giaccaria. 2017. 'Economic Valuation of Ecosystem Services at
609 Local Level for Policy Makers and Planners. The Case of the Island of St. Erasmo in the Lagoon of Venice'.
610 *Environmental Economics* 2 (3). [https://www.businessperspectives.org/index.php/journals/environmental-
611 economics/issue-212/economic-valuation-of-ecosystem-services-at-local-level-for-policy-makers-and-planners-
612 the-case-of-the-island-of-st-erasmo-in-the-lagoon-of-venice](https://www.businessperspectives.org/index.php/journals/environmental-economics/issue-212/economic-valuation-of-ecosystem-services-at-local-level-for-policy-makers-and-planners-the-case-of-the-island-of-st-erasmo-in-the-lagoon-of-venice).
- 613 Pérez-Ruzafa, Angel, Francesca de Pascalis, Michol Ghezze, Jhoni Ismael Quispe-Becerra, Raquel Hernández-García,
614 Irene Muñoz, Carlos Vergara, Isabel María Pérez-Ruzafa, Georg Umgiesser, and Concepción Marcos. 2019.
615 'Connectivity between Coastal Lagoons and Sea: Asymmetrical Effects on Assemblages' and Populations'
616 Structure'. *Estuarine, Coastal and Shelf Science* 216 (January): 171–86.
617 <https://doi.org/10.1016/J.ECSS.2018.02.031>.
- 618 QGIS Association: QGIS Geographic Information System. 2022. 'QGIS'. 2022. <http://www.qgis.org>.
- 619 R Core team. 2022. 'R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing,
620 Vienna, Austria.' 2022. <https://www.r-project.org/>.
- 621 Ravagnan, Gino. 1982. 'Vallicoltura Moderna. Proposte Operative per La Ristrutturazione e Lo Sviluppo Della Itticultura
622 Salmastra Italiana'. Edagricole. 1982. [https://www.libreriauniversitaria.it/vallicoltura-moderna-proposte-
623 operative-ristrutturazione/libro/9788820617769](https://www.libreriauniversitaria.it/vallicoltura-moderna-proposte-operative-ristrutturazione/libro/9788820617769).
- 624 Roner, M., A. D'Alpaos, M. Ghinassi, M. Marani, S. Silvestri, E. Franceschinis, and N. Realdon. 2015. 'Spatial Variation
625 of Salt-Marsh Organic and Inorganic Deposition and Organic Carbon Accumulation: Inferences from the Venice
626 Lagoon, Italy'. *Advances in Water Resources* 000: 1–12. <https://doi.org/10.1016/j.advwatres.2015.11.011>.
- 627 Rova, Silvia, Felix Müller, Patrick Meire, and Fabio Pranovi. 2019. 'Sustainability Perspectives and Spatial Patterns of
628 Multiple Ecosystem Services in the Venice Lagoon: Possible Roles in the Implementation of the EU Water
629 Framework Directive'. *Ecological Indicators* 98 (March): 556–67. <https://doi.org/10.1016/j.ecolind.2018.11.045>.

- 630 Rova, Silvia, and Fabio Pranovi. 2017. 'Analysis and Management of Multiple Ecosystem Services within a Social-
631 Ecological Context'. *Ecological Indicators* 72: 436–43. <https://doi.org/10.1016/j.ecolind.2016.07.050>.
- 632 Rova, Silvia, Fabio Pranovi, and Felix Müller. 2015. 'Provision of Ecosystem Services in the Lagoon of Venice (Italy):
633 An Initial Spatial Assessment'. *Ecohydrology and Hydrobiology* 15 (1): 13–25.
634 <https://doi.org/10.1016/j.ecohyd.2014.12.001>.
- 635 Rova, Silvia, Alice Stocco, and Fabio Pranovi. 2022. 'Ecosystem Services' Capacity and Flow in the Venice Lagoon and
636 the Relationship with Ecological Status'. *One Ecosystem* 7 (September): e79715-
637 <https://doi.org/10.3897/oneeco.7.e79715>.
- 638 RStudio Team. n.d. 'RStudio'.
- 639 Salvetti, Roberta, Marco Acutis, Arianna Azzellino, Marta Carpani, Carlo Giupponi, Paolo Parati, Manfredi Vale, and
640 Renato Vismara. 2008. 'Modelling the Point and Non-Point Nitrogen Loads to the Venice Lagoon (Italy): The
641 Application of Water Quality Models to the Dese-Zero Basin'. *Desalination* 226 (1–3): 81–88.
642 <https://doi.org/10.1016/J.DESAL.2007.01.236>.
- 643 Scapin, Luca, Matteo Zuchetta, Fabio Pranovi, and Piero Franzoi. 2022. 'Influence of Seascape on Coastal Lagoon
644 Fisheries: The Role of Habitat Mosaic in the Venice Lagoon'. *Estuaries and Coasts* 45 (3): 793–811.
645 <https://doi.org/10.1007/S12237-021-00986-3/TABLES/3>.
- 646 Scarton, Francesco. 2017. 'Long-Term Trend of the Waterbird Community Breeding in a Heavily Man-Modified Coastal
647 Lagoon: The Case of the Important Bird Area "Lagoon of Venice"'. *Journal of Coastal Conservation* 21 (1): 35–
648 45. <https://doi.org/10.1007/S11852-016-0470-8/FIGURES/5>.
- 649 Scarton, Francesco, and Mauro Bon. 2009. 'Gli Uccelli Acquatici Svernanti Nella Laguna Di Venezia Nel Periodo 1993-
650 2007: Analisi Delle Dinamiche Temporali e Spaziali'. *Avocetta*.
- 651 Schröter, Matthias, David N. Barton, Roy P. Remme, and Lars Hein. 2014. 'Accounting for Capacity and Flow of
652 Ecosystem Services: A Conceptual Model and a Case Study for Telemark, Norway'. *Ecological Indicators* 36: 539–
653 51. <https://doi.org/10.1016/j.ecolind.2013.09.018>.
- 654 Seitzinger, S., J.A. Harrison, A.F. Bohlke, A.F. Bouwman, R. Lowrance, B. Peterson, C. Tobias, and G. van Drecht. 2006.
655 'Denitrification Across Landscapes and Waterscapes: A Synthesis'. *Ecological Applications* 16 (6): 2064–90.
656 [https://doi.org/10.1890/1051-0761\(2006\)016\[2064:DALAWA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[2064:DALAWA]2.0.CO;2).
- 657 Sfriso, Adriano, and Chiara Facca. 2007a. 'Distribution and Production of Macrophytes and Phytoplankton in the Lagoon
658 of Venice: Comparison of Actual and Past Situation'. *Hydrobiologia*. <https://doi.org/10.1007/s10750-006-0418-3>.
- 659 Sfriso, Adriano, Chiara Facca, and Sonia Ceoldo. 2004. 'Growth and Production of *Cymodocea Nodosa* (Ucria)
660 *Ascherson* in the Venice Lagoon'. *Scientific Research and Safeguarding of Venice* 96: 229–36.
- 661 Sfriso, Adriano, and Pier Francesco Ghetti. 1998a. 'Seasonal Variation in Biomass, Morphometric Parameters and
662 Production of Seagrasses in the Lagoon of Venice'. *Aquatic Botany*. [https://doi.org/10.1016/S0304-3770\(98\)00064-](https://doi.org/10.1016/S0304-3770(98)00064-3)
663 3.
- 664 Sfriso, Adriano, and Antonio Marcomini. 1999. 'Macrophyte Production in a Shallow Coastal Lagoon. Part II: Coupling
665 with Sediment, SPM and Tissue Carbon, Nitrogen and Phosphorus Concentrations'. *Marine Environmental*
666 *Research*. [https://doi.org/10.1016/S0141-1136\(98\)00122-6](https://doi.org/10.1016/S0141-1136(98)00122-6).
- 667 Shang, Yung Cheng. 1982. *Aquaculture Economics: Basic Concepts and Methods of Analysis*. *Journal of the Marine*
668 *Biological Association of the United Kingdom*. Vol. 62. Cambridge University Press.
669 <https://doi.org/10.1017/S0025315400057519>.
- 670 Solidoro, Cosimo, Vinko Bandelj, Fabrizio Aubry Bernardi, Elisa Camatti, Stefano Ciavatta, Gianpiero Cossarini, Chiara
671 Facca, et al. 2010. 'Response of the Venice Lagoon Ecosystem to Natural and Anthropogenic Pressures over the
672 Last 50 Years'. *Coastal Lagoons: Critical Habitats of Environmental Change*, no. January: 483–511.
673 <https://doi.org/10.1201/EBK1420088304>.
- 674 Solidoro, Cosimo, Roberto Pastres, and Gianpiero Cossarini. 2005. 'Nitrogen and Plankton Dynamics in the Lagoon of
675 Venice'. *Ecological Modelling* 184 (1): 103–23. <https://doi.org/10.1016/j.ecolmodel.2004.11.009>.
- 676 Umgiesser, Georg. 2020. 'The Impact of Operating the Mobile Barriers in Venice (MOSE) under Climate Change'.
677 *Journal for Nature Conservation* 54 (April). <https://doi.org/10.1016/j.jnc.2019.125783>.

- 678 Umgiesser, Georg, Donata Melaku Canu, Andrea Cucco, and Cosimo Solidoro. 2004. 'A Finite Element Model for the
679 Venice Lagoon. Development, Set up, Calibration and Validation'. *Journal of Marine Systems* 51 (1-4 SPEC. ISS.):
680 123–45. <https://doi.org/10.1016/j.jmarsys.2004.05.009>.
- 681 Villamagna, Amy M., Paul L. Angermeier, and Elena M. Bennett. 2013. 'Capacity, Pressure, Demand, and Flow: A
682 Conceptual Framework for Analyzing Ecosystem Service Provision and Delivery'. *Ecological Complexity* 15: 114–
683 21. <https://doi.org/10.1016/j.ecocom.2013.07.004>.
- 684 Walton, M. E.M., C. Vilas, J. P. Cañavate, E. Gonzalez-Ortegon, A. Prieto, S. A. van Bergeijk, A. J. Green, M. Librero,
685 N. Mazuelos, and L. le Vay. 2015. 'A Model for the Future: Ecosystem Services Provided by the Aquaculture
686 Activities of Veta La Palma, Southern Spain'. *Aquaculture* 448 (November): 382–90.
687 <https://doi.org/10.1016/j.aquaculture.2015.06.017>.
- 688 Walton, M. E.M., C. Vilas, C. Coccia, A. J. Green, J. P. Cañavate, A. Prieto, S. A. van Bergeijk, et al. 2015. 'The Effect
689 of Water Management on Extensive Aquaculture Food Webs in the Reconstructed Wetlands of the Doñana Natural
690 Park, Southern Spain'. *Aquaculture* 448 (November): 451–63.
691 <https://doi.org/10.1016/J.AQUACULTURE.2015.06.011>.
- 692 Weitzman, Jenny. 2019. 'Applying the Ecosystem Services Concept to Aquaculture: A Review of Approaches,
693 Definitions, and Uses'. *Ecosystem Services* 35 (February): 194–206. <https://doi.org/10.1016/j.ecoser.2018.12.009>.
- 694 Zucchetta, Matteo, Fabrizio Capoccioni, Piero Franzoi, Eleonora Ciccotti, and Chiara Leone. 2021. 'Fish Response to
695 Multiple Anthropogenic Stressors in Mediterranean Coastal Lagoons: A Comparative Study of the Role of Different
696 Management Strategies'. *Water* 13 (2): 130. <https://doi.org/10.3390/w13020130>.
- 697
698

699

Supplementary materials I

700

Scheme of a valle da pesca




Figure S.M. I - 1 Sketch map of a valle da pesca structure. a) brackish water inlet; b) brackish lake; c) winter fishponds; d) freshwater lakes; e) freshwater intake.

701

Supplementary materials II

702

Remote sensing images

703

The following tables list the remote sensing images analyzed in this work. Very High-Resolution (VHR) satellite imageries have been granted by the European Space Agency (ESA) via the ESA On-Demand Restricted Data Access program (Tab. S. M. II - 1). Aerial photograms has been granted by Regione Veneto, “Direzione Pianificazione Territoriale, U.O. Pianificazione territoriale strategica e cartografia” (Tab. S. M. II - 2).

707

Collection data	Satellite/sensor	Satellite scenes			
		bands	resolution	clouds coverage	off-nadir:
2019/07/01	WV 02	8-bands	40 cm	0.0%	13.0°
2019/01/10	WV 02	8-bands	40 cm	1.0%	12.6°
2019/11/07	WV 03	8-bands	32 cm	0.0%	8.1°
2018/11/29	GE 01	4-bands	51 cm	0.0%	26.7°

Table S. M. II - 1 VHR satellite imageries characteristics

Collection data	Flight name	bands	resolution
2018	REVEN 2018	3-bands (RGB)	20 cm
2019	AGEA REVEN	3-bands (RGB)	20 cm

Table S.M. II - 2 Aerial photograms characteristics

708 **Supplementary materials III**

709 **ESs assessment methodology**

ES category	ES	Assessment methodology	
		Valli da pesca	Lagoon
Regulating & maintenance	<p>Climate regulation Carbon sequestration rate by saltmarshes and seagrasses [gC m⁻² y⁻¹]</p>	<p><i>Capacity/flow methodology</i> The spatial distribution of saltmarshes in the valli da pesca has been achieved with a machine learning classification of very high-resolution satellite and aerial imageries (listed in table S. M. II - 1), with calibration points collected during field work. Average saltmarshes' carbon sequestration rate based on literature data (Day et al. 1998; Roner et al. 2015)</p> <p>The spatial distribution of seagrasses meadows has been achieved by photointerpretation of very high-resolution satellite imageries (WV-2A, GeoEye), with the support of UAV aerial imageries and visual census field surveys data collected during periodical visits within the valli da pesca. Seagrasses' C sequestration rate based on species-specific belowground production and organic C content (Sfriso, Facca, and Ceoldo 2004; Sfriso and Francesco Ghetti 1998b; Sfriso, Facca, and Ceoldo 2007; Sfriso and Facca 2007; 2007)</p> <p>According to the literature, capacity and flow indicators of regulating ESs have been considered coincident (Schröter et al. 2014; Hein et al. 2006b).</p>	<p><i>Capacity/flow methodology</i> The saltmarshes spatial distribution in the open, not privately managed part of the lagoon has been retrieved from the saltmarshes map achieved in 2003 by Magistrato alle Acque. Average saltmarshes' carbon sequestration rate based on literature data (Day et al. 1998; Roner et al. 2015)</p> <p>The seagrasses distribution refers to the seagrasses map in 2017 (Provveditorato OO. PP. del Triveneto; SELC, 2018). Seagrasses' C sequestration rate based on species-specific belowground production and organic C content (Sfriso and Facca 2007; Sfriso, Facca, and Ceoldo 2007; 2004; Sfriso and Francesco Ghetti 1998b)</p> <p>According to the literature, capacity and flow indicators of regulating ESs have been considered coincident (Schröter et al. 2014; Hein et al. 2006b).</p>
Regulating & maintenance	<p>Water purification Percentage of Nitrogen load removed by denitrification process in brackish water [%]</p>	<p><i>Capacity/flow methodology</i> Nitrogen removal capacity of the brackish lakes estimated in proportion to the lake extension and the turnover time of the volume of water in the saltwater basin, based on experimental data about denitrification measured by Ravagnan (1982) in other valli da pesca managed in the same way as the valli da pesca considered in this study. According to the literature, capacity and flow indicators of regulating ESs have been considered coincident (Schröter et al. 2014; Hein et al. 2006a).</p>	<p><i>Capacity/flow methodology</i> N load removed through denitrification estimated based on residence time, according to the equation proposed by Seitzinger et al., (2006) for estuarine systems. Residence time calculated with SHYFEM model (Umgiesser et al. 2004) referred to the year 2014 (courtesy of G. Umgiesser, ISMAR-CNR).</p> <p>According to the literature, capacity and flow indicators of regulating ESs have been considered coincident (Schröter et al. 2014; Hein et al. 2006a).</p>

ES category	ES	Assessment methodology	
		Valli da pesca	Lagoon
Regulating & maintenance	Lifecycle support 0-1 scale	<p>Capacity methodology Attractiveness for migratory waterbirds related to the presence of waterbirds lifecycle support factors (saltmarshes, freshwater presence, shrubs, and herbaceous vegetation) that enhance the suitability for resting and molt changing (Korschgen et al., 1985; Havera et al., 1992; Arzel et al., 2006; Hatzziordanou et al., 2019), retrieved from aerial imageries photo interpretation and geospatial layers data. Estimated distribution in the most confined part of the lagoon of juveniles fish biomass [kg ha⁻¹y⁻¹] of the foodweb group “Mugilidae”, retrieved from a spatialized foodweb model of the Venice lagoon (Anelli Monti et al. 2021a) Fish and waterbirds lifecycle support capacities were normalized to a 0-1 scale, then aggregated.</p> <p>Flow methodology Waterbirds: Average number of migratory waterbirds [n ha⁻¹y⁻¹] wintering in the valli da pesca, according to the last ten years’ waterbirds annual censuses (AFV 2020). Fish: Fry biomass [kg ha⁻¹y⁻¹] sown in the valli da pesca for the species Mugil cephalus, Chelon labrosus, Chelon aurata, Chelon saliens and Chelon ramada according to the official management data.</p> <p>Waterbirds and fish lifecycle support capacities were normalized to a 0-1 scale, then aggregated.</p>	<p>Capacity methodology Attractiveness for migratory waterbirds related to the presence of waterbirds lifecycle support factors (saltmarshes, freshwater presence, shrubs, and herbaceous vegetation) that enhance the suitability for resting and molt changing (Korschgen et al., 1985; Havera et al., 1992; Arzel et al., 2006; Hatzziordanou et al., 2019), retrieved from aerial imageries photo interpretation and geospatial layers data. Estimated distribution of juveniles fish biomass [kg ha⁻¹y⁻¹] of the foodweb group “Mugilidae”, retrieved from a spatialized foodweb model of the Venice lagoon (Anelli Monti et al. 2021a) Fish and waterbirds lifecycle support capacities were normalized to a 0-1 scale, then aggregated.</p> <p>Flow methodology Waterbirds: Average number of migratory waterbirds [n ha⁻¹y⁻¹] wintering in the open and not privately managed part of the lagoon, according to the last ten years’ waterbirds annual censuses (AFV 2020). Fish: Fry biomass [kg ha⁻¹y⁻¹] distribution estimated in the lagoon, retrieved from a spatialized foodweb model of the Venice lagoon (Anelli Monti et al., 2021).</p> <p>Waterbirds and fish lifecycle support capacities were normalized to a 0-1 scale, then aggregated.</p>
Provisioning	Fish production Fish biomass [kg ha ⁻¹ y ⁻¹]	<p>Capacity methodology Potential fish biomass hosted by each <i>valle da pesca</i> (kg ha⁻¹ y⁻¹), estimated basing on an Ecopath-Ecosim with Ecospace spatialized foodweb model (Anelli Monti et al. 2021a).</p> <p>Flow methodology Average fish catches per hectare of brackish water surfaces per year (kg ha⁻¹ y⁻¹) in the valli da pesca, according to the 2010-2019 official data (Regione Veneto).</p>	<p>Capacity methodology Sum of the biomass of fish functional groups targeted by artisanal fishing, calculated by an Ecopath-Ecosim with Ecospace spatialized foodweb model (Anelli Monti et al. 2021).</p> <p>Flow methodology Average fish catches from artisanal fishing per hectare of water per year (kg ha⁻¹ y⁻¹), calculated by an Ecopath-Ecosim with Ecospace spatialized foodweb model (Anelli Monti et al. 2021).</p>
Provisioning	Waterfowl hunting Number of huntable waterbirds [n ha ⁻¹ y ⁻¹]	<p>Capacity methodology Mean number of huntable waterbirds per hectare per year wintering in the valli da pesca, according to the time series of waterbirds censuses from 2010 to 2019 (Associazione Faunisti Veneti, 2020), spatially interpolated by Dirichlet tessellation.</p> <p>Flow methodology Average number of waterbirds catches calculated on 2010 - 2020 hunting registries data of the <i>valli da pesca</i>, spatialized considering the effective hunting reserve area.</p>	<p>Capacity methodology Mean number of huntable waterbirds per hectare per year wintering in the open part of the lagoon, according to the time series of waterbirds censuses from 2010 to 2019 (Associazione Faunisti Veneti, 2020), spatially interpolated by Dirichlet tessellation</p> <p>Flow methodology Estimated considering the catches per capita (n. birds/person/hunting trip), the hunting effort (n. of hunting trips/person/year) and the proportion of active hunters in the lagoon outside the hunting farms. Data were gathered through 84 interviews to hunters and by asking the total number of hunters active in the lagoon to the “Ambito Territoriale di caccia VE5” in 2020. Spatial distribution based on the location of the hunting blinds in the lagoon (Regione Veneto, 2019), weighted by the average effective shotgun range.</p>
Provisioning	Wild herbs & honey	<p>Capacity methodology Kg of <i>Salicornia spp.</i> biomass that could be harvested in the <i>valli da pesca</i>, per year, estimated from the vegetational patches distribution in the valli da pesca retrieved with both visual census method and through the</p>	<p>Capacity methodology Kg of <i>Salicornia</i> biomass that could be harvested in the lagoon, per year, estimated from the vegetational patches distribution in the open lagoon retrieved with both visual census method and through the identification of their</p>


ES category	ES	Assessment methodology	
		Valli da pesca	Lagoon
		<p>identification of their predictive range in R, G, B bands reflectance values, filtered through a cut-off NDVI value calculated from aerial and satellite imageries (listed in Supplementary Materials II).</p> <p>Kg of honey potentially produced per unit area in the valli da pesca, considering the <i>Limonium sp.</i> inflorescence cover in the patches (Fantinato and Buffa 2019) and the ratio between the number of sea lavender flowers and the grams of honey that can be obtained from them.</p> <p><i>Flow methodology</i> Kg of harvested edible plants reported by 2020 market data (courtesy of Veneto Agricoltura), by interviewed local people and restaurant chefs. Kg of produced sea-lavender honey witnessed by 5 interviewed beekeepers.</p>	<p>predictive range in R, G, B bands reflectance values, filtered through a cut-off NDVI value calculated from aerial and satellite imageries (listed in Supplementary Materials II).</p> <p>Kg of honey potentially produced per unit area in the lagoon, considering the <i>Limonium</i> inflorescence cover in the patches (Fantinato and Buffa 2019) and the ratio between the number of sea lavender flowers and the grams of honey that can be obtained from them.</p> <p><i>Flow methodology</i> Kg of harvested edible plants reported by 2020 market data (courtesy of Veneto Agricoltura), by interviewed local people and restaurant chefs. Kg of produced sea-lavender honey witnessed by 5 interviewed beekeepers.</p>
Cultural	Tourism	<p><i>Capacity methodology</i> Tourism attractiveness map, assessed by a survey carried out in the second half of 2019. Spatialization achieved by considering the attractiveness factors evaluated by tourists in the questionnaires, namely saltmarshes presence, the possibility to observe birds and fauna, good water quality, and the chance to contemplate natural terrestrial habitats in the valli da pesca. Each element was weighted to depict the interest declared by people and normalized to a 0-1 scale index.</p> <p><i>Flow methodology</i> Number of people who, during a year, have passed at least one night in one of the accommodation facilities within the valli da pesca.</p>	<p><i>Capacity methodology</i> Tourism attractiveness map, assessed by a survey carried out in the second half of 2019. Spatialization achieved by considering the attractiveness factors evaluated by tourists in the questionnaires, namely saltmarshes presence, the possibility to observe birds and fauna, good water quality, and the chance to contemplate natural terrestrial habitats in the lagoon. Each element was weighted to depict the interest declared by people and normalized to a 0-1 scale index.</p> <p><i>Flow methodology</i> Number of people visiting the lagoon in the year 2019, excluding the historical center of Venice, obtained from tourism and transportation operators (public transport company AVM-ACTV S.p.a., 17 private navigation companies, 9 ecotourism associations).</p>
Cultural	Information for cognitive development	<p><i>Capacity methodology</i> The natural factors that enhance touristic attractiveness assessed by a survey carried out in the second half of 2019 have been mapped along with the inclusivity for the necessities of people with disabilities. The resulting map was normalized to a 0-1 scale.</p> <p><i>Flow methodology</i> Number of persons that yearly attend outdoor educational activities or guided one-day trip in the valli da pesca, as reported by the valli da pesca managers, 6 touristic guides and 6 major ecotourism associations.</p>	<p><i>Capacity methodology</i> The natural factors that enhance touristic attractiveness assessed by a survey carried out in the second half of 2019 have been mapped along with the inclusivity for the necessities of people with disabilities. The resulting map was normalized to a 0-1 scale.</p> <p><i>Flow methodology</i> Number of persons that yearly attend outdoor educational activities or guided one-day trip in the lagoon, as reported by 6 touristic guides and 6 major ecotourism associations.</p>
Cultural	Birdwatching	<p><i>Capacity methodology</i> Birdwatching attractiveness normalized map, scaled 0-1, based on the factors reported as important by 30 interviewed birdwatchers, namely the presence of pedestrian paths, saltmarshes presence, birds nesting areas in the visual field, and high probability for observing birds. All these factors were summed up to obtain a normalized attractiveness map.</p> <p><i>Flow methodology</i> Average number of active birdwatchers [$n \cdot y^{-1}$] calculated from the observers' activity trend, recorded from 2010 to 2020 in the Italian birdwatchers' database (https://www.ornitho.it/).</p>	<p><i>Capacity methodology</i> Birdwatching attractiveness normalized map, scaled 0-1, based on the factors reported as important by 30 interviewed birdwatchers, namely the presence of pedestrian paths, saltmarshes presence, birds nesting areas in the visual field, and high probability for observing birds. All these factors were summed up to obtain a normalized attractiveness map.</p> <p><i>Flow methodology</i> Average number of active birdwatchers [$n \cdot y^{-1}$] calculated from the observers' activity trend, recorded from 2010 to 2020 in the Italian birdwatchers' database (https://www.ornitho.it/).</p>

711 **S. M. III references**


- 712 AFV, A.F.V. (2020) CENSIMENTO DEGLI UCCELLI ACQUATICI SVERNANTI IN PROVINCIA DI VENEZIA.
- 713 Anelli Monti, M. et al. (2021) ‘Ecosystem functioning and ecological status in the Venice lagoon, which relationships?’,
714 *Ecological Indicators*, 133, p. 108461. Available at: <https://doi.org/10.1016/j.ecolind.2021.108461>.
- 715 Arzel, C., Elmberg, J. and Guillemain, M. (2006) ‘Ecology of spring-migrating Anatidae: A review’, *Journal of*
716 *Ornithology*. Springer, pp. 167–184. Available at: <https://doi.org/10.1007/s10336-006-0054-8>.
- 717 Day, J.W. et al. (1998) ‘Relative sea level rise and Venice lagoon wetlands’, *Journal of Coastal Conservation*, 4(1), pp.
718 27–34. Available at: <https://doi.org/10.1007/BF02806486>.
- 719 Fantinato, E. and Buffa, G. (2019) ‘Animal-mediated interactions for pollination in saltmarsh communities’, *Plant*
720 *Sociology*, 56(2), pp. 35–42. Available at: <https://doi.org/10.7338/pls2019562/02>.
- 721 Hatziiordanou, L. et al. (2019) ‘Indicators for mapping and assessment of ecosystem condition and of the ecosystem
722 service habitat maintenance in support of the EU biodiversity strategy to 2020’, *One Ecosystem*, 4. Available at:
723 <https://doi.org/10.3897/oneeco.4.e32704>.
- 724 Havera, S.P. et al. (1992) ‘HUMAN DISTURBANCE OF WATERFOWL ON KEOKUK POOL’, undefined [Preprint].
- 725 Hein, L. et al. (2006a) ‘Spatial scales, stakeholders and the valuation of ecosystem services’, *Ecological Economics*,
726 57(2), pp. 209–228. Available at: <https://doi.org/10.1016/J.ECOLECON.2005.04.005>.
- 727 Hein, L. et al. (2006b) ‘Spatial scales, stakeholders and the valuation of ecosystem services’, *Ecological Economics*,
728 57(2), pp. 209–228. Available at: <https://doi.org/10.1016/J.ECOLECON.2005.04.005>.
- 729 Korschgen, C., George, L. and Green, W.L. (1985) ‘Disturbance of diving ducks by boaters on a migrational staging
730 area’, undefined [Preprint].
- 731 Ravagnan, G. (1982) Vallicoltura moderna. Proposte operative per la ristrutturazione e lo sviluppo della itticoltura
732 salmastra italiana, Edagricole. Available at: [https://www.libreriauniversitaria.it/vallicoltura-moderna-proposte-](https://www.libreriauniversitaria.it/vallicoltura-moderna-proposte-operative-ristrutturazione/libro/9788820617769)
733 [operative-ristrutturazione/libro/9788820617769](https://www.libreriauniversitaria.it/vallicoltura-moderna-proposte-operative-ristrutturazione/libro/9788820617769) (Accessed: 27 March 2021).
- 734 Roner, M. et al. (2015) ‘Spatial variation of salt-marsh organic and inorganic deposition and organic carbon
735 accumulation: Inferences from the Venice lagoon, Italy’, *Advances in Water Resources*, 000, pp. 1–12. Available at:
736 <https://doi.org/10.1016/j.advwatres.2015.11.011>.
- 737 Schröter, M. et al. (2014) ‘Accounting for capacity and flow of ecosystem services: A conceptual model and a case
738 study for Telemark, Norway’, *Ecological Indicators*, 36, pp. 539–551. Available at:
739 <https://doi.org/10.1016/j.ecolind.2013.09.018>.
- 740 Seitzinger, S. et al. (2006) ‘Denitrification Across Landscapes and Waterscapes : a Synthesis’, *Ecological Applications*,
741 16(6), pp. 2064–2090. Available at: [https://doi.org/10.1890/1051-0761\(2006\)016\[2064:DALAWA\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[2064:DALAWA]2.0.CO;2).
- 742 Sfriso, A. and Facca, C. (2007) ‘*Nanozostera noltii* growth and production in the lagoon of Venice’, *Biologia Marina*
743 *Mediterranea*, 14, pp. 326–327.
- 744 Sfriso, A., Facca, C. and Ceoldo, S. (2004) ‘Growth and production of *Cymodocea nodosa* (Ucria) ascherson in the
745 Venice Lagoon’, *Scientific Research and Safeguarding of Venice*, 96, pp. 229–236.
- 746 Sfriso, A., Facca, C. and Ceoldo, S. (2007) *Scientific Research and Safeguarding of Venice*.
- 747 Sfriso, A. and Francesco Ghetti, P. (1998) ‘Seasonal variation in biomass, morphometric parameters and production of
748 seagrasses in the lagoon of Venice’, *Aquatic Botany*, 61(3), pp. 207–223. Available at: [https://doi.org/10.1016/S0304-](https://doi.org/10.1016/S0304-3770(98)00064-3)
749 [3770\(98\)00064-3](https://doi.org/10.1016/S0304-3770(98)00064-3).
- 750 Umgiesser, G. et al. (2004) ‘A finite element model for the Venice Lagoon. Development, set up, calibration and
751 validation’, *Journal of Marine Systems*, 51(1-4 SPEC. ISS.), pp. 123–145. Available at:
752 <https://doi.org/10.1016/j.jmarsys.2004.05.009>
753

Supplementary materials IV


a) Climate regulation




b) Denitrification




c) Lifecycles support - capacity




d) Lifecycles support - flow




e) Aquaculture - capacity




f) Aquaculture - flow




g) Waterfowls hunting - capacity




h) Waterfowls hunting - flow




i) Edible halophytes & honey production - cap



j) Edible halophytes & honey production - flow





755

756

757

758

Fig. S.M. IV Results of the ESs assessment. a), b), c), d) refer to regulating and maintenance ESs. e), f), g), h), i), j) refer to provisioning ESs. k), l), m), n), o), p) refer to cultural ESs.