

38 **The effects of preformed vitamin A and provitamin A carotenoid supplementation on**
39 **tadpoles of the poison frog *Phyllobates vittatus***

40
41 **Abstract**

42
43 Understanding the nutritional requirements of captive animals is necessary for proper
44 animal husbandry, however, the specific dietary requirements for many amphibian species
45 commonly kept in captivity are unknown. Like most vertebrates, frogs cannot synthesize
46 carotenoids or vitamin A, and must therefore obtain these essential nutrients through diet. It is
47 unclear if amphibians can cleave provitamin A carotenoids to form vitamin A metabolically
48 within the body, so common practice is to supplement their captive diets with both preformed
49 vitamin A and provitamin A carotenoids. We carried out a feeding experiment in tadpoles of
50 *Phyllobates vittatus*, a commonly kept poison frog species, to test the effects of supplementing a
51 fish flake diet with a provitamin A carotenoid (2.5mg/g β -carotene) and vitamin A (0.5 mg/g
52 retinyl acetate), both individually and in combination. Contrary to our expectations,
53 supplementation had either no effect or adverse effects on tadpole growth and survivorship.
54 Tadpoles reared under supplemented diets showed higher mortality rates, coupled with
55 symptoms of hypervitaminosis A. Survivors had a smaller body size and mass at metamorphosis.
56 The vitamin A and β -carotene levels in our supplemented diet have been shown to benefit
57 tadpoles of other species, yet our results indicate that adding these amounts to what is found in a
58 generalist fish flake mix (5.466 μ g/g and 1.88 μ g/g, respectively) can have detrimental effects on
59 *P. vittatus* tadpoles. More broadly, this study highlights the importance of creating husbandry
60 guidelines based on the specific physiological needs of the species (or species groups) being kept
61 in captivity, rather than general ones for all amphibians, as is often done.

62
63
64
65 **Introduction**

66
67 As wild populations of a considerable number of amphibian species decline (Campbell et
68 al., 2020) , captive populations of some of them are increasing. This is due to both their
69 increasing popularity in the pet trade and captive breeding programs (Edmonds, 2021; Tapley et
70 al., 2015). Proper husbandry is required for all captive animals to ensure that their physiological
71 needs are met. An important aspect of proper husbandry is animal nutrition. However, the
72 nutritional requirements of most amphibians are unknown (Tapley et al., 2015). Breeders of
73 captive frogs are aware that feeder insects, like *Drosophila* species, do not contain all of the
74 nutrients that amphibians require and, thus, enhance diets with commercially available
75 supplements. Despite the wide variety of products available for amphibian diet supplementation,
76 there is very little research to inform their use in particular species.

77 Carotenoids are powerful antioxidants that quench and stabilize reactive oxygen species
78 produced during periods of growth and development (Goodwin, 1984; Keogh et al., 2018;

79 McInerney et al. 2019). This limits the damage that reactive oxygen species cause to cells, which
80 is believed to result in increased cell division and differentiation (Keogh et al., 2018; McInerney
81 et al. 2019). In amphibians, antioxidants are especially important during the larval stage and
82 metamorphic climax because this is when the production of reactive oxygen species is greatest
83 (Keogh et al., 2018; McInerney et al. 2019). Furthermore, carotenoids and their metabolism are
84 involved in coloration in a variety of species across the tree of life (Goodwin, 1984; Toews et al.
85 2017). Since frogs, like most animals, are unable to produce carotenoids, they must be obtained
86 exclusively through their diet (Densmore and Green 2007; Keogh et al., 2018; McInerney et al.
87 2019). It is therefore common practice amongst frog breeders to add carotenoid supplements to
88 the captive diets of most species. However, breeders must be careful not to over supplement with
89 carotenoids as they can have negative effects when given in excess (Keogh et al., 2018; Palozza
90 et al. 2003). For instance, one of the most commonly used carotenoids, β -carotene, can be toxic
91 at high concentrations and has been found to have a pro-oxidant effect that can cause a reduced
92 growth rate (Keogh et al., 2018; Palozza et al. 2003).

93

94 For many animals, carotenoids also act as vitamin A precursors (Brenes-Soto and
95 Dierenfeld, 2014; Clugston and Blaner, 2014). This means that provitamin A carotenoids can be
96 cleaved to form vitamin A metabolically within the body. Whether this is the case for
97 amphibians remains unclear. The few studies that have addressed the problem *in vivo* have not
98 found evidence for conversion of specific carotenoids to vitamin A (McComb 2010; Collins et
99 al., 1952; Wright, 2006), although there is evidence that at least some species may be able to
100 perform this conversion (Baruah and Goswami 2012, Brenes-Soto and Dierenfeld 2014).
101 Vitamin A plays important roles in gene expression, vision, limb regeneration and embryonic
102 development (McInerney et al. 2019; Clugston and Blaner 2014), and in frogs its deficiency has
103 been linked to squamous metaplasia of the mouth, which is commonly referred to as “short
104 tongue syndrome” (Clugston and Blaner 2014). Frogs affected by short tongue syndrome have
105 difficulty catching prey, resulting in lethargy, weight loss, and eventually death if untreated
106 (Clugston and Blaner 2014). To prevent hypovitaminosis A, a common practice amongst
107 breeders today is to co-supplement with carotenoids and preformed vitamin A.

108

109 Nevertheless, since vitamin A is fat-soluble, excessive intake poses a risk for toxicity
110 (Densmore and Green 2007). Hypervitaminosis A has been recorded in African clawed frogs
111 (*Xenopus laevis*) at different stages of their life cycle (Clugston and Blaner 2014). This condition
112 causes weight loss and negatively affects the skin and liver of adult frogs (Pessier 2013), and has
113 been reported to cause abnormal development in tadpoles, with notable observations of
114 hyperpigmented, hemorrhagic skin and chronic diarrhea (Weissmann 1961; Weissman et al.
115 1963). Proper dosages of preformed vitamin A have not been determined for most species, even
116 if under or over-supplementation can lead to hypovitaminosis A or hypervitaminosis A, both of
117 which can be life threatening if untreated (Clugston and Blaner 2014).

118

119 With the goal of improving husbandry practices of captive amphibians, this study
120 investigated the effects of supplementing feed with β -carotene (a commonly used provitamin A
121 carotenoid), vitamin A, and their combination on tadpoles of the poison-dart frog *Phyllobates*
122 *vittatus*, a species commonly kept in captivity. Through observation and measurement of
123 tadpoles during development, we evaluated the effect of the above supplementation regimes on
124 tadpole survivorship, growth rates, final size at metamorphosis, and time to metamorphosis. Our
125 results provide somewhat unexpected initial insights into the dietary carotenoid requirements of
126 *Phyllobates* poison-dart frog tadpoles, and highlight the need for further species-specific research
127 on the diets of captive amphibians.

128 **Methods**

129 *Study Species*

130 *P. vittatus* is a diurnal, terrestrial species that is endemic to the Golfo Dulce Region of
131 Costa Rica, where it can be found in lowland moist and wet forests. Adults range in size from
132 22.5mm to 31mm, with females being larger than the males. *P. vittatus* is black with a broad
133 gold, red-orange, or orange dorsolateral stripe that runs continuously from the dorsal base of
134 each thigh to the tip of the snout, and blue-green speckling on its limbs. Their diet is made up of
135 a variety of small arthropods. Females will lay clutches of 7-21 eggs every two to three weeks.
136 The male cares for the eggs by providing them with moisture, and once the eggs hatch, males
137 will transport the tadpoles to a pool of water (Silverstone 1976), where they likely maintain an
138 omnivorous diet (Grant et al. 2006). Although this has not been studied in *P. vittatus*, predatory
139 behaviors, including tadpole cannibalism, have been observed in multiple closely related species
140 (Caldwell and de Araujo 1998, Márquez *in review*). Tadpoles are uniformly dark brown on the
141 dorsal side, and are lighter brown ventrally, and gradually develop orange dorsolateral stripes,
142 starting around Gosner stage 37 (Gosner 1960). Metamorphosis occurs approximately ten weeks
143 after hatching (Savage 2002; Paitz and Dugas 2021, R. Márquez pers. obs. R. Arkin pers. obs.).

144 *Experimental Animals*

145 A total of 35 *P. vittatus* eggs, from 6 different clutches, were collected from a single
146 captive colony of 7 adults (4M, 3F) maintained at the University of Michigan. Adults are fed
147 *Drosophila hydei* and *D. melanogaster* fruit flies three times a week, supplemented with
148 Repashy Calcium Plus, Rep-Cal Herptivite multivitamin, and Rep-Cal Calcium with Vitamin D
149 (one supplement per feeding). All eggs were laid between 1/20/22 and 2/11/22. They were
150 removed from the terrarium and reared in petri dishes until hatching. Thirteen eggs from 2
151 clutches had been de-jellied shortly after laying. We found no differences between these and
152 other eggs in any of the measured parameters (Table S1). All eggs hatched within the time frame

153 typical of our lab colony (mean 16.4 days, s.d. 2.65, range 14-23 days). One tadpole died within
154 2 days of hatching and before ever being fed, so it was removed from the study.

155 *Experimental diets*

156 All four diets were composed of of 99.7% ground fish flakes (75:25 mixture of
157 SeraFlora/SeraSans; SERA, Germany), which was previously analyzed for nutrient content
158 (Byrne and Silla, 2017), and contributes 0.01454 mg g⁻¹ total carotenoids and 0.00188 mg g⁻¹
159 preformed vitamin A The remaining 0.03% varied between four treatments as follows: Diet 1
160 (control) was unsupplemented. Diet 2 was supplemented with 2.5mg g⁻¹ β-carotene (Sigma
161 Aldrich, Cat. No. C9750), Diet 3 was supplemented with 0.5 mg g⁻¹ vitamin A acetate (i.e. retinyl
162 acetate; Sigma-Aldrich Cat. No. R4632), and Diet 4 was supplemented with 2.5mg g⁻¹ β-carotene
163 and 0.5 mg g⁻¹ vitamin A. Cellulose powder (Sigma Aldrich Cat. No. 435236), which is odorless
164 and has been found to have no impact on the growth, survivorship, or development of tadpoles
165 (Keogh et al., 2018; Dias et al., 1998; Deng et al., 2006), was added to treatments 1, 2, and 3 to
166 ensure that all experimental diets contained the same amount of feed. Table 1 shows the total
167 amounts of carotenoids and vitamin A in each diet treatment, and Table S2 summarizes the
168 recipes used in each diet. β-carotene was selected as the provitamin A carotenoid due to its
169 commercial availability and presence in popular supplements. Retinyl acetate was chosen
170 because it is the form of vitamin A commonly found in commercial vitamin A supplements for
171 both animals and humans (e.g. Repashy vitamin A plus). All diet mixes were suspended to a 1:10
172 ratio in deionized H₂O, wrapped in aluminum foil and mixed on a rocker for 48 hr at room
173 temperature. Once mixed, diets were loaded onto 3ml syringes, wrapped in foil, and stored at 0
174 to -4°C until used.

175 *Feeding experiment*

176 After hatching, individuals were randomly assigned to one of four treatments, and housed
177 in individual 230 ml plastic cups filled with approximately 150ml of reverse-osmosis (RO)
178 water. They were fed triweekly: Monday, Wednesday, and Friday. Before each feeding, syringes
179 were thawed and the feed was homogenized using a blunt tip needle. Tadpoles received 2 drops
180 of food (~0.1 ml) for the first month. After that time, individuals received 4 drops of feed (~0.2
181 ml). Tadpoles were photographed using a Canon EOS R6 camera with a Canon RF24-105mm F4
182 L IS USM lens after each feeding. The body (from tip of snout to base of tail) and tail length of
183 each individual were measured using image analysis software (ImageJ, version 1.53; Schneider
184 et al. 2012). Feeding and photographing stopped once tadpoles began the metamorphic climax at
185 Gosner's stage 42, as evidenced by the emergence of forelimbs (Gosner, 1960). Once both arms

186 emerged, tadpoles were moved to a tilted 1L cup with ~75ml of new RO water to facilitate
187 leaving the water. Once out of the water, froglets were moved to plastic containers (10 cm x 20
188 cm x 10 cm) lined with sphagnum moss and dried magnolia leaves. Froglets were monitored
189 every two days until they reached Gosner's stage 46, marked by the full absorption of their tail
190 (Gosner, 1960). At that time, individuals were photographed (NIKON D5100 camera with an AF
191 Micro-Nikkor 60mm f/2.8D lens) and their body length and weight were recorded. Two tadpoles
192 had not yet developed arms after 81 days, so they were put into new 230 ml plastic cups and a
193 partial water change was performed. This was done to clear the algae buildup that made it
194 difficult to observe and photograph the tadpole.

195 *Statistical analyses*

196 To compare mortality over time between treatments, we computed survival curves using the
197 Kaplan-Meier method (Kaplan & Meier 1958) and tested for differences between survival curves
198 using the chi-squared approach described by Harrington & Fleming (1982). Survival curves were
199 fitted using the `survfit()` function and comparisons were made using the `survdiff()` function, both
200 implemented in the R package `survival` (Therneau 2022).

201 We then evaluated the effect of diet treatment of tadpole growth as a function of age by fitting
202 logistic curves to each tadpole's growth trajectory and comparing the best-fitting growth rate and
203 maximum size parameters between treatments. The logistic equation was used in the following
204 form:

$$205 \quad S_t = \frac{M}{1 + \left(\frac{M - m}{m}\right) e^{-rt}}$$

206 where S_t is the size at a given age, M is the maximum size asymptote, m is the minimum size
207 asymptote, r is the growth rate, and t is the age. Growth curves were fit to each individual using
208 non-linear least squares as implemented in the function `nlsLM()` of the `minpack.lm` R package
209 (Elzhov et al. 2022). We tested for differences in maximum size and growth parameter estimates
210 between treatments using bi-variate anovas with treatment and whether tadpoles survived to
211 metamorphosis as fixed factors. To further explore differences between treatments and
212 survivorship categories, we conducted post-hoc Tukey-Kramer tests. Finally, we tested for
213 between-treatment differences in our empirical measurements of size, weight and age at
214 metamorphosis also using anovas and Tukey-Kramer tests. The data and code used for all
215 analyses and visualizations are available on the dryad repository associated with this paper.

216 Results

217 *Effects of diet on survivorship*

218 Overall, 82.35% (28/34) of tadpoles survived to complete metamorphosis. Tadpoles that
219 were fed diets supplemented with vitamin A (i.e. diets 3 and 4) had higher mortality rates,
220 especially the treatment group that was supplemented with both vitamin A and β -carotene (i.e.
221 Diet 4), which showed a clear excess of observed deaths relative to the expectation under the
222 Kaplan-Meier model ($\chi^2 = 9.67$, $df = 3$, $p = 0.02$; Fig 1A-B). Remarkably, tadpoles that died in
223 the treatment groups supplemented with vitamin A (1 in treatment 3, 4 in treatment 4), were
224 observed to have skin and gastrointestinal symptoms (persistent diarrhea) that reflect what was
225 described in *X. laevis* tadpoles with hypervitaminosis A (Fig 1C; Weissmann 1961; Weissman et
226 al. 1963).

227 *Effects of diet on growth*

228 All of the tested tadpoles displayed growth trajectories characterized by initial rapid
229 growth that gradually slows down towards metamorphosis (Fig. 2A, S1). We found significant
230 effects of treatment, survivorship, and their interaction on both maximum body size (Fig. 2B;
231 treatment: $F_{3,27} = 32.5$, $p = 4.2e-9$; survivorship: $F_{1,27} = 171.3$, $p = 3.3e-13$; treatment \times
232 survivorship: $F_{2,27} = 16.9$, $p = 1.7e-5$) and growth rate (Fig. 2C; treatment: $F_{3,27} = 15.9$, $p = 3.7e-$
233 6 ; survivorship: $F_{1,27} = 67.4$, $p = 8.2e-9$; treatment \times survivorship: $F_{2,27} = 12.4$, $p = 1.5e-4$).
234 However, most of the signal in these tests is due to the tadpoles that did not survive to
235 metamorphosis, most of which displayed faster initial growth spurts but stopped growing soon
236 after, at considerably smaller sizes, where they remained for several days before dying (Fig. 2A-
237 C; Fig. S1). The only significant post-hoc comparisons were either between non-surviving
238 tadpoles of different diet treatments, or between surviving and non-surviving tadpoles, both
239 between and within treatments (Table 2).

240 Among the tadpoles that survived to metamorphosis, diet treatment had moderate effects
241 on body length ($F_{3,24} = 3.41$, $p = 0.034$) and body mass ($F_{3,24} = 2.68$, $p = 0.069$) at metamorphosis,
242 but not on age at metamorphosis ($F_{3,24} = 1.06$, $p = 0.38$;) (Fig 2D-F). Individuals fed the control
243 diet (Diet 1) were on average the largest and heaviest at metamorphosis, while those
244 supplemented with just vitamin A (Diet 3) were smaller and weighed less than those in other
245 treatments. Individuals supplemented with β -carotene (Diet 2) and β -carotene combined with
246 vitamin A (Diet 4) were found to have intermediate sizes and weights (Fig 2D-E). However, the
247 only statistically significant pairwise comparisons in both metamorphic size and weight were

248 between the vitamin A and control treatments (Diet 3 vs. Diet 1; Table 3), possibly due to our
249 moderate sample sizes.

250

251 **Discussion**

252 Our goal in this study was to examine the effects of supplementing the diets of *P. vittatus*
253 tadpoles with additional carotenoids (β -carotene) and preformed vitamin A. This practice is
254 commonly carried out by captive amphibian owners and caretakers, but the safety, benefits, and
255 appropriate dosages of these supplements have only been studied in a limited number of species.
256 The results of our feeding experiment suggest that, at least in the case of *P. vittatus*,
257 supplementing with additional β -carotene and/or vitamin A than what is found in our base fish
258 flake feed (5.466 $\mu\text{g/g}$ β -carotene and 0.00188 mg/g preformed vitamin A, respectively) does not
259 provide any benefits, and can have detrimental effects in terms of tadpole mortality, and
260 probably also on pre-metamorphic growth.

261 The most severe negative effects were observed in the tadpoles fed Diet 4, which
262 contained both supplemental β -carotene and vitamin A. This treatment resulted in a significantly
263 lower survival rate than the other tested diets, and the affected individuals displayed clearly
264 different growth trajectories, as well as symptoms reminiscent of descriptions of
265 hypervitaminosis A in tadpoles of other frog species (Fig 1C; Weissmann 1961; Weissman et al.
266 1963), suggesting that these tadpoles died due to a diet-associated disease, most likely
267 hypervitaminosis A. The same was true for the single tadpole that died under Diet 3 (only
268 vitamin A supplement). The fact that tadpoles supplemented with only vitamin A were much less
269 likely to die or show symptoms than those supplemented with both β -carotene and vitamin A
270 could be due to some of the β -carotene being converted endogenously to vitamin A. While it is
271 commonly assumed that amphibians are not capable of cleaving provitamin A carotenoids to
272 form vitamin A, there is evidence that suggests that at least some species can do so (Brenes-Soto
273 and Dierenfeld 2014, Baruah and Goswami 2012). For instance, false tomato frogs (*Dyscophus*
274 *guineti*) supplemented with a mixture of carotenoids showed increased vitamin A plasma levels,
275 indicating they are likely able to convert carotenoids to vitamin A (Brenes-Soto and Dierenfeld
276 2014). In vitro experiments on the β -carotene metabolism of Asian common toad (*Duttaphrynus*
277 *melanostictus*) and Indian bullfrog (*Haplobatrachus tigerinus*) tadpoles also suggest that they are
278 able to convert β -carotene to vitamin A (Baruah and Goswami 2012). However, it is unknown if
279 this is the case for *P. vittatus* tadpoles, so further research is necessary to isolate the
280 physiological causes of this result. In the end, if a species is able to convert carotenoids into

281 vitamin A, then proper carotenoid supplementation could eliminate the need for additional
282 preformed vitamin A supplements, highlighting the importance of species-specific research for
283 husbandry practices.

284 We based the composition of the experimental diets on previous work with other
285 (distantly related) frog species (e.g. *Pseudophryne corroboree*, Byrne and Silla 2017; Keogh et
286 al. 2018; Weissmann 1961; Weissman et al. 1963), in an attempt to avoid over supplementation .
287 Yet, it seems clear that tadpoles in this study did experience adverse effects associated with over
288 supplementation. This highlights the importance of knowledge on the biology of focal species for
289 adequate captive breeding. The lack of research on the nutritional requirements of most
290 amphibian species kept in captivity (or their close relatives) is concerning, considering the
291 increasing number of species kept in the pet trade, in zoos, and in ex-situ conservation efforts.
292 Understanding the nutritional requirements of captive amphibians is especially important for
293 endangered species since adequate husbandry is the keystone of captive breeding programs.
294 Implementing the proper diet with proper levels of supplementation leads to larger, healthier,
295 more genetically diverse populations for release, which is essential for reintroduction success
296 (McInerney et al. 2019).

297 More generally, our findings highlight that the dietary requirements of frogs vary
298 between species. This is not surprising considering the clade made up by anurans shared a
299 common ancestor approximately 221.8 million years ago (Kumar et al. 2022). Despite their
300 degree of evolutionary divergence, it is common for all frog species to be grouped as a single
301 category in husbandry contexts, which leads to and this belief is reflected by assumed similarities
302 in husbandry practices, which may be unwarranted. This frame of thinking is not usually applied
303 to more “mainstream” species kept in captivity. For instance cats (*Felis catus*) and dogs (*Canis*
304 *lupus familiaris*), for which clearly distinct husbandry practices are in place, diverged
305 approximately 55 million years ago (Kumar et al. 2022). In contrast, dendrobatids and
306 mantellids, two families of frogs commonly kept in captivity, diverged approximately 150
307 million years ago (Kumar et al. 2022) but are assumed to have very similar dietary and other
308 husbandry needs. Considering the phylogenetic diversity of amphibians there is a clear need for
309 further research to establish adequate husbandry and veterinary guidelines and practices that
310 reflect the specific needs of each species (or at least taxonomic group). This information can
311 improve the success of ex-situ conservation efforts, as well as the health of amphibians kept as
312 pets, in zoos, and in laboratories. Our study is a step in this direction for *Phyllobates* poison-dart
313 frogs, and possibly other dendrobatid frogs kept in captivity.

314

315 **Author Contributions**

316 RA and RM conceived and designed the study. RA obtained the data. RA and RM analyzed the
317 data. RA wrote the manuscript with contributions from RM. RM obtained funding. Both authors
318 read and approved the manuscript.

319 **Acknowledgements**

320 We would like to thank members of the Márquez group for comments and suggestions. This
321 work was funded by the Michigan Society of Fellows and the Department of Ecology and
322 Evolutionary Biology at the University of Michigan.

323 **Tables**

324 Table 1. Total carotenoid, β -carotene, and Vitamin A contents of the four tested diets. Baseline contents
 325 were obtained from Byrne and Silla’s (2017) liquid chromatography analyses of the same commercially-
 326 available feeds used for the control diet (i.e. Diet 1).

	Total Carotenoids	β-carotene	Vitamin A
Treatment 1	0.01454 mg/g	0.005466 mg/g	0.00188 mg/g
Treatment 2	2.51454 mg/g	2.505466 mg/g	0.00188 mg/g
Treatment 3	0.01454 mg/g	0.005466 mg/g	0.50188 mg/g
Treatment 4	2.51454 mg/g	2.505466 mg/g	0.50188 mg/g

327
328

329 Table 2. Pairwise comparisons between diet treatments and survivorship categories in growth curve
 330 parameters obtained through a Tukey-Kramer post-hoc test. Maximum size and growth rate comparisons
 331 correspond to panels B and C of Figure 2, respectively.

Comparison	Maximum size (mm)				Growth Rate			
	Difference	Lower 2.5%	Upper 97.5%	p-value	Difference	Lower 2.5%	Upper 97.5%	p-value
D2 Surv. – D1 Surv	-0.03	-0.68	0.62	1.00	0.01	-0.04	0.06	1.00
D3 Surv. – D1 Surv	-0.37	-1.03	0.28	0.58	0.01	-0.04	0.06	1.00
D4 Surv. – D1 Surv	-0.12	-0.86	0.62	1.00	0.01	-0.05	0.06	1.00
D3 Surv. – D2 Surv	-0.34	-0.98	0.29	0.64	0.00	-0.05	0.04	1.00
D4 Surv. – D2 Surv	-0.09	-0.81	0.63	1.00	0.00	-0.06	0.05	1.00
D4 Surv. – D3 Surv	0.26	-0.46	0.98	0.93	0.00	-0.05	0.05	1.00
D3 Died – D1 Died	-1.84	-3.63	-0.05	0.04	0.05	-0.08	0.18	0.86
D4 Died – D1 Died	-2.89	-4.31	-1.48	8.34e-6	0.17	0.06	0.27	3.20e-4
D4 Died – D3 Died	-1.05	-2.47	0.36	0.26	0.11	0.01	0.21	0.026
D1 Surv. – D1 Died	0.65	-0.70	2.00	0.76	-0.02	-0.11	0.08	1.00
D2 Surv. – D1 Died	0.62	-0.72	1.96	0.79	-0.01	-0.10	0.09	1.00
D3 Surv. – D1 Died	0.28	-1.06	1.62	1.00	-0.01	-0.10	0.09	1.00
D4 Surv. – D1 Died	0.54	-0.85	1.92	0.90	-0.01	-0.11	0.09	1.00
D1 Surv. – D3 Died	2.49	1.14	3.85	4.56e-5	-0.07	-0.17	0.03	0.31
D2 Surv. – D3 Died	2.46	1.12	3.81	4.87e-5	-0.06	-0.16	0.04	0.49

D3 Surv. – D3 Died	2.12	0.78	3.46	4.36e-4	-0.06	-0.16	0.04	0.45
D4 Surv. – D3 Died	2.38	0.99	3.76	1.37e-4	-0.06	-0.17	0.04	0.44
D1 Surv. – D4 Died	3.55	2.75	4.34	5.89e-13	-0.18	-0.24	-0.12	1.89e-9
D2 Surv. – D4 Died	3.52	2.74	4.29	4.12e-13	-0.17	-0.23	-0.12	3.80e-9
D3 Surv. – D4 Died	3.17	2.40	3.95	4.86e-12	-0.17	-0.23	-0.12	2.99e-9
D4 Surv. – D4 Died	3.43	2.58	4.28	6.64e-12	-0.18	-0.24	-0.11	1.52e-8

332


333 Table 3. Pairwise comparisons between diet treatments in size and weight of surviving tadpoles at
 334 metamorphosis. Weight and length comparisons correspond to panels E and F of Figure 2, respectively.

	Weight at Metamorphosis (g)				Body Length at Metamorphosis (mm)			
	Difference	Lower 2.5%	Upper 97.5%	p-value	Difference	Lower 2.5%	Upper 97.5%	p-value
D2 – D1	0.00	-0.02	0.01	0.82	-0.10	-0.53	0.33	0.93
D3 – D1	-0.02	-0.03	0.0001	0.052	-0.45	-0.88	-0.02	0.038
D4 – D1	-0.01	-0.03	0.01	0.55	-0.32	-0.80	0.17	0.31
D3 – D2	-0.01	-0.03	0.005	0.24	-0.35	-0.77	0.06	0.11
D4 – D2	-0.004	-0.02	0.01	0.94	-0.22	-0.69	0.26	0.59
D4 – D3	0.01	-0.01	0.02	0.68	0.14	-0.34	0.61	0.86

335

336


337 **Figures**



338

339 **Figure 1.** The effect of dietary supplementation on tadpole survivorship. A) Kaplan-Meier (KM)
340 survivorship curves for each of the four treatments. Translucent polygons represent 95% confidence
341 intervals. B) Observed and expected mortality by treatment under a KM model with equal survivorship by
342 across treatments. C) Examples of a healthy tadpole (left), and one exhibiting the symptoms associated
343 with A hypervitaminosis (right). Photos were taken around Gosner stage 30, 24 days after eclosion.

344



345

346 **Figure 2.** The effect of diet supplementation on tadpole growth. A) Best-fitting logistic growth curves for the 34
 347 tadpoles used in the experiment. Tadpoles that did not survive to metamorphosis are marked with a red diamond. B-
 348 C) Best-fitting maximum size and growth rate parameters for the curves in panel A grouped by treatment and
 349 survivorship status. D-F) Metamorphic age, mass, and body length of individuals that successfully survived past
 350 metamorphosis, grouped by treatments. Colors in all figures match those in Figure 1.

351

352 **Supplementary Materials**

353 Table S1. Survivorship and growth parameters compared between tadpoles born from jellied and de-
 354 jellied eggs. Mortality was compared using Fisher’s exact test, and all other parameters were compared
 355 using one-way anovas. OR = Odds Ratio.

356

Parameter	Test Statistic	df	p-value
Mortality	OR = 0.565	-	0.642
Maximum Size	F = 0.512	1, 32	0.479
Growth Rate	F = 1.699	1, 32	0.202
Time to Metamorphosis	F = 0.095	1, 26	0.760
Mass at Metamorphosis	F = 0.714	1, 26	0.406
Body Length at Metamorphosis	F = 0.664	1, 26	0.422


357

358 Table S2. Recipes for the four tested diets. The flake mix was a 75:25 by weight mixture of Sera Flora
 359 and Sera Sans fish flakes.

Prepared Diet	Flake Mix	β -carotene	Vitamin A	Cellulose	Total
Control (Diet 1)	3.988 g	0 mg	0 mg	12.0 mg	4 g
β-carotene (Diet 2)	3.988 g	10.0 mg	0 mg	2.0 mg	4 g
Vitamin A (Diet 3)	3.988 g	0 mg	2.0 mg	10 mg	4 g
β-carotene and Vitamin A (Diet 4)	3.988 g	10.0 mg	2.0 mg	0 mg	4 g

360

361



362

363 **Figure S1.** Individual growth trajectories for all tadpoles used in the study. Points represent raw
 364 measurements, and red lines are the best-fitting logistic growth curves for each individual.

365

References

- 366 Baruah, P., & Goswami, U. C. (2012). In vitro metabolism of carotenoids , β carotene and
367 lutein into retinoids in amphibians. *Journal of Ecobiotechnology*, 4(1), 46–50.
- 368 Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects
369 Models Using lme4. *Journal of Statistical Software*, 67(1 SE-Articles), 1–48.
370 <https://doi.org/10.18637/jss.v067.i01>
- 371 Brenes-Soto, A., & Dierenfeld, E. S. (2014). Effect of dietary carotenoids on vitamin A
372 status and skin pigmentation in false tomato frogs (*Dyscophus guineti*). *Zoo Biology*,
373 33(6), 544–552. <https://doi.org/10.1002/zoo.21175>
- 374 Byrne, P. G., & Silla, A. J. (2017). Testing the effect of dietary carotenoids on larval
375 survival, growth and development in the critically endangered southern corroboree
376 frog. *Zoo Biology*, 36(2), 161–169. <https://doi.org/10.1002/zoo.21352>
- 377 Caldwell, J. P., & de Araújo, M. C. (1998). Cannibalistic Interactions Resulting from
378 Indiscriminate Predatory Behavior in Tadpoles of Poison Frogs (Anura:
379 Dendrobatidae). *Biotropica*, 30(1), 92–103.
380 <https://doi.org/https://doi.org/10.1111/j.1744-7429.1998.tb00372.x>
- 381 Campbell Grant, E.H., Miller, D.A.W. & Muths, E. (2020) A Synthesis of evidence of
382 drivers of amphibian declines. *Herpetologica* 76, 101–107.
383 <https://doi.org/10.1655/0018-0831-76.2.101>
- 384 Clugston, R. D., & Blaner, W. S. (2014). Vitamin A (retinoid) metabolism and actions:
385 What we know and what we need to know about amphibians. *Zoo Biology*, 33(6), 527–
386 535. <https://doi.org/10.1002/zoo.21140>
- 387 Collins, F. D., Love, R. M., & Morton, R. A. (1953). Studies in vitamin A. 23. Vitamin A
388 and its occurrence in *Amblystoma tigrinum*. *The Biochemical Journal*, 53(4), 626–629.
389 <https://doi.org/10.1042/bj0530626>
- 390 Deng, J., Mai, K., Ai, Q., Zhang, W., Wang, X., Xu, W., & Liufu, Z. (2006). Effects of
391 replacing fish meal with soy protein concentrate on feed intake and growth of juvenile
392 Japanese flounder, *Paralichthys olivaceus*. *Aquaculture*, 258(1–4), 503–513.

393 Densmore, C. L., & Green, D. E. (2007). Diseases of Amphibians. *ILAR Journal*, 48(3),
394 235–254. <https://doi.org/10.1093/ilar.48.3.235>

395 Denver, R. J. (2021). Stress hormones mediate developmental plasticity in vertebrates with
396 complex life cycles. *Neurobiology of Stress*, 14, 100301.
397 <https://doi.org/10.1016/j.ynstr.2021.100301>

398 Dias, J., Huelvan, C., Dinis, M. T., & Métailler, R. (1998). Influence of dietary bulk agents
399 (silica, cellulose and a natural zeolite) on protein digestibility, growth, feed intake and
400 feed transit time in European seabass (*Dicentrarchus labrax*) juveniles. *Aquatic Living
401 Resources*, 11(4), 219–226.

402 Edmonds, D. (2021). Poison frogs traded and maintained by U.S. private breeders.
403 *Herpetological Review*, 52, 779–786.

404 Elzhov, T. V., Mullen, K. M., Spiess, A.-N., & Bolker, B. (2022). *R Interface to the
405 Levenberg-Marquardt Nonlinear Least-Squares Algorithm Found in MINPACK, Plus
406 Support for Bounds* (1.2-2).
407 <https://cran.r-project.org/web/packages/minpack.lm/minpack.lm.pdf>

408 Goodwin, T.W. (1984) II: Animal *The Biochemistry of the Carotenoids*. Springer
409 Netherlands, Dordrecht.

410 Gosner, K. L. (1960). A Simplified Table for Staging Anuran Embryos and Larvae with
411 Notes on Identification. *Herpetologica*, 16(3), 183–190.
412 <http://www.jstor.org/stable/3890061>

413 Grant, T., Frost, D. R., Caldwell, J. P., Gagliardo, R., Haddad, C. F. B., Kok, P. J. R.,
414 Means, D. B., Noonan, B. P., Schargel, W. E., & Wheeler, W. C. (2006). Phylogenetic
415 systematics of dart-poison frogs and their relatives (Amphibia: Athesphatanura:
416 Dendrobatidae). *Bulletin of the American Museum of Natural History*, 299, 1–262.
417 [https://doi.org/10.1206/0003-0090\(2006\)299\[1:PSODFA\]2.0.CO;2](https://doi.org/10.1206/0003-0090(2006)299[1:PSODFA]2.0.CO;2)

418 Harrington, D. P., & Fleming, T. R. (1982). A Class of Rank Test Procedures for Censored
419 Survival Data. *Biometrika*, 69(3), 553–566. <https://doi.org/10.2307/2335991>

420 Kaplan, E. L., & Meier, P. (1958). Nonparametric Estimation from Incomplete
421 Observations. *Journal of the American Statistical Association*, 53(282), 457–481.
422 <https://doi.org/10.1080/01621459.1958.10501452>

- 423 Keogh, L. M., Silla, A. J., McFadden, M. S., & Byrne, P. G. (2018). Dose and life stage-
424 dependent effects of dietary beta-carotene supplementation on the growth and
425 development of the Booroolong frog. *Conservation Physiology*, 6(1), coy052.
426 <https://doi.org/10.1093/conphys/coy052>
- 427 Kumar, S., Suleski, M., Craig, J. M., Kasprowicz, A. E., Sanderford, M., Li, M., Stecher,
428 G., & Hedges, S. B. (2022). TimeTree 5: An Expanded Resource for Species
429 Divergence Times. *Molecular Biology and Evolution*, 39(8).
430 <https://doi.org/10.1093/molbev/msac174>
- 431 Márquez, R. *In review*. Tadpole cannibalism in Phyllobates poison frogs. *Evolutionary*
432 *Ecology*.
- 433 McComb, A. (2010). *Evaluation of Vitamin A Supplementations for Captive Amphibian*
434 *Species* [North Carolina State University].
435 <http://www.lib.ncsu.edu/resolver/1840.16/6567>
- 436 McInerney, E. P., Silla, A. J., & Byrne, P. G. (2019). Effect of carotenoid class and dose on
437 the larval growth and development of the critically endangered southern corroboree
438 frog. *Conservation Physiology*, 7(1), coz009. <https://doi.org/10.1093/conphys/coz009>
- 439 Ogilvy, V., Preziosi, R. F., & Fidgett, A. L. (2012). A brighter future for frogs? The
440 influence of carotenoids on the health, development and reproductive success of the
441 red-eye tree frog. *Animal Conservation*, 15(5), 480–488.
442 <https://doi.org/https://doi.org/10.1111/j.1469-1795.2012.00536.x>
- 443 Paitz, R. T., & Dugas, M. B. (2022). Steroid levels in frog eggs: Manipulations,
444 developmental changes, and implications for maternal steroid effects. *Journal of*
445 *Experimental Zoology Part A: Ecological and Integrative Physiology*, 337(4), 293–
446 302. <https://doi.org/https://doi.org/10.1002/jez.2566>
- 447 Palozza, P., Serini, S., Di Nicuolo, F., Piccioni, E., & Calviello, G. (2003). Prooxidant
448 effects of beta-carotene in cultured cells. *Molecular Aspects of Medicine*, 24(6), 353–
449 362. [https://doi.org/10.1016/s0098-2997\(03\)00031-1](https://doi.org/10.1016/s0098-2997(03)00031-1)
- 450 Pessier, A. P. (2014). Short Tongue Syndrome and Hypovitaminosis A. In D. R. Mader &
451 S. J. Divers (Eds.), *Current Therapy in Reptile Medicine and Surgery* (pp. 271–276).
452 Saunders. <https://doi.org/https://doi.org/10.1016/B978-1-4557-0893-2.00023-5>

453 Savage, J. M. (2002). Frogs and Toads (Order Anura). In *The Amphibians and Reptiles of*
454 *Costa Rica: a herpetofauna between two continents, between two seas* (pp. 390–391).
455 University of Chicago Press.

456 Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years
457 of image analysis. *Nature Methods*, 9(7), 671–675. <https://doi.org/10.1038/nmeth.2089>

458 Silverstone, P. A. (1976). A revision of the poison-arrow frogs of the genus *Phyllobates*
459 *Bibron* in Sagra (Family Dendrobatidae). Natural History Museum of Los Angeles
460 County, Science Bulletin, 27,1–53.

461 Tapley, B., Bradfield, K. S., Michaels, C., & Bungard, M. (2015). Amphibians and
462 conservation breeding programmes: do all threatened amphibians belong on the ark?
463 *Biodiversity and Conservation*, 24(11), 2625–2646. [https://doi.org/10.1007/s10531-](https://doi.org/10.1007/s10531-015-0966-9)
464 [015-0966-9](https://doi.org/10.1007/s10531-015-0966-9)

465 Therneau, T. M. (2022). *A Package for Survival Analysis in R* (3.3-1). [https://cran.r-](https://cran.r-project.org/package=survival)
466 [project.org/package=survival](https://cran.r-project.org/package=survival)

467 Toews, D.P.L., Hofmeister, N.R. & Taylor, S.A. (2017) The Evolution and Genetics of
468 Carotenoid Processing in Animals. *Trends in Genetics* 33, 171–182.
469 <https://doi.org/10.1016/j.tig.2017.01.002>

470 Weissmann, G. (1961). Changes in connective tissue and intestine caused by vitamin A in
471 amphibia, and their acceleration by hydrocortisone. *The Journal of Experimental*
472 *Medicine*, 114(4), 581–592. <https://doi.org/10.1084/jem.114.4.581>

473 Weissmann, G., Bell, E., & Thomas, L. (1963). Prevention by hydrocortisone of changes in
474 connective tissue induced by an excess of vitamin A acid in Amphibia. *The American*
475 *Journal of Pathology*, 42(5), 571–585.

476 Wright, K. M. (2006). Overview of Amphibian Medicine. In D. R. Mader (Ed.), *Reptile*
477 *Medicine and Surgery* (Second, pp. 941–971). W.B. Saunders.
478 <https://doi.org/https://doi.org/10.1016/B0-72-169327-X/50079-1>

479

480

481
482