

# 1 Landscape changes in the “valli da pesca” of the Venice lagoon 2 and possible effects on the Ecosystem Services supply

3  
4 Alice Stocco<sup>1\*</sup>, Lorenzo Duprè<sup>1</sup>, Fabio Pranovi<sup>1</sup>

5  
6 <sup>1</sup> Environmental Sciences, Informatics and Statistics Department, Ca’ Foscari University of Venice,

7 Via Torino 155, 30170 Venezia Mestre, Italy

8 \* corresponding author: [alice.stocco@unive.it](mailto:alice.stocco@unive.it)

## 9 Highlights

- 10 • Aquaculture and hunting reserves of the Venice lagoon changed through the years
- 11 • Remote sensing data allowed for multi-temporal landscape analyses
- 12 • Landscape indicators were effective in describing the valli da pesca evolution
- 13 • Human interventions influenced the landscape structure for maximizing different  
14 ecosystem services
- 15 • Fish production and hunting arise different landscape arrangements

## 16 Abstract


17 Coastal lagoons have long been subject to continuous changes caused by mutual interactions with  
18 human activities. Monitoring such changes becomes critical, particularly when modifications in  
19 landscape and land cover classes can affect their capacity to ensure Ecosystem Services (ESs). In the  
20 Venice lagoon, some confined areas called “valli da pesca” supply provisioning ESs, namely  
21 aquaculture and hunting, but also other ESs that are important for the entire lagoon, such as regulating  
22 and cultural ones. Being heavily modified ecosystems under human control, valli da pesca underwent  
23 considerable morphological evolution depending on the maximized ES and the applied management.

This preprint represents an older version of this paper. The final version has been published in *Estuarine, Coastal and Shelf Science* with the title “Exploring the interplay of landscape changes and ecosystem services maximization in man-managed lagoon areas” (DOI: <https://doi.org/10.1016/j.ecss.2023.108597>). Please refer to the published version for citation purposes.

24 Using remote sensing data from different sources, we reconstructed changes in land cover and  
25 landscape elements in valli da pesca over the last century. By calculating landscape indicators related  
26 to land, saltmarshes, and water, we found that landscape features were initially similar for all the valli  
27 da pesca. Then, a process began between 1975 and 1987, in which management devoted to  
28 maximizing different ESs shaped the land cover in specific patterns. This study confirms the  
29 importance of these areas in the context of the entire lagoon and require for monitoring their land  
30 cover changes. Remote sensing data represent an important source of historical data to deepen the  
31 knowledge about human-Nature interactions, for keeping trace not only of the landscape evolution  
32 but also of the dynamics in the ESs supply in response to human interventions.

33 **Keywords:** ecosystem services, landscape, land cover changes, landscape indicators, Venice  
34 lagoon, managed ecosystems, valli da pesca, remote sensing

35 **Graphical abstract**



36

37

## 38 **1. Introduction**

39 Coastal areas have long been known worldwide for their productivity. They represented the cradle of  
40 different civilizations, becoming some of the most important areas that ensure contributions to human  
41 well-being in the form of Ecosystem Services (Barbier et al., 2011; Duarte et al., 2008). Since ancient  
42 times, humans have learned how to exploit the resources of lagoons, estuaries, and coasts, modifying  
43 the environment and the landscape to their own advantage, building structures, and altering habitats  
44 (Halpern et al., 2008).

45 The most common anthropogenic modifications of coastal areas relate to land reclamation (Gaglio et  
46 al., 2017; Sousa et al., 2020), conversion into croplands (Tian et al., 2021; Zhan et al., 2022),  
47 deployment of aquaculture (Dias et al., 2013), and urbanization (Floerl et al., 2021; Gedan et al.,  
48 2009).

49 In the Venice lagoon, the larger coastal lagoon of the Mediterranean region, the long-standing  
50 interactions between human society and the ecosystem achieved many similar modifications to the  
51 landscape, through a multifaceted process of co-evolution (Gatto and Carbognin, 1981). Today, we  
52 have a fairly clear picture of this evolution thanks to numerous works focusing on the Venice lagoon  
53 from the perspective of various disciplines (Ravera, 2000; UNESCO, 1987). However, the most  
54 confined areas at the interface between the mainland and the lagoon water, called in Italian “valli da  
55 pesca”, are not as well-known.

56 The valli da pesca are peculiar areas confined by levees and embankments, where fresh and brackish  
57 water inputs are entirely regulated by humans, who also intentionally shaped the landscape. Born  
58 during the XIV Century to exploit fishing and waterfowl hunting, in the following decades they  
59 underwent a complex evolution that changed their numerosity, surface extent, and structure (Bullo,  
60 1940; D’Alpaos, 2011; Laffaille, 2016).

61 Even if the valli da pesca are today artificial ecosystems, managed to maximize one or a few  
62 Ecosystem Services (ESs), they still provide ESs belonging to different categories along with the

63 maximized one. Recently, they have been shown to contribute substantially to the ESs budget of the  
64 whole lagoon and act as decisive conservational areas (Stocco et al., 2023).

65 Despite being such an essential part of the Venice lagoon, the land cover evolution in valli da pesca  
66 has not been studied thoroughly. Indeed, all the works reconstructing the changes in the lagoon  
67 landscape on a multi-temporal scale mainly focused on the open lagoon, excluding the valli da pesca  
68 (Brivio and Zilioli, 2014; Gačić and Solidoro, 2004; Molinaroli et al., 2009).

69 One of the main reasons for this exclusion is that these areas include landscape features that can  
70 be smaller than 10 meters wide, resulting in difficulty in monitoring them remotely. Moreover, since  
71 the valli da pesca are privately managed, entering them for field surveys has always been difficult,  
72 making it even more unlikely to overcome this lack of data. Consequently, following their evolution  
73 has always been challenging: they can suddenly undergo artificial changes in the landscape because  
74 of management choices (Wang et al., 2021).

75 Even if cost-effective satellites data with a high temporal resolution, such as Sentinel and Landsat  
76 imageries, have an insufficient spatial resolution to distinguish such features on the ground (Anderson  
77 and Gaston, 2013; Casella et al., 2016), data of adequate spatial resolution come from highly detailed  
78 historical maps and aerial photos, taken during flights scheduled about 10-15 years apart. Such aerial  
79 photos are gathered and made available by the Veneto Region archive; this is an excellent opportunity  
80 to depict most interactions between human society and natural resources in the valli da pesca. This  
81 work aims to evaluate if changes in the management strategy (that is, the ES which is mainly  
82 maximized) have impacted, and how, the morphology of the valli da pesca during the last century. In  
83 particular, we assessed the land cover changes in the valli da pesca in the last century (from 1901 to  
84 2021) and searched for possible relationships between landscape evolution and the ESs maximization.

85

86

## 87 **2. Materials and methods**


### 88 *2.1 Study area*

89 The 31 valli da pesca of the Venice lagoon collectively cover 97 km<sup>2</sup> and are located in both the  
90 Northern and Southern parts of the lagoon (Fig. 1); of these, 27 are still productive under private  
91 management.

92 The managers of these 27 managed valli da pesca provided information about the principal ES on  
93 which their main business relied, as well as about the periodic anthropogenic interventions and the  
94 access rules in the valle da pesca. In addition, aquaculture, hunting, and touristic activities related  
95 data were collected through 54 interviews carried out during our periodical visits to the valli da pesca.  
96 In particular, we retrieved yearly data about fish seeding, fish production, hunting catches, herbs and  
97 honey harvesting, and tourist visits. Literature review and historical maps collection were the primary  
98 sources of information for the valli da pesca that are no longer managed. We also collected data on  
99 abandoned valli da pesca through 12 interviews with the Veneto Region, local police, and ecotourism  
100 guides.

101 The ESs assessment and the collected data allowed for the classification of the valli da pesca into five  
102 different management groups, depending on the maximized ES (Stocco & Pranovi, submitted):

- 103 • valli da pesca devoted to fish production through extensive aquaculture (F)
- 104 • valli da pesca devoted to waterfowl hunting (H)
- 105 • valli da pesca devoted to both fish production and hunting (M)
- 106 • valli da pesca devoted to recreational ES, e.g. nature-based tourism and environmental  
107 education activities (R)
- 108 • valli da pesca no longer managed (N).



109

110 **Figure 1** The Venice lagoon and the valli da pesca. The colors of the polygons indicate the management style aiming  
111 to maximize ESS: blue stands for fish production, red for hunting, violet for multiple ESS, orange for recreational ESS,  
112 green for lack of management. 1 = Valle Dogà, 2 = Valle Grassabò, 3 = Vallesina, 4 = Valle Fosse, 5 = Valle Lio  
113 Maggiore, 6 = Valle Bianca, 7 = Valle Dragojesolo, 8= Valle Cavallino, 9 = Valle Falconera, 10 = Valle Liona, 11 =  
114 Valle Olivara, 12 = Valli Saline-Manciane-Sparasera, 13 = Valle Paleazza, 14 = Valle Sacchettina, 15 = Valle Sacchetta,  
115 16 = Valle Ca' Zane, 17 = Santa Cristina island, 18 = Valle Perini, 19 = Valle Miana-Serraglia, 20 = Valle Avertò, 21 =  
116 Valle A.M.A., 22 = Valle Contarina, 23 = Valle Cornio Alto e Cornio Basso, 24 = Valle Zappa, 25 = Valle Figheri, 26 =  
117 Valle Pierimpiè, 27 = Valle Morosina-Ghebo Storto, 28 = Valle Baseggia, 29 = Valle delle Mesole, 30 = La Cura, 31 =  
118 Valle Millecampi.

## 119 2.2 Land cover and landscape indicators

120 Hydrographic maps of the Venice lagoon dating back to 1901 and 1932  
121 (<http://cigno.atlantedellalaguna.it/geoserver/wms>) were digitized to create categorical raster layers  
122 with three land cover classes, namely land, saltmarshes, and water.

123 For the following time steps, we used 148 aerial photograms from the geo-topographic database of  
124 the Veneto Region, selecting among the aerial images taken in 1954-1955, 1975-1978, 1987, 1999,  
125 2010, and 2018 (Tab. SM1 Supplementary Materials). We georeferenced the images with the QGIS  
126 3.16 core GDAL Georeferencer (QGIS Association: QGIS Geographic Information System, 2022),  
127 considering 6 to 12 ground control points per frame projected on the national coordinate system  
128 Monte Mario – Italy zone 2 (EPSG:3004, 2021 CRS revision). All the aerial photographs were  
129 rectified on the most recent orthophoto mosaics of the study area. A 2<sup>nd</sup>-degree polynomial  
130 transformation algorithm was applied.

131 The georeferenced photos from 1954, 1975, and 1987 had the imprint frame of the on-board camera  
132 holder, which represented an issue for the mosaicking of the photograms. To address this problem, it  
133 was necessary to pre-process the photos using the software GIMP 2.10.30. The pre-processing  
134 workflow required cropping the border of the photograms, masking the clouds and over-exposure  
135 errors, and finally desaturating them to a homogeneous value to obtain reflectance values falling in a  
136 narrow range.

137 Pre-processed photographs were then mosaicked together in QGIS, resulting in seven unique scenes  
138 covering the extent of the study area, with one scene for each considered year. For layers referred to  
139 years 1954, 1975, 1987, and 1999 a post-processing raster correction was performed using a pixel-  
140 based approach, limited to areas that were previously masked because they were affected by clouds  
141 or sunlight halos.

142 Subsequently, we obtained for each one of these scenes a land cover map with 5 meters per pixel side,  
143 in which class 1 corresponds to land, class 2 corresponds to salt marshes, and class 3 to areas covered  
144 by water, coherently with the layers referred to 1901 and 1932. To do so, we chose to apply a random  
145 forest classification algorithm (Liaw and Wiener, 2002), starting by identifying and labeling a  
146 minimum of 50 polygons of interest from which to extract the reflectance values. Obtained reflectance  
147 values of the pixels that fell within each polygon result in a matrix input for each model run. Since

148 mosaics were mono-band or tri-band in the visible, depending on the year and the original aerial  
149 photos, we modified the classification algorithm accordingly. Based on the random forest models  
150 testing results, only the output maps that resulted in at least 86% accuracy were considered.

151 Finally, we obtained the land cover data for 2021 from very-high-resolution satellite scenes collected  
152 by Worldview-02, Worldview-03, and GeoEye-01 on-board sensors, updated through field surveys  
153 carried out in 2021 (Stocco et al., 2023).

154 Geostatistical analyses were performed on the land cover raster maps to calculate the area covered by  
155 each class within the area of each valle da pesca.

156 Three landscape indicators were calculated:

157 
$$\text{land ratio (LR)} = \frac{\text{land area}}{\text{total area}}$$

158 
$$\text{saltmarshes ratio (SR)} = \frac{\text{saltmarshes area}}{\text{total area}}$$

159 
$$\text{water ratio (WR)} = \frac{\text{water covered area}}{\text{total area}}$$


160 where the area is expressed in m<sup>2</sup> and “total area” refers to the total area of the valle da pesca for  
161 which the calculation is made. Obtained data were analyzed with R software (R Core, 2022).

### 162 **3. Results**

163 All the groups of valli da pesca showed similar trends of LR from 1901 to 1975 but then started to  
164 follow different trends depending on the management group (Figs. 2 – 6, a). The statistical analyses  
165 showed that trends followed by the indicators were quite gradual and did not mark significant  
166 differences between contiguous periods.

167 In the valli da pesca belonging to group F, the LR was the lowest in 1901 (0.02±0.005 s.e.). After  
168 that, LR increased until 1975; then, it remained around the same value (0.089 ± 0.007 s.e.) in the last  
169 decades (Fig. 2a). The SR, on the contrary, decreased until 1987, then slowed down towards 2021


170 (Fig. 2b). The WR increases with an opposite trend (Fig. 2c). For SR and WR, the heterogeneity of  
171 data decreases along time.  
172



173  
174 **Figure 2** Landscape indicators for the valli da pesca of group F, maximizing the fish production ES. The dots represent  
175 the means, the error bars show the standard errors. a) land area to total area ratio; b) saltmarshes area to total area ratio;  
176 c) water area to total area ratio.

177 In the valli da pesca belonging to group H, LR is higher in 1932 than in 1901 and remained almost  
178 constant in the following years (Fig. 3a). SR decreases, reaching the lowest values in 1975, then  
179 slightly increases (Fig. 3b). WR showed a specular trend in comparison with the previous indicator  
180 SR (Fig. 3c). All indicators showed higher heterogeneity than that F group.

181



182

183

184

185

**Figure 3** Landscape indicators for the valli da pesca of group H, maximizing hunting ES. The dots represent the means, the error bars show the standard errors. a) land area to total area ratio; b) saltmarshes area to total area ratio; c) water area to total area ratio.

186

187


188

189

190

191

Group M showed trends similar to H for all the indicators but with lower heterogeneity (Figs. 4a-4c). SR in group M recorded the lowest value in 1975, as showed in group H. From the following time step in 1987, SR increased until reaching a steady value in both the group H and M where hunting is practiced. However, group M presented a slightly lower SR value than group H in the last two decades.



192


193 **Figure 4** Landscape indicators for the valli da pesca of group M, maximizing both fish production and hunting ESs.

194 a) land area to total area ratio; b) saltmarshes area to total area ratio; c) water area to total area ratio. The dots represent  
195 the means, and the error bars show the standard errors.

196 In general, group R showed higher LR than the values of groups F, H, and M. An initial increase  
197 is noticed in 1932, followed by a sort of stabilization (Fig. 5a). The SR abruptly decreased between  
198 1901 and 1932 (from  $0.38 \pm 0.14$  to  $0.13 \pm 0.06$ ), and in 2010 when the SR reaches the lowest value  
199 (Fig. 5b). On the contrary, WR increased slightly going towards 2021 (Fig. 5c).

200


201



202

203 **Figure 5** Landscape indicators for the valli da pesca of group R, that abandoned fishing and hunting while maximizing  
204 recreational ESs. a) land area to total area ratio; b) saltmarshes area to total area ratio; c) water area to total area ratio. The  
205 dots represent the means, and the error bars show the standard errors.

206 In group N, an increase in LR can be seen from 1901 to the following years, when LR seems to  
207 stabilize (Fig. 6a). Although showing a wide heterogeneity, SR followed a decreasing trend especially  
208 from 1901 to 1975; after a momentary increase in 1987 SR continued on its decreasing trend (Fig.  
209 6b). The WR shows a quite specular path to SR, with an increase toward 1954-1975 followed by a  
210 decrease in 1987, when the upward trend of WR occurs again. The heterogeneity slightly decreased  
211 with time (Fig. 6c).



212

213

214

215

216

217

## 4. Discussion

218

219

220

221

222

223

224

225

226

227

Figure 6 Landscape indicators for the valli da pesca of group N, which are not managed. a) land area to total area ratio; b) saltmarshes area to total area ratio; c) water area to total area ratio. The dots represent the means, and the error bars show the standard errors.

In this work, we used remote sensing data from different sources and based on different technologies to assess the evolution of the land cover in the valli da pesca of the Venice lagoon in the last century, focussing on the possible relationship between ESs maximization and the landscape arrangement.

By using historical maps, aerial photographs, and high-resolution satellite optical imageries, it was possible to demonstrate that at the beginning of the 20th century, the morphological structure, that is the areas covered by land, saltmarshes, and water, was similar for all the valli da pesca of the Venice lagoon.

The abrupt change in LR between 1901 and 1932 is explainable mainly by the land reclamation process, dating back to 1920-1927 (D’Alpaos, 2010; Sarretta et al., 2010) that caused a deep

228 modification in land use in the study area, confirming that land reclamation is one of the processes  
229 that mainly affected northern Italy coastal lagoons in the past (Fontolan et al., 2012; Gaglio et al.,  
230 2017).

231 In the following years, the building of terrain levees and embankments changed the landscape  
232 proportions in all the valli da pesca; from 1922 to 1943, the land area increased due to a decree law  
233 that allowed, and even financed, the transformation in terrain levees of the thin fences made of swamp  
234 reeds (Bullo, 1940; D’Alpaos, 2010; Gatto and Carbognin, 1981). In that period, however, noticeable  
235 portions of saltmarshes were lost in all the valli da pesca until 1975, regardless of the management  
236 group they belonged, following the same decreasing trends described for the open lagoon  
237 (Madricardo and Donnici, 2014; Molinaroli et al., 2009; Sarretta et al., 2010).

238 A change in the law in 1973 started a process for which the managers of the valli da pesca could  
239 implement substantial changes to the landscape elements, so the development trends have become  
240 distinctive of each management group. In a long process, started during the last years of the Seventies,  
241 all the valli da pesca were provided with bold perimeter structures reinforced with ripraps well above  
242 the high tide elevation (Basurco, 2001; D’Alpaos, 2010; Finotello et al., 2019). This change explains  
243 the little spike in 1975 for the LR. Moreover, the freshwater and brackish water inputs began to be  
244 fully regulated by employing outlet works. Thus, all the water inputs fell completely under human  
245 control, and the valli da pesca became “regulated artificial ecosystems”.

246 In the 80’s, different management strategies of the valli da pesca were established (Laffaille, 2016),  
247 according to the need to maximize different ESs, highly influencing the evolution of the landscape,  
248 especially after 1987.

249 For instance, the increase of land in the valli da pesca of group F is linked to the construction of  
250 artificial, rectangular boundaries of the fishponds, built for protecting fish during winter. The  
251 managers of the valli da pesca of group F seemed to lose interest in conserving saltmarshes: such  
252 observation is concordant with the urgency to maximize the volume of water where to carry out

253 aquaculture. Therefore, SR decreased due to a continuous loss of saltmarshes as time passed under  
254 this type of management.

255 On the other hand, in groups H and M, where hunting is preferred to aquaculture, the saltmarshes  
256 have been restored and preserved between 1987 and 2021, being regarded as crucial elements for  
257 attracting waterfowl in terms of places to rest and shelter (Arzel et al., 2006; Cherkaoui et al., 2017;  
258 Liang et al., 2015; Rizzo and Battisti, 2009). Therefore, the need to preserve suitable habitats to attract  
259 waterfowl to shoot at has led to maintaining these landscape elements. However, the higher  
260 heterogeneity shown by groups H and M indicates that each valle da pesca underwent different  
261 interventions and that no shared management was applied, even if they all aim to maximize the same  
262 ES. This heterogeneity could be linked to the willingness to attract huntable species that require  
263 different habitat characteristics (Adair et al., 1996; Ma et al., 2010; Velasquez, 1992). For instance,  
264 dabbling ducks and diving ducks need different water levels, the former preferring shallow lakes  
265 punctuated with bare shoals and saltmarshes, the latter preferring deeper and wider lakes surrounded  
266 by reed beds (Colwell and Taft, 2000; Isola et al., 2000).

267 A different situation can be identified for group R, where the land-covered area was already higher  
268 than in the other group since 1901. After the land reclamation in 1920-1940, it increased to twice its  
269 initial value. We could hypothesize that such a high ratio has driven the future evolution of the valli  
270 da pesca belonging to group R, perhaps leading their managers to maximize the recreational ESs for  
271 which these valli da pesca were suitable instead of fishing and hunting. Consequently, the cessation  
272 of the need to maintain habitats to attract fish and birds has led to overlooking the importance of  
273 saltmarshes (Fontolan et al., 2012).

274 Finally, in group N, the land cover trend continued to be the same as in the open lagoon (Carniello  
275 et al., 2016; Sarretta et al., 2010). In this case, the heterogeneity among data is more remarkable than  
276 that of the managed groups because each valle da pesca belonging to group N has a different origin  
277 and has seen the ceasing of the management in different time steps.

278 Our findings show that the trends in landscape indicators in the valli da pesca of the Venice lagoon  
279 are related to the type of management implemented and, consequently, to the maximized ES. The  
280 maximization of one (or a few) ES(s) turns out to be effective in making the valli da pesca landscape  
281 more similar to each other when belonging to the same management group, and rather different when  
282 compared to the ones belonging to a different management strategy group. Indeed, as soon as the  
283 management started to modify the landscape to maximize the potential of its own valle da pesca to  
284 provide the desired ES, this caused the same changes in the landscape for all the valli da pesca that  
285 are managed to maximize that specific ES.

286 This result concurs with other authors who assessed the effects of wetland management on the  
287 landscape, especially when dwelling on aquaculture ES maximization (Kelly, 2001; Maltby, 1991;  
288 Saha and Paul, 2021; Schneiders et al., 2012).

289 In the Venice lagoon, such an approach did work appropriately for each one of the valli da pesca,  
290 when analyzed alone. However, they have performed well in providing ESs as long as they shared  
291 the same ecological gradients and processes timeline, in terms of structure and functionality. When,  
292 around the 80'ies, each one of the valli da pesca began to be regulated and partially isolated from the  
293 lagoon, it likewise began to be self-reliant and to detach its dynamics from the one of the other valli  
294 da pesca that surrounded it. This turned out to be quite counterproductive because it was thanks to  
295 the saltmarshes and the synchronized freshwater inputs, initially found in all the confined areas of the  
296 lagoon, that the fish fry and the waterfowl migrate towards the valli da pesca (Cavraro et al., 2017;  
297 Flaherty et al., 2013; Fortibuoni et al., 2014; Zucchetta et al., 2021).

298 The “de-synchronization” of the interventions, and the reduction of the fish stocks, could be among  
299 the reasons why, in the last decades, only a low quantity of fry spontaneously migrated inside the  
300 valli da pesca, as reported by the managers. In a situation where the lagoon fish struggle to overcome  
301 the pressures on the lagoon ecosystems (Zucchetta et al., 2021), the ecological connectivity and the  
302 movements of the fish may have been affected even more by anthropogenic modifications in the

303 lagoon borders. The presence of levees and barriers in front of the fringing saltmarshes, as well as the  
304 loss of plentiful freshwater inputs, could have negatively influenced the ability of fingerlings to reach  
305 the confined areas where they can find refuge and nourishment (Cavraro et al., 2017; Flaherty et al.,  
306 2013; Huisman et al., 1979; Zucchetto et al., 2021). Consequently, the valli da pesca managers needed  
307 to increase the artificial sowing of fish fry to maintain the aquaculture ES, because their management  
308 caused unexpected feedback on the life-cycle support ES.

309 Also, other feedback may represent a matter of concern because the loss of landscape elements  
310 might result in the risk of losing ESs related to them (Grizzetti et al., 2019; Rova et al., 2022). For  
311 example, the tendency to conserve saltmarshes in the groups H and M can help in maintaining a high  
312 supply for regulating ESs (Stocco et al., 2023; Stürck et al., 2015) and provisioning ESs as well. On  
313 the contrary, the replacement of saltmarshes with consolidated and built-up land, as shown in group  
314 R, may result in the loss of ESs that are significant on a broader scale.

315 In light of the results, monitoring the valli da pesca through multi-temporal remote-sensing data  
316 can be an excellent long-term investment, especially when keeping track of the landscape elements  
317 changes. Landscape management aimed at maximizing one ES may indeed exclude interventions  
318 needed for other ES, thus causing drawbacks in the ESs balance of these areas and, consequently, in  
319 the ESs balance of the whole lagoon system to which the valli da pesca belong.

320 Seen that the valli da pesca management can mitigate the loss of landscape elements in the Venice  
321 lagoon, monitoring these areas is as important as monitoring the whole lagoon to prevent harms to  
322 landscape elements, ecological functions, and the essential ESs related to them.

## 323 **5. Conclusions**

324 The multiple interactions between humans and the Venice lagoon led to centuries of human-driven  
325 manipulation of the ecosystem, which has altered landscape structure and ecosystem functions.  
326 Studying the land cover change since the past century in the valli da pesca helped in understanding

327 their evolution, allowing for the identification of the trends followed by landscape elements and the  
328 related land cover change.

329 Owing to the multi-temporal analyses performed, we found that after a period of similarity in the  
330 landscape indicators trends, the private management came into play to mold the landscape elements,  
331 giving different importance to distinct features based on their relationship to the maximized ES.

332 The mutual relationship between landscape indicators and ESs supply confirms that the valli da  
333 pesca are areas important to the entire lagoon, being capable of conserving landscape elements, such  
334 as the saltmarshes, and habitats supporting the life cycle of several species. These aspects make the  
335 valli da pesca to be highly considerable because they can help the lagoon ecosystem in mitigating the  
336 negative effects of climate change through their contribution in terms of ESs supply.

337 However, due to their being completely managed and under men's control, it is important to  
338 monitor the valli da pesca. Changes in them could trigger the put in place of functional and  
339 morphological adaptations that, on the one hand, can be customized and effective for each one of the  
340 valli da pesca, but on the other hand, could prove to be detrimental to the system as a whole.

341 Much work remains to do to understand the dynamics of such areas since field data are still scarce.  
342 However, since aerial orthophotographs and remote sensing imageries proved to be very informative,  
343 it will be well worth considering putting effort into regularly collecting and analyzing them through  
344 the calculation of the proposed landscape indicators, that were effective in depicting the landscape  
345 structure of these areas and its evolution. They can be of great help for monitoring the landscape  
346 changes in the valli da pesca, especially when related to management choices, in order not to lose  
347 benefits and services related to the consistency of meaningful landscape elements.

348

## References

- 349 Adair, S.E., Moore, J.L., Kiel, W.H., 1996. Wintering Diving Duck Use of Coastal Ponds: An  
350 Analysis of Alternative Hypotheses. *J Wildl Manage* 60, 83. <https://doi.org/10.2307/3802043>
- 351 Anderson, K., Gaston, K.J., 2013. Lightweight unmanned aerial vehicles will revolutionize spatial  
352 ecology. *Front Ecol Environ* 11, 138–146. <https://doi.org/10.1890/120150>
- 353 Arzel, C., Elmberg, J., Guillemain, M., 2006. Ecology of spring-migrating Anatidae: A review. *J*  
354 *Ornithol.* <https://doi.org/10.1007/s10336-006-0054-8>
- 355 Barbier, E.B., Hacker, S.D., Kennedy, C., Koch, E.W., Stier, A.C., Silliman, B.R., 2011. The value  
356 of estuarine and coastal ecosystem services. *Ecol Monogr* 81, 169–193.  
357 <https://doi.org/10.1890/10-1510.1>
- 358 Basurco, B., 2001. Mediterranean Aquaculture: Marine Fish Farming Development.
- 359 Brivio, P.A., Zilioli, E., 2014. Assessing wetland changes in the venice lagoon by means of satellite  
360 remote sensing data. *Journal of Coastal Conservation* 1996 2:1 2, 23–32.  
361 <https://doi.org/10.1007/BF02743034>
- 362 Bullo, G., 1940. Le valli salse da pesca e la vallicultura.
- 363 Carniello, L., D’Alpaos, A., Botter, G., Rinaldo, A., 2016. Statistical characterization of  
364 spatiotemporal sediment dynamics in the Venice lagoon. *J Geophys Res Earth Surf* 121, 1049–  
365 1064. <https://doi.org/10.1002/2015JF003793>
- 366 Casella, E., Rovere, A., Pedroncini, A., Stark, C.P., Casella, M., Ferrari, M., Firpo, M., 2016. Drones  
367 as tools for monitoring beach topography changes in the Ligurian Sea (NW Mediterranean).  
368 *Geo-Marine Letters* 36, 151–163. <https://doi.org/10.1007/S00367-016-0435-9/TABLES/2>
- 369 Cavraro, F., Zucchetta, M., Malavasi, S., Franzoi, P., 2017. Small creeks in a big lagoon: The  
370 importance of marginal habitats for fish populations. *Ecol Eng* 99, 228–237.  
371 <https://doi.org/10.1016/j.ecoleng.2016.11.045>
- 372 Cherkaoui, S.I., Selmi, S., Hanane, S., 2017. Ecological factors affecting wetland occupancy by  
373 breeding Anatidae in the southwestern mediterranean. *Ecol Res* 32, 259–269.  
374 <https://doi.org/10.1007/s11284-017-1436-5>
- 375 Colwell, M.A., Taft, O.W., 2000. Waterbird Communities in Managed Wetlands of Varying Water  
376 Depth. *Waterbirds: The International Journal of Waterbird Biology* 23, 45–55.
- 377 D’Alpaos, L., 2011. Evoluzione dei rapporti idraulici tra la laguna di Venezia e le sue valli da pesca.  
378 Istituto Veneto di Scienze, Lettere ed Arti.
- 379 D’Alpaos, L., 2010. L’evoluzione morfologica della laguna di Venezia attraverso la lettura di alcune  
380 mappe storiche e delle sue carte idrografiche. Comune di Venezia, Venezia.
- 381 Dias, J.A., Cearreta, A., Isla, F.I., de Mahiques, M.M., 2013. Anthropogenic impacts on  
382 Iberoamerican coastal areas: Historical processes, present challenges, and consequences for  
383 coastal zone management. *Ocean Coast Manag* 77, 80–88.  
384 <https://doi.org/10.1016/j.ocecoaman.2012.07.025>
- 385 Duarte, C.M., Dennison, W.C., Orth, R.J.W., Carruthers, T.J.B., 2008. The charisma of coastal  
386 ecosystems: Addressing the imbalance. *Estuaries and Coasts* 31, 233–238.  
387 <https://doi.org/10.1007/S12237-008-9038-7/FIGURES/3>
- 388 Finotello, A., Canestrelli, A., Carniello, L., Ghinassi, M., D’Alpaos, A., 2019. Tidal Flow Asymmetry  
389 and Discharge of Lateral Tributaries Drive the Evolution of a Microtidal Meander in the Venice

- 390 Lagoon (Italy). *J Geophys Res Earth Surf* 124, 3043–3066.  
391 <https://doi.org/10.1029/2019JF005193>
- 392 Flaherty, K.E., Matheson, R.E., McMichael, R.H., Perry, W.B., 2013. The Influence of Freshwater  
393 on Nekton Community Structure in Hydrologically Distinct Basins in Northeastern Florida Bay,  
394 FL, USA 36, 918–939. <https://doi.org/10.1007/sl>
- 395 Floerl, O., Atalah, J., Bugnot, A.B., Chandler, M., Dafforn, K.A., Floerl, L., Zaiko, A., Major, R.,  
396 2021. A global model to forecast coastal hardening and mitigate associated socioecological risks.  
397 *Nature Sustainability* 2021 4:12 4, 1060–1067. <https://doi.org/10.1038/s41893-021-00780-w>
- 398 Fontolan, G., Pillon, S., Bezzi, A., Villalta, R., Lipizer, M., Triches, A., D’Aietti, A., 2012. Human  
399 impact and the historical transformation of saltmarshes in the Marano and Grado Lagoon,  
400 northern Adriatic Sea. *Estuar Coast Shelf Sci* 113, 41–56.  
401 <https://doi.org/10.1016/j.ecss.2012.02.007>
- 402 Fortibuoni, T., Gertwagen, R., Giovanardi, O., Raicevich, S., 2014. The progressive deregulation of  
403 fishery management in the Venetian Lagoon after the fall of the Repubblica Serenissima: food  
404 for thought on sustainability. *Global Bioethics* 25, 42–55.  
405 <https://doi.org/10.1080/11287462.2014.894707>
- 406 Gačić, M., Solidoro, C., 2004. Lagoon of Venice: Circulation, water exchange and ecosystem  
407 functioning. *Journal of Marine Systems* 51, 1–3. <https://doi.org/10.1016/j.jmarsys.2004.06.001>
- 408 Gaglio, M., Aschonitis, V.G., Gissi, E., Castaldelli, G., Fano, E.A., 2017. Land use change effects on  
409 ecosystem services of river deltas and coastal wetlands: case study in Volano–Mesola–Goro in  
410 Po river delta (Italy). *Wetl Ecol Manag* 25, 67–86. <https://doi.org/10.1007/s11273-016-9503-1>
- 411 Gatto, P., Carbognin, L., 1981. The lagoon of venice: Natural environmental trend and man-induced  
412 modification. *Hydrological Sciences Bulletin* 26, 379–391.  
413 <https://doi.org/10.1080/02626668109490902>
- 414 Gedan, K.B., Silliman, B.R., Bertness, M.D., 2009. Centuries of Human-Driven Change in Salt Marsh  
415 Ecosystems. *Ann Rev Mar Sci* 1, 117–141.  
416 <https://doi.org/10.1146/annurev.marine.010908.163930>
- 417 Grizzetti, B., Liqueste, C., Pistocchi, A., Vigiak, O., Zulian, G., Bouraoui, F., de Roo, A., Cardoso,  
418 A.C., 2019. Relationship between ecological condition and ecosystem services in European  
419 rivers, lakes and coastal waters. *Science of the Total Environment* 671, 452–465.  
420 <https://doi.org/10.1016/j.scitotenv.2019.03.155>
- 421 Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C. v., Micheli, F., D’Agrosa, C., Bruno, J.F.,  
422 Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin, E.M.P.,  
423 Perry, M.T., Selig, E.R., Spalding, M., Steneck, R., Watson, R., 2008. A global map of human  
424 impact on marine ecosystems. *Science (1979)* 319, 948–952.  
425 <https://doi.org/10.1126/SCIENCE.1149345>
- 426 Huisman, E.A., FAO, R., Hague, W. on M.R. of F. and F. of F.W.F. eng 8 M. 1979 T., Huisman,  
427 E.A., Hogendoorn, H., Ministerie van Landbouw en Visserij, T.H., 1979. The role of fry and  
428 fingerlings in (freshwater) fish culture. <https://doi.org/10.3/JQUERY-UIJS>
- 429 Isola, C.R., Colwell, M.A., Taft, O.W., Safran, R.J., 2000. Interspecific Differences in Habitat Use  
430 of Shorebirds and Waterfowl Foraging in Managed Wetlands of California’s San Joaquin Valley.  
431 *Waterbirds: The International Journal of Waterbird Biology* 23, 196–203.

- 432 Kelly, N.M., 2001. Changes to the landscape pattern of coastal North Carolina wetlands under the  
433 Clean Water Act, 1984-1992. *Landsc Ecol* 16, 3–16.
- 434 Laffaille, F., 2016. Biens publics, biens communs, fonction sociale de l’État. La lagune de Venise et  
435 le statut des Valli salse di pesca. *Revue internationale de droit comparé* 68, 681–698.  
436 <https://doi.org/10.3406/RIDC.2016.20688>
- 437 Liang, J., Hua, S., Zeng, G., Yuan, Y., Lai, X., Li, X., Li, F., Wu, H., Huang, L., Yu, X., 2015.  
438 Application of weight method based on canonical correspondence analysis for assessment of  
439 anatidae habitat suitability: A case study in east dongting lake, middle china. *Ecol Eng* 77, 119–  
440 126. <https://doi.org/10.1016/j.ecoleng.2015.01.016>
- 441 Liaw, A., Wiener, M., 2002. Classification and Regression by randomForest. *R news* 2/3, 18–22.
- 442 Ma, Z., Cai, Y., Li, B., Chen, J., 2010. Managing Wetland Habitats for Waterbirds: An International  
443 Perspective. *Wetlands* 30, 15–27. <https://doi.org/10.1007/s13157-009-0001-6>
- 444 Madricardo, F., Donnici, S., 2014. Mapping past and recent landscape modifications in the Lagoon  
445 of Venice through geophysical surveys and historical maps. *Anthropocene* 6, 86–96.  
446 <https://doi.org/10.1016/J.ANCENE.2014.11.001>
- 447 Maltby, E., 1991. Wetland management goals: wise use and conservation. *Landsc Urban Plan* 20, 9–  
448 18. [https://doi.org/10.1016/0169-2046\(91\)90085-Z](https://doi.org/10.1016/0169-2046(91)90085-Z)
- 449 Molinaroli, E., Guerzoni, S., Sarretta, A., Masiol, M., Pistolato, M., 2009. Thirty-year changes (1970  
450 to 2000) in bathymetry and sediment texture recorded in the Lagoon of Venice sub-basins, Italy.  
451 *Mar Geol* 258, 115–125. <https://doi.org/10.1016/J.MARGEO.2008.12.001>
- 452 QGIS Association: QGIS Geographic Information System, 2022. QGIS [WWW Document]. URL  
453 <http://www.qgis.org>
- 454 R Core team, 2022. R: A language and environment for statistical computing. R Foundation for  
455 Statistical Computing, Vienna, Austria. [WWW Document]. URL <https://www.r-project.org/>
- 456 Ravera, O., 2000. The Lagoon of Venice : the result of both natural factors and human influence. *J*  
457 *Limnol* 59, 19. <https://doi.org/10.4081/jlimnol.2000.19>
- 458 Rizzo, E., Battisti, C., 2009. Habitat preferences of anatidae (Aves , Anseriformes) in a Mediterranean  
459 patchy wetland (Central Italy). <https://doi.org/10.4149/ekol>
- 460 Rova, S., Stocco, A., Pranovi, F., 2022. Ecosystem services’ capacity and flow in the Venice Lagoon  
461 and the relationship with ecological status. *One Ecosystem* 7.  
462 <https://doi.org/10.3897/oneeco.7.e79715>
- 463 Saha, J., Paul, S., 2021. An insight on land use and land cover change due to tourism growth in coastal  
464 area and its environmental consequences from West Bengal, India. *Spatial Information Research*  
465 29, 577–592. <https://doi.org/10.1007/S41324-020-00368-0/FIGURES/8>
- 466 Sarretta, A., Pillon, S., Molinaroli, E., Guerzoni, S., Fontolan, G., 2010. Sediment budget in the  
467 Lagoon of Venice, Italy. *Cont Shelf Res* 30, 934–949.  
468 <https://doi.org/10.1016/J.CSR.2009.07.002>
- 469 Schneiders, A., van Daele, T., van Landuyt, W., van Reeth, W., 2012. Biodiversity and ecosystem  
470 services: Complementary approaches for ecosystem management? *Ecol Indic* 21, 123–133.  
471 <https://doi.org/10.1016/J.ECOLIND.2011.06.021>
- 472 Sousa, C.A.M., Cunha, M.E., Ribeiro, L., 2020. Tracking 130 years of coastal wetland reclamation  
473 in Ria Formosa, Portugal: Opportunities for conservation and aquaculture. *Land use policy* 94,  
474 104544. <https://doi.org/10.1016/j.landusepol.2020.104544>

- 475 Stocco, A., Basconi, L., Rova, S., Pranovi, F., 2023. Like Little Lagoons: The Contribution of Valli  
476 da Pesca to the Ecosystem Services Supply of the Venice Lagoon. *Estuaries and Coasts* 1, 1–14.  
477 <https://doi.org/10.1007/s12237-023-01168-z>
- 478 Stürck, J., Schulp, C.J.E., Verburg, P.H., 2015. Spatio-temporal dynamics of regulating ecosystem  
479 services in Europe- The role of past and future land use change. *Applied Geography* 63, 121–  
480 135. <https://doi.org/10.1016/j.apgeog.2015.06.009>
- 481 Tian, P., Li, J., Cao, L., Pu, R., Gong, H., Liu, Y., Zhang, H., Chen, H., 2021. Impacts of reclamation  
482 derived land use changes on ecosystem services in a typical gulf of eastern China: A case study  
483 of Hangzhou bay. *Ecol Indic* 132, 108259. <https://doi.org/10.1016/j.ecolind.2021.108259>
- 484 UNESCO, 1987. Venice and its Lagoon – UNESCO World Heritage List Advisory Body Evaluation  
485 (ICOMOS).
- 486 Velasquez, C.R., 1992. Managing Artificial Saltpans as a Waterbird Habitat: Species’ Responses to  
487 Water Level Manipulation. *Colonial Waterbirds* 15, 43. <https://doi.org/10.2307/1521353>
- 488 Wang, J., Chen, J., Wen, Y., Fan, W., Liu, Q., Tarolli, P., 2021. Monitoring the coastal wetlands  
489 dynamics in Northeast Italy from 1984 to 2016. *Ecol Indic* 129.  
490 <https://doi.org/10.1016/j.ecolind.2021.107906>
- 491 Zhan, L., Xin, P., Chen, J., 2022. Subsurface salinity distribution and evolution in low-permeability  
492 coastal areas after land reclamation: Field investigation. *J Hydrol (Amst)* 612.  
493 <https://doi.org/10.1016/j.jhydrol.2022.128250>
- 494 Zucchetta, M., Capoccioni, F., Franzoi, P., Ciccotti, E., Leone, C., 2021. Fish Response to Multiple  
495 Anthropogenic Stressors in Mediterranean Coastal Lagoons: A Comparative Study of the Role  
496 of Different Management Strategies. *Water (Basel)* 13, 130. <https://doi.org/10.3390/w13020130>  
497
- 498

This preprint represents an older version of this paper. The final version has been published in *Estuarine, Coastal and Shelf Science* with the title “Exploring the interplay of landscape changes and ecosystem services maximization in man-managed lagoon areas” (DOI: <https://doi.org/10.1016/j.ecss.2023.108597>). Please refer to the published version for citation purposes.

499 **Author Contributions:** (according to CRediT taxonomy)

500 Stocco Alice: Conceptualization, Data curation, Formal analysis, Investigation, Methodology,  
501 Coding, Writing - original draft, Writing - review & editing.

502 Duprè Lorenzo: Data curation, Formal analysis, Investigation, Methodology, Writing - review &  
503 editing.

504 Pranovi Fabio: Conceptualization, Funding acquisition, Project administration, Supervision,  
505 Validation, Writing - review & editing.

506 All authors have read and agreed to submit this version of the manuscript.

507

508 **Funding:** This scientific work has been developed in the context of the “Venezia 2021” project,  
509 thanks to a doctoral fellowship co-funded by Ca’ Foscari University of Venice with the contribution  
510 of the Provveditorato Interregionale Opere Pubbliche per il Veneto, Trentino Alto Adige e Friuli  
511 Venezia Giulia through the Consorzio Venezia Nuova and coordinated by CORILA.

512

513 **Data Availability Statement:** The data supporting this study's findings are available upon request  
514 from the corresponding author.

515

516 **Acknowledgments:** The authors want to acknowledge the Centre for Cartography of the Regione  
517 Veneto.

518

519 **Conflicts of Interest:** The authors have no conflicts of interest to disclose.

520 **Supplementary material**

521 **Table SM1** Data of the flights whose aerial images have been used in this work upon request to the Aerophotography

522 Collection of the Veneto Region.

<b>Flight related code</b>	<b>year</b>	<b>cruising altitude [m]</b>	<b>band(s)</b>	<b>Study area coverage [%]</b>
<b>1954-1955 GAI</b>	1954-1955	5000	1	100
<b>1975 ReVen Benedetti</b>	1975	2600	1	75
<b>1978 ReVen Benedetti</b>	1978	6000	1	100
<b>1987 ReVen</b>	1987	3000	3, R-G-B	100
<b>1999 ReVen Veneto Centrale</b>	1999	2500	3, R-G-B	100
<b>2010 ReVen Area Venezia VA 18</b>	2010	1680 - 3030	3, R-G-B	100
<b>AGEA orthoimage 20 cm [2018]</b>	2018	NA (likely 1600-3000)	3, R-G-B	100

523