

# 1 Building a Portuguese Coalition for Biodiversity Genomics



2  
3 João P. Marques<sup>a,1,2</sup>, Paulo C. Alves<sup>a,1,2,3,4</sup>, Isabel R. Amorim<sup>a,5</sup>,  
4 Ricardo J. Lopes<sup>a,1,2,6</sup>, Mónica Moura<sup>a,2,7,8</sup>, Eugene W. Myers<sup>a,9,10</sup>,  
5 Manuela Sim-Sim<sup>a,11,12,13</sup>, Carla Sousa-Santos<sup>a,14</sup>, M. Judite Alves<sup>11,12</sup>, Paulo A. V. Borges<sup>5</sup>,  
6 Thomas Brown<sup>9</sup>, Miguel Carneiro<sup>1,2</sup>, Carlos Carrapato<sup>15,16</sup>, Luís M. P. Cerfaco<sup>1,12,17</sup>, Cláudio  
7 Ciofi<sup>18</sup>, Luís P. da Silva<sup>1,2</sup>, Genevieve Diedericks<sup>19</sup>, Maria A. Diroma<sup>18</sup>, Liliana Farelo<sup>1,2</sup>,  
8 Giulio Formenti<sup>20</sup>, Fátima Gil<sup>21</sup>, Miguel Grilo<sup>14,22</sup>, Henrique G. Leitão<sup>19</sup>, Alessio Ianucci<sup>18</sup>,  
9 Cristina Máguas<sup>11,13</sup>, Ann M. Mc Cartney<sup>23</sup>, Sofia L. Mendes<sup>11,24</sup>, João M. Moreno<sup>11,25</sup>, Marco  
10 Morselli<sup>26</sup>, Alice Mouton<sup>18,27</sup>, Chiara Natali<sup>18</sup>, Fernando Pereira<sup>5</sup>, Rúben M. C. Rego<sup>7</sup>,  
11 Roberto Resendes<sup>7</sup>, Guilherme Roxo<sup>7</sup>, Hannes Svardal<sup>19,28</sup>, Helena Trindade<sup>11,13</sup>, Sara  
12 Vicente<sup>11,29</sup>, Sylke Winkler<sup>9,30</sup>, Marcela Alvarenga<sup>1,2,3</sup>, Andreia J. Amaral<sup>31,32,33</sup>, Agostinho  
13 Antunes<sup>3,34</sup>, Paula F. Campos<sup>34</sup>, Adelino V. M. Canário<sup>35</sup>, Rita Castilho<sup>35,36</sup>, L. Filipe C.  
14 Castro<sup>3,34</sup>, Angelica Crottini<sup>1,2,3</sup>, Mónica V. Cunha<sup>11,37</sup>, Gonçalo Espregueira Themudo<sup>11,34</sup>,  
15 Pedro J. Esteves<sup>1,2,3</sup>, Rui Faria<sup>1,2</sup>, Carlos Rodríguez Fernandes<sup>11,38</sup>, Jean-Baptiste Ledoux<sup>34</sup>,  
16 Bruno Louro<sup>35</sup>, Sara Magalhães<sup>11,24</sup>, Octávio S. Paulo<sup>11,24</sup>, Gareth Pearson<sup>35</sup>, João  
17 Pimenta<sup>1,2</sup>, Francisco Pina-Martins<sup>11,24</sup>, Teresa L. Santos<sup>11,24</sup>, Ester Serrão<sup>35</sup>, José Melo-  
18 Ferreira<sup>b,1,2,3</sup> and Vítor C. Sousa<sup>b,11,24</sup>

19  
20 <sup>a</sup> shared first authorship

21 <sup>b</sup> shared senior authorship

22

23 **Corresponding authorship:** João P. Marques ([joao.marques@cibio.up.pt](mailto:joao.marques@cibio.up.pt)), José Melo-

24 Ferreira ([jmeloferreira@cibio.up.pt](mailto:jmeloferreira@cibio.up.pt)) and Vitor C. Sousa ([vmsousa@ciencias.ulisboa.pt](mailto:vmsousa@ciencias.ulisboa.pt))

25

## 26 Affiliations

27 <sup>1</sup> CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado,

28 Campus de Vairão, Universidade do Porto, 4485-661 Vairão, Portugal

29 <sup>2</sup> BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão, 4485-661 Vairão,  
30 Portugal

31 <sup>3</sup> Departamento de Biologia, Faculdade de Ciências, Universidade do Porto, 4099-002 Porto, Portugal

32 <sup>4</sup> EBM - Estação Biológica de Mértola,, 7750-329 Mértola, Portugal

33 <sup>5</sup> cE3c-Centre for Ecology, Evolution and Environmental Changes, Azorean Biodiversity Group, CHANGE –  
34 Global Change and Sustainability Institute, and University of the Azores, Rua Capitão João d'Ávila, Pico da Urze,  
35 9700-042, Angra do Heroísmo, Portugal

36 <sup>6</sup> MHNC-UP, Natural History and Science Museum of the University of Porto, 4099-002 Porto, Portugal

37 <sup>7</sup> CIBIO, Research Centre in Biodiversity and Genetic Resources, InBIO Associated Laboratory, Azores Unit,  
38 University of the Azores, Campus of Ponta Delgada, Rua da Mãe de Deus 13A, 9500-321 Ponta Delgada,  
39 Azores, Portugal

40 <sup>8</sup> UNESCO Chair – Land Within Sea: Biodiversity & Sustainability in Atlantic Islands

41 <sup>9</sup> Max Planck Institute of Molecular Cell Biology and Genetics Pfothenhauerstraße 108 01307 Dresden, Germany

42 <sup>10</sup> Okinawa Institute of Science and Technology, Okinawa, Japan

43 <sup>11</sup> cE3c - Centre for Ecology, Evolution and Environmental Changes & CHANGE - Global Change and  
44 Sustainability Institute, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa,  
45 Portugal

46 <sup>12</sup> MUHNAC - Museu Nacional de História Natural e da Ciência, Universidade de Lisboa, Rua da Escola  
47 Politécnica, 58, 1250-102 Lisboa, Portugal

48 <sup>13</sup> Departamento de Biologia Vegetal, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-  
49 016 Lisboa, Portugal

50 <sup>14</sup> MARE-Marine and Environmental Sciences Centre/ARNET-Aquatic Research Network, ISPA-Instituto Superior  
51 de Ciências Psicológicas, Sociais e da Vida, Rua Jardim do Tabaco, 34, 1149-041 Lisbon, Portugal.

52 <sup>15</sup> Instituto da Conservação da Natureza e das Florestas, IP, Portugal

53 <sup>16</sup> DCNF-Alentejo/DLAP/Parque Natural do Vale do Guadiana

54 <sup>17</sup> Universidade Federal do Rio de Janeiro, Museu Nacional, Departamento de Vertebrados, Quinta da Boa Vista,  
55 São Cristóvão, 20940-040 Rio de Janeiro, RJ, Brasil

56 <sup>18</sup> Department of Biology, University of Florence, 50019 Sesto Fiorentino (FI), Italy

57 <sup>19</sup> Department of Biology, University of Antwerp, 2610 Antwerp, Belgium

58 <sup>20</sup> The Vertebrate Genome Laboratory, The Rockefeller University, 1230 York Ave, New York City, 10021 NY,  
59 USA

60 <sup>21</sup> Aquário Vasco da Gama, R. Direita do Dafundo, 1495-718 Cruz Quebrada – Dafundo, Portugal

61 <sup>22</sup> Egas Moniz Center for Interdisciplinary Research (CiiEM), Egas Moniz School of Health and Science, 2829-  
62 511 Caparica, Almada, Portugal

- 63 <sup>23</sup> Genomics Institute, University of California, Santa Cruz, CA 95060 USA
- 64 <sup>24</sup> Departamento de Biologia Animal, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016
- 65 Lisboa, Portugal
- 66 <sup>25</sup> MARE – Centro de Ciências do Mar e do Ambiente (MARE), ARNET Aquatic Research Network , Faculdade
- 67 de Ciências , Universidade de Lisboa, Lisboa, Portugal
- 68 <sup>26</sup> Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, Parco Area
- 69 delle Scienze 23/A, 43124 Parma, Italy
- 70 <sup>27</sup> InBios/Conservation Genetics Laboratory, University of Liege, Chemin de la Vallée 4, 4000 Liege, Belgium
- 71 <sup>28</sup> Naturalis Biodiversity Center, 2333 CR Leiden, The Netherlands
- 72 <sup>29</sup> ERISA – Escola Superior de Saúde Ribeiro Sanches, Instituto Politécnico da Lusofonia (IPLUSO), Rua do
- 73 Telhal aos Olivais nº 8 – 8a, 1950-396 Lisboa, Portugal
- 74 <sup>30</sup> DRESDEN concept genome center, c/o Max Planck Institute of Molecular Cell Biology and Genetics
- 75 Pfotenhauerstraße 108 01307 Dresden, Germany
- 76 <sup>31</sup> Departamento de Zootecnia, Escola Superior de Ciência e Tecnologia, Universidade de Évora, 7000-671
- 77 Évora, Portugal
- 78 <sup>32</sup> CIISA - Centre for Interdisciplinary Research in Animal Health, Faculty of Veterinary Medicine, University of
- 79 Lisbon, Alto da Ajuda, 1300-477 Lisbon, Portugal
- 80 <sup>33</sup> AL4AnimalS - Associate Laboratory for Animal and Veterinary Sciences, Portugal
- 81 <sup>34</sup> CIIMAR/CIMAR, Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Porto,
- 82 Portugal
- 83 <sup>35</sup> CCMAR/CIMAR LA Centre for Marine Sciences, Campus de Gambelas, University of Algarve, 8005-139 Faro,
- 84 Portugal
- 85 <sup>36</sup> Universidade do Algarve, Campus de Gambelas, 8005-139 Faro, Portugal
- 86 <sup>37</sup> BiolSI - Biosystems and Integrative Sciences Institute, Faculdade de Ciências da Universidade de Lisboa,
- 87 Lisboa, Portugal.
- 88 <sup>38</sup> Faculdade de Psicologia, Universidade de Lisboa, Alameda da Universidade, 1649-013 Lisboa, Portugal

89

## 90 **Abstract**

91 The diverse physiography of the Portuguese land and marine territory, spanning from

92 continental Europe to the Atlantic archipelagos, has made it an important repository of

93 biodiversity throughout the Pleistocene glacial cycles, leading to a remarkable diversity of

94 species and ecosystems. This rich biodiversity is under threat from anthropogenic drivers,

95 such as climate change, invasive species, land use changes, overexploitation or pathogen  
96 (re)emergence. The inventory, characterization and study of biodiversity at inter- and intra-  
97 specific levels using genomics is crucial to promote its preservation and recovery by informing  
98 biodiversity conservation policies, management measures and research. The participation of  
99 researchers from Portuguese institutions in the European Reference Genome Atlas (ERGA)  
100 initiative, and its pilot effort to generate reference genomes for European biodiversity, has  
101 reinforced the establishment of Biogenome Portugal. This nascent institutional network will  
102 connect the national community of researchers in genomics. Here, we describe the  
103 Portuguese contribution to ERGA's pilot effort, which will generate high-quality reference  
104 genomes of six species from Portugal that are endemic, iconic and/or endangered, and include  
105 plants, insects and vertebrates (fish, birds and mammals) from mainland Portugal or the  
106 Azores islands. In addition, we outline the objectives of Biogenome Portugal, which aims to (i)  
107 promote scientific collaboration, (ii) contribute to advanced training, (iii) stimulate the  
108 participation of institutions and researchers based in Portugal in international biodiversity  
109 genomics initiatives, and (iv) contribute to the transfer of knowledge to stakeholders and  
110 engaging the public to preserve biodiversity. This initiative will strengthen biodiversity  
111 genomics research in Portugal and fuel the genomic inventory of Portuguese eukaryotic  
112 species. Such efforts will be critical to the conservation of the country's rich biodiversity and  
113 will contribute to ERGA's goal of generating reference genomes for European species.

114

## 115 **Keywords**

116 Biodiversity, Genomics, Conservation, Portugal

117

## 118 **Sumário**

119 A diversa fisiografia dos territórios terrestre e marinho portugueses, que se estende desde a  
120 Europa continental até aos arquipélagos do Atlântico, promoveu a sua importância como  
121 repositório de biodiversidade ao longo dos ciclos glaciares do Pleistoceno, resultando numa  
122 notável diversidade de espécies e ecossistemas. No entanto, esta biodiversidade enfrenta

123 ameaças decorrentes de atividades humanas, tais como as alterações climáticas, invasão de  
124 espécies, mudanças no uso do solo, exploração excessiva e (re)emergência de agentes  
125 patogénicos. A fim de promover a preservação e recuperação da biodiversidade, é crucial  
126 inventariar, caracterizar e estudar a biodiversidade a níveis inter e intraespecíficos, utilizando  
127 técnicas genómicas. Essa informação é essencial para informar políticas de conservação da  
128 biodiversidade, medidas de gestão e pesquisas futuras. A participação de investigadores de  
129 instituições portuguesas na iniciativa European Reference Genome Atlas (ERGA), e no  
130 projeto-piloto para gerar genomas de referência para a biodiversidade europeia, promoveu a  
131 criação do Biogenoma Portugal. Esta rede institucional em desenvolvimento visa conectar a  
132 comunidade nacional de investigadores em genómica. Neste contexto, descrevemos a  
133 contribuição portuguesa para o projeto-piloto do ERGA, que tem como objetivo gerar  
134 genomas de referência de alta qualidade de seis espécies de Portugal. Estas espécies são  
135 endémicas, icónicas e/ou ameaçadas e incluem plantas, insetos e vertebrados (peixes, aves  
136 e mamíferos) do território continental de Portugal e das ilhas dos Açores. Adicionalmente,  
137 delineamos os objetivos do Biogenoma Portugal que visa: (i) promover a colaboração  
138 científica, (ii) contribuir para a formação avançada, (iii) estimular a participação de instituições  
139 e investigadores baseados em Portugal em iniciativas internacionais de genómica da  
140 biodiversidade e (iv) contribuir para a transferência de conhecimento e envolvimento do  
141 público em geral na preservação da biodiversidade. Esta iniciativa fortalecerá a investigação  
142 em genómica da biodiversidade em Portugal e contribuirá para o objetivo do ERGA de gerar  
143 genomas de referência para espécies europeias. Estes esforços terão um papel fundamental  
144 na conservação da biodiversidade do país e na promoção da conscientização sobre a  
145 importância de preservar a diversidade biológica.

146

## 147 **Introduction**

148 The remarkable diversity of life on Earth is essential for maintaining ecosystem stability,  
149 fostering ecological interactions among species, regulating the climate, and providing  
150 fundamental resources that sustain human well-being and promote the development of

151 society. Today, Earth's biodiversity, ecological and evolutionary heritage, and the ecosystem  
152 services they provide are under intense and extensive strain as a result of many direct and  
153 indirect anthropogenic activities. These major activities include habitat destruction and  
154 fragmentation caused by agricultural expansion and other forms of land conversion,  
155 overexploitation of natural resources, overharvesting, illegal killing and trading of wild species,  
156 climate changes, environmental pollution, and global spread of exotic species (Ceballos et al.,  
157 2017; Cowie et al., 2022; Habel et al., 2019; Karger et al., 2021; O'Hara et al., 2021). These  
158 pressures not only result in the decline and extinction of populations, species and ecosystems,  
159 but also have cascading effects on various dimensions of biodiversity, including genetic,  
160 functional or phenological diversity and the interactions within biodiversity networks. These  
161 impacts harm the health of natural populations and their survival, and reduce their potential to  
162 adapt to environmental challenges (Dauphin et al., 2021). Characterising and understanding  
163 in detail the genetic variation underlying biodiversity at all levels provides essential modern  
164 tools to delineate sound and efficient strategies to mitigate the impacts of human activities,  
165 design management plans for conservation and restoration, and support data-driven  
166 biodiversity policy development (Browne et al., 2019; Hohenlohe et al., 2021; Ralls et al., 2020;  
167 Segelbacher et al., 2022). Regional biodiversity genomics initiatives that promote research  
168 and applications under the umbrella of international coalitions, can play an important role in  
169 this global challenge.

170 Europe hosts a remarkable biodiversity richness despite its temperate climate and historically  
171 anthropogenic landscapes (Etté & Geburek, 2021), and has an important biodiversity hotspot  
172 along the Mediterranean basin (Habel et al., 2019; Myers et al., 2000). Most of the continental  
173 territory of Portugal, located in south-western Europe in the Iberian Peninsula, is part of the  
174 Mediterranean biodiversity hotspot (Myers et al., 2000). The country also includes the volcanic  
175 Atlantic archipelagos of Azores, Madeira and Selvagens (Figure 1). Even though Portugal is  
176 the 19<sup>th</sup> largest European country (land area ~92000 km<sup>2</sup>), it holds the 5<sup>th</sup> largest marine territory  
177 (~1.7 million km<sup>2</sup>) (Pauly et al., 2020). As a result of its unique geographic location, climate  
178 and geophysical features, Portugal comprises a variety of landscapes and ecosystems and

179 harbours a large diversity of species. The Iberian Peninsula provided refugia for species  
180 throughout the glacial periods and remains one of the most important repositories of  
181 biodiversity in Europe, both at inter- and intraspecific level (Gómez & Lunt, 2007; Hewitt,  
182 2004). In addition, the geographic isolation of the archipelagos of Azores, Madeira and  
183 Selvagens, which together with the Canary and Cabo Verde archipelagos form the  
184 biogeographical unit of Macaronesia, has led to the divergence and speciation of many  
185 lineages that resulted in many endemisms (Florencio et al., 2021). According to the  
186 Portuguese National Strategy for the Conservation of Nature and Biodiversity 2030 (ENCNB  
187 2030; <https://dre.pt/dre/detalhe/resolucao-conselho-ministros/55-2018-115226936>), Portugal  
188 holds ~35,000 animal and plant species, representing a relevant proportion of these species  
189 groups present in Europe. The IUCN Red List database (<https://www.iucnredlist.org>; 27 April  
190 2023) lists ~800 of the assessed species as endemic to the continental mainland or  
191 archipelagos of Portugal. There are 414 protected areas in Portugal encompassing 22.28% of  
192 the land and 2.46% of marine water, and 404 species and 102 habitats are protected under  
193 EU law (Biodiversity Information System for  
194 Europe: <https://biodiversity.europa.eu/countries/portugal>). The ENCNB 2030 recognizes the  
195 importance of a systematic inventory and characterization of biodiversity at all levels –  
196 ecosystems, species and genetic diversity – to properly anchor conservation strategies.  
197 Genomics provides unprecedented power to understand biodiversity at the inter- and  
198 intraspecific levels (Formenti et al., 2022; Theissinger et al., 2023). Thus, gathering scientific  
199 expertise in the area, coordinating research efforts and promoting genomic studies of native  
200 Portuguese species, can give an invaluable contribution to preserving the rich biodiversity of  
201 Portugal and its ecosystem services.

202

### 203 **ERGA-Portugal community and the Pilot effort**

204 The European Reference Genome Atlas (ERGA; <https://www.erga-biodiversity.eu/>) is an  
205 ambitious and ground-breaking initiative that aims to revolutionise our understanding of  
206 biodiversity by creating a comprehensive genomic resource for European species. ERGA is

207 implemented under the umbrella of the Earth Biogenome Project (EBP;  
208 <https://www.earthbiogenome.org/>), the global network of networks that aims to catalogue the  
209 eukaryotic life diversity of Earth through genome sequencing. ERGA gathers researchers,  
210 scientists, and biodiversity enthusiasts to collect and sequence the genomes of species across  
211 Europe, providing a valuable tool for studying and preserving biodiversity, understanding  
212 evolutionary processes, and addressing pressing environmental challenges. The network  
213 leverages the power of genomics to unlock the secrets of European biodiversity and foster  
214 collaborative efforts towards its conservation and sustainable management.

215 ERGA includes more than 50 members in Portugal, distributed across seven research units,  
216 five universities and two natural history museums (May 31<sup>st</sup>, 2023; Figure 1). This community  
217 of ERGA members based in Portugal is called ERGA-Portugal, and covers a wide range of  
218 expertise that is relevant for biodiversity genomics research, from taxonomy and systematics  
219 across diverse biological groups (e.g., microorganisms, plants, insects, fish, amphibians, birds  
220 or mammals), to comparative and population genomics and bioinformatics. This community  
221 communicates through open meetings, mailing lists and message sharing channels to  
222 facilitate the exchange of biodiversity genomic expertise across Portugal. Many Portuguese  
223 community members are also active participants in ERGA's scientific committees.

224 To contribute to the establishment and mission of ERGA, ERGA-Portugal participated in the  
225 pilot test of ERGA (Cartney et al., 2023). This pilot aimed at testing and developing the  
226 biodiversity genomic networks across Europe through the sequencing of several European  
227 species, with the contribution of expert ERGA members and European institutions and without  
228 a centralised source of funding. In Portugal, the goal of this project was to consolidate the  
229 ERGA-Portugal community, and initiate collective efforts to generate reference genomes for  
230 Portuguese species. This pilot effort in Portugal focused on endemic, endangered and iconic  
231 species from several taxonomic groups and different ecosystems in both the mainland and  
232 the Azores islands. From an initial list of 53 species proposed by members of ERGA-Portugal,  
233 11 were prioritised based on ERGA's feasibility criteria defined by the ERGA Sampling and

234 Sample Processing Committee (Böhne et al., 2023). These species were then ranked based  
235 on an anonymous online survey open to all members of ERGA-Portugal at the time. The six  
236 species with the highest score were included in the pilot effort. This shortlist included one fish,  
237 one mammal, one bird, one invertebrate and two plants (Figure 2).

238 Each species was championed by a genome team, led by one or two species ambassadors,  
239 and included ERGA-Portugal members and other ERGA members with transdisciplinary skills,  
240 who were responsible for successfully implementing each step for generating a reference  
241 genome. These steps included, e.g., sampling, taxonomic identification, vouchering,  
242 laboratory work, sequencing, assembly, annotation and downstream analysis. These six  
243 genome teams, detailed below for each species, promoted new national and international  
244 collaborations. The implementation of this project also promoted interactions with national  
245 authorities, for example, to obtain permits for capture, storage and export of samples, in  
246 coordination with the Portuguese National authority on Nature Conservation (Institute for  
247 Nature Conservation and Forests - ICNF), and the Azorean Regional Directorate for the  
248 Environment and Climate Change (DRAAC). For mainland Portugal, the capture and  
249 collection of samples of wildlife specimens were authorised by the ICNF, while for the  
250 Autonomous regions of Azores permission was given by the DRAAC. For the Azores,  
251 collections complied with the Access and Benefit Sharing provisions codified in the Nagoya  
252 Protocol (Convention on Biological Diversity, 2010). For all species, the sampling steps  
253 followed the guidelines provided by the ERGA SSP committee (Böhne et al., 2023). Sampling  
254 was optimised to ensure that High Molecular Weight (HMW) DNA and RNA could be obtained  
255 from at least one individual and ideally from multiple tissues (see details for each species  
256 below). The immediate preservation and transport of tissues in liquid nitrogen was prioritised  
257 and tissues were subsequently maintained at -80°C conditions.

258

### 259 **ERGA-Portugal pilot species**

260 Iberian minnowcarp - *Anaecypris hispanica* (Steindachner, 1866)

261 Common name in Portuguese: saramugo

262 The **Iberian minnowcarp** (*Anaocypris hispanica*) is a strictly freshwater fish with a short  
263 lifespan that inhabits a restricted geographical area in two river basins of the Southern Iberian  
264 Peninsula (Cardoso, 2023; De Miguel et al., 2010), a region which is severely affected by the  
265 effects of global climate change. This endemic species is considered to be the most  
266 endangered strictly freshwater fish in Iberian rivers, and is listed as Endangered on the IUCN  
267 Red List (Crivelli, 2006) and as Critically Endangered on the Portuguese Red List (Cabral et  
268 al., 2005). These fish live in vulnerable freshwater ecosystems and are exposed to multiple  
269 threats, including increased temperatures and propensity for drought, pollution, habitat  
270 fragmentation, dams, intensive water use and invasive species proliferation (Cabral et al.,  
271 2005; Cardoso, 2023). The Iberian minnowcarp is considered an iconic species as it is the  
272 only living member of the genus *Anaocypris*, representing a phylogenetically unique old  
273 lineage that persisted in the Iberian Peninsula (Perea et al., 2010).

274 *Sampling and sample processing:* An adult female was collected by hand net from an open-  
275 air breeding tank at Aquário Vasco da Gama (Cruz Quebrada-Dafundo, Portugal), where an  
276 *ex-situ* conservation program is ongoing with captive-born fish derived from 37 individuals of  
277 River Chança (Guadiana river basin, Portugal). The individual was sampled, euthanized and  
278 dissected on the same day, according to permits from the ICNF (Permit P-026382/2021) and  
279 animal welfare regulations of the host institution (ORBEA-MARE 02/2021). The head of the  
280 fish was flash-frozen and used for biobanking. Fin clips were used for barcoding and preserved  
281 in ethanol. Fin clips were also used to derive cell lines, which were flash-frozen and can be  
282 used for karyotyping and further genetic studies in the future. Tissues were collected  
283 immediately and flash-frozen in liquid nitrogen. Liver, spleen, muscle, eyes and ovary tissues  
284 were collected.

285 *Sequencing and assembly:* Aliquots of these tissues were shipped to the Max Planck Institute  
286 of Molecular Cell Biology and Genetics (MPI-CBG, Germany), where the remaining steps for  
287 DNA extraction and sequencing were performed. The HMW DNA extraction and library  
288 preparation for PacBio HiFi was done using the spleen, whereas Hi-C library preparation for  
289 Illumina was done using muscle. RNA-Sequencing and Iso-Seq libraries from eye and gonad

290 tissues were produced and sequenced. Sequencing data was used to obtain a genome  
291 assembly, which was done and completed by the genome team members from MPI-CBG  
292 (Germany) using the VGP standards. A manually curated assembly is completed, and the  
293 annotation of the genome is ongoing in a collaborative effort involving researchers from  
294 ERGA-Portugal, as well as institutions and national computational resources.

295 *Expected results:* A high-quality reference genome will open the door to reconstruct the  
296 evolutionary history of this monospecific genus, elucidating its relationship with other fish  
297 lineages. Given that the Iberian minnowcarp is fragmented into small populations, inhabiting  
298 a region highly impacted by drought and water scarcity (Sousa-Santos et al., 2014), the  
299 availability of its reference genome will be crucial to understand adaptation to intermittent  
300 environments, to inform conservation management, and predict responses to increasing water  
301 temperatures. The Iberian minnowcarp is included in ongoing *ex-situ* conservation programs,  
302 hence these genomic resources will allow quantifying the loss of genetic diversity due to  
303 consecutive generations bred from small initial stocks of wild breeders, as well as optimise  
304 and monitor future *ex-situ* efforts.

305

### 306 Iberian hare - *Lepus granatensis* (Rosenhauer, 1856)

307 Common name in Portuguese: lebre Ibérica

308 The **Iberian hare** (*Lepus granatensis*), also known as Granada hare, is a lagomorph species  
309 endemic to the Iberian Peninsula, and the only hare species present in Portugal (Alves et al.,  
310 2023). It is an iconic species of major ecological and economic importance in the Iberian  
311 ecosystems, acting as an important prey to apex predators and a relevant small game species  
312 in Portugal and Spain. Genetic studies have promoted this species as an outstanding model  
313 to understand diverse evolutionary questions, such as the impact of glacial cycles in repeated  
314 changes in species ranges, and the influence of introgressive hybridization on their adaptive  
315 potential (Seixas et al., 2018). It is currently classified as Least Concern in the IUCN Red List  
316 (Soriguer & Carro, 2018), but a marked decline over the last decade led to its classification as  
317 Vulnerable in the last review of the Red Book of Mammals of Portugal (Mathias et al., 2023).

318 The rate of population decline has increased due to the emergence of a natural recombinant  
319 derived from the rabbit *Myxoma virus*, the *hare Myxoma virus* (ha-MYXV) (Águeda-Pinto et  
320 al., 2019; Dalton et al., 2019).

321 *Sampling and Sample Processing.* One male individual Iberian hare was sampled in Mértola,  
322 southern Portugal, during the regular permitted hunting season, duly authorised by ICNF  
323 (Permit 012/2022/CAPT). Tissues were collected shortly after the animal's death and flash  
324 frozen in liquid nitrogen. Tissues from several organs were collected during the necropsy and  
325 kept at -80°C for biobanking: liver, kidney, spleen, lung and testis. The whole body will be  
326 prepared for deposition at the Natural History and Science Museum of the University of Porto  
327 (MHNCUP\_MAM 0812).

328 *Sequencing and Assembly.* Tissue samples were shipped to the University of Antwerp for  
329 Omni-C and RNA-Sequencing library preparation, and to the University of Florence for PacBio  
330 DNA library preparation. Final Omni-C libraries were generated using kidney tissue, while  
331 RNA-Sequencing libraries were produced for liver, kidney, spleen, lung and testis tissues.  
332 These libraries along with PacBio DNA libraries were sequenced at the University of Florence.  
333 Long-read DNA sequencing was performed in HiFi mode in a PacBio Sequel IIe platform using  
334 five 8 million ZMW SMRT cells. Sequencing data was used to produce a genome assembly  
335 following the gold standard instructions implemented in the VGP Galaxy Pipeline (Batut et al.,  
336 2018; Lariviere et al., accessed June 2023). The genome manual curation and annotation is  
337 ongoing, and involves ERGA-Portugal members, as well as institutions and national  
338 computational resources.

339 *Expected results:* Generating a high-quality reference genome will improve genomic studies  
340 in this species, upgrading the genomic resources for this and related species. This will open  
341 the door for fundamental research in speciation and adaptation genomics. Furthermore, it will  
342 anchor research aiming to quantify the impact of emerging pathogens on the naïve host hare  
343 populations, which can be used to inform conservation measures.

344

345 Azores laurel - *Laurus azorica* (Seub.) Franco

346 Common name in Portuguese: louro-da-terra and louro-bravo

347 The **Azores laurel** (*Laurus azorica*) is a micro or mesophanerophyte dioecious perennial tree  
348 endemic to the Azores Islands. It can be commonly found in submontane *Laurus* forests (Elias  
349 et al., 2016) but also occupies habitats such as coastal scrublands, among other native  
350 Azorean vegetation (Elias et al., 2016). It is listed as Least Concern in the IUCN Red List (Silva  
351 & Beech, 2016), yet it is facing habitat loss due to competition with invasive species and  
352 alteration of habitats resulting from the implementation of production forests and pastures  
353 (Borges Silva et al., 2018; Dutra Silva et al., 2019; Hortal et al., 2010; Lourenço et al., 2011).  
354 The genus *Laurus* L. (Lauraceae) is currently restricted to isolated refugia in the southern  
355 Black Sea area, Mediterranean Basin, Northern Africa, and the Macaronesian archipelagos.  
356 Its taxonomic classification remains uncertain, with some recognizing the existence of two  
357 species, *Laurus azorica* (Seub.) Franco, endemic from the Azores and *Laurus nobilis* L.  
358 However, while some molecular studies contradict this distinction (Rodríguez-Sánchez et al.,  
359 2009), others even recognise a third species, *Laurus novo-canariensis* Rivas Mart., Lousã,  
360 Fern.Prieto, E.Dias, J.C.Costa & C.Aguiar (Rivas-Martínez et al., 2002).

361 *Sampling and sampling processing:* Fresh branches with healthy leaves were collected using  
362 gardening scissors from one mature male individual in the Azores islands, specifically, São  
363 Miguel Island, in Lombadas. The branches were loosely wrapped in moist absorbent paper,  
364 placed inside an open plastic bag and sent on the same day through express mail to the  
365 University of Lisbon, where they were kept at -20°C until DNA extraction. The corresponding  
366 voucher duplicate was dried using standard herbarium procedures and stored in the AZB  
367 herbarium at the Biology Department of the Faculty of Sciences and Technology of the  
368 University of the Azores, with voucher AZB4382, and collector code LZ001. A second  
369 sampling was conducted during March 2023 and followed the same methodology. The Nagoya  
370 certificate has CCIR number 61/2021/DRCTD.

371 *Sequencing:* The HMW DNA extraction was performed at the Department of Plant Biology of  
372 the Faculty of Sciences of the University of Lisbon (FCUL), using a modified  
373 cetyltrimethylammonium bromide (CTAB) protocol (Doyle & Doyle, 1987, later modified by

374 Weising et al., 1994) adjusted based on the exchange of protocols within the ERGA  
375 community. DNA extraction was challenging, as extractions did not pass the quality control for  
376 PacBio Hi-Fi library preparation and sequencing. Leaf tissue was thus shipped in dry ice to  
377 the University of Florence, where the libraries for Pac-Bio were prepared and sequenced. The  
378 assembly of the genome is ready to start, and will involve ERGA-Portugal members, as well  
379 as institutions and national computational resources.

380 *Expected results:* A high-quality reference genome will allow performing phylogenetic and  
381 population genetic studies to reconstruct the evolutionary history of this group, contributing to  
382 clarify the diversity and taxonomy of the *Laurus* genus.

383

384 Black wheatear - *Oenanthe leucura* (Gmelin, 1789)

385 Common name in Portuguese: chasco-preto

386 The **black wheatear** (*Oenanthe leucura*) is a passerine bird species confined to North Africa  
387 and southwest Europe, a region where global environmental changes (namely desertification  
388 and agriculture abandonment) are impacting the distribution of many bird species (Fernández-  
389 Nogueira & Corbelle-Rico, 2018, 2020; Fusco et al., 2021). Black wheatears are no exception,  
390 with findings supporting that some edge populations have already vanished (France) whilst  
391 others continue to decrease (Western populations in Portugal and Spain). Despite the decline  
392 and fragmentation of these populations, an isolated population located in northern Portugal  
393 (~100 pairs) still subsists, confined mainly to cultivated lands (primarily vineyards and olive  
394 orchards) of the Portuguese section of the Douro River, internationally known for the Port  
395 wine. For this reason, despite being classified globally as Least Concern by the IUCN Red  
396 List, Portuguese populations are classified as Critically Endangered in the Portuguese Red  
397 List (Cabral et al., 2005).

398 *Sampling and sample processing:* Due to the small size of the black wheatear population in  
399 Portugal, the sacrifice of a bird would be unjustified and, therefore, ICNF authorised the  
400 collection of blood samples from live birds (Permit 720/2021/REC). Three individuals (one  
401 female and two males) were caught using spring traps at the Douro Valley, near the Tua river

402 mouth. Each bird was ringed, photographed and measured, and a blood sample (~100  
403 microliters) was taken by venepuncture at the ulnar vein with a microcapillary. Each blood  
404 sample was immediately stored in liquid nitrogen and transferred to a -80°C freezer on the  
405 same day.

406 *Sequencing and assembly.* Samples were sent to the ERGA partners in dry ice, to generate  
407 Omni-C (University of Antwerp) and PacBio HiFi data (University of Florence) from the female  
408 (C96097) and RNA-Sequencing (University of Antwerp) from one of the males (E29638). All  
409 DNA libraries were sequenced at the University of Florence. Long-read DNA sequencing was  
410 performed using two 8 million ZMW SMRT cells. Genome assembly was performed following  
411 the gold standard instructions implemented in the VGP Galaxy Pipeline (Batut et al., 2018;  
412 Lariviere et al., accessed June 2023). The genome manual curation and annotation is ongoing,  
413 involving ERGA-Portugal members, as well as institutions and national computational  
414 resources.

415 *Expected results:* A high-quality reference genome, combined with population genomics data,  
416 will allow to characterise the structure and history of this population, which is closely  
417 associated with anthropogenic agricultural lands in the Douro valley, yet geographically  
418 isolated from other Iberian populations that occur mostly in natural habitats. This isolation and  
419 fragmentation could have irreversible effects on the survival of these populations and their  
420 conservation may require the implementation of specific conservation measurements.  
421 Moreover, it will establish the baseline for genomic analysis of the Portuguese populations,  
422 providing information on individuals' relatedness and dispersal capacity.

423

#### 424 Portuguese crowberry - *Corema album* (L.) D. Don

425 Common name in Portuguese: Camarina and Camarinha

426 The **Portuguese crowberry** (*Corema album*) is a dioecious perennial woody shrub endemic  
427 to the Atlantic coast of the Iberian Peninsula (ssp. *album*), and to the Azores Islands (ssp.  
428 *azoricum*). In the Iberian Peninsula it inhabits coastal areas from Galicia to Gibraltar, and is  
429 an important species in sand dune habitats which are highly valuable for conservation

430 purposes (Council Directive 92/43/EEC of 21 May 1992 on the Conservation of Natural  
431 Habitats and of Wild Fauna and Flora, 1992). The dynamic nature of these coastal ecosystems  
432 provides a vast variety of habitats with unique floristic and animal richness. However, these  
433 dune systems face increasing disturbances as they support various economic and leisure  
434 activities, associated with the growth of the coastal population (Antunes et al., 2018). Because  
435 of habitat loss, *C. album* has been classified as Vulnerable on the Red List of Andalusia, Spain  
436 (Cabezudo et al., 2005). In the Azores Islands, it inhabits volcanic lava and ash fields (de  
437 Oliveira & Dale, 2012). The fruits of the Portuguese crowberry are edible, producing bioactive  
438 compounds that have been associated with chemoprotective activity and potential health-  
439 benefits (de Oliveira & Dale, 2012; Jacinto et al., 2021; Zunzunegui et al., 2006).

440 *Sampling and sample processing:* Two male and two female adult plants from the same  
441 population were sampled during two field trips to Azeitão, near Arrábida Natural Park. Young  
442 expanding leaves and fruits were collected after 48 hours of dark treatment (plant covered  
443 with light-opaque paper sheets with a few holes that allowed air flow), according to permits  
444 from ICNF (21PTLX00657D). Samples from leaves and fruits for DNA and RNA extraction  
445 were flash frozen at -20°C and -80°C, respectively. Voucher specimens were deposited at the  
446 Herbarium (LISU 270092) of the MUHNAC, Universidade de Lisboa.

447 *Sequencing:* The HMW DNA extraction was very challenging. First, it was performed at the  
448 Department of Plant Biology of FCUL using a CTAB protocol (Doyle & Doyle, 1987 later  
449 modified by Weising et al., 1994), which resulted in low DNA quality even after purification.  
450 Taking advantage of the ERGA network, we used a nuclei isolation protocol adapted from the  
451 ARIMA-HiC kit, shared by Narjes Yousefi (pers. comm.), a member of ERGA. This resulted in  
452 higher quality and quantity of extracted DNA, but it still did not pass the quality control for  
453 PacBio Hi-Fi library preparation and sequencing at the University of Florence. After several  
454 attempts, to prevent DNA degradation during shipping, leaf tissue was directly shipped in dry  
455 ice so that DNA extraction could be performed at the University of Florence, where the libraries  
456 for Pac-Bio were prepared and sequenced. The assembly of the genome is ongoing, involving  
457 ERGA-Portugal members and Biogenome Portugal institutions.

458 *Expected results:* Building a high-quality annotated reference genome will support ongoing  
459 projects on Portuguese crowberry and related plant species from coastal areas, and it will  
460 provide the opportunity to explore evolutionary biology questions associated with adaptations  
461 to coastal environments under a scenario of global changes. Moreover, the annotated  
462 reference genome will allow to uncover the genetic basis of the production of Portuguese  
463 crowberry's bioactive compounds associated with health benefits.

464

465 Cave ground beetle - *Trechus terceiranus* (Machado, 1988)

466 Common name in Portuguese: escaravelho-cavernícola-da-Ilha-Terceira

467 The **cave ground beetle** *Trechus terceiranus* (Coleoptera) is endemic to Terceira Island in  
468 the Azores and is restricted to subterranean habitats, including lava tubes. These tiny beetles  
469 have up to 4.3 mm in length, do not fly and have several morphological traits associated with  
470 their life in subterranean habitats (Machado, 1988). This species is listed as Vulnerable in the  
471 IUCN Red List (Borges & Amorim, 2018). Interestingly, several species of the genus *Trechus*  
472 occur in the Azores islands, most of which inhabit subterranean habitats (seven species), but  
473 two occur in surface habitats. One epigeal species, *T. terrabravensis* Borges, Serrano &  
474 Amorim, 2004, co-exists on Terceira Island with *T. terceiranus*.

475 *Sampling and sample processing:* Six adult individuals were sampled in a subterranean  
476 habitat (volcanic pit, lava tube) at 45-70m deep, using pitfall traps. Sampling was done  
477 according to permits issued by the Regional Government of the Azores (IRCC  
478 23/2021/DRCTD). Adult individuals were flash frozen with liquid nitrogen. Dried vouchers were  
479 stored at the Dalberto Teixeira Pombo entomological collection at the University of the Azores,  
480 Terceira (DTP-MF1091).

481 *Sequencing:* The frozen whole organisms were shipped in dry ice to the University of Lisbon,  
482 from where they were subsequently sent to Centro Nacional de Análisis Genómico (CNAG) –  
483 Barcelona for RNA extraction and sequencing; and to the University of Florence where DNA  
484 extraction was attempted. Due to constraints to obtain high quality HMW DNA for PacBio,

485 additional samples were sent in order to repeat the extraction to obtain enough DNA amount  
486 and quality for sequencing, which is ongoing.

487 *Expected results:* A high-quality reference genome will be crucial to pursue comparative  
488 genomics, phylogenetic and population genetic studies to reconstruct the evolutionary history  
489 of this group and to implement adequate conservation strategies. Furthermore, it will open the  
490 door to detect genes involved in adaptation to subterranean habitats and to determine the  
491 genetic architecture of traits related with such habitats (e.g., eye development, pigmentation  
492 and biological clock).

493

494

## 495 **Building a national network for biodiversity genomics: Biogenome Portugal**

### 496 Objectives and structure of Biogenome Portugal

497 While ERGA and ERGA-Portugal are a community of researchers, their collaboration in ERGA  
498 activities and in the pilot test has played a significant role in forming a national institutional  
499 network for biodiversity genomics, called Biogenome Portugal (BGP). This network is being  
500 formally established with the objective of fostering scientific collaboration, facilitating the  
501 exchange of expertise and infrastructure, organising advanced training programs, promoting  
502 Portugal's active participation in international biodiversity genomics initiatives, and  
503 contributing to knowledge transfer and outreach efforts. The network will comprise members  
504 with a diverse range of expertise, encompassing fields such as genomics, ecology, taxonomy,  
505 evolution, and more, spanning across various taxonomic groups.

506 By bringing together experts from different fields, the BGP network aims to facilitate the  
507 exchange of ideas, data, and resources that will lead to the development of new research  
508 projects centred on generating and using high quality reference genomes for Portuguese  
509 species, and address a plethora of biodiversity applications. Such collaboration also implies  
510 sharing institutional genomic and computational infrastructures and articulating with the  
511 national network of infrastructures related to BGP's areas of activity. Cooperation among

512 members is crucial for promoting the exchange of expertise, a key objective of the network.  
513 By sharing knowledge and skills, the network members will strengthen their collective  
514 expertise and coordinate efforts. Activities towards the transfer of skills and expertise include  
515 organising advanced training in the field of biodiversity genomics. Through these initiatives,  
516 BGP aims to foster the training of specialised human resources able to accompany the fast  
517 development of genomic data analysis tools. To broaden its scope, the network intends to  
518 facilitate the involvement of individual researchers, even if their institution is not formally  
519 affiliated with the network.

520 BGP also aims to disseminate the results of its activities to stakeholders and the general  
521 public, thus promoting knowledge transfer and outreach. These activities are important not  
522 only to enrich scientific literacy about the importance of genomics for biodiversity conservation,  
523 but also to promote and encourage the use of the genomic research findings across  
524 stakeholders.

525 Finally, BGP also aims to coordinate and synchronise Portuguese participation in global  
526 biodiversity genomics initiatives. By reinforcing and sharing the opportunities for institutional  
527 representation in internationally funded projects, BGP can promote the contribution of  
528 Portuguese expertise to important regional, continental or worldwide initiatives.

529

530 Articulation with the Portuguese National Strategy for the Conservation of Nature and  
531 Biodiversity 2030

532 The Portuguese National Strategy for the Conservation of Nature and Biodiversity (ENCNB  
533 2030; <https://dre.pt/dre/detalhe/resolucao-conselho-ministros/55-2018-115226936>) is the  
534 main legal Portuguese document that is used to frame all national policies until 2030. It  
535 recognizes the importance of Portugal's national biodiversity at both the European and global  
536 scale and acknowledges how the nation's biogeographic specificities have led to high levels  
537 of endemic and relict species, each with a unique evolutionary history and genetic  
538 composition, which is important to preserve. The national strategy has several main  
539 objectives, including planning and executing action promoting the conservation and recovery

540 of species and habitats at the national level and promoting the conservation of plant and  
541 animals' genetic diversity. These objectives align with the aims of the nascent BGP. The  
542 network can provide the knowledge-based capacities necessary for the conservation and  
543 sustainable use of the national marine, terrestrial and freshwater genetic resources. BGP can  
544 reinforce a science-based approach for species conservation action plans and guide complex  
545 conservation strategies, including *in situ* and *ex situ* conservation.

546

#### 547 Fostering national research infrastructures

548 In 2014, the national public agency for science, technology and innovation (Fundação para a  
549 Ciência e a Tecnologia, FCT) created a *National Roadmap for Research Infrastructures of*  
550 *Strategic Interest*. This roadmap aimed at promoting cooperation, excellence and  
551 internationalisation to strengthen national scientific infrastructures. A total of 56 Research  
552 Infrastructures (RIs) have been supported during the first funding cycle in key areas, such as  
553 the Environment, Health and Food, Social and Cultural Innovation, among others. BGP  
554 identified several RIs of interest, intersecting its area of intervention. Among these are  
555 GenomePT (the National Laboratory for Genome Sequencing and Analysis), Biodata.pt  
556 (ELIXIR PT - Portuguese Distributed Infrastructure for Biological Data), PORBIOTA  
557 (Portuguese E-Infrastructure for Information and Research on Biodiversity), EMBRC.PT  
558 (European Marine Biological Resource Centre – Portugal), and PRISC (Portuguese Research  
559 Infrastructure for Scientific Collections). Future action implies fostering the establishment of  
560 strategic cooperation with the active infrastructures, which can be facilitated by the current  
561 presence of several institutions of BGP in the RIs, pending the continuation of the current  
562 roadmap.

563

#### 564 Engaging the community: establishing training programs, supporting the generation of 565 additional reference genomes and expanding funding opportunities

566 The analysis of reference genomes at scale requires standardised procedures for the  
567 sampling and storing of the biological material, as well as sequencing and analysis of the

568 genomic data. Researchers working to analyse and utilise these cutting-edge genomic  
569 resources need specific training, to maintain quality standards and streamline procedures.  
570 The development of bioinformatic pipelines by the larger ERGA community using open-source  
571 platforms such as Galaxy (The Galaxy Community, 2022) allows quick implementation of  
572 analyses across research centres. Within the BGP network, an online training programme is  
573 being implemented to cover different topics across pipelines of genomic analyses, such as  
574 genome and transcriptome assembly, annotation, comparative and population genomics. This  
575 programme aims to expand and facilitate training to enhance expertise in genomic analyses  
576 across the Portuguese research community. The first introductory course to Galaxy and VGP  
577 assembly pipeline took place in the first semester of 2023.

578 Further expertise sharing will come from the expansion of reference genomes projects under  
579 the BGP umbrella, which meets the technical and quality standards set by ERGA. For  
580 example, two reference genomes of Anthozoan species belonging to octocorals are being  
581 generated under the BGP initiative: the pink sea fan (*Eunicella verrucosa*) and the dead man's  
582 fingers (*Alcyonium digitatum*). Octocoral genomes remain poorly studied, with only ~10  
583 reference genomes available from the more than 3,000 described species. These two species  
584 belong to a pool of octocorals for which the sequencing of reference genomes was supported  
585 by EASI-Genomics (H2020 824110; Project ID 10240, CoGeCo). The two species are widely  
586 distributed along the Portuguese coast in sublittoral rocky habitats, and sequencing, genome  
587 assembly and annotations are underway.

588 Attracting national and international funding will be crucial to support the implementation of  
589 BGP's activities. Working as a network will lead to stronger research proposals, more  
590 successful grant applications, and increase collective participation in international consortia to  
591 foster biodiversity genomics cutting-edge research and knowledge transfer. This strategy will  
592 allow consolidating, strengthening and expanding the BGP network and its impact on  
593 fundamental and applied science and innovation.

594

595 Producing high quality chromosome-level assemblies of Portuguese species

596 The alignment of BGP with ERGA can foster the multiplication of opportunities for sequencing  
597 projects for Portuguese biodiversity. This aligns with the aim of ERGA of “propagating  
598 guidelines for state-of-the-art genome establishment through training and knowledge transfer”.  
599 Produced genomes can therefore take advantage of the standard sampling and molecular  
600 protocols and bioinformatics pipelines for sequencing, genome assembly and annotation  
601 established by ERGA, and be included under the ERGA umbrella. While promoting the  
602 visibility and accessibility of the genomes to the ERGA network and beyond, those genomes  
603 will contribute to the Reference Genome Atlas of European biodiversity. Such parallel  
604 initiatives under the BGP and ERGA umbrella, coordinated with ERGA projects, will expedite  
605 the production of a genomic inventory for all Portuguese eukaryotic species. Those efforts are  
606 invaluable to enable the use of genomic applications to preserve and protect the country’s rich  
607 biodiversity.

608

609


610 **Figures**



611

612 Fig.1 - a) Spatial context of the Portuguese territory, including the two Autonomous Regions (Azores  
613 and Madeira) b) Distribution of the Portugal-based members of ERGA (ERGA-Portugal), including their  
614 host institutions (May 31, 2023).

615



616

617 Fig.2 – Approximate distributions and sampling localities of ERGA-Portugal species sequenced in

618 ERGA's pilot effort: a) Iberian minnowcarp - *Anaecypris hispanica*; b) Iberian hare - *Lepus granatensis*;

619 c) Azores laurel - *Laurus azorica*; d) Black wheatear - *Oenanthe leucura*; e) Portuguese crowberry -

620 *Corema album*; f) Cave ground beetle - *Trechus terceiranus*. Picture credits: a) Carlos Carrapato; b)

621 Pedro Moreira; c) Mónica Moura; d) Ricardo Jorge Lopes; e) Manuela Sim-Sim; f) Javier Torrent

622 (Azorean Biodiversity Group). Species distributions coloured according to terrestrial (green) or aquatic

623 (blue) territories. Sampling localities marked with a black dot ● except for the Iberian minnowcarp (a)

624 that is double marked, as it was sampled in the Aquário Vasco da Gama – Lisbon ⬠, but it is originally

625 from a population of River Chança ●. Distributions adjusted from the IUCN Red List of Threatened

626 Species Website (<https://www.iucnredlist.org>), except for the Portuguese crowberry, which was based

627 on information from the Anthos geographical information system for Spanish Flora ([www.anthos.es](http://www.anthos.es)).

628

629

630

630 **Acknowledgements**

631 **Co-authors:** This work was supported by National Funds through FCT-Fundação para a  
632 Ciência e a Tecnologia in the scope of the project UIDP/50027/2020. Strategic funding from  
633 FCT to cE3c and BioISI Research Units (UIDB/00329/2020 and UIDB/04046/2020) and to the  
634 associate laboratory CHANGE (LA/P/0121/2020) are gratefully acknowledged. Financing by  
635 Azorean government regional funds, SRCCTD/DRCTD, project ERGA TAXA AÇORES  
636 (M2.2.A/PROJ.INT/A/001/2021 – ERGA taxa Açores). J. Melo-Ferreira acknowledges support  
637 from FCT – contract 2021.00150.CEECIND and project grant HybridChange PTDC/BIA-  
638 EVL/1307/2020, via Portuguese national funds, and GenomePT for supporting CIBIO's  
639 genomic infrastructure. P.C. Alves acknowledges support by NORTE-01-0246-FEDER-  
640 000063, supported by Norte Portugal Regional Programme. T. Brown was supported by DFG  
641 (INST 269/768-1). FCT/MEC support the research contract to da L. P. Silva  
642 (CEEIND/02064/2017). R.J. Lopes was supported by a FCT – Transitory Norm contract  
643 (DL57/2016/CP1440/CT0006). P.A.V. Borges was funded by project FCT-UIDB/00329/2020-  
644 2024 (Thematic Line 1 – integrated ecological assessment of environmental change on  
645 biodiversity). S. L. Mendes is supported by an FCT PhD studentship  
646 (SFRH/BD/145153/2019). J. M. Moreno is supported by an FCT PhD studentship  
647 (SFRH/BD/143199/2019). This study received Portuguese national funds from FCT through  
648 project UIDB/04326/2020. H. Svardal was supported by the Flemish University Research  
649 Fund (BOF). A. Mouton was supported by a Postdoctoral Researcher Fellowship of the Fonds  
650 de la Recherche Scientifique – FNRS. Portuguese National Funds through FCT support the  
651 research contract to A. Crottini (2020.00823.CEECIND/CP1601/CT0003) and R. Faria  
652 (contract 2020.00275.CEECIND; grant Invcontinuum – PTDC/BIA-EVL/1614/2021). C.  
653 Fernandes thanks the support of cE3c through an assistant researcher contract (Fciência.ID  
654 contract #366) and FCT for Portuguese National Funds attributed to cE3c within the strategic  
655 project UID/BIA/00329/2020; C. Rodriguez Fernandes also thanks FPUL for a contract of  
656 invited assistant professor. Operational Programme (NORTE2020), under the PORTUGAL  
657 2020 Partnership Agreement, through the European Regional. Project 24rant HYBRIDOMICS

658 – PTDC/BIA-EVL/4345/2021, via Portuguese national funds. We gratefully acknowledge the  
659 generous support of the Max Planck Society for this research. Additional support was obtained  
660 from Horizon Europe under the Biodiversity, Circular Economy and Environment (REA.B.3);  
661 co-funded by the Swiss State Secretariat for Education, Research and Innovation (SERI)  
662 under contract number 22.00173; and by the UK Research and Innovation (UKRI) under the  
663 Department for Business, Energy and Industrial Strategy’s Horizon Europe Guarantee  
664 Scheme.

665 **ERGA Hubs:** Illumina and PacBio HiFi sequencing (RNA + DNA) was supported by the  
666 sequencing facility of the Department of Biology, University of Florence through the  
667 Departments of Excellence programme funded by the Italian Ministry for University and  
668 Research. We thank the Antwerp University Hospital Center of Medical Genetics and Jarl  
669 Bastianen for access to sequencing library quality control equipment. The Hi-C library  
670 preparation for *Trechus terceiranus* is being done at the Metazoa Phylogenomics Lab, Institute  
671 of Evolutionary Biology (CSIC-University Pompeu Fabra) through a collaboration with Judit  
672 Salces-Ortiz, Nuria Escudero and Rosa Fernández.

673 **ERGA infrastructure:** We acknowledge access to the storage resources at Barcelona  
674 Supercomputing Center, which are partially funded from European Union H2020-  
675 INFRAEOSC-2018-2020 programme through the DICE project (Grant Agreement no.  
676 101017207). We would like to thank Alisha Ahamed, Josephine Burgin, Joana Paupério,  
677 Jeena Rajan and Guy Cochrane from the European Nucleotide Archive (ENA) for their support  
678 regarding data coordination and submission of genomic data.

679 **Commercial Partners:** We would like to acknowledge and thank all supplier partners that  
680 have kindly donated kits and reagents to the ERGA pilot Library Preparation Hubs to support  
681 species without funding to produce the generation of high-quality genomes and annotations.  
682 This support has been key to embedding a culture of diversity, equity, inclusion, and justice in  
683 the Pilot Project. Specifically, we want to thank Dovetail Genomics, Part of Cantata Bio LLC,  
684 especially Mark Daly, Thomas Swale and Lily Shuie; Arima Genomics; PacBio; Integrated

685 DNA Technologies (IDT); MagBio Genomics Europe GmbH; Zymo Research; Agilent  
686 Technologies; Fisher Scientific Spain; Illumina Inc.

687

688 **Author contributions**

689 **Species Ambassador or Executive coordination:** Marques JP, Alves PC, Amorim IR,  
690 Lopes RJ, Moura M, Myers EW, Sim-Sim M, Sousa-Santos C and Melo-Ferreira J

691 **Genome Team Member:** Marques JP, Alves PC, Amorim IR, Lopes RJ, Moura M, Myers EW,  
692 Sim-Sim M, Sousa-Santos C, Alves MJ, Borges PAV, Brown T, Carneiro M, Carrapato C,  
693 Ceríaco LMP, Ciofi C, da Silva LP, Diedericks G, Diroma MA, Farelo L, Formenti G, Gil F,  
694 Grilo ML, Ianucci A, Leitão HG, Máguas C, Mc Cartney AM, Mendes SL, Moreno JM, Morselli  
695 M, Mouton A, Natali C, Pereira F, Rego RMC, Resendes R, Roxo G, Svardal H, Trindade H,  
696 Vicente S, Winkler S, Sousa VC and Melo-Ferreira J

697 **ERGA-Portugal Member:** Marques JP, Alves PC, Amorim IR, Lopes RJ, Moura M, Sim-Sim  
698 M, Sousa-Santos C, Alves MJ, Carneiro M, Ceríaco LMP, Trindade H, Vicente S, Alvarenga  
699 M, Amaral AJ, Antunes A, Campos PF, Canário AVM, Castilho R, Castro LFC, Crottini A,  
700 Cunha MV, Espregueira Themudo G, Esteves PJ, Faria R, Rodríguez Fernandes C, Ledoux  
701 JB, Louro B, Magalhães S, Paulo OS, Pearson GA, Pimenta J, Santos TL, Serrao EA, Pina-  
702 Martins F, Sousa VC and Melo-Ferreira J

703 **ERGA-Portugal Coordination:** Sousa VC and Melo-Ferreira J

704 **Write Manuscript:** Marques JP, Amorim IR, Lopes RJ, Moura M, Sim-Sim M, Sousa-Santos  
705 C, Trindade H, Espregueira Themudo G, Rodríguez Fernandes C, Ledoux JB, Paulo OS,  
706 Sousa VC and Melo-Ferreira J

707 **Revise Manuscript:** Marques JP, Alves PC, Amorim IR, Lopes RJ, Moura M, Sim-Sim M,  
708 Sousa-Santos C, Ciofi C, Mc Cartney AM, Mouton A, Trindade H, Alvarenga M, Amaral AJ,  
709 Antunes A, Canário AVM, Castilho R, Castro LFC, Crottini A, Cunha MV, Espregueira  
710 Themudo G, Esteves PJ, Faria R, Rodríguez Fernandes C, Ledoux JB, Louro B, Magalhães  
711 S, Paulo OS, Santos TL, Serrao EA, Pina-Martins F, Sousa VC and Melo-Ferreira J

712 **Read and Approve Manuscript:** all authors

713 **References**

- 714 Águeda-Pinto, A., Lemos de Matos, A., Abrantes, M., Kraberger, S., Rivalde, M. A., Gortázar, C.,  
715 McFadden, G., Varsani, A., & Esteves, P. J. (2019). Genetic Characterization of a Recombinant  
716 *Myxoma virus* in the Iberian Hare (*Lepus granatensis*). *Viruses*, 11(6), 530.  
717 <https://doi.org/10.3390/v11060530>
- 718 Alves, P. C., Acevedo, P., & Melo-Ferreira, J. (2023). Iberian Hare *Lepus granatensis* Rosenhauer,  
719 1856. In K. Hackländer & F. E. Zachos (Eds.), *Handbook of the Mammals of Europe* (pp. 1–23).  
720 Springer International Publishing. [https://doi.org/10.1007/978-3-319-65038-8\\_8-1](https://doi.org/10.1007/978-3-319-65038-8_8-1)
- 721 Antunes, C., Pereira, A. J., Fernandes, P., Ramos, M., Ascensão, L., Correia, O., & Máguas, C.  
722 (2018). Understanding plant drought resistance in a Mediterranean coastal sand dune ecosystem:  
723 Differences between native and exotic invasive species. *Journal of Plant Ecology*, 11(1), 26–38.  
724 <https://doi.org/10.1093/jpe/rtx014>
- 725 Batut, B., Hiltemann, S., Bagnacani, A., Baker, D., Bhardwaj, V., Blank, C., Bretaudeau, A., Brillet-  
726 Guéguen, L., Čech, M., Chilton, J., Clements, D., Doppelt-Azeroual, O., Erxleben, A., Freeberg, M. A.,  
727 Gladman, S., Hoogstrate, Y., Hotz, H.-R., Houwaart, T., Jagtap, P., ... Grüning, B. (2018).  
728 Community-Driven Data Analysis Training for Biology. *Cell Systems*, 6(6), 752-758.e1.  
729 <https://doi.org/10.1016/j.cels.2018.05.012>
- 730 Böhne, A., Fernández, R., Leonard, J. A., McCartney, A. M., McTaggart, S., Melo-Ferreira, J.,  
731 Monteiro, R., Oomen, R. A., Pettersson, O. V., & Struck, T. H. (2023). Contextualising samples:  
732 Supporting reference genomes for European biodiversity through sample and associated metadata  
733 collection (p. 2023.06.28.546652). bioRxiv. <https://doi.org/10.1101/2023.06.28.546652>
- 734 Borges, P.A.V. & Amorim, I.R. 2018. *Trechus terceiranus*. The IUCN Red List of Threatened  
735 Species 2018: e.T97125072A99166594. <https://dx.doi.org/10.2305/IUCN.UK.2018->  
736 1.RLTS.T97125072A99166594.en. Accessed on 22 June 2023.
- 737 Borges Silva, L., Lourenço, P., Teixeira, A., Azevedo, E. B., Alves, M., Elias, R. B., & Silva, L. (2018).  
738 Biomass valorization in the management of woody plant invaders: The case of *Pittosporum*  
739 *undulatum* in the Azores. *Biomass and Bioenergy*, 109, 155–165.  
740 <https://doi.org/10.1016/j.biombioe.2017.12.025>
- 741 Browne, L., Wright, J. W., Fitz-Gibbon, S., Gugger, P. F., & Sork, V. L. (2019). Adaptational lag to  
742 temperature in valley oak (*Quercus lobata*) can be mitigated by genome-informed assisted gene flow.

743 *Proceedings of the National Academy of Sciences*, 116(50), 25179–25185.  
744 <https://doi.org/10.1073/pnas.1908771116>

745 Cabezudo, B., Talavera, S., Blanca, G., Salazar, C., Cueto, M., Valdés, B., Hernández-Bermejo, J. E.,  
746 Herrera, C. M., Rodríguez-Hiraldo, C., & Navas, D. (2005). *Lista Roja de la flora vascular de*  
747 *Andalucía*. Consejería de Medio Ambiente, Junta de Andalucía, Sevilla.

748 Cabral, M. J., Almeida, J., Almeida, P. R., Dellinger, T., Ferrand de Almeida, N., Oliveira, M. E.,  
749 Palmeirim, J. M., Queirós, A. I., Rogado, L., & Santos-Reis, M. (2005). *Livro Vermelho dos*  
750 *Vertebrados de Portugal*. Instituto da Conservação da Natureza.  
751 <http://dspace.uevora.pt/rdpc/handle/10174/6006>

752 Cardoso, A. (2023, February 7). Updating *Anaocypris hispanica* distribution and conservation status  
753 in Portugal. *FiSHMED Journal*. <https://www.fishmedjournal.sibic.org/FiSHMED.2022.001/>

754 Cartney, A. M. M., Formenti, G., Mouton, A., Panis, D. D., Marins, L. S., Leitão, H. G., Diedericks, G.,  
755 Kirangwa, J., Morselli, M., Salces-Ortiz, J., Escudero, N., Iannucci, A., Natali, C., Svoldal, H.,  
756 Fernández, R., Pooter, T. D., Joris, G., Strazisar, M., Wood, J., ... Mazzoni, C. J. (2023). The  
757 European Reference Genome Atlas: Piloting a decentralised approach to equitable biodiversity  
758 genomics (p. 2023.09.25.559365). *bioRxiv*. <https://doi.org/10.1101/2023.09.25.559365>

759 Ceballos, G., Ehrlich, P. R., & Dirzo, R. (2017). Biological annihilation via the ongoing sixth mass  
760 extinction signaled by vertebrate population losses and declines. *Proceedings of the National*  
761 *Academy of Sciences*, 114(30), E6089–E6096. <https://doi.org/10.1073/pnas.1704949114>

762 Council Directive 92/43/EEC of 21 May 1992 on the conservation of natural habitats and of wild fauna  
763 and flora, CONSIL, 206 OJ L (1992). <http://data.europa.eu/eli/dir/1992/43/oj/eng>

764 Cowie, R. H., Bouchet, P., & Fontaine, B. (2022). The Sixth Mass Extinction: Fact, fiction or  
765 speculation? *Biological Reviews*, 97(2), 640–663. <https://doi.org/10.1111/brv.12816>

766 Crivelli, A. J. (2006). IUCN Red List of Threatened Species: *Anaocypris hispanica*. *IUCN Red List of*  
767 *Threatened Species*. <https://www.iucnredlist.org/en>

768 Dalton, K. P., Martín, J. M., Nicieza, I., Podadera, A., de Llano, D., Casais, R., Gimenez, S., Badiola,  
769 I., Agüero, M., Duran, M., Buitrago, D., Romero, L. J., García, E., & Parra, F. (2019). *Myxoma virus*  
770 jumps species to the Iberian hare. *Transboundary and Emerging Diseases*, 66(6), 2218–2226.  
771 <https://doi.org/10.1111/tbed.13296>

772 Dauphin, B., Rellstab, C., Schmid, M., Zoller, S., Karger, D. N., Brodbeck, S., Guillaume, F., &  
773 Gugerli, F. (2021). Genomic vulnerability to rapid climate warming in a tree species with a long  
774 generation time. *Global Change Biology*, 27(6), 1181–1195. <https://doi.org/10.1111/gcb.15469>

775 De Miguel, R., Pino, E., Ramiro, A., Aranda, F., Peña, J. P., Doadrio, I., & Fernández-Delgado, C.  
776 (2010). On the occurrence of *Anaecypris hispanica*, an extremely endangered Iberian endemism, in  
777 the Guadalquivir River basin. *Journal of Fish Biology*, 76(6), 1454–1465.  
778 <https://doi.org/10.1111/j.1095-8649.2010.02574.x>

779 de Oliveira, P. B., & Dale, A. (2012). *Corema album* (L.) D. Don, the white crowberry – a new crop.  
780 *Journal of Berry Research*, 2(3), 123–133. <https://doi.org/10.3233/JBR-2012-033>

781 Doyle, J. J., & Doyle, J. L. (1987). A rapid DNA isolation procedure for small quantities of fresh leaf  
782 tissue. *Phytochemical Bulletin*, 19: 11–15.

783 Dutra Silva, L., Brito de Azevedo, E., Vieira Reis, F., Bento Elias, R., & Silva, L. (2019). Limitations of  
784 Species Distribution Models Based on Available Climate Change Data: A Case Study in the Azorean  
785 Forest. *Forests*, 10(7), Article 7. <https://doi.org/10.3390/f10070575>

786 Elias, R. B., Gil, A., Silva, L., Fernández-Palacios, J. M., Azevedo, E. B., & Reis, F. (2016). Natural  
787 zonal vegetation of the Azores Islands: Characterization and potential distribution. *Phytocoenologia*,  
788 107–123. <https://doi.org/10.1127/phyto/2016/0132>

789 Ette, J.-S., & Geburek, T. (2021). Why European biodiversity reporting is not reliable. *Ambio*, 50(4),  
790 929–941. <https://doi.org/10.1007/s13280-020-01415-8>

791 Fernández-Nogueira, D., & Corbelle-Rico, E. (2018). Land Use Changes in Iberian Peninsula 1990–  
792 2012. *Land*, 7(3), Article 3. <https://doi.org/10.3390/land7030099>

793 Fernández-Nogueira, D., & Corbelle-Rico, E. (2020). Determinants of Land Use/Cover Change in the  
794 Iberian Peninsula (1990–2012) at Municipal Level. *Land*, 9(1), Article 1.  
795 <https://doi.org/10.3390/land9010005>

796 Florencio, M., Patiño, J., Nogué, S., Traveset, A., Borges, P. A. V., Schaefer, H., Amorim, I. R.,  
797 Arnedo, M., Ávila, S. P., Cardoso, P., de Nascimento, L., Fernández-Palacios, J. M., Gabriel, S. I., Gil,  
798 A., Gonçalves, V., Haroun, R., Illera, J. C., López-Darias, M., Martínez, A., ... Santos, A. M. C. (2021).  
799 Macaronesia as a Fruitful Arena for Ecology, Evolution, and Conservation Biology. *Frontiers in*  
800 *Ecology and Evolution*, 9, 718169. <https://doi.org/10.3389/fevo.2021.718169>

801 Formenti, G., Theissingner, K., Fernandes, C., Bista, I., Bombarely, A., Bleidorn, C., Ciofi, C., Crottini,  
802 A., Godoy, J. A., Höglund, J., Malukiewicz, J., Mouton, A., Oomen, R. A., Paez, S., Palsbøll, P. J.,  
803 Pampoulie, C., Ruiz-López, M. J., Svardal, H., Theofanopoulou, C., ... Zammit, G. (2022). The era of  
804 reference genomes in conservation genomics. *Trends in Ecology & Evolution*, 37(3), 197–202.  
805 <https://doi.org/10.1016/j.tree.2021.11.008>

806 Fusco, J., Walker, E., Papaix, J., Debolini, M., Bondeau, A., & Barnagaud, J.-Y. (2021). Land Use  
807 Changes Threaten Bird Taxonomic and Functional Diversity Across the Mediterranean Basin: A  
808 Spatial Analysis to Prioritize Monitoring for Conservation. *Frontiers in Ecology and Evolution*, 9.  
809 <https://www.frontiersin.org/articles/10.3389/fevo.2021.612356>

810 Gómez, A., & Lunt, D. H. (2007). Refugia within Refugia: Patterns of Phylogeographic Concordance  
811 in the Iberian Peninsula. In S. Weiss & N. Ferrand (Eds.), *Phylogeography of Southern European*  
812 *Refugia: Evolutionary perspectives on the origins and conservation of European biodiversity* (pp. 155–  
813 188). Springer Netherlands. [https://doi.org/10.1007/1-4020-4904-8\\_5](https://doi.org/10.1007/1-4020-4904-8_5)

814 Habel, J. C., Rasche, L., Schneider, U. A., Engler, J. O., Schmid, E., Rödder, D., Meyer, S. T., Trapp,  
815 N., Sos del Diego, R., Eggermont, H., Lens, L., & Stork, N. E. (2019). Final countdown for biodiversity  
816 hotspots. *Conservation Letters*, 12(6), e12668. <https://doi.org/10.1111/conl.12668>

817 Hewitt, G. M. (2004). Genetic consequences of climatic oscillations in the Quaternary. *Philosophical*  
818 *Transactions of the Royal Society B: Biological Sciences*, 359(1442), 183–195.

819 Hohenlohe, P. A., Funk, W. C., & Rajora, O. P. (2021). Population genomics for wildlife conservation  
820 and management. *Molecular Ecology*, 30(1), 62–82. <https://doi.org/10.1111/mec.15720>

821 Hortal, J., Borges, P. A. V., Jiménez-Valverde, A., de Azevedo, E. B., & Silva, L. (2010). Assessing  
822 the areas under risk of invasion within islands through potential distribution modelling: The case of  
823 *Pittosporum undulatum* in São Miguel, Azores. *Journal for Nature Conservation*, 18(4), 247–257.  
824 <https://doi.org/10.1016/j.jnc.2009.11.002>

825 Jacinto, J., Giovanetti, M., Oliveira, P. B., Valdivieso, T., Máguas, C., & Alegria, C. (2021). Quality  
826 attributes of cultivated white crowberries (*Corema album* (L.) D. Don) from a multi-origin clonal field.  
827 *Euphytica*, 217(3), 40. <https://doi.org/10.1007/s10681-021-02767-2>

828 Karger, D. N., Kessler, M., Lehnert, M., & Jetz, W. (2021). Limited protection and ongoing loss of  
829 tropical cloud forest biodiversity and ecosystems worldwide. *Nature Ecology & Evolution*, 5(6), Article  
830 6. <https://doi.org/10.1038/s41559-021-01450-y>

831 Lariviere, D., Ostrovsky, A., Gallardo, C., Syme, A., Abueg, L., Pickett, B., Formenti, G., & Sozzoni, M.  
832 VGP assembly pipeline. (Galaxy Training Materials); <https://training.galaxyproject.org/training->  
833 [material/topics/assembly/tutorials/vgp\\_genome\\_assembly/tutorial.html](https://training.galaxyproject.org/training-material/topics/assembly/tutorials/vgp_genome_assembly/tutorial.html), Online; accessed June 2023

834 Lourenço, P., Medeiros, V., Gil, A., & Silva, L. (2011). Distribution, habitat and biomass of  
835 *Pittosporum undulatum*, the most important woody plant invader in the Azores Archipelago. *Forest*  
836 *Ecology and Management*, 262(2), 178–187. <https://doi.org/10.1016/j.foreco.2011.03.021>

837 Machado, A. (1988). Two new cavernicolous species of the genus *Trechus* Clairv. From the Azores  
838 (Coleoptera, Carabidae). *Bocagiana*, 119, 1-8.

839 Mathias, M. L., Fonseca, C., Rodrigues, L., Grilo, C., Lopes-Fernandes, M., Palmeirim, J. M., Santos-  
840 reis, M., Alves, P. C., Cabral, J. A., Ferreira, M., Mira, A., Eira, C., Negrões, N., Paupério, J., Pita, R.,  
841 Rainho, A., Rosalino, L. M., Tapisso, J. T., & Vingada, J. (2023). *Livro Vermelho dos Mamíferos de*  
842 *Portugal Continental*. FCIências.ID / ICNF.

843 Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B., & Kent, J. (2000). Biodiversity  
844 hotspots for conservation priorities. *Nature*, 403(6772), Article 6772. <https://doi.org/10.1038/35002501>

845 O'Hara, C. C., Frazier, M., & Halpern, B. S. (2021). At-risk marine biodiversity faces extensive,  
846 expanding, and intensifying human impacts. *Science*, 372(6537), 84–87.  
847 <https://doi.org/10.1126/science.abe6731>

848 Pauly, D., Zeller, D., & Palomares, M. L. D. (2020). Sea around us. Concepts, Design and Data.

849 Perea, S., Böhme, M., Zupančič, P., Freyhof, J., Šanda, R., Özuluğ, M., Abdoli, A., & Doadrio, I.  
850 (2010). Phylogenetic relationships and biogeographical patterns in Circum-Mediterranean subfamily  
851 Leuciscinae (Teleostei, Cyprinidae) inferred from both mitochondrial and nuclear data. *BMC*  
852 *Evolutionary Biology*, 10(1), 265. <https://doi.org/10.1186/1471-2148-10-265>

853 Ralls, K., Sunnucks, P., Lacy, R. C., & Frankham, R. (2020). Genetic rescue: A critique of the  
854 evidence supports maximizing genetic diversity rather than minimizing the introduction of putatively  
855 harmful genetic variation. *Biological Conservation*, 251, 108784.  
856 <https://doi.org/10.1016/j.biocon.2020.108784>

857 Rivas-Martínez, S., Díaz, T. E., Fernández-González, F., Izco, J., Loidi, J., Lousa, M., & Penas, A.  
858 (2002). Vascular plant communities of Spain and Portugal: Addenda to the syntaxonomical checklist  
859 of 2001. 2. *Itinera Geobotanica (España)*.  
860 [https://scholar.google.com/scholar\\_lookup?title=Vascular+plant+communities+of+Spain+and+Portuga](https://scholar.google.com/scholar_lookup?title=Vascular+plant+communities+of+Spain+and+Portuga)

861 %3A+addenda+to+the+syntaxonomical+checklist+of+2001.+2&author=Rivas-  
862 Mart%C3%ADnez%2C+S.+%28Universidad+Complutense+de+Madrid.+Facultad+de+Farmacia%29  
863 &publication\_year=2002

864 Rodríguez-Sánchez, F., Guzmán, B., Valido, A., Vargas, P., & Arroyo, J. (2009). Late Neogene  
865 history of the laurel tree (*Laurus* L., Lauraceae) based on phylogeographical analyses of  
866 Mediterranean and Macaronesian populations. *Journal of Biogeography*, 36(7), 1270–1281.  
867 <https://doi.org/10.1111/j.1365-2699.2009.02091.x>

868 Segelbacher, G., Bosse, M., Burger, P., Galbusera, P., Godoy, J. A., Helsen, P., Hvilsom, C., Iacolina,  
869 L., Kahric, A., Manfrin, C., Nonic, M., Thizy, D., Tsvetkov, I., Veličković, N., Vilà, C., Wisely, S. M., &  
870 Buzan, E. (2022). New developments in the field of genomic technologies and their relevance to  
871 conservation management. *Conservation Genetics*, 23(2), 217–242. [https://doi.org/10.1007/s10592-](https://doi.org/10.1007/s10592-021-01415-5)  
872 [021-01415-5](https://doi.org/10.1007/s10592-021-01415-5)

873 Seixas, F. A., Boursot, P., & Melo-Ferreira, J. (2018). The genomic impact of historical hybridization  
874 with massive mitochondrial DNA introgression. *Genome Biology*, 19(1), 91.  
875 <https://doi.org/10.1186/s13059-018-1471-8>

876 Silva, L., & Beech, E. (2016). IUCN Red List of Threatened Species: *Laurus azorica*. *IUCN Red List of*  
877 *Threatened Species*. <https://www.iucnredlist.org/en>

878 Soriguer, R., & Carro, F. (2018). IUCN Red List of Threatened Species: *Lepus granatensis*. *IUCN Red*  
879 *List of Threatened Species*. <https://www.iucnredlist.org/en>

880 Sousa-Santos, C., Robalo, J. I., Francisco, S. M., Carrapato, C., Cardoso, A. C., & Doadrio, I. (2014).  
881 Metapopulations in temporary streams – The role of drought–flood cycles in promoting high genetic  
882 diversity in a critically endangered freshwater fish and its consequences for the future. *Molecular*  
883 *Phylogenetics and Evolution*, 80, 281–296. <https://doi.org/10.1016/j.ympev.2014.08.007>

884 The Galaxy Community. (2022). The Galaxy platform for accessible, reproducible and collaborative  
885 biomedical analyses: 2022 update. *Nucleic Acids Research*, 50(W1), W345–W351.  
886 <https://doi.org/10.1093/nar/gkac247>

887 Theissinger, K., Fernandes, C., Formenti, G., Bista, I., Berg, P. R., Bleidorn, C., Bombarely, A.,  
888 Crottini, A., Gallo, G. R., Godoy, J. A., Jentoft, S., Malukiewicz, J., Mouton, A., Oomen, R. A., Paez,  
889 S., Palsbøll, P. J., Pampoulie, C., Ruiz-López, M. J., Secomandi, S., ... Zammit, G. (2023). How

890 genomics can help biodiversity conservation. *Trends in Genetics*.  
891 <https://doi.org/10.1016/j.tig.2023.01.005>  
892 Weising, K., Nybom, H., Pfenninger, M., Wolff, K., & Meyer, W. (1994). DNA Fingerprinting in Plants  
893 and Fungi. CRC Press, Boca Raton, 338 pp. Taylor & Francis.  
894 <https://books.google.pt/books?id=AtkHllks64YC>  
895 Zunzunegui, M., Díaz Barradas, M. C., Clavijo, A., Alvarez Cansino, L., Ain Lhout, F., & García Novo,  
896 F. (2006). Ecophysiology, growth timing and reproductive effort of three sexual forms of *Corema*  
897 *album* (Empetraceae). *Plant Ecology*, 183(1), 35–46. <https://doi.org/10.1007/s11258-005-9004-4>