1	Cumulative	impacts	of invasiv	ve plant	t species i	in British
---	------------	---------	------------	----------	-------------	------------

2 Columbia's riparian ecosystems: a systematic map protocol

3 4	
5	*Fabio Mologni [,]
6	Chandra E. Moffat ^{2,1}
7	Jason Pither
8	
9	1. Department of Biology, and Institute for Biodiversity, Resilience, and Ecosystem Services
10	I.K. Barber Faculty of Science,
11	University of British Columbia Okanagan
12	1177 Research Road, Kelowna, BC V1V 1V7, Canada
13	
14	2. Agriculture and Agri-Food Canada
15	Summerland Research and Development Centre
16	4200 Highway #97, Summerland, BC V0H 1Z0
17	
18	* Corresponding author
19	Postal address: Department of Biology, University of British Columbia Okanagan, 1177
20	Research Road, Kelowna, BC V1V 1V7, Canada
21	Phone: +12369709033

22 Email: fabio.mologni@ubc.ca

23 ORCiD: https://orcid.org/0000-0003-4750-9974

24

25 Other authors email:

26 Chandra E. Moffat (chandra.moffat@agr.gc.ca)

27 Jason Pither (jason.pither@ubc.ca)

28

29 Abstract

30 Background

31 Globally, the structure and functioning of foreshore and riparian ecosystems are being 32 dramatically impacted by non-native invasive plant species. Invasive species can 33 outcompete and replace native species, modify geochemical and hydraulic cycles, alter 34 trophic processes and change the composition and structure of communities above and 35 below ground. However, these impacts are often investigated in isolation, even though one 36 invasive species might increase or mitigate the impacts of others (i.e. cumulative impacts), 37 potentially with cascading effects. Although cumulative impacts have long been studied within other environmental contexts, research on the cumulative impacts of invasive species 38 is comparatively scarce. We aim to (1) develop a protocol for systematically assessing the 39 cumulative impacts of invasive species and (2) conduct a test of this protocol using a suite of 40 non-native plants that are invasive in foreshore and riparian ecosystems of British Columbia. 41 Canada. The protocol itself aims to standardize future evaluations of the cumulative impacts 42 of invasive species. In addition, our systematic map will identify the strengths and gaps in 43 44 knowledge pertaining to invasive plant species impacts in foreshore and riparian 45 ecosystems, with the ultimate goal of facilitating the development of evidence-based 46 management strategies.

47

48 Methods

We identified the research topic and the primary and secondary questions with the support 49 of stakeholders. We then devised a flexible string that allows for searching target invasive 50 51 species. Using this string, we searched the literature for pilot species that aided the iterative 52 development of the protocol. Once all target species are identified, we will carry out a 53 systematic literature search on their impacts. We will search Web of Science and the CABI 54 compendium for invasive species. We will include studies if they (i) refer to the target 55 invasive species, (ii) focus on its environmental impacts and (iii) investigate such impacts in 56 riparian ecosystems (iv) within North America (i.e. Canada & U.S.A.). We will use a two-57 stage screening process: titles and abstracts first, then the full manuscript. From each 58 source, we will extract impact description, ecosystem component impacted, and magnitude 59 and directionality of impacts. We will include a publicly available database of studies, descriptive statistics and a narrative summary within our synthesis outcomes. 60 61

Keywords: Cumulative impacts, British Columbia, Invasive species, Impacts, Riparian
ecosystems, Plant invasions, Foreshore ecosystems, Protocol, Systematic maps

- 64
- 65

66 Background

67 Biological invasions in foreshore and riparian ecosystems

Foreshore and riparian ecosystems are vitally important from ecological, cultural, and
economic standpoints. Although their spatial extent is small, they are often hotspots of
biodiversity, hosting rare species and serving as refugia and corridors essential to many
others (1–3). Riparian ecosystems also provide essential functions and services such as
improving water quality, flood mitigation, and minimizing erosion (2,4,5). As such, foreshore

and riparian ecosystems are the focus of targeted management and conservation strategies
in many countries (6–9).

75 Despite their recognized importance, foreshore and riparian ecosystems are being 76 impacted by many anthropogenic stressors (10). Infrastructures (e.g. dams, dyking, 77 channelization) and water management (e.g. water diversion, irrigation, dredging) can 78 radically modify water levels and flow and disrupt natural fluvial dynamics (1,5,11,12). 79 Contamination and nutrient additions can alter water quality, reduce biodiversity, and 80 bioaccumulate (1,13). Habitat loss through agriculture, deforestation and development 81 disproportionately impacts foreshore and riparian zones (1,14–16), and was estimated to be 82 up to two-thirds in the U.S. alone (17). Additionally, freshwater ecosystems are oftentimes highly invaded by non-native species due to their proximity to human settlements and their 83 84 function as dispersal corridors (14,18–21).

Invasive species can impact riparian ecosystems in various ways, but invasive plants 85 have particularly pervasive impacts to ecosystem structure and functioning. By spreading 86 aggressively, they displace both plant and animal native species (22-25), modify 87 geochemical and hydraulic cycles (26,27), alter trophic processes (28) and change the 88 89 composition and structure of communities above and below ground (2,29). Additionally, invasive plants alter traditional practices and resource use by Indigenous peoples (28). The 90 91 cumulative impacts of invasive plants on riparian ecosystems are potentially profound, but research to quantify such effects remains limited (2,31). 92

Here, we aim to develop a framework for quantifying the cumulative impacts of invasive species. We test the protocol using a set of plant species invasive in British Columbia's foreshore and riparian ecosystems, that are also invasive in other regions.

96

97 Cumulative impacts: definitions, examples and previous work

In Invasion ecology, impacts are defined as measurable changes caused by non-native
species on a target ecosystem (32,33). They can vary greatly in type, magnitude, and

directionality. For instance, some impacts might be barely detectable, while others can
produce pronounced, observable effects. Impacts can be direct, but also mediated through
other factors (32). Finally, while nonnative species have been investigated in large part
because of their negative effects, impacts can vary along a continuum from negative to
positive (33,34), and can be ecosystem or context-dependent.

Identifying an impact's directionality presents some challenges. Negative impacts are typically equated to unfavourable outcomes for humans (33). However, this approach is strongly biased by the value system and worldview of the researcher (34,35). In an effort to minimize subjectivity and value-based identifications of impact directionality, we define as negative or positive any quantifiable reduction or increase in ecosystem properties or attributes (33). For instance, we define as positive an increase in the fitness or number of individuals of a native species but as negative its reduction.

112 The combination and interaction of multiple impacts are referred to as cumulative 113 impacts and many definitions of this concept exist. For the Canadian Environmental 114 Assessment Act (CEAA), they are "changes to the environment that are caused by an action in combination with other past, present and future human actions" (36). The Council on 115 116 Environmental Quality (CEQ) suggests impacts have to be incremental (37). The most well-117 articulated definition is that of the European Environmental Agency (EEA), which defines 118 them as: 'the impacts (positive or negative, direct and indirect, long-term and short-term 119 impacts) arising from a range of activities throughout an area or region, where each 120 individual effect may not be significant if taken in isolation. Such impacts can arise from the 121 growing volume of traffic, the combined effect of a number of agriculture measures leading to more intensive production and use of chemicals, etc. Cumulative impacts include a time 122 123 dimension, since they should calculate the impact on environmental resources resulting from 124 changes brought about by past, present and reasonably foreseeable future actions." (38). Consistent elements among these definitions are (1) the combination of multiple individual 125 impacts, (2) a time component and (3) the human agency. While not explicitly stated in the 126

previous definitions, cumulative impacts also have a spatial dimension, or they canaccumulate in space as well as temporally (39).

We define cumulative impacts in biological invasions as the combined effect of multiple impacts when at least one is generated by an invasive species. Cumulative impacts include recurrent impacts of a single species and the combined effect of multiple invaders, but also the compounded impact of invading species and other anthropogenic stressors (12). Our definition incorporates all the elements of previous definitions; however, it is more restrictive, as the primary focus is the impacts of invasive species. Conversely, it includes impacts of any magnitude, type or directionality.

The term 'cumulative' might imply that the total effect of multiple impacts is always 136 greater than that of individual impacts. Multiple invaders can collectively increase native 137 species displacement, or enhance topsoil nutrient concentration (additive impacts, 29,30). 138 139 An N-fixer might increase soil nitrogen, facilitating invasions by more competitive nitrophilous species, which in turn will displace natives (multiplicative impacts, 29). However, additive or 140 multiplicative impacts are not the only potential outcomes. Competition between two 141 invaders might instead reduce their impact per capita. For example, an allopathic species 142 143 might negatively affect both native and non-native species. In this case, one invader 144 mitigates the impacts of another invader (39).

145 Despite a long history of research on cumulative impacts within environmental 146 contexts, (39), the literature on the cumulative impacts of invasive species is relatively 147 scarce. Most work in biological invasions focuses on a single species or single direct impact 148 (41–46). Even when multiple impacts are identified, their cumulative effect is rarely considered (31,40). This is despite previously proposed theoretical frameworks share some 149 150 conceptual overlap. One such example is the invasion meltdown, which posits that 151 interactions among invaders might increase their impacts (47). Critically for our work, little research effort explored the cumulative impacts of invasive plant species in riparian and 152 foreshore ecosystems. 153

154

155 Topic Identification and Stakeholder Input

156 There is a clear need for work assessing the cumulative impacts of invasive species in riparian ecosystems. The Province of British Columbia, Ministry of Forests Invasive Plant 157 158 Program, highlighted the need to synthesize current evidence on the impacts of invasive 159 plant species in riparian and foreshore ecosystems within the province, in order to inform 160 research and management needs. British Columbia's riparian and foreshore ecosystems are 161 invaded by numerous highly destructive invasive plant species, such as Russian Olive (Elaeagnus angustifolia), Phragmites (Phragmites australis), Knotweeds (Reynoutria spp., 162 syn. Fallopia), Tree of Heaven (Ailanthus altissima) and Canary reed grass (Phalaris 163 164 arundinacea). While the impacts of these species have been extensively investigated (43,48–52), there is no comprehensive assessment of their cumulative impacts. 165

166 Stakeholders in the provincial government played a pivotal role in shaping the research topic and refining the scope of the systematic map. Based on their expert 167 168 knowledge and the available data, they provided a list of 10-15 plant species that are invasive in the target ecosystems and geographic areas, thereby aiding in the identification 169 of specific research questions and objectives. Input from practitioners and other researchers 170 helped refine the approach and the methodology. Through ongoing dialogue and feedback, 171 172 stakeholders were able to establish clear expectations, develop a robust methodology, and identify appropriate outcomes for the systematic map. In addition to quantifying the 173 174 cumulative impacts of plant species invasive to riparian ecosystems, stakeholders have identified two additional aspects as essential. First is the development of a reproducible 175 176 protocol that can be employed in future systematic studies of invasive species impacts. 177 Second is the investigation of how the cumulative impacts of invasive species will vary under current climate change scenarios. 178

Protocols are a crucial aspect of developing a project, particularly in the case of systematic work (53). Good protocols need to be transparent, detailed and reproducible, allowing other researchers to replicate their work (53–56). In this case, we do not simply want to describe our procedure for mapping the existing literature, but we specifically aim to
provide a tool that is sufficiently flexible and reproducible to be applied in the investigation of
other invasive species or ecosystems.

185 Climate change is a key contributor to the cumulative impacts of invasive species 186 across both terrestrial and aquatic ecosystems. However, the nature and magnitude of these 187 are often unclear. Interactions between particular invasive plants and the diverse facets of climate change are challenging to predict and likely species- and context-dependent (57). 188 189 For instance, while the ranges of many non-native invasive species may expand as 190 temperature rises (58), others may contract or shift in response to both abiotic and biotic factors (57,59). Nevertheless, strategies for mitigating negative impacts are sorely needed. A 191 key first step is synthesizing the diverse and extensive research on this topic. 192

Here, we propose to first devise and publish a reproducible systematic map protocol 193 (53) for screening, collating, and describing research on the impacts of priority invasive 194 plants in riparian and foreshore ecosystems in British Columbia. We will develop and refine 195 196 our systematic map protocol using an iterative approach to pilot invasive species. Next, we aim to publish the findings of our systematic map. Given their efficacy and 197 198 comprehensiveness, systematic maps are increasingly common in environmental 199 management (54). Through the systematic map process, we will identify knowledge clusters 200 and gaps (i.e. areas of high and low concentration of the research effort), and synthesize 201 results within the context of current climate change scenarios. Key outputs will include (1) a 202 robust analytical framework for qualitatively predicting – based on the best available 203 evidence - the cumulative impacts of invasive plants under changing climates and followed 204 by (2) a more detailed assessment for a selection of priority invasive plant species (identified 205 by the BC Ministry of Forests Invasive Alien Plant Program). These outputs will have high 206 utility for policy, planning and strategic, evidence-based decision management of 207 ecosystems impacted by priority invasive plant species in British Columbia.

208 **Objective of the review**

209	We aim to systematically identify and map studies assessing the impacts of a selection of		
210	plant species invasive to riparian ecosystems in British Columbia. We will combine the		
211	results to quantify the individual and collective cumulative impacts of these species.		
212	Primary question		
213	What are the cumulative impacts of invasive plants in the riparian and foreshore ecosystems		
214	of British Columbia?		
215	Components of the primary question		
216	Population: Riparian ecosystems in British Columbia		
217	• Exposure: Impacts of a set of non-native plant species invasive to riparian		
218	ecosystems of British Columbia		
219	• Comparator : No impact or absence of invasive plant species.		
220	• Outcome : A synthesis of both the individual and collective cumulative impacts of the		
221	selected invasive plant species		
222	Secondary question		
223	We will describe variations in the research effort with regard to:		
224 225 226	Geography and fluvial systems investigated		
227	Invasive species		
228	 Impacts and their directionality (negative, positive, or neutral) 		
229	Impacted ecosystem components		
230	• Type of study (e.g. correlational, experimental, etc.)		
231	Time (did the level of knowledge change over time?)		
232 233	Additionally, we will delineate potential changes in impact magnitude by species under		
234	current climate change scenarios based on the available literature.		

235 Methods

236 Search string

We will conduct multiple systematic searches, one for each of our focus species. For each search, we will use as keywords the scientific name of a species and "impact", formatted for Web of Science (WOS). For example:

240

241 Elaeagnus angustifolia AND impact*

242

The selected search string is purposely broad. Searches including keywords associated with the target ecosystem (riparian, foreshore, freshwater, wetland, aquatic, etc) and geographic area (British Columbia, Canada, North America, etc.) were deemed to be too restrictive. A broader search allows for capturing also studies that either use different keywords or investigate impacts in different circumstances and yet might be relevant to the target ecosystem. Using this string, we searched the literature for pilot species that aided the iterative development of the protocol. Pilot species will be included in the systematic map.

251 Bibliographic sources

We will conduct searches in WOS, accessing the core database. The core database assigns 252 metadata to a study based exclusively on the information provided by the publisher and 253 journal. Since other databases assign additional metadata to a study, some material might 254 go undetected despite meeting our criteria. We will expand our search to all databases and 255 then refine it to the core collection. This will identify studies that match our keywords across 256 257 all databases but are only present in the core collection, and thus accessible to the authors 258 (Mathew Vis-Dunbar, pers. comm. 2023). Additionally, we will screen all references in the 259 CABI Invasive Species Compendium factsheet for each species, except for references in the 260 Distribution References section. Review studies that fit the criteria for inclusion will be used as sources as well, and references extracted and screened. We will detail exceptions in the 261

supplementary material. Accessing multiple databases will help reduce location and index
biases (i.e.not all journals are indexed in all databases, incomplete or poor indexing, 46).

265 Screening and inclusion criteria

266 The screening process will include two stages. First, we will screen titles and abstracts. If the information is insufficient to make a decision, we will assess the full manuscript as well. 267 These steps will be applied to all studies, regardless of the source they were extracted from. 268 A single reviewer will conduct the screening (FM). A random subset of studies (10%) will 269 270 also be assessed by a second reviewer (JP). We will appraise consistency using Cohen's 271 kappa statistics and set 0.6 as a threshold (60,61). If consistency is below the cut-off limit, 272 screening and inclusion criteria will be adjusted for clarity. All disagreements will be discussed and resolved. Any study authored by one of the systematic reviewers that meet 273 274 the criteria for inclusion will be assessed by the other reviewer at every stage of the 275 process.

We will screen published and unpublished material, but not personal communications 276 or expert opinions. Including unpublished work reduces the risk of publication and citation 277 biases (i.e. significant results are more likely to be published and cited than non-significant 278 279 results, 46,48). We will consider only material in English. To minimize language bias (i.e. significant results are more likely to be published in English, 46,48), we will assess the title 280 and abstract if translated into English. Studies were included irrespective of the magnitude, 281 282 type or directionality of the impact (negative, positive or neutral), and irrespective of the statistical significance of reported results. This will help reduce the prevailing paradigm bias 283 (i.e. a bias towards studies supporting the prevailing paradigm; in this case, invasive 284 species' impacts are extensive and negative, 26,46,48). Currently, the time span includes all 285 286 studies up to the day the search was conducted (09 January 2023), countering temporal bias (i.e. older studies might be overlooked, 46,54). 287

288	We will include studies if they (i) refer to the invasive species searched, (ii) focus on	
289	its environmental impacts and (iii) investigate such impacts in riparian ecosystems (iv) within	
290	North America (i.e. Canada & U.S.A.). We will justify all exceptions (63).	
291		
292	Meta-data extraction	
293	Studies included in the systematic literature map will undergo a full-manuscript screening to	
294	identify the investigated impact (or impacts). We will provide a description of the investigated	
295	impacts and the ecosystem component impacted. Then, we will categorize impacts by their	
296	magnitude and directionality. Impacts magnitude will be assessed following previous work,	
297	modified to include both positive and negative impacts (31-33):	
298 299 300	• Minimal : The impact is unlikely or negligible.	
301	• Minor : It causes changes in the fitness of individuals in the native biota, but no	
302	changes in native population densities.	
303	• Moderate: It causes changes in the population densities of native species, but no	
304	changes to the structure of communities or to the abiotic or biotic components of	
305	ecosystems.	
306	• Major: It causes the local or population extinction/introduction of at least one native	
307	species, and leads to reversible/transient changes in the structure of communities	
308	and the abiotic or biotic components of ecosystems.	
309	• Massive: It leads to the replacement and local extinction/introduction of multiple	
310	native species, and produces irreversible changes in the structure of communities	
311	and the abiotic or biotic components of ecosystems.	
312		
313	Data coding	

314 For each study at the full-text screening stage, we will provide the following information:

315 316 317	1. Bibliographic information
318	1. Authors list
319	2. Article title
320	3. Publication year
321	4. Bibliographic source
322	2. Inclusion criteria
323	1. Exposure: Focuses on target species (Y/N)
324	2. Exposure: Focuses on environmental impacts (Y/N)
325	3. Population: Focuses on riparian ecosystems (Y/N)
326	4. Population: Within North America (Y/N)
327	3. Screening stage
328	1. Excluded at full-text stage
329	2. Included
330	3. Exceptions
331	4. Additional information
332	1. Duplicate (Y/N)
333	2. Notes
334 335	For included studies only, we will provide also the following information:
336 337 338	1. Bibliographic information
339	1. Authors list
340	2. Article title
341	3. Publication year
342	2. Information on impacts
343	c. Impact description
344	c. Ecosystem component impacted (e.g. species, soil, etc.)

345	c. Magnitude of impact	
346	c. Impact direction (negative, positive, neutral)	
347	3. Additional information	
348	1. Geographic region	
349	2. Notes	
350 351	We will compile subsection 3c. <i>Exceptions</i> on a case-by-case basis. For included studies,	
352	we will provide information by impact so that if a study investigated more than one, there will	
353	be a number of entries equivalent to the number of impacts assessed.	
354		
355	Synthesis and presentation	
356	For each species, we will provide a first database with all studies included at the full-text	
357	screening stage and a second database with the studies included in the review, along with a	
358	graphical representation of the screening process. Both databases will contain	
359	corresponding coded metadata (see Data Coding section). We will import studies included in	
360	the review into a reference manager and share them as a public library to facilitate	
361	accessibility. We will develop a graphical representation of riparian ecosystems,	
362	representing identified impacts and their magnitude and directionality for each individual	
363	species. Then, we will create a matrix combining multiple species (as rows) and impacts (as	
364	columns) to assess the collective cumulative impacts. Descriptive statistics will be used to	
365	answer secondary questions, and co-occurrence matrices to identify research effort biases	
366	(64). Lastly, we will provide a narrative synthesis of results for both main and secondary	
367	questions. The narrative synthesis will focus on (i) species and impact prioritization, (ii)	
368	clusters and gaps in present knowledge, (iii) predicted variations in impact magnitude and	
369	direction under current climate change scenarios and (iv) avenues for future research.	

370

371 Ethics approval and consent to participate

- 372 Not applicable.
- 373 Consent for publication
- Not applicable.
- 375 Availability of data and materials
- 376 Data sharing is not applicable to this article as no datasets were generated or analyzed
- 377 during the current study.
- 378 Competing interests
- 379 The authors declare that they have no competing interests.
- 380 Funding
- 381 FM is funded by the Ministry of Forests, British Columbia, Canada, and by the Irving K.
- 382 Faculty of Science at the Okanagan campus of UBC.
- 383 Authors' contributions
- FM drafted the protocol with input from JP and CM. All authors read and approved the finalmanuscript.
- 386 Acknowledgements
- 387 We thank Mathew Vis-Dunbar for support and guidance in developing search strings and
- extracting information from bibliographic databases. We thank all the stakeholders for the
- 389 feedback provided throughout the process.
- 390 References

Poff B, Koestner KA, Neary DG, Henderson V. Threats to Riparian Ecosystems in
 Western North America: An Analysis of Existing Literature. JAWRA Journal of the American
 Water Resources Association. 2011;47(6):1241–54.

Stella JC, Rodríguez-González PM, Dufour S, Bendix J. Riparian vegetation research
 in Mediterranean-climate regions: common patterns, ecological processes, and

considerations for management. Hydrobiologia. 2013 Nov;719(1):291–315.

Ragan K, Schipper J, Bateman HL, Hall SJ. Mammal use of riparian corridors in
 semi-arid Sonora, Mexico. The Journal of Wildlife Management. 2023;87(1):e22322.

4. Koundouri P, Boulton AJ, Datry T, Souliotis I. Ecosystem Services, Values, and

Societal Perceptions of Intermittent Rivers and Ephemeral Streams. In: Intermittent Rivers
and Ephemeral Streams. Elsevier; 2017. p. 455–76.

402 5. Patten DT. Riparian ecosytems of semi-arid North America: Diversity and human
403 impacts. Wetlands. 1998 Dec 1;18(4):498–512.

Knopf FL, Johnson RR, Rich T, Samson FB, Szaro RC. Conservation of Riparian
Ecosystems in the United States. The Wilson Bulletin. 1988;100(2):272–84.

406 7. Erős T, Kuehne L, Dolezsai A, Sommerwerk N, Wolter C. A systematic review of
407 assessment and conservation management in large floodplain rivers – Actions postponed.
408 Ecological Indicators. 2019;98:453–61.

8. Fryirs KA, Zhang N, Duxbury E, Ralph T, Fryirs KA, Zhang N, et al. Rivers up in
smoke: impacts of Australia's 2019–2020 megafires on riparian systems. Int J Wildland Fire.
2022 Jun 16;31(7):720–7.

412 9. Brierley GJ, Hikuroa D, Fuller IC, Tunnicliffe J, Allen K, Brasington J, et al.

413 Reanimating the strangled rivers of Aotearoa New Zealand. WIREs Water.

414 2023;10(2):e1624.

415 10. Stella JC, Bendix J. Chapter 5 - Multiple Stressors in Riparian Ecosystems. In:

416 Sabater S, Elosegi A, Ludwig R, editors. Multiple Stressors in River Ecosystems. Elsevier;

417 2019. p. 81–110.

Airoldi L, Beck M. Loss, Status and Trends for Coastal Marine Habitats of Europe. In:
Gibson R, Atkinson R, Gordon J, editors. Oceanography and Marine Biology. CRC Press;
2007. p. 345–405. (Oceanography and Marine Biology - An Annual Review; vol. 20074975).
Johnson PT, Olden JD, Vander Zanden MJ. Dam invaders: impoundments facilitate
biological invasions into freshwaters. Frontiers in Ecology and the Environment.

423 2008;6(7):357–63.

13. Chiu MC, Leigh C, Mazor R, Cid N, Resh V. Anthropogenic Threats to Intermittent
Rivers and Ephemeral Streams. In: Intermittent Rivers and Ephemeral Streams. Elsevier;
2017. p. 433–54.

427 14. Greene SL. A Roadmap for Riparian Invasion Research. River Research and
428 Applications. 2014;30(5):663–9.

429 15. Post GC, Chang H, Banis D. The spatial relationship between patterns of
430 disappeared streams and residential development in Portland, Oregon, USA. Journal of

431 Maps. 2022 Dec 1;18(2):210–8.

432 16. Sweeney BW, Bott TL, Jackson JK, Kaplan LA, Newbold JD, Standley LJ, et al.

433 Riparian deforestation, stream narrowing, and loss of stream ecosystem services.

434 Proceedings of the National Academy of Sciences. 2004;101(39):14132–7.

435 17. Swift BL. Status of Riparian Ecosystems in the United States. JAWRA Journal of the
436 American Water Resources Association. 1984;20(2):223–8.

437 18. Aguiar FCF, Ferreira MT. Plant invasions in the rivers of the Iberian Peninsula, south438 western Europe: A review. Plant Biosystems. 2013;147(4):1107–19.

439 19. Arianoutsou M, Delipetrou P, Celesti-Grapow L, Basnou C, Bazos I, Kokkoris Y, et al.

440 Comparing naturalized alien plants and recipient habitats across an east-west gradient in

the Mediterranean Basin. Journal of Biogeography. 2010;37(9):1811–23.

442 20. Kowarik I, Säumel I. Water dispersal as an additional pathway to invasions by the

443 primarily wind-dispersed tree Ailanthus altissima. Plant Ecol. 2008;198(2):241–52.

444 21. Pysek P, Prach K. Plant Invasions and the Role of Riparian Habitats: A Comparison

of Four Species Alien to Central Europe. Journal of Biogeography. 1993;20(4):413–20.

Barnes WJ. The Rapid Growth of a Population of Reed Canarygrass (Phalaris
arundinacea L.) and Its Impact on Some Riverbottom Herbs. Journal of the Torrey Botanical
Society. 1999;126(2):133–8.

Price EPF, Spyreas G, Matthews JW. Biotic homogenization of regional wetland
plant communities within short time-scales in the presence of an aggressive invader. Journal
of Ecology. 2018;106(3):1180–90.

452 24. Bateman HL, Chung-MacCoubrey A, Snell HL. Impact of Non-Native Plant Removal
453 on Lizards in Riparian Habitats in the Southwestern United States. Restoration Ecology.
454 2008;16(1):180–90.

Pendleton R, Pendleton B, Finch D. Displacement of Native Riparian Shrubs by
Woody Exotics: Effects on Arthropod and Pollinator Community Composition. Natural
Resources and Environmental Issues. 2011;16(1).

458 26. Martinez AE, McDowell PF. Invasive Reed Canarygrass (Phalaris arundinacea) and
459 Native Vegetation Channel Roughness. Invasive Plant Science and Management.
460 2016;9(1):12–21.

27. Simons SB, Seastedt TR. Decomposition and Nitrogen Release from Foliage of
Cottonwood (Populus deltoides) and Russian-olive (Elaeagnus angustifolia) in a Riparian
Ecosystem. The Southwestern Naturalist. 1999;44(3):256–60.

464 28. Mineau MM, Baxter CV, Marcarelli AM. A Non-Native Riparian Tree (Elaeagnus
465 angustifolia) Changes Nutrient Dynamics in Streams. Ecosystems. 2011;14(3):353–65.
466 29. Jacinthe PA, Bills JS, Tedesco LP. Size, activity and catabolic diversity of the soil
467 microbial biomass in a wetland complex invaded by reed canary grass. Plant Soil.

468 2010;329(1):227–38.

469 30. Collier N, Austin BJ, Bradshaw CJA, McMahon CR. Turning Pests into Profits:

470 Introduced Buffalo Provide Multiple Benefits to Indigenous People of Northern Australia.

471 Hum Ecol. 2011;39(2):155–64.

- 472 31. Katsanevakis S, Tempera F, Teixeira H. Mapping the impact of alien species on
- 473 marine ecosystems: the Mediterranean Sea case study. Diversity and Distributions.

474 2016;22(6):694–707.

- 475 32. Blackburn TM, Essl F, Evans T, Hulme PE, Jeschke JM, Kühn I, et al. A Unified
- 476 Classification of Alien Species Based on the Magnitude of their Environmental Impacts.
- 477 PLOS Biology. 2014;12(5):e1001850.
- 478 33. Vimercati G, Probert AF, Volery L, Bernardo-Madrid R, Bertolino S, Céspedes V, et
- al. The EICAT+ framework enables classification of positive impacts of alien taxa on native
- 480 biodiversity. PLOS Biology. 2022;20(8):e3001729.
- 34. Sax DF, Schlaepfer MA, Olden JD. Identifying key points of disagreement in nonnative impacts and valuations. Trends in Ecology & Evolution. 2023;
- 483 35. Cardou F, Vellend M. Stealth advocacy in ecology and conservation biology.
- 484 Biological Conservation. 2023;280:109968.
- 485 36. Canadian Environmental Assessment Agency, editor. Cumulative effects assessment
 486 practitioners guide. Ottawa; 1999.
- 487 37. Council on Environmental Quality. National Environmental Policy Act Cumulative488 Effects.
- 489 38. European Environment Agency. https://www.eea.europa.eu/help/glossary/eea490 glossary/cumulative-impacts. Cumulative Impacts.
- 491 39. Spaling H, Smit B. Cumulative environmental change: Conceptual frameworks,
- 492 evaluation approaches, and institutional perspectives. Environmental Management.
- 493 1993;17(5):587–600.
- 494 40. Vanderhoeven S, Dassonville N, Meerts P. Increased Topsoil Mineral Nutrient
- 495 Concentrations Under exotic invasive plants in Belgium. Plant Soil. 2005;275(1):169–79.
- 496 41. Follstad Shah JJ, Harner MJ, Tibbets TM. Elaeagnus angustifolia Elevates Soil
- 497 Inorganic Nitrogen Pools in Riparian Ecosystems. Ecosystems. 2010;13(1):46–61.

498 42. Spyreas G, Wilm BW, Plocher AE, Ketzner DM, Matthews JW, Ellis JL, et al.

Biological consequences of invasion by reed canary grass (Phalaris arundinacea). Biol
Invasions. 2010;12(5):1253–67.

43. Rojas IM, Zedler JB. An invasive exotic grass reduced sedge meadow species
richness by half. Wetlands Ecol Manage. 2015;23(4):649–63.

503 44. Katz GL, Tuttle GM, Denslow MW, Norton AP. Legacy Effects of Russian Olive

(Elaeagnus angustifolia L.) in a Riparian Ecosystem Three Years Post-Removal. Wetlands.
2020;40(6):1897–907.

506 45. Gallego-Tévar B, Grewell BJ, Whitcraft CR, Futrell JC, Bárcenas-Moreno G, Castillo

507 JM. Contrasted Impacts of Yellow Flag Iris (Iris pseudacorus) on Plant Diversity in Tidal

508 Wetlands within Its Native and Invaded Distribution Ranges. Diversity. 2022;14(5):326.

509 46. Rodgers VL, Scanga SE, Kolozsvary MB, Garneau DE, Kilgore JS, Anderson LJ, et

al. Where Is Garlic Mustard? Understanding the Ecological Context for Invasions of Alliaria

511 petiolata. BioScience. 2022;72(6):521–37.

512 47. Simberloff D, Von Holle B. Positive Interactions of Nonindigenous Species:

513 Invasional Meltdown? Biological Invasions. 1999 Mar 1;1(1):21–32.

48. Benoit LK, Askins RA. Impact of the spread of Phragmites on the distribution of birds

in Connecticut tidal marshes. Wetlands. 1999;19(1):194–208.

516 49. Brooks RK, Barney JN, Salom SM. The invasive tree, Ailanthus altissima, impacts

517 understory nativity, not seedbank nativity. Forest Ecology and Management.

518 2021;489:119025.

519 50. Claeson SM, LeRoy CJ, Barry JR, Kuehn KA. Impacts of invasive riparian knotweed

on litter decomposition, aquatic fungi, and macroinvertebrates. Biol Invasions.

521 2014;16(7):1531–44.

522 51. Katz GL, Shafroth PB. Biology, ecology and management of Elaeagnus angustifolia

523 L. (Russian olive) in western North America. Wetlands. 2003;23(4):763–77.

524 52. Medina-Villar S, Rodríguez-Echeverría S, Lorenzo P, Alonso A, Pérez-Corona E,

525 Castro-Díez P. Impacts of the alien trees Ailanthus altissima (Mill.) Swingle and Robinia

pseudoacacia L. on soil nutrients and microbial communities. Soil Biology and Biochemistry.
2016;96:65–73.

528 53. Collaboration for Environmental Evidence. Guidelines and Standards for Evidence
529 synthesis in Environmental Management. Version 5.1 (AS Pullin, GK Frampton, B Livoreil &
530 G Petrokofsky, Eds) [Internet]. 2022. Available from:

531 https://environmentalevidence.org/information-for-authors/

532 54. Haddaway NR, Bernes C, Jonsson BG, Hedlund K. The benefits of systematic

533 mapping to evidence-based environmental management. Ambio. 2016;45(5):613–20.

534 55. Haddaway NR, Macura B, Whaley P, Pullin AS. ROSES RepOrting standards for

535 Systematic Evidence Syntheses: pro forma, flow-diagram and descriptive summary of the

plan and conduct of environmental systematic reviews and systematic maps. Environ Evid.

537 2018;7(1):7.

538 56. Palacio FX, Callaghan CT, Cardoso P, Hudgins EJ, Jarzyna MA, Ottaviani G, et al. A

protocol for reproducible functional diversity analyses. Ecography. 2022;2022(11):e06287.

540 57. Bellard C, Thuiller W, Leroy B, Genovesi P, Bakkenes M, Courchamp F. Will climate 541 change promote future invasions? Glob Chang Biol. 2013;19(12):3740–8.

542 58. Perry LG, Reynolds LV, Shafroth PB. Divergent effects of land-use, propagule

543 pressure, and climate on woody riparian invasion. Biol Invasions. 2018;20(11):3271–95.

544 59. Poland TM, Patel-Weynand T, Finch DM, Miniat CF, Hayes DC, Lopez VM, editors.

545 Invasive Species in Forests and Rangelands of the United States: A Comprehensive

546 Science Synthesis for the United States Forest Sector. Cham: Springer International

547 Publishing; 2021.

60. Ridley FA, McGowan PJ, Mair L. The scope and extent of literature that maps threats
to species: a systematic map protocol. Environmental Evidence. 2020;9(1):23.

550 61. Cohen J. A coefficient of agreement for nomimal scales. Educ Psychol Meas.

551 1960;20:2746.

552 62. Bayliss HR, Beyer FR. Information retrieval for ecological syntheses. Research
553 Synthesis Methods. 2015;6(2):136–48.

- 63. O'Dea RE, Lagisz M, Jennions MD, Koricheva J, Noble DWA, Parker TH, et al.
- 555 Preferred reporting items for systematic reviews and meta-analyses in ecology and
- evolutionary biology: a PRISMA extension. Biol Rev. 2021;96(5):1695–722.
- 557 64. Ridley FA, Hickinbotham EJ, Suggitt AJ, McGowan PJK, Mair L. The scope and
- 558 extent of literature that maps threats to species globally: a systematic map. Environ Evid.
- 559 2022;11(1):1–26.

560