

1 Collating existing evidence on cumulative impacts of invasive
2 plant species in riparian ecosystems of British Columbia,
3 Canada: a systematic map protocol

4
5

6 *Fabio Mologni¹

7 Chandra E. Moffat^{2,1}

8 Jason Pither¹

9

10 1. Department of Biology, and Institute for Biodiversity, Resilience, and Ecosystem Services

11 I.K. Barber Faculty of Science,

12 University of British Columbia Okanagan

13 1177 Research Road, Kelowna, BC V1V 1V7, Canada

14

15 2. Agriculture and Agri-Food Canada

16 Summerland Research and Development Centre

17 4200 Highway #97, Summerland, BC V0H 1Z0

18

19 * Corresponding author

20 *Postal address:* Department of Biology, University of British Columbia Okanagan, 1177

21 Research Road, Kelowna, BC V1V 1V7, Canada

22 *Phone: +12369709033*

23 *Email: fabio.mologni@ubc.ca*

24 *ORCID: <https://orcid.org/0000-0003-4750-9974>*

25
26 Other authors email:

27 Chandra E. Moffat (chandra.moffat@agr.gc.ca)

28 Jason Pither (jason.pither@ubc.ca)

29

30 **Abstract**

31 **Background**

32 Globally, the structure and functioning of foreshore and riparian ecosystems are being
33 dramatically impacted by non-native invasive plant species. Invasive species can
34 outcompete and replace native species, modify geochemical and hydraulic cycles, alter
35 trophic processes, and change the composition and structure of communities above and
36 below ground. However, these impacts are often investigated in isolation, even though one
37 invasive species might increase or mitigate the impacts of others (i.e. cumulative impacts),
38 potentially with cascading effects. Although cumulative impacts have long been studied
39 within other environmental contexts, research on the cumulative impacts of invasive species
40 is comparatively scarce. We aim to develop a protocol to systematically identify and collate
41 evidence on the individual and cumulative impacts of a set of plant species invasive in
42 foreshore and riparian ecosystems of British Columbia, Canada. Our primary question is:
43 What evidence is available on the individual and cumulative impacts of invasive plants in the
44 riparian and foreshore ecosystems of British Columbia, Canada? In addition, our systematic
45 map will identify the strengths and gaps in knowledge pertaining to invasive plant species

46 impacts in foreshore and riparian ecosystems, with the ultimate goal of facilitating the
47 development of evidence-based management strategies.

48
49 **Methods**

50 We identified the research topic and the primary and secondary questions with the support
51 of stakeholders. We then devised a flexible string that allows for searching target invasive
52 species. Using this string, we searched the literature for pilot species that aided the iterative
53 development of the protocol. Once all target species are identified, we will carry out a
54 systematic literature search on their impacts. We will search Web of Science and the CABI
55 compendium for invasive species. We will include studies if they (i) refer to the target
56 invasive species, (ii) focus on its environmental impacts and (iii) investigate such impacts in
57 riparian ecosystems (iv) within North America (i.e. Canada & U.S.A.). We will use a two-
58 stage screening process: titles and abstracts first, then the full manuscript. From each
59 source, we will extract impact description, ecosystem component impacted, and magnitude
60 and directionality of impacts. We will include a publicly available database of studies,
61 descriptive statistics, and a narrative summary within our synthesis outcomes.

62
63 **Keywords:** *Cumulative impacts, British Columbia, Invasive species, Impacts, Riparian*
64 *ecosystems, Plant invasions, Foreshore ecosystems, Protocol, Systematic maps*

65
66
67 **Background**

68 *Biological invasions in foreshore and riparian ecosystems*

69 Foreshore and riparian ecosystems are vitally important from ecological, cultural, and
70 economic standpoints. Although their spatial extent is small, they are often hotspots of
71 biodiversity, hosting rare species, and serving as refugia and corridors essential to many
72 others (1–3). These ecosystems also provide essential functions and services such as

73 improving water quality, flood mitigation, and minimizing erosion (2,4,5). As such, foreshore
74 and riparian habitats are the focus of targeted management and conservation strategies in
75 many countries (6–9).

76 Despite their recognized importance, foreshore and riparian ecosystems are being
77 impacted by many anthropogenic stressors (10). Infrastructures (e.g. dams, dyking,
78 channelization) and water management (e.g. water diversion, irrigation, dredging) can
79 radically modify water levels and flow and disrupt natural fluvial dynamics (1,5,11,12).
80 Contamination and nutrient additions can alter water quality, reduce biodiversity, and
81 promote bioaccumulation (1,13). Habitat loss through agriculture, deforestation, and
82 development disproportionately impacts foreshore and riparian zones (1,14–16), and was
83 estimated to be up to two-thirds in the U.S. alone (17). Additionally, freshwater ecosystems
84 are oftentimes highly invaded by non-native species due to their proximity to human
85 settlements and their function as dispersal corridors (14,18–21).

86 Invasive species can impact riparian ecosystems in various ways, but invasive plants
87 have particularly pervasive impacts on ecosystem structure and functioning. By spreading
88 aggressively, they displace both plant and animal native species (22–25), modify
89 geochemical and hydraulic cycles (26,27), alter trophic processes (28), and change the
90 composition and structure of communities above and below ground (2,29). Additionally,
91 invasive plants alter traditional practices and resource use by Indigenous peoples (28). The
92 cumulative impacts of invasive plants on riparian ecosystems are potentially profound, but
93 research to quantify such effects remains limited (2,31).

94 Here, we aim to develop a framework for systematically collating and mapping
95 evidence on the individual and cumulative impacts of plant species that are invasive within
96 foreshore and riparian ecosystems, and we will apply our protocol to systems in British
97 Columbia, Canada.

98 *Individual and cumulative impacts: definitions, examples and previous work*

99 In invasion ecology, individual impacts are defined as measurable changes caused by non-
100 native species on a target ecosystem (32,33). They can vary greatly in type, magnitude, and
101 directionality. For instance, some impacts might be barely detectable (e.g. gene flow through
102 hybridization), while others can produce pronounced, observable effects (e.g. ecosystem
103 dominance). Impacts can be direct (e.g. displacement of native species), but also mediated
104 through other factors (e.g. competition for resources, 32). Finally, while non-native species
105 have been investigated in large part because of their negative effects, impacts can vary
106 along a continuum from negative to positive (33,34), and can be ecosystem or context-
107 dependent.

108 Identifying an impact's directionality presents some challenges. Negative impacts are
109 typically equated to unfavourable outcomes for humans (33). However, this approach is
110 strongly biased by the value system and worldview of the researcher (34,35). In an effort to
111 minimize subjectivity and value-based identifications of impact directionality, we define as
112 negative or positive any quantifiable reduction or increase in ecosystem properties or
113 attributes (e.g. native species richness and abundance, nutrient cycling, water quality, etc.,
114 33). For instance, we define as positive an increase in the fitness or number of individuals of
115 a native species but as negative its reduction.

116 The combination and interaction of multiple individual impacts are referred to as
117 cumulative impacts and many definitions of this concept exist. For the Canadian
118 Environmental Assessment Act (CEAA), they are "*changes to the environment that are
119 caused by an action in combination with other past, present and future human actions*" (36).
120 The Council on Environmental Quality (CEQ) suggests impacts have to be incremental (37).
121 The most well-articulated definition is that of the European Environmental Agency (EEA),
122 which defines them as: '*the impacts (positive or negative, direct and indirect, long-term and
123 short-term impacts) arising from a range of activities throughout an area or region, where
124 each individual effect may not be significant if taken in isolation. Such impacts can arise from
125 the growing volume of traffic, the combined effect of a number of agriculture measures
126 leading to more intensive production and use of chemicals, etc. Cumulative impacts include*

127 *a time dimension, since they should calculate the impact on environmental resources*
128 *resulting from changes brought about by past, present and reasonably foreseeable future*
129 *actions.” (38). Consistent elements among these definitions are (1) the combination of*
130 *multiple individual impacts, (2) a time component and (3) the human agency. While not*
131 *explicitly stated in the previous definitions, cumulative impacts also have a spatial*
132 *dimension, or they can accumulate in space as well as temporally (39).*

133 We define cumulative impacts in biological invasions as the combined effect of
134 multiple impacts when at least one is generated by an invasive species. Cumulative impacts
135 include recurrent impacts of a single species and the combined effect of multiple invaders,
136 but also the compounded impact of invading species and other anthropogenic stressors (12).
137 Our definition incorporates all the elements of previous definitions; however, it is more
138 restrictive, as the primary focus is the impacts of invasive species. Conversely, it includes
139 impacts of any magnitude, type or directionality.

140 The term ‘cumulative’ might imply that the total effect of multiple impacts is always
141 greater than that of individual impacts. Multiple invaders can collectively increase native
142 species displacement, or enhance topsoil nutrient concentration (additive impacts, 29,30).
143 An N-fixer might increase soil nitrogen, facilitating invasions by more competitive nitrophilous
144 species, which in turn will displace natives (multiplicative impacts, 29). However, additive or
145 multiplicative impacts are not the only potential outcomes. Competition between two
146 invaders might instead reduce their impact per capita. For example, an allopathic species
147 might negatively affect both native and non-native species. In this case, one invader
148 mitigates the impacts of another invader (39).

149 Despite a long history of research on cumulative impacts within environmental
150 contexts (39), the literature on the cumulative impacts of invasive species is relatively
151 scarce. Most work in biological invasions focuses on a single species or single direct impact
152 (41–46). Even when multiple impacts are identified, their cumulative effect is rarely
153 considered (31,40). This is despite previously proposed theoretical frameworks share some
154 conceptual overlap. One such example is the invasion meltdown, which posits that

155 interactions among invaders might increase their impacts (47). Critically for our work, little
156 research effort explored the cumulative impacts of invasive plant species in riparian and
157 foreshore ecosystems. Therefore, anticipating a lack of studies on cumulative impacts, we
158 will also include individual impacts in this systematic map.

159

160 *Topic Identification and Stakeholder Input*

161 There is a clear need for work identifying the cumulative impacts of invasive species in
162 riparian ecosystems. The Province of British Columbia, Ministry of Forests Invasive Plant
163 Program, highlighted the need to synthesize current evidence on the impacts of invasive
164 plant species in riparian and foreshore ecosystems within the province, to inform research
165 and management needs. British Columbia's riparian and foreshore ecosystems are invaded
166 by numerous highly destructive invasive plant species, such as Russian Olive (*Elaeagnus*
167 *angustifolia*), Phragmites (*Phragmites australis*), Knotweeds (*Reynoutria* spp., syn. *Fallopia*),
168 Tree of Heaven (*Ailanthus altissima*) and Canary reed grass (*Phalaris arundinacea*). While
169 the impacts of these species have been extensively investigated (43,48–52), there is no
170 comprehensive assessment of their cumulative impacts.

171 Stakeholders in the provincial government played a pivotal role in shaping the
172 research topic and refining the scope of the systematic map. Stakeholders include the British
173 Columbia Ministry of Forests, Agriculture and Agri-Food Canada, and the University of
174 British Columbia. Based on their expert knowledge and the available data, they provided a
175 list of 10-15 plant species that are invasive in the target ecosystems and geographic areas,
176 thereby aiding in the identification of specific research questions and objectives. Input from
177 practitioners and other researchers helped refine the approach and the methodology.
178 Through ongoing dialogue and feedback, stakeholders were able to establish clear
179 expectations, develop a robust methodology, and identify appropriate outcomes for the
180 systematic map. In addition to quantifying the cumulative impacts of plant species invasive to
181 riparian ecosystems, stakeholders have identified two additional aspects as essential. First is

182 the development of a reproducible protocol that can be employed in future systematic
183 studies of invasive species impacts. Second is the investigation of how the cumulative
184 impacts of invasive species will vary under current climate change scenarios.

185 Protocols are a crucial aspect of developing a project, particularly in the case of
186 systematic work (53). Good protocols need to be transparent, detailed and reproducible,
187 allowing other researchers to replicate their work (53–56). In this case, we do not simply
188 want to describe our procedure for mapping the existing literature, but we specifically aim to
189 provide a tool that is sufficiently flexible and reproducible to be applied in the investigation of
190 other invasive species or ecosystems.

191 Climate change is a key contributor to the cumulative impacts of invasive species
192 across both terrestrial and aquatic ecosystems. However, the nature and magnitude of its
193 effect of invasive species' impacts is often unclear. Interactions between particular invasive
194 plants and the diverse facets of climate change are challenging to predict and likely species-
195 and context-dependent (57). For instance, while the ranges of many non-native invasive
196 species may expand as temperature rises (58), others may contract or shift in response to
197 both abiotic and biotic factors (57,59). Nevertheless, strategies for mitigating negative
198 impacts are sorely needed. A key first step is synthesizing the diverse and extensive
199 research on this topic.

200 Here, we present a reproducible systematic map protocol (53) for screening,
201 collating, and describing research on the impacts of priority invasive plants in riparian and
202 foreshore ecosystems, and we will apply it to systems in British Columbia. Given their
203 efficacy and comprehensiveness, systematic maps are increasingly common in
204 environmental management (54). Through the systematic map process, we will identify
205 knowledge clusters and gaps (i.e. areas of high and low concentration of the research effort),
206 and synthesize results within the context of current climate change scenarios. Key outputs
207 will include (1) a robust analytical framework for qualitatively predicting – based on the best
208 available evidence – the cumulative impacts of invasive plants under changing climates and
209 followed by (2) a more detailed assessment for a selection of priority invasive plant species

210 (identified by the BC Ministry of Forests Invasive Alien Plant Program). These outputs will
211 have high utility for policy, planning and strategic, evidence-based decision management of
212 ecosystems impacted by priority invasive plant species in British Columbia.

213 **Objective of the review**

214 We aim to systematically collate and map evidence on the individual and cumulative impacts
215 of a selection of plant species invasive to riparian ecosystems in British Columbia, Canada.

216 *Primary question*

217 What evidence is available on the individual and cumulative impacts of invasive plants in the
218 riparian and foreshore ecosystems of British Columbia, Canada?

219 *Components of the primary question*

- 220 • **Population:** Riparian and foreshore ecosystems in British Columbia
- 221 • **Exposure:** non-native plant species invasive to riparian and foreshore ecosystems
222 of British Columbia
- 223 • **Comparator:** No impact or absence of invasive plant species.
- 224 • **Outcome:** A synthesis of both the individual and collective cumulative impacts of the
225 selected invasive plant species

226 *Secondary question*

227 We will describe variations in the research effort with regard to:

- 228 • Geography and fluvial systems investigated
- 229 • Invasive species
- 230 • Impacts and their directionality (negative, positive, or neutral)
- 231 • Impacted ecosystem components
- 232 • Type of study (e.g. correlational, experimental, etc.)
- 233

234 • Time (did the level of knowledge change over time?)

235

236 Additionally, we will delineate potential changes in impact magnitude by species under

237 current climate change scenarios based on the available literature.

238 **Methods**

239 *Search string*

240 We will conduct multiple systematic searches, one for each of our focus species. For each

241 search, we will use as keywords the scientific name of a species and “impact”, formatted for

242 Web of Science (WOS). For example:

243

244 *Elaeagnus angustifolia* AND impact*

245

246 The selected search string is purposely broad. Searches including keywords associated with

247 the target ecosystem (riparian, foreshore, freshwater, wetland, aquatic, etc.) and geographic

248 area (British Columbia, Canada, North America, etc.) were deemed to be too restrictive. A

249 broader search allows for capturing additional studies that either use different keywords or

250 investigate impacts in different circumstances and yet might be relevant to the target

251 ecosystem.

252 We tested the comprehensiveness of searches using two pilot species, the Russian

253 Olive (*Elaeagnus angustifolia*) and the Canary Reed Grass (*Phalaris arundinacea*). For each

254 species, we selected 5 primary articles, which used a variety of keywords (e.g. impact,

255 effect, alter, change, consequence, see Appendix 1 for the full list). Then, we used the

256 search strings to extract studies from WOS and we extracted references from CABI and

257 review studies for pilot species. All studies were detected by search strings. These two

258 species aided the iterative development of the protocol and will be included in the systematic

259 map.

260

261 *Bibliographic sources*

262 We will conduct searches in WOS, accessing the core database using an institutional licence
263 (University of British Columbia). The core database assigns metadata to a study based
264 exclusively on the information provided by the publisher and journal. Since other databases
265 assign additional metadata to a study, some material might go undetected despite meeting
266 our criteria. We will expand our search to all databases and then refine it to the core
267 collection. This will identify studies that match our keywords across all databases but are
268 only present in the core collection, and thus accessible to the authors (Mathew Vis-Dunbar,
269 UBC librarian, pers. comm. 2023). Additionally, we will screen all references in the CABI
270 Invasive Species Compendium factsheet for each species, except for references in the
271 *Distribution References* section. Review studies that fit the criteria for inclusion will be used
272 as sources as well, and references extracted and screened. These sources will allow for
273 capturing also the grey literature. WOS identifies dissertations and conference proceedings,
274 especially if expanding searches to all databases, while the CABI and other review papers
275 identify technical reports. We will detail exceptions in the supplementary material. Accessing
276 multiple databases will help reduce location and index biases (i.e. not all journals are
277 indexed in all databases, incomplete or poor indexing, 46).

278

279 *Screening and inclusion criteria*

280 The screening process will include two stages. First, we will screen titles and abstracts. If the
281 information is insufficient to make a decision, we will assess the full manuscript as well.
282 These steps will be applied to all studies, regardless of the source they were extracted from.
283 A single reviewer will conduct the screening (FM). A random subset of studies (10%) will
284 also be assessed by a second reviewer (JP) at the full-manuscript stage. We will appraise
285 consistency using Cohen's kappa statistics and set 0.6 as a threshold (60,61). If consistency
286 is below the cut-off limit, screening and inclusion criteria will be adjusted for clarity. All
287 disagreements will be discussed and resolved. Any study authored by one of the systematic

288 reviewers that meets the criteria for inclusion will be assessed by the other reviewer at every
289 stage of the process.

290 We will screen both commercially published and grey literature, but not personal
291 communications or expert opinions. Including grey literature reduces the risk of publication
292 and citation biases (i.e. significant results are more likely to be published and cited than non-
293 significant results, 46,48). We will consider only material in English. To minimize language
294 bias (i.e. significant results are more likely to be published in English, 46,48), we will assess
295 the title and abstract if translated into English. Studies were included irrespective of the
296 magnitude, type or directionality of the impact (negative, positive or neutral), and irrespective
297 of the statistical significance of reported results. This will help reduce the prevailing paradigm
298 bias (i.e. a bias towards studies supporting the prevailing paradigm; in this case, invasive
299 species' impacts are extensive and negative, 26,46,48). The time span includes all studies
300 up to the day the search will be conducted, countering temporal bias (i.e. older studies might
301 be overlooked, 46,54). Finally, we will include studies regardless of study design (e.g.
302 experimental, observational, etc.).

303 We will include studies if they:

- 304 (1) Refer to the non-native invasive plant species searched. We defined as
305 invasive widespread, impactful non-native species.
- 306 (2) Focus on its abiotic and biotic impacts. We defined impacts as measurable
307 changes caused by non-native species on a target ecosystem.
- 308 (3) Investigate such impacts in riparian and foreshore ecosystems. Riparian
309 ecosystems are defined as areas adjacent to streams or rivers (flowing
310 water), while foreshore ecosystems are defined as the land adjacent to still
311 (non-flowing) water bodies.
- 312 (4) within North America (i.e. Canada & U.S.A.).

313 We will include all studies in North America because many environmental conditions and
314 invasive species will be shared between British Columbia and other regions within Canada
315 and the U.S. However, including all studies in North America might capture information not

316 relevant to British Columbia. For instance, studies might investigate the impacts of invasive
317 plant species on abiotic and biotic components absent in our study system. Such cases will
318 be excluded, and exclusions justified. Similarly, we will justify all other exceptions (63).

319

320 *Study Validity Assessment*

321 We assessed the validity of each study based only on the eligibility criteria.

322

323 *Data coding*

324 For each study at the full-text screening stage, we will provide the following information:

325

326

327 1. Bibliographic information

328 a) Authors list

329 b) Article title

330 c) Publication year

331 d) Bibliographic source

332 2. Inclusion criteria

333 a) Exposure: Focuses on target species (Y/N)

334 b) Exposure: Focuses on abiotic and biotic impacts (Y/N)

335 c) Population: Focuses on riparian and foreshore ecosystems (Y/N)

336 d) Population: Within North America (Y/N)

337 3. Screening stage

338 a) Excluded at full-text stage

339 b) Included

340 c) Exceptions

341 4. Additional information

342 a) Duplicate (Y/N)

343 b) Notes

344

345 For included studies only, we will provide also the following information:

346

347

348 1. Bibliographic information

349 a) Authors list

350 b) Article title

351 c) Publication year

352 2. Information on impacts

353 a) Impact description

354 b) Ecosystem component impacted (e.g. species, soil, etc.)

355 c) Magnitude of impact

356 d) Impact direction (negative, positive, neutral)

357 3. Additional information

358 a) Geographic region

359 b) Study Design (i.e. field or laboratory experiment, correlation or direct
360 observation)

361 c) Notes

362

363 We will compile subsection 3c. *Exceptions* on a case-by-case basis. For included studies,

364 we will provide information by impact so that if a study investigated more than one, there will

365 be a number of entries equivalent to the number of impacts assessed.

366

367 *Meta-data extraction*

368 Studies included in the systematic literature map will undergo a full-manuscript screening to

369 identify the investigated impact (or impacts). We will provide a description of the investigated

370 impacts and the ecosystem component impacted. Then, we will categorize impacts by their

371 magnitude and directionality. Impacts magnitude will be assessed following previous work,
372 modified to include both positive and negative impacts (31–33):

373

374

375

- **Minimal:** The impact is unlikely or negligible.

376

- **Minor:** It causes changes in the fitness of individuals in the native biota, but no changes in native population densities.

377

378

- **Moderate:** It causes changes in the population densities of native species, but no changes to the structure of communities or the abiotic or biotic components of ecosystems.

379

380

381

- **Major:** It causes the local or population extinction/introduction of at least one native species, and leads to reversible/transient changes in the structure of communities and the abiotic or biotic components of ecosystems.

382

383

384

- **Massive:** It leads to the replacement and local extinction/introduction of multiple native species, and produces irreversible changes in the structure of communities and the abiotic or biotic components of ecosystems.

385

386

387

388 *Synthesis and presentation*

389 For each species, we will provide a first database with all studies included at the full-text

390 screening and a reason for exclusions at this stage. A second database with the studies

391 included in the map, along with a graphical representation of the screening process. Both

392 databases will contain corresponding coded metadata (see *Data Coding* section). We will

393 import studies included in the review into a reference manager and share them as a public

394 library to facilitate accessibility. We will develop a graphical representation of riparian

395 ecosystems, representing identified impacts and their magnitude and directionality for each

396 species. Then, we will create a matrix combining multiple species (as rows) and impacts (as

397 columns) to illustrate the collective impacts of the focus species. Descriptive statistics will be

398 used to answer secondary questions. We will provide the geographic distribution of studies,

399 visualize publication trends over time, and illustrate differences in species and impacts
400 research effort. We will use co-occurrence matrices to identify research effort biases (64).
401 Lastly, we will provide a narrative synthesis of results for both main and secondary
402 questions. The narrative synthesis will focus on (i) species and impact prioritization, (ii)
403 clusters and gaps in present knowledge, (iii) predicted variations in impact magnitude and
404 direction under current climate change scenarios, and (iv) avenues for future research.

405

406 *Ethics approval and consent to participate*

407 Not applicable.

408 *Consent for publication*

409 Not applicable.

410 *Availability of data and materials*

411 Data sharing is not applicable to this article as no datasets were generated or analyzed
412 during the current study.

413 *Competing interests*

414 The authors declare that they have no competing interests.

415 *Funding*

416 FM is funded by the Ministry of Forests, British Columbia, Canada, and by the Irving K.
417 Faculty of Science at the Okanagan campus of UBC. JP acknowledges financial support
418 from the Natural Science and Engineering Council of Canada (Discovery Grant 2020-
419 06543).

420 *Authors' contributions*

421 FM drafted the protocol with input from JP and CM. All authors read and approved the final
422 manuscript.

423 *Acknowledgements*

424 We thank Mathew Vis-Dunbar (UBC librarian) for support and guidance in developing search
425 strings and extracting information from bibliographic databases. We thank all the
426 stakeholders for the feedback provided throughout the process.

427 *References*

- 428 1. Poff B, Koestner KA, Neary DG, Henderson V. Threats to Riparian Ecosystems in
429 Western North America: An Analysis of Existing Literature. JAWRA Journal of the American
430 Water Resources Association. 2011;47(6):1241–54.
- 431 2. Stella JC, Rodríguez-González PM, Dufour S, Bendix J. Riparian vegetation research
432 in Mediterranean-climate regions: common patterns, ecological processes, and
433 considerations for management. Hydrobiologia. 2013 Nov;719(1):291–315.
- 434 3. Ragan K, Schipper J, Bateman HL, Hall SJ. Mammal use of riparian corridors in
435 semi-arid Sonora, Mexico. The Journal of Wildlife Management. 2023;87(1):e22322.
- 436 4. Koundouri P, Boulton AJ, Datry T, Souliotis I. Ecosystem Services, Values, and
437 Societal Perceptions of Intermittent Rivers and Ephemeral Streams. In: Intermittent Rivers
438 and Ephemeral Streams. Elsevier; 2017. p. 455–76.
- 439 5. Patten DT. Riparian ecosystems of semi-arid North America: Diversity and human
440 impacts. Wetlands. 1998 Dec 1;18(4):498–512.
- 441 6. Knopf FL, Johnson RR, Rich T, Samson FB, Szaro RC. Conservation of Riparian
442 Ecosystems in the United States. The Wilson Bulletin. 1988;100(2):272–84.
- 443 7. Erős T, Kuehne L, Dolezsai A, Sommerwerk N, Wolter C. A systematic review of
444 assessment and conservation management in large floodplain rivers – Actions postponed.
445 Ecological Indicators. 2019;98:453–61.

- 446 8. Fryirs KA, Zhang N, Duxbury E, Ralph T, Fryirs KA, Zhang N, et al. Rivers up in
447 smoke: impacts of Australia's 2019–2020 megafires on riparian systems. *Int J Wildland Fire*.
448 2022 Jun 16;31(7):720–7.
- 449 9. Brierley GJ, Hikuroa D, Fuller IC, Tunnicliffe J, Allen K, Brasington J, et al.
450 Reanimating the strangled rivers of Aotearoa New Zealand. *WIREs Water*.
451 2023;10(2):e1624.
- 452 10. Stella JC, Bendix J. Chapter 5 - Multiple Stressors in Riparian Ecosystems. In:
453 Sabater S, Elozegi A, Ludwig R, editors. *Multiple Stressors in River Ecosystems*. Elsevier;
454 2019. p. 81–110.
- 455 11. Airoldi L, Beck M. Loss, Status and Trends for Coastal Marine Habitats of Europe. In:
456 Gibson R, Atkinson R, Gordon J, editors. *Oceanography and Marine Biology*. CRC Press;
457 2007. p. 345–405. (*Oceanography and Marine Biology - An Annual Review*; vol. 20074975).
- 458 12. Johnson PT, Olden JD, Vander Zanden MJ. Dam invaders: impoundments facilitate
459 biological invasions into freshwaters. *Frontiers in Ecology and the Environment*.
460 2008;6(7):357–63.
- 461 13. Chiu MC, Leigh C, Mazor R, Cid N, Resh V. Anthropogenic Threats to Intermittent
462 Rivers and Ephemeral Streams. In: *Intermittent Rivers and Ephemeral Streams*. Elsevier;
463 2017. p. 433–54.
- 464 14. Greene SL. A Roadmap for Riparian Invasion Research. *River Research and
465 Applications*. 2014;30(5):663–9.
- 466 15. Post GC, Chang H, Banis D. The spatial relationship between patterns of
467 disappeared streams and residential development in Portland, Oregon, USA. *Journal of
468 Maps*. 2022 Dec 1;18(2):210–8.
- 469 16. Sweeney BW, Bott TL, Jackson JK, Kaplan LA, Newbold JD, Standley LJ, et al.
470 Riparian deforestation, stream narrowing, and loss of stream ecosystem services.
471 *Proceedings of the National Academy of Sciences*. 2004;101(39):14132–7.
- 472 17. Swift BL. Status of Riparian Ecosystems in the United States. *JAWRA Journal of the
473 American Water Resources Association*. 1984;20(2):223–8.

- 474 18. Aguiar FCF, Ferreira MT. Plant invasions in the rivers of the Iberian Peninsula, south-
475 western Europe: A review. *Plant Biosystems*. 2013;147(4):1107–19.
- 476 19. Arianoutsou M, Delipetrou P, Celesti-Gradow L, Basnou C, Bazos I, Kokkoris Y, et al.
477 Comparing naturalized alien plants and recipient habitats across an east–west gradient in
478 the Mediterranean Basin. *Journal of Biogeography*. 2010;37(9):1811–23.
- 479 20. Kowarik I, Säumel I. Water dispersal as an additional pathway to invasions by the
480 primarily wind-dispersed tree *Ailanthus altissima*. *Plant Ecol*. 2008;198(2):241–52.
- 481 21. Pysek P, Prach K. Plant Invasions and the Role of Riparian Habitats: A Comparison
482 of Four Species Alien to Central Europe. *Journal of Biogeography*. 1993;20(4):413–20.
- 483 22. Barnes WJ. The Rapid Growth of a Population of Reed Canarygrass (*Phalaris*
484 *arundinacea* L.) and Its Impact on Some Riverbottom Herbs. *Journal of the Torrey Botanical*
485 *Society*. 1999;126(2):133–8.
- 486 23. Price EPF, Spyreas G, Matthews JW. Biotic homogenization of regional wetland
487 plant communities within short time-scales in the presence of an aggressive invader. *Journal*
488 *of Ecology*. 2018;106(3):1180–90.
- 489 24. Bateman HL, Chung-MacCoubrey A, Snell HL. Impact of Non-Native Plant Removal
490 on Lizards in Riparian Habitats in the Southwestern United States. *Restoration Ecology*.
491 2008;16(1):180–90.
- 492 25. Pendleton R, Pendleton B, Finch D. Displacement of Native Riparian Shrubs by
493 Woody Exotics: Effects on Arthropod and Pollinator Community Composition. *Natural*
494 *Resources and Environmental Issues*. 2011;16(1).
- 495 26. Martinez AE, McDowell PF. Invasive Reed Canarygrass (*Phalaris arundinacea*) and
496 Native Vegetation Channel Roughness. *Invasive Plant Science and Management*.
497 2016;9(1):12–21.
- 498 27. Simons SB, Seastedt TR. Decomposition and Nitrogen Release from Foliage of
499 Cottonwood (*Populus deltoides*) and Russian-olive (*Elaeagnus angustifolia*) in a Riparian
500 Ecosystem. *The Southwestern Naturalist*. 1999;44(3):256–60.

- 501 28. Mineau MM, Baxter CV, Marcarelli AM. A Non-Native Riparian Tree (*Elaeagnus*
502 *angustifolia*) Changes Nutrient Dynamics in Streams. *Ecosystems*. 2011;14(3):353–65.
- 503 29. Jacinthe PA, Bills JS, Tedesco LP. Size, activity and catabolic diversity of the soil
504 microbial biomass in a wetland complex invaded by reed canary grass. *Plant Soil*.
505 2010;329(1):227–38.
- 506 30. Collier N, Austin BJ, Bradshaw CJA, McMahon CR. Turning Pests into Profits:
507 Introduced Buffalo Provide Multiple Benefits to Indigenous People of Northern Australia.
508 *Hum Ecol*. 2011;39(2):155–64.
- 509 31. Katsanevakis S, Tempera F, Teixeira H. Mapping the impact of alien species on
510 marine ecosystems: the Mediterranean Sea case study. *Diversity and Distributions*.
511 2016;22(6):694–707.
- 512 32. Blackburn TM, Essl F, Evans T, Hulme PE, Jeschke JM, Kühn I, et al. A Unified
513 Classification of Alien Species Based on the Magnitude of their Environmental Impacts.
514 *PLOS Biology*. 2014;12(5):e1001850.
- 515 33. Vimercati G, Probert AF, Volery L, Bernardo-Madrid R, Bertolino S, Céspedes V, et
516 al. The EICAT+ framework enables classification of positive impacts of alien taxa on native
517 biodiversity. *PLOS Biology*. 2022;20(8):e3001729.
- 518 34. Sax DF, Schlaepfer MA, Olden JD. Identifying key points of disagreement in non-
519 native impacts and valuations. *Trends in Ecology & Evolution*. 2023;
- 520 35. Cardou F, Vellend M. Stealth advocacy in ecology and conservation biology.
521 *Biological Conservation*. 2023;280:109968.
- 522 36. Canadian Environmental Assessment Agency, editor. Cumulative effects assessment
523 practitioners guide. Ottawa; 1999.
- 524 37. Council on Environmental Quality. National Environmental Policy Act - Cumulative
525 Effects.
- 526 38. European Environment Agency. [https://www.eea.europa.eu/help/glossary/eea-](https://www.eea.europa.eu/help/glossary/eea-glossary/cumulative-impacts)
527 [glossary/cumulative-impacts](https://www.eea.europa.eu/help/glossary/eea-glossary/cumulative-impacts). Cumulative Impacts.

- 528 39. Spaling H, Smit B. Cumulative environmental change: Conceptual frameworks,
529 evaluation approaches, and institutional perspectives. *Environmental Management*.
530 1993;17(5):587–600.
- 531 40. Vanderhoeven S, Dassonville N, Meerts P. Increased Topsoil Mineral Nutrient
532 Concentrations Under exotic invasive plants in Belgium. *Plant Soil*. 2005;275(1):169–79.
- 533 41. Follstad Shah JJ, Harner MJ, Tibbets TM. *Elaeagnus angustifolia* Elevates Soil
534 Inorganic Nitrogen Pools in Riparian Ecosystems. *Ecosystems*. 2010;13(1):46–61.
- 535 42. Spyreas G, Wilm BW, Plocher AE, Ketzner DM, Matthews JW, Ellis JL, et al.
536 Biological consequences of invasion by reed canary grass (*Phalaris arundinacea*). *Biol*
537 *Invasions*. 2010;12(5):1253–67.
- 538 43. Rojas IM, Zedler JB. An invasive exotic grass reduced sedge meadow species
539 richness by half. *Wetlands Ecol Manage*. 2015;23(4):649–63.
- 540 44. Katz GL, Tuttle GM, Denslow MW, Norton AP. Legacy Effects of Russian Olive
541 (*Elaeagnus angustifolia* L.) in a Riparian Ecosystem Three Years Post-Removal. *Wetlands*.
542 2020;40(6):1897–907.
- 543 45. Gallego-Tévar B, Grewell BJ, Whitcraft CR, Futrell JC, Bárcenas-Moreno G, Castillo
544 JM. Contrasted Impacts of Yellow Flag Iris (*Iris pseudacorus*) on Plant Diversity in Tidal
545 Wetlands within Its Native and Invaded Distribution Ranges. *Diversity*. 2022;14(5):326.
- 546 46. Rodgers VL, Scanga SE, Kolozsvary MB, Garneau DE, Kilgore JS, Anderson LJ, et
547 al. Where Is Garlic Mustard? Understanding the Ecological Context for Invasions of *Alliaria*
548 *petiolata*. *BioScience*. 2022;72(6):521–37.
- 549 47. Simberloff D, Von Holle B. Positive Interactions of Nonindigenous Species:
550 Invasional Meltdown? *Biological Invasions*. 1999 Mar 1;1(1):21–32.
- 551 48. Benoit LK, Askins RA. Impact of the spread of *Phragmites* on the distribution of birds
552 in Connecticut tidal marshes. *Wetlands*. 1999;19(1):194–208.
- 553 49. Brooks RK, Barney JN, Salom SM. The invasive tree, *Ailanthus altissima*, impacts
554 understory nativity, not seedbank nativity. *Forest Ecology and Management*.
555 2021;489:119025.

- 556 50. Claeson SM, LeRoy CJ, Barry JR, Kuehn KA. Impacts of invasive riparian knotweed
557 on litter decomposition, aquatic fungi, and macroinvertebrates. *Biol Invasions*.
558 2014;16(7):1531–44.
- 559 51. Katz GL, Shafroth PB. Biology, ecology and management of *Elaeagnus angustifolia*
560 L. (Russian olive) in western North America. *Wetlands*. 2003;23(4):763–77.
- 561 52. Medina-Villar S, Rodríguez-Echeverría S, Lorenzo P, Alonso A, Pérez-Corona E,
562 Castro-Díez P. Impacts of the alien trees *Ailanthus altissima* (Mill.) Swingle and *Robinia*
563 *pseudoacacia* L. on soil nutrients and microbial communities. *Soil Biology and Biochemistry*.
564 2016;96:65–73.
- 565 53. Collaboration for Environmental Evidence. Guidelines and Standards for Evidence
566 synthesis in Environmental Management. Version 5.1 (AS Pullin, GK Frampton, B Livoreil &
567 G Petrokofsky, Eds) [Internet]. 2022. Available from:
568 <https://environmentalevidence.org/information-for-authors/>
- 569 54. Haddaway NR, Bernes C, Jonsson BG, Hedlund K. The benefits of systematic
570 mapping to evidence-based environmental management. *Ambio*. 2016;45(5):613–20.
- 571 55. Haddaway NR, Macura B, Whaley P, Pullin AS. ROSES RepOrting standards for
572 Systematic Evidence Syntheses: pro forma, flow-diagram and descriptive summary of the
573 plan and conduct of environmental systematic reviews and systematic maps. *Environ Evid*.
574 2018;7(1):7.
- 575 56. Palacio FX, Callaghan CT, Cardoso P, Hudgins EJ, Jarzyna MA, Ottaviani G, et al. A
576 protocol for reproducible functional diversity analyses. *Ecography*. 2022;2022(11):e06287.
- 577 57. Bellard C, Thuiller W, Leroy B, Genovesi P, Bakkenes M, Courchamp F. Will climate
578 change promote future invasions? *Glob Chang Biol*. 2013;19(12):3740–8.
- 579 58. Perry LG, Reynolds LV, Shafroth PB. Divergent effects of land-use, propagule
580 pressure, and climate on woody riparian invasion. *Biol Invasions*. 2018;20(11):3271–95.
- 581 59. Poland TM, Patel-Weynand T, Finch DM, Miniati CF, Hayes DC, Lopez VM, editors.
582 *Invasive Species in Forests and Rangelands of the United States: A Comprehensive*

583 Science Synthesis for the United States Forest Sector. Cham: Springer International
584 Publishing; 2021.

585 60. Ridley FA, McGowan PJ, Mair L. The scope and extent of literature that maps threats
586 to species: a systematic map protocol. *Environmental Evidence*. 2020;9(1):23.

587 61. Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas*.
588 1960;20:2746.

589 62. Bayliss HR, Beyer FR. Information retrieval for ecological syntheses. *Research*
590 *Synthesis Methods*. 2015;6(2):136–48.

591 63. O’Dea RE, Lagisz M, Jennions MD, Koricheva J, Noble DWA, Parker TH, et al.
592 Preferred reporting items for systematic reviews and meta-analyses in ecology and
593 evolutionary biology: a PRISMA extension. *Biol Rev*. 2021;96(5):1695–722.

594 64. Ridley FA, Hickenbotham EJ, Suggitt AJ, McGowan PJK, Mair L. The scope and
595 extent of literature that maps threats to species globally: a systematic map. *Environ Evid*.
596 2022;11(1):1–26.

597