

# PHILOSOPHICAL TRANSACTIONS OF THE ROYAL SOCIETY B

BIOLOGICAL SCIENCES

## Amazonian soundscapes: unravelling the secrets of insect acoustic niches in diverse habitats

Journal:	<i>Philosophical Transactions B</i>
Manuscript ID	RSTB-2023-0112
Article Type:	Research
Date Submitted by the Author:	05-Nov-2023
Complete List of Authors:	Do Nascimento, Leandro A.; biometrio.earth, Science Department Pérez-Granados, Cristian; Universidad de Alicante, Ecology Department Alencar, Janderson; Instituto Nacional de Pesquisas da Amazônia, Programa de Pós-Graduação em Ciências Biológicas (Entomologia) Beard, Karen; Utah State University, Wildland Resources and Ecology Center
Issue Code (this should have already been entered and appear below the blue box, but please contact the Editorial Office if it is not present):	INSECTS
Subject:	Ecology < BIOLOGY
Keywords:	Acoustic Evenness Index, acoustic indices, crickets, habitat structure, passive acoustic monitoring, acoustic niche

SCHOLARONE™  
Manuscripts

**Author-supplied statements**

Relevant information will appear here if provided.

**Ethics**

*Does your article include research that required ethical approval or permits?:*

Yes

*Statement (if applicable):*

Data collection complied with all Brazilian legal requirements (SISBIO license #56539-2).

**Data**

*It is a condition of publication that data, code and materials supporting your paper are made publicly available. Does your paper present new data?:*

Yes

*Statement (if applicable):*

Raw datasets and codes can be found at [REDACTED]

**Conflict of interest**

I/We declare we have no competing interests

*Statement (if applicable):*

CUST\_STATE\_CONFLICT :No data available.

**Use of AI**

*Please provide a statement of any use of AI technology in the preparation of the paper.*

No, we have not used AI-assisted technologies in creating this article

CUST\_IF\_YES\_DECLARATION\_OF\_AI\_USE :No data available.

1  
2  
3 Amazonian soundscapes: unravelling the secrets of insect acoustic niches in diverse habitats  
4  
5  
6  
7

8 Authors: Leandro A. Do Nascimento<sup>1\*</sup>, Cristian Pérez-Granados<sup>2</sup>, Janderson B. Rodrigues Alencar<sup>3</sup>,  
9 Karen H. Beard<sup>4</sup>

10 <sup>1</sup>Science Department, biometrio.earth, 66123 Saarbrücken, Germany. ORCID: 0000-0001-8254-2900

11 <sup>2</sup>Ecology Department, Alicante University, 03690 Alicante, Spain. ORCID: 0000-0003-3247-4182

12 <sup>3</sup>Instituto Nacional de Pesquisas da Amazônia, Programa de Pós-Graduação em Ciências Biológicas  
13 (Entomologia), 69067-375 Manaus, AM, Brazil. ORCID: 0000-0001-9482-7866

14 <sup>4</sup>Department of Wildland Resources and Ecology Center, Utah State University, 84322-5230 Logan, UT,  
15 USA. ORCID: 0000-0003-4997-2495

16  
17  
18  
19 \*Correspondence: Leandro A. Do Nascimento, Email: le\_nasc@hotmail.com  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Abstract

Insects are the most diverse animal taxon on Earth and play a key role in ecosystem functioning. However, they are often neglected from biodiversity surveys due to the difficulties of monitoring this small and hyper-diverse taxon. With technological advancements in biomonitoring and analytical methods, these shortcomings may finally be addressed. Here, we performed passive acoustic monitoring at 141 sites (eight habitats) in the Amazon to advance insect surveys using acoustic methods. We first describe the frequency range occupied by three soniferous insect groups (cicadas, crickets, and katydids) to calculate the Acoustic Evenness Index (AEI). Then, we assess the temporal and spatial variations of AEI among the insect categories, and finally we investigate the relationship among vegetation structure variables and AEI for each category. Overall, crickets occupied lower and narrower frequency bands than cicadas and katydids. AEI values varied among insect categories and across space and time. There was higher acoustic activity before sunrise and lower acoustic activity was recorded at pastures. Canopy cover was positively associated with crickets' acoustic activity but not with katydids. Our findings contribute to a better understanding of the role of time, habitat, and vegetation structure in shaping insect activity within the diverse Amazonian ecosystem.

## 1. Introduction

Global diversity has declined worldwide in recent decades in what is often considered the sixth major extinction event (Cowie et al. 2022). However, the biodiversity crisis is not homogeneous and varies among areas and taxonomic groups. For example, biodiversity loss is greater in tropical forests, where deforestation is causing extinctions at unprecedented rates (Brooks et al. 1999) and is also greater for insects than for other taxa (van Klink et al. 2020). Insects are the most diverse group of animals, comprising around 80% of all described animal life (Stork 2018), which together with their small size and our limited knowledge about their ecology (c. 80% of insect species remains undescribed, Cardoso et al. 2011), have made them challenging to study. However, technological advances during the last decades have provided new scalable, automated, and non-invasive methods for monitoring insects at large spatial and temporal scales (van Klink et al. 2022, Van Klink et al. 2024).

Among the recently developed automated and non-invasive techniques for biomonitoring is passive acoustic monitoring (PAM, Sugai et al. 2019). PAM requires the placement of programmed autonomous sound recorders in the field and followed by recording interpretation (manually or using automated software). This technique has proven to be a suitable method for monitoring wildlife across many regions and taxa (Sugai et al. 2019). However, invertebrates are, by far, the biological group least monitored using this technique (<5% of the studies, Sugai et al. 2019). Likewise, most research has been concentrated in the northern temperate zone (65% of the studies, Sugai et al. 2019), even though one of the main advantages of using PAM relies on its ability to monitor wildlife in ecosystems difficult to reach for ecologists, such as tropical

1  
2  
3 forests (Deichman et al. 2018, Pérez-Granados and Schuchmann 2020, Do Nascimento et al.  
4  
5 2021).

6  
7  
8  
9 A diverse range of insects produce sounds and therefore PAM enables a scalable and  
10  
11 standardized methodology for monitoring soniferous insects, which dominate nocturnal  
12  
13 soundscapes in the tropical region (Aide et al. 2017, Müller et al. 2023). Indeed, insects can be  
14  
15 acoustically detected far away from the recorders (e.g., up to 100 m), which is a much longer  
16  
17 distance than insect detection range using pitfalls or nets (Jeliazkov et al. 2016). Moreover, PAM  
18  
19 surveys can be coupled with automated species recognition, thus allowing for species-specific  
20  
21 monitoring with less effort than traditional survey methods (van Klink et al. 2022, Symes et al.  
22  
23 2024). However, this approach is limited by the lack of available training data for the algorithms,  
24  
25 and this is especially problematic for insects (ter Hofstede et al. 2020, Symes et al. 2022).

26  
27  
28  
29  
30 PAM-based surveys collect soundscape recordings, which contain available information  
31  
32 about the state of biological assemblages. A soundscape can be described as the sonic structure  
33  
34 of a given location, accounting for signal composition, diversity, intensity and temporal patterns  
35  
36 (Schafer 1977). Soundscapes are usually divided into biotic and abiotic components according to  
37  
38 main sound sources, with the biotic component, usually defined as biophony, composed of the  
39  
40 collective sounds produced by living organisms (Pijanowski et al. 2011). Several acoustic indices  
41  
42 have been developed to aid in the interpretation of soundscapes and have often proven their  
43  
44 ability to assess species richness, composition, or whether acoustic activity varies within  
45  
46 different ecosystems (Buxton et al. 2018, Bradfer-Lawrence et al. 2020, Do Nascimento et al.  
47  
48 2020, Allen-Ankins et al. 2023, Müller et al. 2023; but see Alcocer et al. 2022 for limitations).  
49  
50 Several groups of insects produce sounds at very specific (e.g., higher than birds) and narrow  
51  
52 frequencies (Metcalf et al. 2021), and therefore acoustic indices could be an effective alternative  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 for monitoring the relationship between insect acoustic activity and other variables of interest.  
4  
5 Indeed, some of the existing indices already have been used as a surrogate of insect richness or  
6  
7 composition (Aide et al. 2017; Müller et al. 2023) and to describe temporal patterns of insect  
8  
9 communities (Ferreira et al. 2018, Burivalova et al. 2022). However, our current knowledge  
10  
11 about the use of acoustic indices for monitoring insects is still limited and their effectiveness is  
12  
13 expected to vary among regions owing to different background noise and species compositions  
14  
15 (Müller et al. 2023, Scarpelli et al. 2023).  
16  
17  
18

19 Here, we draw upon the acoustic adaptation hypothesis (Morton 1975) to test the effects  
20  
21 of habitat type and vegetation structure on the acoustic signals produced by insect communities.  
22  
23 We recorded soundscapes and collected vegetation data at 141 sites across eight habitat types  
24  
25 (from natural to anthropogenic) representing dominant habitat types found in the Amazon basin.  
26  
27 Our approach comprises three main goals: (1) to describe manually the frequency range occupied  
28  
29 by three soniferous insects groups (cicadas, crickets, and katydids) in our study area, and to  
30  
31 calculate acoustic indices targeting these groups to semi-automatically study their ecology  
32  
33 (Burivalova et al. 2022, Metcalf et al. 2021), (2) to assess the temporal (hourly) and spatial  
34  
35 (habitat type) variations of acoustic indices among insect groups, and (3) to investigate the  
36  
37 relationship between vegetation structure variables and acoustic indices on targeted insect groups  
38  
39 to better understand relationships between insect activity and habitat structure. We predict that:  
40  
41 (a) insect acoustic activity will vary among habitat types because habitat type influences insect  
42  
43 community composition as well as their acoustic communication and behaviour (Römer and  
44  
45 Lewalds 1992, Schmidt and Balakrishnan 2015). More specifically, we expect forested habitats  
46  
47 (and those with higher vegetation complexity) to have more diverse insect communities and thus  
48  
49 a more diverse acoustic environment, while open (more homogeneous) habitat types would have  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 more homogeneous insect communities and thus less diverse acoustic environments, due to more  
4 favourable microenvironment conditions in forested habitats that may sustain greater insect  
5 richness and abundance. We also predict that (b) insect acoustic activity will vary among  
6 recording hours because several environmental factors (e.g., moonlight, temperature) may  
7 influence their activity and sound production throughout the night in tropical forests (Oliveira et  
8 al. 2021, Gomez-Morales and Acevedo-Charry 2022, Scarpelli et al. 2023, Symes et al. 2024).  
9 Our findings will contribute to a better understanding of the role of time, habitat and vegetation  
10 structure in shaping insect acoustic activity within the diverse Amazonian ecosystem and might  
11 be useful to guide future research on other tropical regions.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24


## 25 **2. Methods**

### 26 (a) Study sites

27  
28  
29  
30 We conducted this research in and around Viruá National Park (VNP), Roraima, Brazil,  
31 in the north of the Brazilian Amazon (Figure 1). VNP was established in 1998 and is 240,000 ha  
32 (ICMBio, 2014). The climate in VNP is warm and wet with mean annual temperature of 26°C  
33 and mean annual precipitation of ~2,000 mm (ICMBio, 2014). Rainfall is mostly concentrated  
34 from May to September (ICMBio, 2014). VNP is regulated by floods that create a vegetation  
35 mosaic ranging from dense forests to grasslands and has most major habitats found across the  
36 Amazon biome (ICMBio, 2014). These habitats share common species, but also harbour unique  
37 fauna and flora. The study was carried out in eight different habitat types, separated according to  
38 their vegetation characteristics and flooding regime (Figure 1). The eight habitats surveyed can  
39 be grouped into three main categories: i) open habitats (Burned campina, Campina, and  
40 Pastures), ii) flooded forests (Igapó, Islands, and Várzea), and iii) non-flooded forests  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



(Campinarana and Terra-firme). In general, open habitats have lower vertebrate species richness than forested habitats, and Campiranana, Igapó, and Island forests are not as diverse as Terra-firme and Várzea forests, yet it is unclear if these patterns hold for invertebrates (see Do Nascimento et al. 2020 for additional information on habitats).



**Figure 1.** Location of the surveyed sites (141) across the eight habitat types studied in the Viruá National Park, Roraima, Brazil. The inset shows the location of the study area in northern Brazil.

(b) Passive acoustic monitoring

We used Arbimon acoustic recorders (Campos-Cerqueira & Aide, 2016) to collect acoustic data during the dry season, from November 2016 to February 2017, in the eight habitat types. We deployed recorders at 20 replicate sites in each habitat, except for Pastures (six replicates because they comprised a small area in the VNP); and Campinarana and Terra-firme

1  
2  
3 (18 and 17 replicates, respectively, because of recorder malfunctions). Therefore, the final  
4  
5 acoustic dataset was collected in 141 sites. Recorders were separated by at least 500 m to  
6  
7 minimize the risk of overlapping detections across recorders (vocalizations of birds and frogs  
8  
9 recorded with the employed recorder can be detected up to ~100 m, Campos-Cerqueira et al.  
10  
11 2020). We attached recorders to trees or fixed poles at the height of 1.5 m. Acoustic devices were  
12  
13 programmed to record 1 min of audio every 10 min (144 1-min recordings per day) for six days  
14  
15 in each sampling site (sample rate = 44.1 kHz; resolution = 16 bit; format = WAV). After six  
16  
17 days, the 20 recorders were moved to a different habitat type and the method repeated. We  
18  
19 inspected recordings for file corruption, heavy rain, and wind and removed those files when any  
20  
21 of these circumstances occurred (details in Do Nascimento et al. 2020). The acoustic dataset we  
22  
23 used for subsequent analysis had 95,141 one-minute recordings (1,585 hours).  
24  
25  
26  
27  
28  
29  
30

### 31 (c) Vegetation structure survey

32  
33 Vegetation structure data were collected within a 20-m radius plot from each acoustic  
34  
35 recorder location (141 sites) after the devices were moved to a different habitat type to avoid  
36  
37 interference with the recordings. For each site, we measured the percent canopy cover, canopy  
38  
39 height, litter depth, number of small trees (Diameter at Breast Height – DBH > 1 cm to < 10 cm),  
40  
41 number of large trees (DBH > 10 cm), and shrub cover. For each vegetation structure variable,  
42  
43 we used the mean value per site for subsequent analysis. Details about the vegetation structure  
44  
45 survey methodology can be found in Do Nascimento et al. 2020.  
46  
47  
48  
49  
50

### 51 (d) Spectral frequencies occupied by insects

1  
2  
3 To characterize the spectral frequencies occupied by insects two experienced researchers  
4 (LAN and JBRA) with insect sounds and the study area manually reviewed 1,152 one-minute  
5 audio files (19.2 hours) from one randomly chosen day/site (24 hours) per habitat type. The  
6 amount of time manually reviewed was similar or higher to that considered in most prior  
7 research using acoustic indices and sonotypes (e.g., 228 minutes in Gasc et al. 2018, 672 minutes  
8 in Duarte et al. 2021, 63 minutes in Burivalova et al. 2022, and 1,342 minutes in Gomez-Morales  
9 and Acevedo-Charry 2022).

10  
11 For each file, we annotated the minimum and maximum frequencies of three main sound-  
12 producing insect groups in the study area using a sonotype approach (details in Aide et al. 2017):  
13 Cicadidae (Cicada), Grylloidea (Crickets), and Tettigoniidae (Katydids). We used Audacity  
14 (Audacity Software 3.3.3) to visually inspect the spectrograms and annotated all selected files  
15 using selection boxes around each sound (like Burivalova et al. 2022). When in doubt about the  
16 identification of a sonotype, we would consult each other or other specialists to reach consensus.  
17 In total, we annotated 1,714 sonotypes across insect groups. We also annotated if any other  
18 animal or sound would overlap in the frequency range occupied by the insects because this  
19 would allow us to focus our analysis on periods of little to no overlapping sounds (163 sonotypes  
20 were annotated). We also registered the beginning and ending time of insect choruses (i.e., two  
21 or more insects producing a very characteristic and continuous acoustic signature that may last  
22 for hours, Figure S1) for each site and habitat reviewed. Additionally, we estimated the chorus  
23 length for each habitat type as the time difference between the ending and beginning time of  
24 insect choruses.

25  
26 Our analysis revealed the cicada sounds were not present across all habitats in our  
27 manually reviewed dataset, therefore this group was excluded from subsequent analysis.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Nonetheless, we have opted for including in the results the spectral frequencies occupied by  
4 cicadas, which might be useful to guide future studies in the Amazon.  
5  
6  
7  
8  
9

10 (e) Acoustic index calculation  
11

12 We used the Python package “scikit-maad” (Ulloa et. al 2021) and Anaconda (Anaconda  
13 Software Distribution, 2016) with Python 3.9.17 to calculate the Acoustic Evenness Index (AEI,  
14 see description in Villanueva-Rivera et al. 2011) on three average frequency ranges mainly  
15 occupied by Crickets (5,000 Hz to 7,500 Hz), Katydid (8,000 Hz to 17,000 Hz), and what we  
16 called “All insects” category - encompassing both crickets and katydids (5,000 Hz to 17,000 Hz)  
17 using the default settings provided in the package (frequency step = 500 Hz). The AEI measures  
18 unevenness among frequency band steps returning a value close to one when sound intensity is  
19 restricted to few frequencies and a value close to zero when soundscapes are saturated with  
20 sounds (Bradfer-Lawrence et al. 2023, Do Nascimento et al., 2020). Therefore, recordings with  
21 lower values of AEI can be interpreted as recordings with higher sound activity, while recordings  
22 with higher AEI values can be interpreted as recordings with limited sound activity (Figures S1–  
23 S8). The AEI was chosen to describe insect acoustic activity based on prior research in the study  
24 area, which showed that the AEI was the acoustic index that better explained ( $R^2 = 81\%$ ) the  
25 relationship between soundscapes and vegetation structure outperforming several other acoustic  
26 indices on this task (Do Nascimento et. al 2020). We focused the index calculation on the  
27 nocturnal period because during this period there was minimal frequency overlap of crickets and  
28 katydids sounds with other animals or sound sources at our manually annotated dataset. In fact,  
29 only three out of 1,152 files reviewed had birds overlapping with crickets which reassures that  
30 the AEI values estimated during the nocturnal period will reflect the acoustic activity of insects.  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 We considered the nocturnal period to be the time elapsed between evening and morning  
4 astronomical twilight (Foote et al. 2017, Pérez-Granados and Schuchmann 2021), as extracted  
5 from [www.timeanddate.com](http://www.timeanddate.com). To standardize the number of recordings analysed per site and due  
6 to the low variation in twilight timing during the study period, we considered as nocturnal those  
7 recordings made between 1900 h to 0500 h, which was always between the evening and morning  
8 twilight times.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18

19 (f) Statistics

20  
21 First, to assess variations of the AEI among hours and habitat type, we fitted generalized  
22 linear mixed models (GLMM, Gaussian family) using the values (log transformed) of the AEI  
23 targeting each insect category (“All insects”, Crickets, and Katydid) as the response variable,  
24 hour of recording (“ten levels”) and habitat type (“eight levels”) as fixed effects and recording  
25 station as a random effect to control for variations within habitats owing to site. One independent  
26 GLMM was fitted for each insect category (“All insects”, Crickets, and Katydid). The Gaussian  
27 family was selected after testing different distribution families and choosing the most appropriate  
28 according to the Akaike information criteria (AIC; Burnham and Anderson, 2004) and visually  
29 inspection of the residuals. Model performance was evaluated by plotting standardized residuals  
30 versus fixed variables, normal Q-Q plots and histograms of residuals. No concrete pattern was  
31 found in any case. When a fixed effect was found to be significant, a Tukey’s post hoc test was  
32 performed to test for differences among levels.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

49 Second, to determine the influence of vegetation structure variables on the AEI, we fitted  
50 linear mixed models (LMM) using the mean values (log transformed) per site of the AEI  
51 targeting each insect category as the response variables (one independent LMM per insect  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 category), six vegetation structure variables (canopy cover, canopy height, litter depth, number  
4 of large trees, number of small trees, and shrub cover) as fixed effects, and habitat type as a  
5 random effect to control for variations owing to habitat groups. Fixed effects were scaled to  
6 make their parameter estimates comparable within models. We performed model selection using  
7 the AICc. We selected the top four performing models based on  $\Delta\text{AICc}$  and considered models  
8 to be similar if  $\Delta\text{AICc} < 2$  (Burnham and Anderson, 2004). We checked for multicollinearity of  
9 predictors and removed canopy height from the analysis. Residuals of the models were checked  
10 for linearity, homoscedasticity, independence, and normality. Finally, we calculated the marginal  
11 and conditional  $R^2$  values (Nakagawa and Schielzeth, 2013) to estimate the proportion of  
12 variance explained by fixed and random effects.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

26 All statistical analyses were performed with R 3.6.2 (R Development Core Team 2021).  
27 We used the packages “lme4” (Bates et al. 2015) for the GLMM construction, “lmerTest”  
28 (Kuznetsova et al. 2014) to calculate the significance of fixed effects, “multcomp” (Hothorn  
29 2008) for post hoc comparison tests, “nlme” (Pinheiro et al. 2019) for the LMM construction,  
30 “car” (Fox and Weisberg, 2019) and “SjPlot” (Lüdecke, 2020) for checking model assumptions,  
31 and “R2glmm” (Jaeger, 2017) to extract  $R^2$  values. The level of significance adopted was  $p <$   
32 0.05.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43

### 44 **3. Results**

#### 45 (a) Mean spectral frequencies occupied by insects and chorus temporal patterns

46  
47  
48 Cicadas were only found in Burned Campinas, Campinaranas, and Terra-firme sites. In  
49 Campinaranas, the mean maximum frequency of Cicadas was much higher (16.21 kHz) than at  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Terra Firme and Burned Campinas (8.64 and 8.95 kHz, respectively), but just one call was  
4 registered in Campinaranas. Crickets were found across all eight habitats surveyed, however, the  
5 variation among habitats between minimum and maximum frequencies was small in magnitude  
6 compared to the other insect groups (range between 5.13-7.73 kHz in all habitats). Katydid were  
7 also found across all habitats surveyed, and higher variations between minimum and maximum  
8 frequencies were identified across habitats (Table 1). For example, mean minimum and  
9 maximum frequencies of Katydid in Burned Campina were of 7.71 and 8.77 kHz, respectively,  
10 while in Island the mean minimum and maximum frequencies were of 12.11 kHz and 15.54 kHz  
11 (Table 1).  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

26 **Table 1.** Minimum and maximum frequencies (kHz, mean  $\pm$  SD) manually measured for  
27 sonotypes (N) of three insect groups across eight Amazon habitat types.  
28

Insect group	Habitat type	N	Minimum frequency	Maximum frequency
Cicadas	Burned campina	3	3.28 $\pm$ 0.35	8.95 $\pm$ 1.88
	Campinarana	1	3.06	16.21
	Terra-firme	77	3.19 $\pm$ 1.43	8.64 $\pm$ 2.84
Crickets	Burned campina	118	5.46 $\pm$ 1.09	6.48 $\pm$ 1.27
	Campina	118	5.27 $\pm$ 1.1	5.62 $\pm$ 1.28
	Campinarana	173	6.75 $\pm$ 1.81	7.73 $\pm$ 1.76
	Igapó	236	5.68 $\pm$ 1.56	7.17 $\pm$ 1.71
	Island	206	5.13 $\pm$ 1.4	7.13 $\pm$ 1.61

	Pasture	147	$5.37 \pm 1.09$	$6.41 \pm 1.36$
	Terra-firme	330	$6.15 \pm 2.61$	$7.26 \pm 2.53$
	Várzea	96	$6.13 \pm 0.83$	$7.02 \pm 1.06$
Katydid	Burned campina	4	$7.71 \pm 1.15$	$8.77 \pm 1.35$
	Campina	12	$9.34 \pm 0.46$	$12.63 \pm 0.47$
	Campinarana	2	$11.51 \pm 2.98$	$16.00 \pm 3.81$
	Igapó	15	$9.02 \pm 2.7$	$17.30 \pm 2.14$
	Island	56	$12.11 \pm 0.64$	$15.54 \pm 1.87$
	Pasture	35	$8.73 \pm 1.25$	$11.75 \pm 2.48$
	Terra-firme	73	$9.29 \pm 1.58$	$17.29 \pm 2.15$
	Várzea	12	$12.83 \pm 0.8$	$14.74 \pm 0.26$

The onset and end of the insect chorus greatly varied across habitats, lasting the longest (18 hours) on Islands and the shortest (7 hours) on Várzea (Table 2).

**Table 2.** Temporal patterns of insect chorus activity across Amazonian habitats based on the analysis of one randomly chosen replicate site for each habitat type during a 24-hour period.

Habitat	Beginning time	Ending time	Chorus length
Burned campina	1710 h	0540 h	12 h 30 m
Campina	1840 h	0310 h	8 h 30 m
Campinarana	1510 h	0810 h	17 h 0 m



Igapó	1740 h	0700 h	13 h 20 m
Island	1600 h	1000 h	18 h 0 m
Pasture	1730 h	0620 h	13 h 20 m
Terra-firme	1240 h	0600 h	17 h 20 m
Várzea	1800 h	0100 h	7 h 0 m

---

(b) The influence of the hour and habitat type on the AEI targeting insect sounds

According to GLMMs, the AEI values of “All insects” (5,000 Hz to 17,000 Hz), Crickets (5,000 to 7,500 Hz), and Katydid (8,000 Hz to 17,000 Hz) significantly varied across recording hours and habitat types (Table 3). Overall, there was a significant decreasing pattern of AEI values from sunset to sunrise, which was very consistent for the three insect categories (Figure 2). According to post-hoc tests, the higher AEI values were obtained from the recordings made at 1900 h and 2000 h for all tree insect categories, while the lowest AEI values were observed at 0200 h for Crickets (Figure 2B), and between 0200 h to 0400 h for “All insects” and Katydid (Figure 2A, 2C).


**Table 3.** Summary table of type-III partitioning of variances testing the effects of recording hour and habitat type on the variations of the Acoustic Evenness Index per insect category.

Recordings were collected at eight habitat types and during ten different recording hours.

Detailed results of the GLMMs per insect category can be found at Table S1–S3 in the online supplementary material.

<b>Insect</b>	<b>Fixed effect</b>	<b>Sum Sq</b>	<b>Mean Sq</b>	<b>DF</b>	<b>F</b>	<b><i>p</i></b>
<b>Category</b>						
All	Recording hour	1090.49	17	416	203.31	<b>&lt;0.001</b>
	Habitat type	11.98	1	33	2.87	<b>0.008</b>
Crickets	Recording hour	73.91	1	401	19.68	<b>&lt;0.001</b>
	Habitat type	28.54	8	33	9.77	<b>&lt;0.001</b>
Katydids	Recording hour	1272.89	43	415	270.71	<b>&lt;0.001</b>
	Habitat type	18.43	3	33	5.04	<b>&lt;0.001</b>


At the habitat level, there were more subtle differences across habitat types compared to the differences found at hourly scale (Figure 2). AEI values for all three insect categories were significantly higher for Pasture than the other habitats (Figure 2), although similar AEI values (non-significant differences according to post-hoc tests) were observed at Pastures and Burned campina for Crickets (Figure 2E), and similar values (non-significant) at Pasture and Islands for Katydids (Figure 2F). Lower AEI values were observed in Campina and Várzea for “All insects” category, in Várzea and Island for Crickets, and in Burned campina and Campina for Katydids.



**Figure 2.** Boxplots showing the variation of AEI values as a function of recording nocturnal hours (A-C) (19:00 to 04:00) and (D-F) habitat type. Results are shown separately for each insect category. Different colours show significant differences in AEI values between recording hours or habitat type from Tukey's post hoc test. B=Burned campina, C=Campina, CAP=Campinarana, IG=Igapó, IL=Island, PAS=Pasture, TF=Terra-firme, and V=Várzea.

(c) The influence of vegetation structure on the AEI targeting insect sounds

Across the top performing models (Table S4), only the AEI calculated in the frequency range dominated by Crickets was significantly and positively associated with canopy cover and negatively associated with shrub cover (Figure 3). The top performing models for AEI targeting Katydid and "All insects" category included null models, therefore we did not consider these groups to be significantly related to any of the vegetation structure variables, although canopy cover and the number of small trees was marginally associated with the "All insects" category. Conditional  $R^2$  of the Cricket top perform model was 41% while the marginal  $R^2$  of canopy cover and shrub cover were, respectively, 41% and 5%.



**Figure 3.** Results of linear mixed models for three insect categories (“All insects”, Crickets, and Katydid) showing the effects of five vegetation structure variables on the AEI values calculated for each insect category. Symbols are the normalized coefficients’ estimate values and horizontal lines represent the 95% confidence intervals. Estimates were normalized by subtracting raw values by the mean and dividing by the standard deviation. Detailed results of the LMMs can be found at Table S5 in the online supplementary material.

#### 4. Discussion

Acoustic indices are useful tools to analyse large quantities of audio data and investigate ecological patterns at broad spatial and temporal scales. Their usefulness is particularly high when coupled with expert ecological knowledge about the studied region and with other analytical methods or available datasets that may help to interpret the underlying ecological

1  
2  
3 patterns. Here, we demonstrate how manual annotation of a subset of recordings was useful in  
4  
5 calculating the Acoustic Evenness Index (AEI) for a time and frequency range sensitive to  
6  
7 different insect groups that dominate nocturnal soundscapes in the Neotropics. Overall, we found  
8  
9 that crickets occupy lower and narrower frequency bands than cicadas and katydids. We also  
10  
11 found that AEI values varied among insect categories across space and time, with a higher  
12  
13 acoustic activity (i.e., lower AEI values) during the hours before sunrise and lower acoustic  
14  
15 activity (higher AEI values) in Pastures across all insect categories analysed. Canopy cover was  
16  
17 positively associated with Cricket acoustic activity but not with Katydid. Our findings  
18  
19 contribute to our understanding of the role of time, habitat, and vegetation structure in shaping  
20  
21 insect acoustic activity within the hyper-diverse habitats of the Amazon using a non-invasive and  
22  
23 semi-automated approach based on soundscapes that is scalable (Müller et al. 2023).  
24  
25  
26  
27

28  
29 The manual annotation process allowed us to verify the absence of other (non-target)  
30  
31 vocally-active taxa during the night, and therefore reduce their influence when assessing changes  
32  
33 in AEI values. Indeed, birds were only detected in three of the 1,152 files reviewed, while  
34  
35 anurans appeared in few recordings, but their vocalizations were always made at frequency  
36  
37 bands lower than the ones used by insects (e.g., Figure S1). The frequency measurements  
38  
39 provided in our study might be useful as a baseline for future research aiming to work with these  
40  
41 groups of insects in the Amazon. However, prior research has demonstrated the existence of  
42  
43 geographical variation in insects' frequencies, partly driven by habitat type, and even of dialects  
44  
45 within the same species, and so we encourage researchers to develop their own group-specific  
46  
47 frequency bands (Claridge and Morgan 1993, Chen et al. 2019, Metcalf et al. 2021).  
48  
49  
50

51  
52 We found a clear pattern of increasing acoustic activity (lower AEI values) from sunset to  
53  
54 sunrise across all three insect categories analysed (Figure 2), in agreement with our prediction  
55  
56  
57  
58  
59  
60

1  
2  
3 that insect acoustic activity will significantly vary throughout the night in Amazonian habitats.  
4  
5 Our results agree with prior research in tropical forests from Panama (Symes et al. 2022) and  
6  
7 Colombia (Gomez-Morales and Acevedo-Charry 2022), where acoustic activity of crickets and  
8  
9 katydids also significantly varied throughout the night. However, the results were species-  
10  
11 dependent. For example, in Colombia two cricket sonotypes were more often detected around  
12  
13 sunset while five katydids sonotypes were detected more homogeneously throughout the night  
14  
15 (Gomez-Morales and Acevedo-Charry 2022). The authors attributed some of this variation  
16  
17 among groups to cricket acoustic activity being influenced by rainfall, while katydids were more  
18  
19 influenced by moon phase. In Panama, Symes et al. (2022) found that 24 katydid species were  
20  
21 detected throughout the night, but some species were reported to call more often right after  
22  
23 sunset and right before sunrise, with no moonlight effect (Symes et al. 2024). It is important to  
24  
25 note that we focused our study during the nocturnal period and excluded the sunset and sunrise  
26  
27 choruses from our analysis to avoid frequency and time overlap with other soniferous species, if  
28  
29 we could easily filter out non-target sounds (e.g. birds) to calculate the AEI (see advances in  
30  
31 source separation in Tolkova 2023), perhaps we would also detect more insect activity right after  
32  
33 sunset and right before sunrise as reported in prior research (Symes et al. 2022, Gomez-Morales  
34  
35 and Acevedo-Charry 2022). Despite this limitation about the use of acoustic indices, a  
36  
37 soundscape analysis approach is likely capturing the activity of much more insect species than  
38  
39 species-specific approach and thus reflecting broader patterns of calling behaviour (Müller et al.  
40  
41 2023, Scarpelli et al. 2023).

42  
43  
44  
45  
46  
47  
48  
49 When analysing the effect of habitat type on AEI values targeting three insect categories,  
50  
51 we did not find any clear pattern emerging from the results (Figure 2). Nonetheless, the variation  
52  
53 across habitats was significant in all insect categories. Pastures displayed the lowest acoustic  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 activity overall (higher AEI values) while forests (e.g., Várzea and Islands) had more acoustic  
4 activity for Crickets (lower AEI values). This result was expected as pastures are human  
5 modified habitats likely hosting lower insect diversity than forests (Do Nascimento 2020). But  
6 only by using fine scale habitat vegetation measurements the patterns in the data became clearer,  
7 unravelling how canopy cover has a positive effect ( $R^2 = 41\%$ ) on Crickets calling activity but  
8 not on Katydid or the broad “All insects” category (Figure 3). Therefore, partially confirming  
9 our prediction that more complex habitats (e.g., Várzea forests) will have higher insect acoustic  
10 activity than less complex habitats (e.g., Campina grasslands). Interestingly, crickets seem to call  
11 more often from sites with less shrub cover (Figure 3), which are more often found in flooded  
12 forests, such as Várzea and Islands, than non-flooded forests, such as Terra-firme. This shrub  
13 cover variation among forests could also be the result of flood pulses that wash out these  
14 ecosystems seasonally (Householder et al. 2021) but in turn may also bring nutrients and other  
15 resources important for cricket survival. Further research could monitor Várzea and Islands  
16 during the flooding period to test if acoustic activity of insects is higher when the herbaceous  
17 vegetation is submerged. To the best of our knowledge, this is the first study to test how major  
18 habitat formations from the Amazon basin may structure the calling activity of different insect  
19 groups using a semi-automated approach based on acoustic indices. The effect of habitat type  
20 and vegetation structure on insect calling activity have been discussed for a long time  
21 (Couldridge and van Staaden 2004) but has rarely been tested on a scale that is relevant for  
22 decision making and conservation (Schmidt and Balakrishnan 2015, Jeliakov et al. 2016).  
23 Soundscapes and acoustic indices could enable these tests on an unprecedented scale (Roe et al.,  
24 2021).

1  
2  
3 In this study, we employed acoustic indices to assess the temporal and spatial variations  
4 of insect acoustic niches. However, the use of acoustic indices for monitoring insects can extend  
5 far beyond and might be useful to answer traditional (and new) ecological questions regarding  
6 insect activity. Currently, our knowledge about how and which drivers shape insect acoustic  
7 activity is very limited, and the use of passive acoustics and indices could clearly contribute to  
8 better understanding this topic. For example, the methodology applied might be useful to assess  
9 the relationship between climate conditions and insect acoustic activity (Scarpelli et al. 2023), as  
10 well as to assess seasonal variations on such relationships (e.g., rainfall is not expected to have  
11 the same impact during the wet and dry season in tropical forests). Likewise, it could also help to  
12 understand how other drivers, such as astronomical (e.g., moonlight), human noise, or predation  
13 pressure (e.g., birds, bats) may have an impact on the acoustic activity of insects (Scarpelli et al.  
14 2023, Belwood and Morris 1987). We are aware that acoustic indices need to be used with  
15 caution (Bradfer-Lawrence et al. 2023), but our results suggest that acoustic indices can greatly  
16 advance insect automated monitoring after carefully selecting the optimal frequency bands for  
17 certain insect groups and with a validation of low interferences in the selected frequencies  
18 (Metcalf et al. 2021).

## 41 42 **5. Conclusions**

43  
44 We have demonstrated that the use of acoustic indices can be a useful tool to accurately  
45 describe the temporal and spatial variation of insect acoustic activity at scale in the Amazon. We  
46 hope that our assessment will encourage researchers and managers to judiciously use this readily  
47 and user-friendly tool. The estimation of acoustic indices is an automated process, which can be  
48 implemented in R (or through friendly interfaces, such as Kaleidoscope Pro) with little  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 informatic knowledge, and as demonstrated here (after being validated through manual  
4  
5 inspection of the sonograms) holds the potential to shed light into the nocturnal ecology of  
6  
7 insects, an aspect of their life cycle difficult to ascertain using traditional surveys methods.  
8  
9  
10 Further research should evaluate the utility of this technique to monitor a wider range of insects,  
11  
12 including those that call in the diurnal period, a time of the day under which the utility of  
13  
14 acoustic indices for monitoring insects may be decreased owing to frequency overlap with other  
15  
16 vocally-active taxa (Scarpelli et al. 2023). Future studies could also focus on applying the  
17  
18 techniques described here to monitor insects calling in the ultrasonic spectrum, which also  
19  
20 presents many challenges (Madhusudhana et al. 2024).  
21  
22  
23  
24  
25

## 26 **Funding**

27  
28 The data collection was supported by grants from: the Ecology Center, Utah State University,  
29  
30 United States; the Rufford Foundation (24612-1), United Kingdom; a fellowship to LAN (Ciências  
31  
32 sem Fronteiras – Conselho Nacional de Desenvolvimento Científico e Tecnológico  
33  
34 203230/2015-9) from Brazil.  
35  
36  
37  
38  
39

## 40 **Acknowledgements**

41  
42 We would like to thank the people who facilitated this work such as the local field assistants Max,  
43  
44 Caçula, Cobra, and Netão. We are grateful to all communities living around the VNP and  
45  
46 especially to Beatriz Ribeiro (former director of VNP) for all her support of this research and data  
47  
48 collection. We also would like to thank Julia Gómez-Catasús for valuable statistical advice and  
49  
50 Sebastian Cadavid for help with Python. LAN acknowledges the support of biometrio.earth, CPG  
51  
52 acknowledges the support from Ministerio de Educación y Formación Profesional through the  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Beatriz Galindo Fellowship (Beatriz Galindo – Convocatoria 2020), and JBRA acknowledges the  
4 support of Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Brazil,  
5 Finance Code 001) and the support from Fundação de Amparo à Pesquisa do Estado do Amazonas  
6 (FAPEAM) through the POSGRAD Programme and EDITAL N. 001/2023 - UNIVERSAL -  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

FAPEAM 20 ANOS.

## References

- Aide, T. M., Hernández-Serna, A., Campos-Cerqueira, M., Acevedo-Charry, O., & Deichmann, J. L. (2017). Species richness (of insects) drives the use of acoustic space in the tropics. *Remote Sensing*, 9(11), 1096. <https://doi.org/10.3390/rs9111096>
- Alcocer, I., Lima, H., Sugai, L. S. M., & Llusia, D. (2022). Acoustic indices as proxies for biodiversity: a meta-analysis. *Biological Reviews*, 97(6), 2209-2236. <https://doi.org/10.1111/brv.12890>
- Allen-Ankins, S., McKnight, D. T., Nordberg, E. J., Hoefer, S., Roe, P., Watson, D. M., ... & Schwarzkopf, L. (2023). Effectiveness of acoustic indices as indicators of vertebrate biodiversity. *Ecological Indicators*, 147, 109937. <https://doi.org/10.1016/j.ecolind.2023.109937>
- Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. lme4: Linear mixed-effects models using Eigen and S4. R package version 1.1\_7. 2014. Institute for Statistics and Mathematics of WU website. Available at <http://cran.r-project.org/package=lme4>.
- Belwood, J. J., & Morris, G. K. (1987). Bat predation and its influence on calling behavior in neotropical katydids. *Science*, 238(4823), 64-67. <https://doi.org/10.1126/science.238.4823.64>

- 1  
2  
3 Bradfer-Lawrence, T., Bunnefeld, N., Gardner, N., Willis, S. G., & Dent, D. H. (2020). Rapid  
4  
5 assessment of avian species richness and abundance using acoustic indices. *Ecological*  
6  
7 *Indicators*, 115, 106400. <https://doi.org/10.1016/j.ecolind.2020.106400>  
8  
9
- 10 Bradfer-Lawrence, T., Desjonqueres, C., Eldridge, A., Johnston, A., & Metcalf, O. (2023).  
11  
12 Using acoustic indices in ecology: Guidance on study design, analyses and interpretation.  
13  
14 *Methods in Ecology and Evolution*, 14(9), 2192-2204..  
15  
16 <https://doi.org/10.1111/2041-210X.14194>  
17  
18
- 19 Brooks, T. M., Pimm, S. L., & Oyugi, J. O. (1999). Time lag between deforestation and bird  
20  
21 extinction in tropical forest fragments. *Conservation Biology*, 13(5), 1140-1150.  
22  
23 <https://doi.org/10.1046/j.1523-1739.1999.98341.x>  
24  
25
- 26 Burivalova, Z., Maeda, T. M., Rayadin, Y., Boucher, T., Choksi, P., Roe, P., ... & Game, E. T.  
27  
28 (2022). Loss of temporal structure of tropical soundscapes with intensifying land use in  
29  
30 Borneo. *Science of The Total Environment*, 852, 158268.  
31  
32 <https://doi.org/10.1016/j.scitotenv.2022.158268>  
33  
34
- 35 Burnham, K. P., & Anderson, D. R. (2004). Multimodel inference: understanding AIC and BIC  
36  
37 in model selection. *Sociological methods & research*, 33(2), 261-304.  
38  
39 <https://doi.org/10.1177/0049124104268644>  
40  
41
- 42 Buxton, R. T., McKenna, M. F., Clapp, M., Meyer, E., Stabenau, E., Angeloni, L. M., ... &  
43  
44 Wittemyer, G. (2018). Efficacy of extracting indices from large-scale acoustic recordings  
45  
46 to monitor biodiversity. *Conservation Biology*, 32(5), 1174-1184.  
47  
48 <https://doi.org/10.1111/cobi.13119>  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Campos-Cerqueira, M., & Aide, T. M. (2016). Improving distribution data of threatened species  
4 by combining acoustic monitoring and occupancy modelling. *Methods in Ecology and*  
5 *Evolution*, 7(11), 1340-1348. <https://doi.org/10.1111/2041-210X.12599>  
6  
7  
8  
9  
10 Campos-Cerqueira, M., Mena, J. L., Tejeda-Gómez, V., Aguilar-Amuchastegui, N., Gutierrez,  
11 N., & Aide, T. M. (2020). How does FSC forest certification affect the acoustically active  
12 fauna in Madre de Dios, Peru?. *Remote Sensing in Ecology and Conservation*, 6(3), 274-  
13 285. <https://doi.org/10.1002/rse2.120>  
14  
15  
16  
17  
18  
19 Cardoso, P., Erwin, T. L., Borges, P. A., & New, T. R. (2011). The seven impediments in  
20 invertebrate conservation and how to overcome them. *Biological Conservation*, 144(11),  
21 2647-2655. <https://doi.org/10.1016/j.biocon.2011.07.024>  
22  
23  
24  
25  
26 Chen, Q., Zhang, X., Wen, M., Wang, Y., & Ren, B. (2019). Dialects of *Metrioptera bicolor*  
27 (Orthoptera: Tettigonioidea): Intraspecific acoustic signal divergence in a small-scale  
28 geographical distribution. *Transactions of the American Entomological Society*, 145(1),  
29 1-18. <https://doi.org/10.3157/061.145.0102>  
30  
31  
32  
33  
34  
35  
36 Claridge, M. F., & Morgan, J. C. (1993). Geographical variation in acoustic signals of the  
37 planthopper, *Nilaparvata bakeri* (Muir), in Asia: species recognition and sexual selection.  
38 *Biological journal of the Linnean Society*, 48(3), 267-281.  
39  
40  
41  
42 <https://doi.org/10.1111/j.1095-8312.1993.tb00891.x>  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60
- Couldridge, V. C., & van Staaden, M. J. (2004). Habitat-dependent transmission of male advertisement calls in bladder grasshoppers (Orthoptera; Pneumoridae). *Journal of Experimental Biology*, 207(16), 2777-2786. <https://doi.org/10.1242/jeb.01092>
- Cowie, R. H., Bouchet, P., & Fontaine, B. (2022). The sixth mass extinction: fact, fiction or speculation? *Biological Reviews*, 97(2), 640-663. <https://doi.org/10.1111/brv.12816>

- 1  
2  
3 Deichmann, J. L., Acevedo-Charry, O., Barclay, L., Burivalova, Z., Campos-Cerqueira, M.,  
4  
5 d'Horta, F., ... & Mitchell Aide, T. (2018). It's time to listen: there is much to be learned  
6  
7 from the sounds of tropical ecosystems. *Biotropica*, 50(5), 713-718.  
8  
9 <https://doi.org/10.1111/btp.12593>  
10  
11  
12 Do Nascimento, L. A. (2020). *Ecoacoustic Methods for Multi-Taxa Animal Surveys in the*  
13  
14 *Amazon* (Doctoral dissertation, Utah State University). [https://doi.org/10.26076/e98b-](https://doi.org/10.26076/e98b-f18c)  
15  
16 [f18c](https://doi.org/10.26076/e98b-f18c)  
17  
18  
19 Do Nascimento, L. A., Campos-Cerqueira, M., & Beard, K. H. (2020). Acoustic metrics predict  
20  
21 habitat type and vegetation structure in the Amazon. *Ecological Indicators*, 117, 106679.  
22  
23 <https://doi.org/10.1016/j.ecolind.2020.106679>  
24  
25  
26 Do Nascimento, L. A., Pérez-Granados, C., & Beard, K. H. (2021). Passive acoustic monitoring  
27  
28 and automatic detection of diel patterns and acoustic structure of Howler Monkey Roars.  
29  
30 *Diversity*, 13(11), 566. <https://doi.org/10.3390/d13110566>  
31  
32  
33 Duarte, M. H. L., Sousa-Lima, R. S., Young, R. J., Vasconcelos, M. F., Bittencourt, E., Scarpelli,  
34  
35 M. D. A., ... & Pieretti, N. (2021). Changes on soundscapes reveal impacts of wildfires in  
36  
37 the fauna of a Brazilian savanna. *Science of the Total Environment*, 769, 144988.  
38  
39 <https://doi.org/10.1016/j.scitotenv.2021.144988>  
40  
41  
42 Ferreira, L. M., Oliveira, E. G., Lopes, L. C., Brito, M. R., Baumgarten, J., Rodrigues, F. H., &  
43  
44 Sousa-Lima, R. S. (2018). What do insects, anurans, birds, and mammals have to say  
45  
46 about soundscape indices in a tropical savanna. *Journal of Ecoacoustics*, 2(1), 1-1.  
47  
48 <https://doi.org/10.22261/jea.pvh6yz>  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Foote, J. R., Nanni, L. K., & Schroeder, R. (2017). Seasonal patterns of nocturnal singing by  
4 ovenbirds and white-throated sparrows. *Behaviour*, 154(13-15), 1275-1295.

5  
6  
7 <https://doi.org/10.1163/1568539X-00003468>

8  
9  
10 Fox, J., & Weisberg, S. (2018). *An R companion to applied regression*. Sage publications.

11  
12 Gasc, A., Gottesman, B. L., Francomano, D., Jung, J., Durham, M., Mateljak, J., & Pijanowski,  
13 B. C. (2018). Soundscapes reveal disturbance impacts: biophonic response to wildfire in  
14 the Sonoran Desert Sky Islands. *Landscape Ecology*, 33, 1399-1415.

15  
16  
17 <https://doi.org/10.1007/s10980-018-0675-3>

18  
19  
20  
21 Gomez-Morales, D. A., & Acevedo-Charry, O. (2022). Satellite remote sensing of  
22 environmental variables can predict acoustic activity of an orthopteran assemblage. *PeerJ*,  
23 10, e13969. <https://doi.org/10.7717/peerj.13969>

24  
25  
26  
27  
28 Householder, J. E., Schöngart, J., Piedade, M. T., Junk, W. J., Ter Steege, H., Montero, J. C., ...  
29 & Wittmann, F. (2021). Modeling the ecological responses of tree species to the flood  
30 pulse of the Amazon Negro River floodplains. *Frontiers in Ecology and Evolution*, 9,  
31 628606. <https://doi.org/10.3389/fevo.2021.628606>

32  
33  
34  
35  
36  
37 Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous inference in general parametric  
38 models. *Biometrical Journal: Journal of Mathematical Methods in Biosciences*, 50(3),  
39 346-363. <https://doi.org/10.1002/bimj.200810425>

40  
41  
42  
43  
44 ICMBio. Instituto Chico Mendes de Conservação da Biodiversidade (2014). *Plano de Manejo do*  
45 *Parque Nacional do Viruá*. ICMBio, Boa Vista, Roraima.

46  
47  
48  
49 Jaeger, B. (2017). *R2glmm: computes R squared for mixed (multilevel) models*. R package  
50 version, 0.1.2. Available at <https://cran.r-project.org/web/packages/r2glmm/index.html>.

1  
2  
3 Jeliaskov, A., Bas, Y., Kerbirou, C., Julien, J. F., Penone, C., & Le Viol, I. (2016). Large-scale  
4 semi-automated acoustic monitoring allows to detect temporal decline of bush-crickets.  
5  
6

7  
8 Global Ecology and Conservation, 6, 208-218.

9  
10 <https://doi.org/10.1016/j.gecco.2016.02.008>  
11

12 Kuznetsova A, P. B. Brockhoff, R. H. B. Christensen. 2014. lmerTest: tests for random and fixed  
13 effects for linear mixed effect models (lmer objects of lme4 package). R package version  
14 2.0\_11. Available at <https://www.CRAN.R-project.org/package=lme4>.  
15  
16

17  
18 Lüdecke, D. (2020). sjPlot: Data visualization for statistics in social science. R package version,  
19 2.8.3. Available at <https://cran.r-project.org/web/packages/sjPlot/index.html>.  
20  
21

22  
23 Madhusudhana, S., Klinck, H., & Symes, L. B. (2024). Rigorous data engineering to the rescue:  
24 building a multi-species katydid detector from unbalanced, atypical training datasets.  
25  
26

27  
28 **This current special issue.**  
29

30  
31 Metcalf, O. C., Barlow, J., Devenish, C., Marsden, S., Berenguer, E., & Lees, A. C. (2021).  
32

33 Acoustic indices perform better when applied at ecologically meaningful time and  
34 frequency scales. *Methods in Ecology and Evolution*, 12(3), 421-431.  
35  
36

37  
38 <https://doi.org/10.1111/2041-210X.13521>  
39

40 Morton, E. S. (1975). Ecological sources of selection on avian sounds. *The American Naturalist*,  
41 109(965), 17-34. <https://doi.org/10.1086/282971>  
42  
43

44 Müller, J., Mitesser, O., Schaefer, H. M., Seibold, S., Busse, A., Kriegel, P., ... & Buřivalová, Z.  
45 (2023). Soundscapes and deep learning enable tracking biodiversity recovery in tropical  
46 forests. *Nature Communications*, 14(1), 6191. <https://doi.org/10.1038/s41467-023-41693->  
47  
48  
49

50  
51 [w](#)  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R<sup>2</sup> from  
4  
5 generalized linear mixed-effects models. *Methods in ecology and evolution*, 4(2), 133-  
6  
7 142. <https://doi.org/10.1111/j.2041-210x.2012.00261.x>  
8  
9
- 10 Oliveira, E. G., Ribeiro, M. C., Roe, P., & Sousa-Lima, R. S. (2021). The Caatinga Orchestra:  
11  
12 Acoustic indices track temporal changes in a seasonally dry tropical forest. *Ecological*  
13  
14 *Indicators*, 129, 107897. <https://doi.org/10.1016/j.ecolind.2021.107897>  
15  
16
- 17 Pérez-Granados, C., & Schuchmann, K. L. (2020). Illuminating the nightlife of two Neotropical  
18  
19 nightjars: vocal behavior over a year and monitoring recommendations. *Ethology*  
20  
21 *Ecology & Evolution*, 32(5), 466-480. <https://doi.org/10.1080/03949370.2020.1753117>  
22  
23
- 24 Pérez-Granados, C., & Schuchmann, K. L. (2021). Nocturnal vocal behaviour of the diurnal  
25  
26 Undulated Tinamou (*Crypturellus undulatus*) is associated with temperature and moon  
27  
28 phase. *Ibis*, 163(2), 684-694. <https://doi.org/10.1111/ibi.12909>  
29  
30
- 31 Pijanowski, B. C., Villanueva-Rivera, L. J., Dumyahn, S. L., Farina, A., Krause, B. L.,  
32  
33 Napoletano, B. M., ... & Pieretti, N. (2011). Soundscape ecology: the science of sound in  
34  
35 the landscape. *BioScience*, 61(3), 203-216. <https://doi.org/10.1525/bio.2011.61.3.6>  
36  
37
- 38 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & Team, R. C. (2019). nlme: Linear and  
39  
40 nonlinear mixed effects models. R package version, 3.1-141. Available at [https://cran.r-](https://cran.r-project.org/web/packages/nlme/index.html)  
41  
42 [project.org/web/packages/nlme/index.html](https://cran.r-project.org/web/packages/nlme/index.html)  
43  
44
- 45 R Development Core Team (2021). A language and environment for statistical computing.  
46  
47 Vienna, Austria. R Foundation for Statistical Computing. Available at [https://www.r-](https://www.r-project.org/)  
48  
49 [project.org/](https://www.r-project.org/).  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



- 1  
2  
3 Roe, P., Eichinski, P., Fuller, R. A., McDonald, P. G., Schwarzkopf, L., Towsey, M., ... &  
4  
5 Watson, D. M. (2021). The Australian acoustic observatory. *Methods in Ecology and*  
6  
7 *Evolution*, 12(10), 1802-1808. <https://doi.org/10.1111/2041-210X.13660>  
8  
9  
10 Romer, H., & Lewald, J. (1992). High-frequency sound transmission in natural habitats:  
11  
12 implications for the evolution of insect acoustic communication. *Behavioral Ecology and*  
13  
14 *Sociobiology*, 29, 437-444. <https://doi.org/10.1007/BF00170174>  
15  
16  
17 Scarpelli, M. D., Roe, P., Tucker, D., & Fuller, S. (2023). Soundscape phenology: The effect of  
18  
19 environmental and climatic factors on birds and insects in a subtropical woodland.  
20  
21 *Science of the Total Environment*, 878, 163080.  
22  
23 <https://doi.org/10.1016/j.scitotenv.2023.163080>  
24  
25  
26 Schafer R. M. (1977). *The tuning of the world*. New York: A. A. Knopf.  
27  
28 Schmidt, A. K., & Balakrishnan, R. (2015). Ecology of acoustic signalling and the problem of  
29  
30 masking interference in insects. *Journal of Comparative Physiology A*, 201, 133-142.  
31  
32 <https://doi.org/10.1007/s00359-014-0955-6>  
33  
34  
35 Stork, N. E. (2018). How many species of insects and other terrestrial arthropods are there on  
36  
37 Earth?. *Annual Review of Entomology*, 63, 31-45.  
38  
39 <https://doi.org/10.1146/annurev-ento-020117-043348>  
40  
41  
42 Sugai, L. S. M., Silva, T. S. F., Ribeiro Jr, J. W., & Llusia, D. (2019). Terrestrial passive acoustic  
43  
44 monitoring: review and perspectives. *BioScience*, 69(1), 15-25.  
45  
46 <https://doi.org/10.1093/biosci/biy147>  
47  
48  
49 Symes, L., ter Hofstede, H. M., Martinson, S. J., Geipel, I., Kernan, C. E., & Madhusudhana, S.  
50  
51 (2022). Using passive acoustic monitoring and machine learning analysis to investigate  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 katydid ecology and behavior. *The Journal of the Acoustical Society of America*, 151(4),  
4 A29-A29. <https://doi.org/10.1121/10.0010553>  
5  
6  
7  
8 Symes, L., Madhusudhana, S., Martinson, S. J., Geipel, I., ter Hofstede, H. M. (2024). Multi-  
9  
10 year soundscape recordings and automated call detection reveals minimal impact of  
11  
12 moonlight on calling activity of Neotropical forest katydids. **This current special issue.**  
13  
14 ter Hofstede, H. M., Symes, L. B., Martinson, S. J., Robillard, T., Faure, P., Madhusudhana, S.,  
15  
16 & Page, R. A. (2020). Calling songs of neotropical katydids (Orthoptera: Tettigoniidae)  
17  
18 from Panama. *Journal of orthoptera research*, 29(2), 137-201.  
19  
20  
21 <https://doi.org/10.3897/jor.29.46371>  
22  
23  
24 Tolkova, I. (2023). *Acoustic Source Separation, Contour Classification, and Trajectory*  
25  
26 *Optimization* (Doctoral dissertation, Harvard University).  
27  
28 Ulloa, J. S., Hauptert, S., Latorre, J. F., Aubin, T., & Sueur, J. (2021). scikit-maad: An  
29  
30 open-source and modular toolbox for quantitative soundscape analysis in Python.  
31  
32 *Methods in Ecology and Evolution*, 12(12), 2334-2340. [https://doi.org/10.1111/2041-](https://doi.org/10.1111/2041-210X.13711)  
33  
34 [210X.13711](https://doi.org/10.1111/2041-210X.13711)  
35  
36  
37 Van Klink, R., Bowler, D. E., Gongalsky, K. B., Swengel, A. B., Gentile, A. & Chase, J. M.  
38  
39 (2020). Meta-analysis reveals declines in terrestrial but increases in freshwater insect  
40  
41 abundances. *Science* 368(6489), 417- 420. <https://doi.org/10.1126/science.aax9931>  
42  
43  
44 van Klink, R., August, T., Bas, Y., Bodesheim, P., Bonn, A., Fossøy, F., ... & Bowler, D. E.  
45  
46 (2022). Emerging technologies revolutionise insect ecology and monitoring. *Trends in*  
47  
48 *Ecology & Evolution*, 37(10): 872-885. <https://doi.org/10.1016/j.tree.2022.06.001>  
49  
50  
51 van Klink, R., Bauer, S., Hoyer, T. T., Sheard, J. K., Do Nascimento, L. A., & Roslin, T. (2024).  
52  
53  
54 Towards a global toolkit for insect biodiversity monitoring. **This current special issue.**  
55  
56  
57  
58  
59  
60

1  
2  
3 Villanueva-Rivera, L. J., Pijanowski, B. C., Doucette, J., & Pekin, B. (2011). A primer of  
4  
5 acoustic analysis for landscape ecologists. *Landscape ecology*, 26, 1233-1246.

6  
7 <https://doi.org/10.1007/s10980-011-9636-9>  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Review Only