



Cryptic species and hybridisation in corals: challenges and opportunities for conservation and restoration

Cynthia Riginos ^{1,2,3}	0000-0002-5485-4197
Iva Popovic ^{1,3}	0000-0001-6582-4236
Zoe Meziere ^{1,3}	0000-0001-6874-8523
Vhon Oliver Garcia ¹	0000-0002-8933-452X
Ilha Byrne ^{1,3}	0000-0003-3909-2902
Samantha M Howitt ¹	0009-0002-6272-5261
Hisatake Ishida ¹	0000-0001-8401-8395
Kevin R Bairos-Novak ¹	0000-0002-0152-1452
Adriana Humanes ⁴	0000-0001-7962-2635
Hugo Scharfenstein ^{3,5}	0000-0003-3408-2350
Thomas Richards ¹	0000-0001-5945-6545
Ethan J. Briggs ¹	0000-0002-3475-5669
Vanessa Clark ¹	0000-0001-6125-9399
Chuan Lei ¹	0000-0002-5691-8545
Mariam Khan ¹	0009-0008-8530-1465
Katharine E Prata ^{1,3}	0000-0001-6679-0066

Corresponding author: Riginos

¹ School of the Environment, The University of Queensland, St Lucia, QLD, Australia

² Department of Marine Sciences – Tjärnö Marine Laboratory, University of Gothenburg, Strömstad, Sweden

³ Australian Institute of Marine Science, Townsville, QLD, Australia

⁴ School of Natural and Environmental Sciences, Newcastle University, Newcastle upon Tyne, UK

⁵ School of BioSciences, The University of Melbourne, Melbourne, VIC, Australia

Keywords

adaptation, conservation, coral, cryptic species, ecotype, gene flow, hybridization, introgression, conservation, restoration, speciation

Abstract

The conservation and management of coral reef ecosystems will benefit from accurate assessments of reef-building coral species diversity. However, the true diversity of corals may be obfuscated by cryptic yet genetically distinct groups, which are likely more pervasive than currently recognised. Here, we investigate the prevalence of cryptic coral groups and assess evidence for their permeability to gene flow (hybridisation) via a structured literature review of genomic studies. Using reproducible criteria to detect distinct genetic groups that are sympatric, we find that 68% of nominal species represented in population genomic studies show evidence for comprising partially reproductively isolated groups and that these distinct groups are often linked by gene flow. Cryptic genetic groups frequently segregate by environment, especially depth, and may differ by phenotypic characteristics including resilience to heat stress. This hidden biodiversity creates challenges for coral conservation and restoration planning that are not well appreciated, including hiding true population declines, biasing estimates for species' phenotypic breadth, overestimating the resilience of species to stressors, yielding uncertainty in evolutionary dynamics inferred from past studies, and implying that reproductive barriers may limit mating between local and translocated corals. Incorporating the expectation that coral cryptic taxa with incomplete species boundaries will frequently be encountered is critical to the long-term success of coral conservation and restoration programs. Studying these phenomena in more detail will directly benefit conservation and restoration goals. Thus, we detail recommendations for best practice and strategies for identifying cryptic taxa and hybridisation. In addition, cryptic coral taxa present an untapped resource for studying speciation which could provide rich opportunities for collaboration among coral and speciation biologists and fill key knowledge gaps relevant to conservation and restoration.

1 Introduction: Hidden dimensions of coral biodiversity

Coral reefs are highly biodiverse and productive ecosystems (Reaka-Kudla, 1997; Fisher *et al.*, 2015) that substantially contribute to human well-being (Moberg & Folke, 1999; Adey, 2000). Yet, reef-building corals are imperilled by rising temperatures and other anthropogenic stressors worldwide (Hughes *et al.*, 2017; Knowlton *et al.*, 2021; Souter *et al.*, 2021). Thus, there is great urgency to inventory coral biodiversity and to deepen knowledge of biodiversity generating processes to guide conservation and restoration actions.

For corals, it has long been recognised that morphological variation is unlikely to align well with genetic delineation of biological units (Knowlton, 1993; Oury *et al.*, 2023). Supporting this notion, Grupstra *et al.* (2024) found over 100 examples across 24 genera of genetic studies that self-reported discovering morphologically cryptic coral species. There is also evidence that many coral taxa have incomplete reproductive barriers and are connected by

occasional gene flow or hybridisation (reviewed by van Oppen & Gates, 2006; Willis *et al.*, 2006; Mao & Satoh, 2019; González, Rivera-Vicéns & Schizas, 2021; Hobbs *et al.*, 2021; Pinsky, Clark & Bos, 2023). Consistent with hybridisation, morphologically intermediate individuals between taxonomically-recognised species are often encountered in the field (Veron, 1995; Richards *et al.*, 2008; DeVantier, Turak & Szava-Kovats, 2020), and many species can be crossed under experimental conditions (Isomura *et al.*, 2016; Chan, Peplow & van Oppen, 2019; Kitanobo *et al.*, 2022). Thus, there is a growing sense that natural units of coral biodiversity may often consist of genetically distinct groups that can be connected by gene flow (hybridisation), even if these groups are not necessarily morphologically distinct. Figure 1 shows examples of such cryptic genetic groups.




Figure 1 – Examples of closely related taxa that are genetically differentiated, yet morphologically cryptic (*sensu lato*). Details can be found in the original studies: *Stylophora pistillata* (Meziere *et al.*, 2024), *Porites* sp. (Starko *et al.*, 2023), and *Agaricia lamarcki* (Prata *et al.*, 2022).

Documenting the prevalence of cryptic genetic groups among corals is an important first step towards learning how differentiated groups may be locally adapted to different environments and understanding the conditions under which their differentiation leads to speciation, fusion, or some in-between state of differentiation-with-gene flow (Abbott *et al.*, 2013; Barraclough, 2024). Scenarios where divergence between groups has been evolutionarily recent and/or accompanied by gene flow present the greatest challenge for recognising

distinct groups and are therefore underpowered to recognise mildly differentiated gene groups, much less to discern the processes contributing to their coexistence when inferences rely on few loci (Seehausen *et al.*, 2014). Thus, in this review, we focus exclusively on population *genomic* studies (supported by whole genome sequencing or reduced representation sequencing of genome-wide variation) and apply clear and reproducible criteria for recognising distinct genetic groups aligned to speciation theory and concepts. This paper builds upon important earlier reviews and syntheses of cryptic coral diversity (Richards, Berry & van Oppen, 2016; Grupstra *et al.*, 2024) and hybridisation in corals (van Oppen & Gates, 2006; Willis *et al.*, 2006; Hobbs *et al.*, 2021; Pinsky *et al.*, 2023) that primarily focused on studies with few markers (microsatellites, few sequenced loci, or allozymes).

We use the term *cryptic* to signify cases where there are distinct genetic groups among sets of colonies that were field-identified as being the same species by experienced researchers (consistent with the definition of Grupstra *et al.*, 2024). Thus, this definition may include both pseudo-cryptic genetic groups and those that truly lack morphological differences (Pante *et al.*, 2015; Cahill, Meglecz & Chenuil, 2024) such that *sensu lato* would be the most appropriate epithet (Chenuil *et al.*, 2019). These genetically distinct coral groups may not have all the contingent properties of species (de Queiroz, 2005), including complete reproductive isolation (as required by the biological species concept: Mayr, 1942). (Indeed, it is possible that some distinct genetic groups represent ecotypes, or genetically distinct populations adapted to specific environments, where their speciation outcome is uncertain: Lowry, 2012.) In this review, we use the terms *genetic groups* or *genetic taxa* to emphasize the genetic coherence (Dobzhansky, 1937) and distinctiveness (following Mallet's operational definition: Mallet, 1995) of genetically defined coral groups, yet acknowledge ambiguity in whether these groups should be taxonomically defined as species (or subspecies). We avoid terms such as clade or lineage, since monophyly across all or many gene trees is unlikely when divergence is recent.

It is likely that many cryptic coral taxa inhabit the “grey zone” of the speciation continuum, where taxa are linked by continuing gene flow that may be variable in strength across the genome (Seehausen *et al.*, 2014; Roux *et al.*, 2016; De Jode *et al.*, 2023), and where taxonomy is likely to be controversial (de Queiroz, 2005; Roux *et al.*, 2016). When reproductive barriers are not complete, the outcome of interbreeding between groups is controlled by the relative strength of divergent selection (when different alleles are advantageous for different groups) and the genomic extent of gene exchange (which promotes homogenisation) (Abbott *et al.*, 2013). However, for genetically distinct groups to coexist within their dispersal range and across time, reproductive barriers of some form must be present (Coyne & Orr, 2004; Seehausen *et al.*, 2014). For corals, reproductive barriers between groups may be due to intrinsic genetic incompatibilities (Levitan *et al.*, 2004), differences in spawning times (Levitan *et al.*, 2004; Rosser, 2015), and strong divergent selection arising from microhabitat differences (Prada & Hellberg, 2013).

Here, we critically assess the prevalence of cryptic genetic taxa among corals. By corals, we refer to benthic Anthozoans including scleractinians (hard corals) as well as octocorals (soft corals and gorgonians). In contrast to previous reviews on corals, we only consider population genomic surveys that have the power to detect cryptic taxa, and we reexamine reported results against reproducible and conservative criteria for detecting cryptic taxa based on population genetic evidence for some degree of reproductive isolation (in the spirit of Mayr's 1942 biological species concept and related concepts stressing evolutionary lineages and populations: de Queiroz, 1998; see section 2.1 for more details). Specifically, we: 1) rigorously quantify the prevalence of cryptic coral taxa and hybridisation, 2) discuss the relevance of cryptic coral taxa and hybridisation for conservation and restoration, and 3) highlight new directions to develop corals as exciting model systems for speciation and adaptation studies that can bring valuable insights to conservation and restoration. The main text is supported by text boxes that (i) provide a worked example of delineating coral taxa, (ii) outline best practices for designing studies when cryptic taxa are likely to be encountered, and (iii) demonstrate how cryptic taxa are commonly overlooked in coral experiments.

2 Closely related coral taxa are common in sympatry and frequently connected by gene flow

To gauge the prevalence and impacts of cryptic coral taxa and hybridisation, we undertook a structured literature search and focus exclusively on population genomic studies of corals. Population genomic surveys have the power to detect subtly differentiated genetic groups when allele frequency differences between groups are small, as we expect for groups that are recently diverged and/or connected by gene flow (Section 2.1). Below, we estimate the frequency of cryptic genetic taxa (2.2). We describe what population genomic results have shown regarding the relationships between cryptic coral taxa and their symbiont partners (2.3) and the environment (2.4). We then investigate how often studies test for and observe hybridisation and gene flow between cryptic taxa (2.5).

2.1 Criteria to detect and delineate cryptic coral taxa

We define cryptic coral taxa as distinct groups of individuals (genotypic clusters, in line with Mallet, 1995) within nominal species that maintain their distinctiveness even when their ranges overlap, and therefore there are no physical barriers to gene exchange. Instances where distinct genetic groups are found together (i.e., sympatric within the scale of dispersal distance) provide the strongest circumstantial evidence for some degree of reproductive isolation between groups, as reduced gene flow due to restricted dispersal cannot be the primary cause of genetic divergence in these cases (Coyne & Orr, 2004; Seehausen *et al.*, 2014). When distinct genetic groups are geographically separated (i.e., allopatric), then

divergence may solely reflect physical dispersal barriers to gene exchange and therefore distinct allopatric groups are not informative for inferring reproductive isolation.

The statistical power of common methods for detecting population genetic structure is determined by the number of loci examined, as well as the extent of genetic covariance among loci, where distinct populations exhibit non-random associations of alleles at various loci across the genome (i.e., *linkage disequilibrium*) (Pritchard, Stephens & Donnelly, 2000; Novembre *et al.*, 2008). While physical linkage on a chromosome alone will cause covariance among loci, genome-wide covariance also arises as a direct consequence of population structure (reflecting distinct gene pools subject to independent outcomes of genetic drift and selection) or selection on interacting loci. Reproductively isolated taxa and geographically separated populations will be differentiated from each other across loci due to genetic drift and selection. In contrast, gene flow between taxa will reduce allele frequency differences and covariances. Statistical power for identifying small allele frequency differences and linkage disequilibrium in empirical surveys is increased by sampling many individuals (both within and between locations) and many loci, where important genomic differences between groups can be missed when few individuals or few loci are sampled. For example, two co-occurring genetic groups of *Montastraea cavernosa* were clearly delineated using thousands of loci and yet were ambiguous with 9 microsatellite loci (Sturm *et al.*, 2020). Similarly, chromosomal inversions have been found in *Acropora kentii* using whole genome sequencing (Zhang *et al.*, 2024) that were missed with single nucleotide polymorphisms from a reduced portion of the genome (Matias *et al.*, 2023).

After adequate sampling and with sufficient genotyped loci, there are two general approaches used for detecting subtle genome-wide differentiation: ordination-based analyses and model-based clustering. Ordination-based analyses, such as principal components analyses (PCA), principal coordinates analysis, and multidimensional scaling describe multidimensional relationships among entities and are based on (and/or visually represent) the genetic covariance matrix (Patterson, Price & Reich, 2006). Model-based clustering analyses – typified by admixture detection analyses such as STRUCTURE (Pritchard *et al.*, 2000), fastSTRUCTURE (Raj, Stephens & Pritchard, 2014), ADMIXTURE (Alexander, Novembre & Lange, 2009) and sNMF (Frichot *et al.*, 2014) – partition groups (K) based on associations among alleles and loci. Ordination and model-based clustering approaches are valuable for exploring relationships between individuals without pre-assigning individuals to “populations”, as is required by *F*-statistics and other population-level metrics. These methods are some of the most common and routinely employed methods in population genomic surveys, perform well at low levels of divergence (i.e. where allele sharing is prevalent), and provide complementary insights into spatial patterns of genetic diversity. We

omit results from discriminant analysis that maximise variance using user-assigned groupings (see Thia, 2022 for extended discussion).

Specifically, we propose three criteria for identifying and delineating taxa based on ordination and model-based clustering:

- 1) Distinct genetic groups occur in sympatry relative to their dispersal ability.
- 2) Ordination analyses (e.g., PCA) strongly cluster these distinct genetic groups based on genotypes of individuals and/or model-based clustering indicates that individuals belong to separate groups.
- 3) Genetic distances between sympatric individuals of provisionally different taxa are greater than the genetic distances among allopatric individuals (when allopatric individuals comprise a single putative taxon). This is evidenced by divergence between sympatric groups across lower ordination axes and/or group numbers (K values) for clustering as compared to axes or groups that describe geographic structure.

2.2 Evaluating published studies for evidence of cryptic taxa

To determine if published genomic surveys of corals typically test for and find evidence of cryptic taxa following the above criteria (2.1, and see Textbox 1 for worked example), we searched the Web of Science Core Collection database for published papers displaying graphical results from ordination (PCA, primarily) and/or model-based clustering analyses (STRUCTURE, fastSTRUCTURE, ADMIXTURE, and sNMF). We focus on studies that genotyped individual cnidarian genomes for 1000s of loci or more and sampled across two or more geographic locations (see Appendix for details of literature search). We additionally collected information on depth ranges and symbiont composition when reported. In our review of published studies, we apply our three criteria (from Section 2.1) conservatively by only looking for sympatric differentiation: i) along the first principal component axis (referred to as 'PC1') for ordination-based analyses, and ii) for model-based clustering, examining outcomes when individuals were allowed to be assigned to one of two groups (i.e. K=2). We consider individuals sampled ≤ 10 km apart as being broadly sympatric. Dispersal distances for most coral species are unknown but there is evidence of spatial population structure at scales of 10 km in brooders (Prata *et al.*, 2024), which guided our choice for sympatric distance.

The literature search uncovered 41 studies describing results for 31 species. Some studies included multiple species, and some species were genotyped multiple times in different studies, thus, our search yielded a total of 51 unique records (available as supplemental data). Although we did not restrict our search by sampling depth, none of the recovered records

included species beyond mesophotic depths (i.e., >150 m), and therefore the results that follow describe shallow water and mesophotic corals.

As shown in Fig. 2, 68% of nominal species with population genomic data showed evidence for distinct cryptic taxa (representing 23 records out of 39 that could be evaluated against all three criteria). While we might expect genetic differentiation to be greater among brooding corals that have less innate dispersal ability relative to broadcast spawning corals (Knowlton, 2001), the relative proportions of sympatric versus non-sympatric groups did not differ between brooding and broadcast spawning corals (2x2 Fisher’s exact test using either ordination or clustering, $P > 0.2$).




Figure 2 – Evidence for cryptic coral taxa is common. For many genomic studies of corals, the greatest axis of genetic differentiation defines groups that are sympatric (e.g., satisfying criteria 1-3 for cryptic taxa). Results by species are summarised as either meeting or not meeting the three criteria for cryptic genetic taxa as applied to ordination or model-based clustering results. For criterion 3, we apply the strictest definition where sympatric differentiation is aligned to the first axis (ordination) or $K = 2$ (model-based clustering). From the 51 studies examined, 39 presented results that could be evaluated against criteria 1-3. Some studies had both ordination and model-based clustering, and some species were included in more than one study; thus, multiple points can appear against each species. Of the 25 nominal species with population genomic data, 17 showed evidence for including cryptic taxa.

Criterion 3 is based on genetically distinct groups co-occurring. This evidence is stronger when these groups co-occur across many geographic locations. Thus, we investigated whether pairs of genetically distinct coral groups were repeated across multiple sites using model-based clustering results based on author-selected K values. Focusing on reported results from principal components analyses and model-based clustering allowed us to evaluate *patterns* across a broad cross-section of published studies. Ideally, the *processes* responsible for

creating such patterns should be further evaluated, where demographic inference can be used to estimate gene flow and thereby provide greater insights on reproductive isolation (see for example, Fraïsse *et al.*, 2021 and section 2.5). Across studies, it was common for cryptic groups to be sympatric at multiple sampled sites (Fig. 2). This observation strengthens the conclusion that closely related, but distinct, taxa can co-occur over extensive geographic areas and implicates some degree of reproductive isolation maintaining the distinctiveness of each group (discussed further in 2.5). In summary, cryptic taxa are common attributes in population genomic studies of corals.

Although there is evidence for cryptic taxa across many studies (Fig. 2), not all studies acknowledged the groupings within their data or partitioned their data appropriately in downstream analyses. For example, some of the studies computed summary statistics such as heterozygosity and *F*-statistics using all individuals from the sampling locations, despite evidence for genetically distinct taxa co-occurring within locations and thereby created inaccurate estimates (discussed further in 3.3). Indeed, many previously published studies that did not have the advantage of detecting cryptic taxa reliably with genomic data (e.g., inference based on microsatellites or allozymes) have likely inadvertently based conclusions on heterogeneous mixes of cryptic taxa. Additionally, analyses using few loci likely lack sufficient power to detect recently differentiated taxa, and thus underpowered studies may conclude the absence of cryptic taxa without sufficient evidence. For these reasons, many published studies – including studies published by authors of this review – may unintentionally base conclusions on heterogeneous mixes of cryptic taxa.

2.3 No clear patterns for symbionts associated with cryptic coral host taxa

An important aspect of coral biodiversity is the diverse microbial community living within the cnidarian host, where mutualistic relationships with endosymbiotic dinoflagellates from the family Symbiodiniaceae are known to affect whole organism physiology (LaJeunesse *et al.*, 2018). A central issue for coral diversification and adaptation is the extent to which symbiotic associations are flexible (see Grupstra *et al.* 2024 for an extended discussion). We found that 43% of surveyed genomic studies had genotyped dinoflagellate symbionts alongside the coral hosts. Symbiont strains were commonly found to be shared across cryptic host taxa regardless of reproductive mode (Bongaerts *et al.*, 2017; van Oppen *et al.*, 2018; Gomez-Corrales & Prada, 2020; Bongaerts *et al.*, 2021a; Prada *et al.*, 2022; Rivera *et al.*, 2022; Buitrago-Lopez *et al.*, 2023; Starko *et al.*, 2023). Yet, some symbiont strains, appeared specific to cryptic taxa (van Oppen *et al.*, 2018; Gomez-Corrales & Prada, 2020; Bongaerts *et al.*, 2021a; Buitrago-Lopez *et al.*, 2023; Starko *et al.*, 2023) so there was no single consistent pattern relating symbionts with their coral hosts, as is commonly observed across nominal species (Bongaerts *et al.*, 2015). However, the variety of methods used to characterise symbionts may contribute to inconsistent patterns.

Moving beyond single marker genotyping of symbionts may provide better resolution of host-symbiont associations (Davies *et al.*, 2023; Ishida *et al.*, 2023; Zhang *et al.*, 2023). For example, Rivera *et al.* (2022) found that symbiont identities among *Porites lobata* did not align to host taxa using internal transcribed spacer (*ITS*) genotyping, but instead were concordant with higher resolution SNP-based analyses. Among the studies examined here, many (e.g., Howells *et al.*, 2016; van Oppen *et al.*, 2018; Gomez-Corrales & Prada, 2020; Bongaerts *et al.*, 2021a; Prata *et al.*, 2022; Buitrago-Lopez *et al.*, 2023; Starko *et al.*, 2023) relied on *ITS* sequences to characterise within-colony symbiont lineages. Some studies used incidentally recorded symbiont sequences retrieved from whole-colony sequencing (either reduced representation or shotgun whole genome sequencing) to make inferences about symbionts, including reconstructing symbiont organelle diversity (Bongaerts *et al.*, 2017; Forsman *et al.*, 2017; Gonzalez-Zapata *et al.*, 2018; Cooke *et al.*, 2020; Bongaerts *et al.*, 2021a; Matias *et al.*, 2023; Zhang *et al.*, 2023) or characterising symbiont genomic diversity with *k*-mer analyses (Zhang *et al.*, 2023). Of these approaches, *k*-mer analysis is the only method that captures genome wide diversity of symbionts and therefore may reveal more nuanced patterns than *ITS* or organelle-based results (which rely on a single marker) (Ishida, Riginos & Chan, 2024).

2.4 Depth can segregate cryptic genetic taxa

For coral hosts, the existence of distinct genetic groups in sympatry implies that differentiation could be preserved by strong divergent selection arising from adaptation to local environments (Richardson *et al.*, 2014). Such divergent selection could maintain ecotypes within species and, in extreme cases, drive reproductive isolation (ecological speciation: Schluter, 2001; Rundle & Nosil, 2005). It has been hypothesised that ecological speciation in corals could be common, resulting from strong environmental gradients on reefs, especially associated with depth (González *et al.*, 2020). Indeed, depth is a predominant structuring aspect in coral reef communities (Knowlton, 1993), with distinct species turnover between shallow (approximately < 30 m) and mesophotic (approximately 30-150 m) depth zones. Importantly, these transitions exist across communities because many environmental factors covary with depth, such as light intensity and spectrum (Lesser, Slattery & Leichter, 2009), temperature (Kahng *et al.*, 2019), nutrients (Leichter, Stokes & Genovese, 2008), water flow (Muir *et al.*, 2015), as well as disturbance frequency and severity (Bongaerts & Smith, 2019), creating highly contrasted habitats often only metres apart. Thus, selective agents could be both strong and multifarious and therefore depth is likely to partition taxa into ecotypes, matching patterns of species turnover.

Among the population genomic studies we surveyed, 11 studies undertook sampling over replicated depth-associated habitat contrasts (i.e., more than one site with two depth habitats sampled within each site). Replicated differentiation by depth was found for *Agaricia fragilis* (shallow vs. mesophotic, Bongaerts *et al.*, 2017), *Agaricia lamarcki* (shallow vs. mesophotic, Prata *et al.*, 2022), *Isopora brueggemanni* (lagoon vs. slope; Thomas *et al.*, 2019), *Pocillopora damicornis* (flat vs. slope, van Oppen *et al.*, 2018) and *Montastraea cavernosa*

(shallow vs. mesophotic, Sturm *et al.*, 2022) but not for *Agaricia grahamae* (upper vs. lower mesophotic, Prata *et al.*, 2022), *Stephanocoenia intersepta* (shallow vs. mesophotic, Bongaerts *et al.*, 2017), *Acropora digitifera* (lagoon vs. slope, Thomas *et al.*, 2019), or *Agaricia undata* (shallow vs. upper mesophotic and upper vs. lower mesophotic, Gonzalez-Zapata *et al.*, 2018). Thus, differentiation by depth frequently, but not always, discriminated cryptic coral taxa as identified by our criteria (see Grupstra *et al.*, 2024 for further discussion and examples).

The structure and composition of coral-associated microbial communities also can vary along environmental gradients, including those associated with depth and disturbance (Klaus *et al.*, 2007; Bongaerts *et al.*, 2013; Howells *et al.*, 2013; Quigley *et al.*, 2022). Shifting compositions of coral-associated microbial communities can expand the environmental niche available to the coral holobiont, mediating adaptation to environmental stress. Indeed, several of the studies examined here reported greater spatial or environmental partitioning among symbionts as compared to hosts (e.g., *Astrangia poculata*: Aichelman & Barshis, 2020; *P. verrucosa*: Buitrago-Lopez *et al.*, 2023; *Stylophora pistillata*: Buitrago-Lopez *et al.*, 2023; *Platygyra daedalea*: Howells *et al.*, 2016; *Acropora tenuis*: Matias *et al.*, 2023; *A. lamarcki*: Prata *et al.*, 2022; but not so for *A. digitifera*: Zhang *et al.*, 2023). Intriguingly, Starko *et al.* (2023) demonstrated that a distinct symbiont community associated with one cryptic taxon of massive *Porites sp.* shifted following a heatwave, such that the post-heatwave composition better matched the symbiont communities living in the other two cryptic taxa. Thus, symbiont communities may shift to track local environments (Baker, 2003), although this flexibility is likely to differ among host taxa (Quigley *et al.*, 2022). How environmentally induced selection shapes genetic and phenotypic variation across biological partners of the coral holobiont remains an open question and would likely benefit from manipulative experiments.

Whereas differentiation by depth and habitat appear to be common in corals (and their associated microbes), sampling strategies for many coral genomic studies are not designed to detect genetic differentiation along these environmental variables. Among the population genomic studies examined here, 25% failed to report sampling depth (or any other relevant habitat, including our own work, e.g. Matias *et al.*, 2023). Presumably, most of the genotyped corals across the studies we reviewed were collected on SCUBA from < 30 m and from a similar depth range across all sites. Among studies that did report depth, many sampling regimes had depth confounded with geography, where each location was sampled at a single depth (Fig. 3). A minority of studies (21%) implemented a structured sampling design where the same depth was sampled at more than one location. More complete reporting on depth and other microenvironmental attributes alongside sampling study designs that replicate environmental contrasts are needed to advance our understanding of how heterogeneous environments, divergent selection, and intrinsic reproductive isolation interact to shape coral biodiversity.




Figure 3 - Summary of depth sampling schemes for studies that reported depth and sampled at more than one depth. Numbers indicate the number of distinct locations that were sampled per depth. Locations were considered distinct if the nearest locations were depth contrasts (e.g., adjacent sites sampled at 5 and 15 m were considered as two locations); otherwise, locations within 10 km and at the same depth were collapsed to a single point. Dotted lines connect locations from the same study and thick grey lines indicate the sampling range (as reported by authors). Citations are as follows: a) Thomas *et al.*, 2019; b) Bongaerts *et al.*, 2017; c) Prata *et al.*, 2022; d) Gonzalez-Zapata *et al.*, 2018; e) Bongaerts *et al.*, 2021a; f) Shilling *et al.*, 2023; g) Rippe *et al.*, 2021; h) Aichelman & Barshis, 2020; i) Drury *et al.*, 2020; j) Sturm *et al.*, 2020; k) Sturm *et al.*, 2022; l) Aurelle *et al.*, 2022; m) van Oppen *et al.*, 2018; n) Meziere *et al.*, 2024.

2.5 Gene flow links coral taxa across divergence histories

Having established the prevalence of cryptic coral taxa (2.2), and their microbial (2.3), and abiotic (2.4) associations, we re-examine coral population genomic studies for evidence of gene flow and contemporary hybridisation between cryptic taxa. Gene flow between coral taxa has long been suspected (Veron, 1995; van Oppen & Gates, 2006; Willis *et al.*, 2006), but

studies using few genetic markers often lack the resolution to appropriately investigate hybridisation in the context of recently diverged taxa, where genetic similarities can result from either shared ancestral diversity or gene flow (discussed in 2.1). Thus, by analysing thousands of genomic SNPs, genomic studies can often resolve the likelihood of divergence with gene flow versus strict divergence scenarios, identify genomic regions that have experienced high or low levels of gene flow, and find regions where introgression patterns have been shaped by selection (Taylor & Larson, 2019).




Figure 4 – Estimated divergence times of coral cryptic taxa vary from 100 thousand to millions of years but are consistently connected by gene flow, as shown by demographic inferences across a variety of methods. Histories of divergence with gene flow were supported in all tested instances and similarly heterogeneous gene flow across genomes was found in all tests (consistent with genomic islands of differentiation). Models follow conventional abbreviations: IM = isolation migration; SC = secondary contact; AM = ancient migration, PER = periodic migration, and SI = strict isolation. The only model that precludes gene flow throughout the entire divergence history is strict isolation. Models with an appended ‘a’ (e.g., ‘IMa’) signify a model with asymmetric migration, otherwise migration was modelled as symmetric. In brackets are alternative models that were tested and discarded. An asterisk in the bracket signifies that there was extensive model testing including the standard scenarios listed above. Dashes indicate information or parameters that we either not tested or not reported.

To gauge if cryptic coral taxa have been linked by gene flow over their divergence history, we focused on the few coral population genomic studies from our literature search that undertook demographic modelling of speciation histories between cryptic genetic groups. Population genetic demographic modelling involves comparing the probability of alternative historical scenarios (e.g., no gene flow versus periodic or ongoing gene flow: Gutenkunst *et al.*, 2009; Beaumont, 2010; Sousa & Hey, 2013; Fraïsse *et al.*, 2021) to resolve the relative contributions of shared ancestral polymorphisms and gene flow to shared genetic variation among taxa. Eight studies used demographic modelling to evaluate competing divergence scenarios between cryptic taxa (e.g., moments and $\delta a\delta i$ methods, representing 11 nominal species: Fig. 4). Strikingly, all 12 evaluated records found the greatest support for models involving periods of divergence with gene flow. Among the variety of divergence with gene flow scenarios supported, only one record (Cooke *et al.*, 2020) found exclusive support for the secondary contact model over other scenarios, while four records had the greatest support for divergence with continuous gene flow (isolation migration model). Other models either could not distinguish between scenarios or supported models that included periods of

isolation after initial divergence with gene flow (Fig 4). In all these examples, gene flow was evolutionarily significant, but divergence was sufficient to overcome the homogenising effects of gene flow. Because divergence is maintained, gene flow between groups cannot be occurring at high enough rates to boost census population sizes, that is, it is not ecologically significant (Waples & Gaggiotti, 2006).

The frequent rejection of secondary contact models can be used as evidence for divergence without physical barriers to gene flow (Prada & Hellberg, 2021; De Jode *et al.*, 2023), consistent with ecological speciation (Schluter, 2001; Rundle & Nosil, 2005). In practice, however, secondary contact and isolation migration models can be difficult to distinguish unless secondary contact is recent and is preceded by a long period of isolation (Roux *et al.*, 2016). Because all studies reported divergence times that predate Holocene reef configurations (< 10,000 years), the genetic groups observed today will have undoubtedly shifted range positions in response to past changes in coastal and reef configurations, and therefore, intermittent periods of isolation cannot be firmly ruled out in the models that support continuous gene flow (see similar arguments in De Jode *et al.*, 2023 for speciation histories across diverse marine taxa). Periods of isolation facilitate the evolution of intrinsic reproductive barriers between groups (Seehausen *et al.*, 2014), which can magnify genotype-by-environment associations such as differentiation by depth (Bierne *et al.*, 2011). Thus, it will be important for future investigations to experimentally measure the strength and nature of reproductive barriers among cryptic coral groups to make inferences about genes associated with local adaptation (discussed further in section 4).

An emerging observation across a diversity of metazoans is that gene flow between closely-related species is variable across genomes (Ravinet *et al.*, 2017) due to reproductive incompatibilities or local adaptations that reduce gene flow within some genomic regions (Martin & Jiggins, 2017; Ravinet *et al.*, 2017). For example, chromosomal inversions and other genome features that lower recombination are thought to assist adaptive divergence when there is ongoing gene flow (Seehausen *et al.*, 2014). However, whether these features maintain existing adaptive differentiation in corals is largely unexplored (Zhang *et al.*, 2024). For pairs of sympatric coral taxa, demographic models that included heterogeneous gene flow rates received the highest support (Fig. 4). These results provide indirect evidence for genomic regions that are resistant to gene flow in the sympatric taxa studied, despite gene flow affecting neutral parts of the genome. For example, in *S. pistillata*, more divergent taxa had a higher proportion of their genomes experiencing reduced gene flow compared to the less divergent taxa, implying that genomic islands of differentiation become wider as speciation proceeds (Meziere *et al.*, 2024). These findings are consistent with *S. pistillata* comprising morphologically similar taxa at various stages of divergence (Roux *et al.*, 2016).

Low levels of gene flow can directly contribute to adaptation via introgression (Martin & Jiggins, 2017; Barraclough, 2024), where alleles derived from a different species can introduce

adaptive traits into the receiving species (e.g. resistance to hypoxia at high altitude in humans: Huerta-Sánchez *et al.*, 2015; and winter coat colour in hares: Giska *et al.*, 2019). In corals, a genomic region of approximately 220 kb appears to contribute to increased bleaching-tolerance for one *Acropora hyacinthus* taxon relative to other cryptic *A. hyacinthus* taxon, which may have been acquired through past hybridisation with *Acropora millepora* (Rose *et al.*, 2021). Supporting this conclusion, there is evidence of historical *Acropora* range expansions coinciding with introgression events, suggesting that ecological opportunities and interspecies competition during range expansions contributed to *Acropora* diversification (Mao, Economo & Satoh, 2018). While these studies implicate a role for hybridisation in adaptive evolution, there have been no comprehensive investigations of adaptive introgression in corals to date.

If interbreeding between divergent groups is ongoing and sufficiently common, then individuals with hybrid or mixed ancestry are likely to be present. Population genomic studies often find coral colonies with possible hybrid ancestries, but only two studies included explicit tests for recent hybridisation (such as tests implemented in NewHybrids: Anderson & Thompson, 2002) that probabilistically assign individuals as putative first-generation and early backcrosses. Early generation hybrid individuals were found among *Agaricia* taxa (Prata *et al.*, 2022) but not among *S. pistillata* taxa (Meziere *et al.*, 2024). Other studies have identified likely hybrid individuals based on the proportion of assignment to different groups from model-based clustering outputs (e.g. Cooke *et al.*, 2020; Kitchen *et al.*, 2020; Bongaerts *et al.*, 2021a; Fifer *et al.*, 2021; Rippe *et al.*, 2021; Rivera *et al.*, 2022; Matias *et al.*, 2023). Using a majority background group assignment score of <0.75 for distinct sympatric genetic groups as an indication of possible recent hybridisation, we identified potential hybrid individuals for 21 of the 34 species surveyed. These mixed ancestry individuals are based on the original author preferred K groups and therefore do not necessarily reflect admixture between partially reproductively isolated groups (see Textbox 1 for further discussion). Nonetheless, the prevalence of individuals with mixed ancestry suggests that hybridisation could be ongoing for many cryptic taxa. Notably, no population genomic study to date has tested whether putatively hybrid individuals were also morphologically or phenotypically distinct from parental individuals (Veron, 1995; Richards *et al.*, 2008).

2.6 Summary: cryptic genetic taxa are common in corals

The preceding reanalysis and review of population genomic studies shows that cryptic coral taxa are common, estimated to be found in 68% of nominal species examined (Fig. 2, Section 2.2), and may be adapted to different microenvironments – especially depth (2.4). These cryptic taxa, however, are often linked by sharing symbiont strains (2.3 and 2.4) and via some gene exchange (2.5). Therefore, cryptic taxa may be distinct in terms of ecology, physiology, and evolution, but how to describe and delineate taxa is not clear, as there may not be obvious morphological characteristics to distinguish them. Even with access to genomic-scale genotyping, taxonomic resolution is affected by sampling and open to interpretation. It is

clear, however, that gross morphology assessed by humans under field conditions is unreliable for recognising closely related taxa (Fig 1). Simply put, any coral investigation that does not genotype the corals under study risks treating a heterogeneous mix of partially reproductively isolated taxa as a single species. Textbox 2 suggests some pathways forward, where collaborations between geneticists and other coral reef scientists will play a key role.

3: Hidden dimensions of coral biodiversity pose conservation and restoration challenges and opportunities

Our literature review shows that cryptic taxa are prevalent, implying that many conclusions related to biodiversity, species traits, and within taxon genetic diversity based at the morphospecies level are likely to be inaccurate. Additionally, there is substantial evidence that such cryptic taxa can be linked by evolutionarily but not ecologically significant levels of gene flow (section 2.5). Failing to recognise cryptic taxa and appropriately adjust interpretations can result in misleading conclusions about fundamentally important aspects of biodiversity measurements in corals, and thus accommodating cryptic taxa will be essential for making informed conservation and restoration decisions into the future (Table 1). Below (3.1-3.6), we discuss the most important manners in which biodiversity can be mischaracterised and its effects on conservation and restoration (see also Chenuil *et al.*, 2019 for similar discussions). We also identify ways in which cryptic taxa and hybridisation could potentially aid restoration (3.5). Figure 5 outlines some possible consequences of cryptic coral taxa to current restoration actions such as direct transplantation and coral gardening (Rinkevich, 1995) and other actions that are being actively researched for managing coral populations to improve their resilience to climate change (National Academies of Sciences, 2018; Anthony *et al.*, 2020; Hein *et al.*, 2020; Bay *et al.*, 2023).

3.1 Underestimates of species diversity & overestimates of population sizes

Biodiversity inventories typically determine counts and abundances of distinct species. Common methods, such as field surveys, are usually based on morphological identification of live organisms ('morphospecies') and often struggle to confidently identify nominal species (DeVantier *et al.*, 2020). If evolutionarily distinct taxa such as cryptic genetic groups are not appropriately recognised and delineated within recognised morphospecies using genotype datasets, then total species counts will be greatly underestimated and will bias coral population sizes, species extinction vulnerabilities, and biodiversity valuations. For example, estimates of census population sizes for present-day corals have been recently debated (Dietzel *et al.*, 2021; Muir *et al.*, 2022), as smaller population sizes would imply greater vulnerability to extinction (Dietzel *et al.*, 2021; Muir *et al.*, 2022). Conversely, some rare morphospecies may be hybrids (Richards *et al.*, 2008) and thus the number of rare species might be reduced. Phenotypic plasticity could sometimes upwardly bias species counts, but underestimation due to cryptic taxa is likely more prevalent. Regardless, management

assessments are often based on population sizes and abundances, and as such, their predictability and accuracy are reduced when estimates rely on species determined solely by morphological characteristics.

Table 1: How undetected cryptic species affect biodiversity metrics and management

Measure	Actual value likely to be	Management implications of undetected, co-occurring cryptic species
Species richness	↑	<ul style="list-style-type: none"> • Number of species under management is underestimated • Management priorities based on current estimates do not reflect evolutionarily distinct groups
Species range	↓	<ul style="list-style-type: none"> • Geographic distributions could be overestimated • Endemic species may not be recognised • Underestimated risk of extinction for cryptic endemic species
Species niche breadth & phenotypic diversity	↓	<ul style="list-style-type: none"> • Perceived generalist species may instead be comprised of multiple specialist species • Translocated cryptic specialist species may fail to establish in new locations due to phenotype-environment mismatch • Species' tolerances to environmental conditions, including temperature, likely overestimated • Extinction risks underestimated
Within-population genetic diversity	↓	<ul style="list-style-type: none"> • In locations where cryptic species co-occur, genetic diversity will be overestimated • Adaptation potential may be overestimated
Inbreeding	↓	<ul style="list-style-type: none"> • In locations where cryptic species co-occur, inbreeding will be overestimated • Baseline estimates of natural inbreeding will be inaccurate
Population structure (e.g., F_{ST})	↑↓	<ul style="list-style-type: none"> • In locations where cryptic species co-occur, population structure measures will be inaccurate • Gene flow (and dispersal) will either be over- or underestimated depending on proportions of cryptic species

3.2 Overestimates of species ranges, niche breadths & generalist phenotypes

Trait diversity assessments will be compromised if trait measurement occurs on an amalgamation of distinct taxa. The combined group is likely to contain more variation than the distinct groups, which will inflate estimates of both genetic and phenotypic variance for natural populations. This phenomenon can affect data interpretation for a diverse array of coral traits. For example, species range estimates would be upwardly biased if cryptic species are geographically restricted within the broader range of a presumed 'species' (as defined by morphology). Environmental niche breadth can also be overestimated: for instance, many coral species that were previously considered depth generalists could resolve into taxa with more restricted depth distributions when cryptic species are considered (e.g., Bongaerts *et al.*, 2021a). Similarly, a presumed species may appear to have a generalist phenotype but actually comprise multiple taxa that are more specialised, as appears to be the case for bleaching responses among *Orbicella faveolata* taxa (Gomez-Corrales & Prada, 2020) and thermal tolerance traits among *A. hyacinthus* taxa (Naugle *et al.*, 2024). This means that cryptic taxa may not be ecologically or functionally equivalent. Thus, restoration actions that

involve coral outplanting (i.e., fragments or sexually propagated colonies) risk mismatching the source taxon's niche with their new destination's environmental conditions, potentially compromising the growth and survival of the outplants (Edwards *et al.*, 2010; Shaver *et al.*, 2020) (Fig. 5). Erroneous estimates of species niches and phenotypes would similarly undermine the accuracy of species distribution models that might be used in planning to predict future locations with suitable environments under climate change. More studies are urgently needed to test how often sets of cryptic coral taxa differ in their phenotypes, including their preferred niches. Current evidence suggests that conclusions drawn from incorrectly identified taxa are likely to be overly optimistic in terms of species ranges and niche breadth, and therefore will likely overestimate coral resilience to environmental change.

To get a sense of how often unrecognised cryptic coral taxa could affect inferences from other fields in coral biology, we investigated whether thermal biology studies are alert to cryptic coral taxa (Textbox 3). If cryptic coral taxa consistently differ in their phenotypes, including response to thermal stress, then experimental outcomes need to be evaluated by taxon. Yet, we find that only 8% of such studies included genotyping that could identify cryptic taxa, suggesting that many studies could be inadvertently evaluating multiple cryptic taxa as single species and thereby producing incorrect or biased conclusions. Although it is outside the scope of this paper to examine the thousands of papers in coral biology that examine phenotypes (including niches and geographic distributions), it is likely that most experimental and ecological studies have not genotyped colonies nor applied any criteria to detect cryptic taxa (such as those presented in section 2.1). Thus, inferences describing species attributes from many studies need to be viewed with scepticism, especially if policy and management actions are based on the conclusions of these studies. (We recognise that incorporating genotyping into experimental designs increases costs and efforts, nonetheless, coral biologists should design their investigations with the assumption that cryptic taxa may be present and consider how interpretation may change if their experimental subjects are a mix of taxa – see Textbox 2 for discussion and suggestions.)

3.3 Inaccurate estimates of gene flow and within-species genetic diversity

A major consequence for ignoring cryptic genetic groupings is that common measures of population genetic variation and gene flow are likely to be biased and incorrect when discrete taxa are not accounted for within the analysis (Pante *et al.*, 2015). Lumping genetically distinct groups will inflate apparent within-population diversity based on measures of allelic diversity or expected heterozygosity and overestimate inbreeding via the Wahlund effect (as discussed in Schmidt, Thia & Hoffmann, 2023). When comparing across geographic locations, measures of population structure (notably F_{ST}) can be biased either upwards or downwards, depending on the mix of cryptic taxa sampled (Pante *et al.*, 2015). These phenomena are neatly illustrated and discussed by Sheets *et al.* (2018) for *A. hyacinthus* in the western Pacific (see also Warner, van Oppen & Willis, 2015). Similarly, common summary statistics such as genetic

diversity, differentiation, inbreeding, and effective population size can be biased when introgression between differentiated taxa is not considered (Hoban *et al.*, 2022). For example, introgression may inflate the measured diversity for populations that include individuals of mixed ancestry, while population structure (F_{ST} and other similar measures that use both between and within population variation) can be biased in either direction depending on admixture proportions in populations being compared. In summary, cryptic taxa are likely to greatly affect the accuracy of studies aiming to assess and monitor genetic diversity.




Figure 5 - How cryptic coral taxa and hybridisation could affect coral reef restoration actions that aim to preserve biological diversity, counter population declines, and/or promote resilience to climate change through biological adaptation. Terminologies follow (van Oppen *et al.*, 2015; National Academies of Sciences, 2018; Hein *et al.*, 2020; Bay *et al.*, 2023) and are not mutually exclusive. For example, assisted movement could be undertaken on fragments (direct transplantation), larvae, or via the progeny of captive, sexually propagated corals. In all graphics, the coral colony shown as a larger size indicates greater fitness (e.g., survival, reproductive output, thermal tolerance) leading to competitive dominance. Different shades of grey lines are used to aid visualising connections but do not convey any specific meaning. Within images, blue coral silhouettes indicate translocated colonies whereas green silhouettes indicate local colonies.

3.4 Diminished reproductive success due to species isolating barriers

The existence of cryptic taxa implies that many coral groups sit somewhere along the speciation continuum with some degree of reproductive isolation between taxa. Intrinsic reproductive incompatibilities would, therefore, be problematic for spawning corals in captivity, as between-taxa crosses could be blocked when potential parents include more than one taxon. In pooled spawning with multiple parents, reproductive blocks between some sets of parents would increase variance in reproductive success among parents and thereby diminish genetic diversity among resultant offspring.

For restoration actions that involve outplanting (regardless of whether outplants are produced sexually or by clonal propagation), intrinsic reproductive incompatibilities between cryptic taxa could be detrimental or beneficial to restoration, depending on the precise goal. For example, a restoration goal may be for outplanted corals (chosen for specific trait attributes such as thermal tolerance) to interbreed with local corals and thus spread advantageous alleles. However, incompatibilities could prevent outplanted colonies from being able to interbreed with colonies in receiving populations or to produce viable and fertile offspring (compounding any effects of phenotype-environment mismatches as described in 3.2). Alternatively, maintaining partial genetic isolation between outplanted and receiving populations of corals could be considered advantageous if the outplanted corals contain co-adapted gene complexes that enhance their fitness or the fitness of F1 hybrids. In this case, diminished effective gene flow – whether genome-wide or associated with inversions that contain co-adapted loci – would protect beneficial combinations of alleles from being separated by recombination (Barton, 2024), thereby allowing the beneficial outplanted gene combinations to persist among the descendants of outplanted corals. Similarly, restoration actions involving manipulated genes – should that technology become feasible for restoration (Cleves, 2022) – would also be affected by reproductive incompatibilities. On the one hand, reproductive isolation between outplants and receiving corals could be viewed as reducing the impact of a restoration action by limiting spread of manipulated genes in the wild, but on the other hand, a natural block to spreading could potentially reduce risks of unconstrained modified gene release in the wild and avoid the spatial dilution of enhanced corals. Thus, it will be important to clarify restoration goals and then consider how interactions between cryptic taxa could modify intended outcomes.

3.5 Evolutionary consequences arising from hybridisation

Because first generation (F1) hybrids often have higher relative fitness (Mackay *et al.*, 2021), cross-breeding species has been proposed as a possible coral restoration strategy (National Academies of Sciences, 2018; Bay *et al.*, 2023). Towards this end, nominal species of *Acropora* have been crossed and the performance of hybrids evaluated against parental species under laboratory (Chan *et al.*, 2018; Chan *et al.*, 2019) and field conditions (Willis *et al.*, 2006; Fogarty, 2012; Lamb *et al.*, 2024). Encouragingly, none of the experiments indicated lower fitness of hybrids, however, hybrids were also not universally superior to parentals

across various fitness-related traits. Experiments based on fragmenting adult colonies morphologically identified as parental and F1 also found no survival difference between groups but did find that putative F1 fragments grew faster (VanWynen *et al.*, 2021). If first-generation hybrids have higher fitness than parental species under field conditions, then their enhanced performance could be viewed as a restoration boon (better surviving coral colonies) or a liability (hybrids outcompete parental species) depending on restoration goals (mirroring considerations under 3.4).

However, it is well-known that backcrosses and all of the possible variations of later generation hybrids can yield much greater phenotypic variation than their parental species (i.e., transgressive segregation, Abbott *et al.*, 2013; Mackay *et al.*, 2021) and recent simulations indicate that such hybrid populations can rapidly adapt to novel environments not previously experienced by either parental species (Kulmuni, Wiley & Otto, 2024). Thus, coral populations that have experienced past hybridisation may be best suited for future rapid adaptation. Additionally, there may be opportunities to develop restoration strategies using naturally occurring hybrids. For example, locating early and late-generation hybrid colonies in wild populations and characterizing their spatial and environmental distributions could potentially help source coral colonies with heat-resistant or other adaptive phenotypes.

3.6 Summary: conservation and restoration research and planning cannot afford to ignore cryptic taxa and hybridisation

Overlooking cryptic taxa can yield inaccurate conclusions about species' abundances, ranges, niches, phenotypic variance, and patterns of within-species gene flow and genetic diversity (Table 1). Conservation and restoration plans based on such erroneous estimates of species attributes are unlikely to achieve their goals (3.1, 3.2). Therefore, we urge coral biologists to acknowledge cryptic taxa as an important source of bias (Textbox 2) and ideally seek to minimise this error source in their investigations (Textbox 3). Development and implementation of coral restoration methods will similarly need to characterise cryptic taxa, where loss of genetic diversity under crossing experiments is especially concerning (3.4).

And yet, the prevalence of cryptic taxa suggests that coral biodiversity is much richer than anticipated and therefore some taxa may be inherently resilient to future conditions and/or contain sufficient genetic variation for rapid adaptation. Gene flow between taxa appears to be a common attribute of coral biology and this characteristic may provide mechanisms for adaptive traits to spread between taxa and to develop new restoration strategies capitalising upon pre-existing advanced generation hybrids (3.5).

4 Corals are untapped systems for studying adaptation and speciation in a changing world

Corals present a largely overlooked system for studying speciation and adaptation, despite extensive evidence for recent cryptic divergence. Studies of coral speciation and diversification processes can advance understanding of how coral biodiversity emerges and is maintained. Such studies would, in turn, characterise biological attributes that are important for reef conservation management decisions and strategies. Here we highlight several topics worthy of focused study.

The emerging consensus that closely related coral taxa are frequently sympatric at coarse spatial scales yet segregate by depth or other microenvironmental characteristics aligns well with models of ecological speciation (Schluter, 2001; Rundle & Nosil, 2005). Furthermore, the presence of distinct cryptic taxa in close geographic proximity suggests that selection for microhabitat matching (such as depth) may be very strong (a high selection to migration ratio: Richardson *et al.*, 2014) and/or that intrinsic reproductive barriers enhance genetic differentiation between habitats (Bierne *et al.*, 2011). To what extent cryptic taxa differ phenotypically or in terms of competitive ability is unknown, although differences in bleaching susceptibility among some cryptic taxa suggest differing vulnerabilities to climate change (Gomez-Corrales & Prada, 2020; Rose *et al.*, 2021; Rivera *et al.*, 2022).

It is likely that intrinsic barriers to reproduction limit gene flow between taxa to some extent. Analyses to date support evolutionary genomic models that allow genomic regions to differ in permeability to gene flow (see section 2.5 and Fig. 4), which may be consistent with chromosomal inversions or other genome features contributing to reproductive isolation. More studies that use chromosomal resolution genotyping will be critical to forming a deeper understanding of how species boundaries are maintained in corals (e.g., Leitwein *et al.*, 2020) and can guide decisions on assisted migration or choosing broodstock for selective breeding (Fig. 5).

Although the relative importance of extrinsic (including environmental) and intrinsic barriers to reproduction are undetermined, as sessile organisms, corals are well-suited to manipulative experiments. Experimental designs based on common gardens and reciprocal translocations can provide some of the strongest tests of local adaptation and therefore provide evidence for environmental selection (Kawecki & Ebert, 2004). Additionally, the clonal nature of corals means that genetically identical fragments from the same colony can be exposed to differing treatments, offering rich opportunities to combine experiments with genomic analyses to holistically investigate the interactions between taxon identity, phenotype, and environment (Pinsky *et al.*, 2023; Richards *et al.*, 2023).

Divergence dates between cryptic taxa often pre-date Holocene reef configurations (Fig. 4), implying that old standing genetic diversity is spread across contemporary reefs that are characterised by spatially complex yet replicated microhabitats and environmental gradients. Thus, corals are ideal for investigations that explore the genetic mechanisms of parallel divergence, especially over depth gradients (e.g. analogous to fishes that have spread into post-glacial lakes: Rougeux, Bernatchez & Gagnaire, 2017; De-Kayne *et al.*, 2022). Such investigations would also provide insights on the geographic distribution of standing genetic variation, which may be under increasing selective pressure due to pervasive anthropogenic environmental changes. For example, knowing whether geographically distant populations do or do not share alleles for advantageous traits can guide decisions regarding the utility of assisted gene flow. This is because evolutionary rescue is only worth considering if donor and recipient populations have different standing genetic diversity for ecologically functional traits.

Individual colonies with genotypes consistent with recent hybrid ancestry have been noted in the current literature (section 2.5). To date, these likely admixed individuals have primarily been documented as anecdotal observations rather than being the focus of detailed studies. However, research on hybridisation and hybrid zones offers valuable insights into speciation and adaptation, highlighting the need for more focused investigation of these individuals. (Hewitt, 1988; Harrison, 1990). Potential restoration interventions based on hybridisation rest on the supposition that hybrid corals differ in their phenotypes relative to parental species – due to some combination of hybrid vigour or transgressive segregation (section 3.5). Yet, aside from the transect studies of Prada & Hellberg (2014), no other study has mapped the spatial and environmental distributions of hybrids relative to parental taxa nor compared their phenotypes. For example, finding and characterising the phenotypes of advanced generation hybrids (not just F1's) would enable robust tests of hybrid fitness and evaluate evidence for transgressive segregation. Integrated field and genetic studies could thus be critical for advancing our understandings of coral hybrid zone dynamics (see Westram *et al.*, 2018 for a marine example).

Throughout this review, we have focused primarily on the cnidarian component of coral genomes to document evidence for cryptic species and hybridisation. However, in considering how future studies could build on these observations to better understand speciation and adaptation processes, it will also be important to integrate genetic analyses of the coral host as well as the associated symbiotic dinoflagellates and microbial communities. An exciting line of investigation would be to try to understand the co-evolutionary dynamics of hosts and symbionts in reference to environmental adaptation and speciation, where environmental heterogeneity likely exerts direct selection on the genomes of both corals and their symbionts (i.e. the coral holobiont) and indirect selection via host-symbiont genetic interactions.

5. Conclusions

In this review, we demonstrate that cryptic coral taxa are extremely common and are often connected by low levels of gene flow. Although our assessments reflect findings for shallow-water corals, we would anticipate that deep-water corals also harbour substantial cryptic diversity. Our failure to locate population genomic studies of deep-water corals indicates that more genetic studies of deep-water corals are needed. The prevalence of cryptic coral taxa among shallow-water corals means that many accepted understandings and conclusions regarding coral biology could be incorrect. In Section 3, we highlight how ignoring cryptic taxa can mislead management decisions by: biasing estimates of spatial biodiversity patterns; inflating species home ranges, trait spaces, and niches; and skewing inferences regarding intraspecific population structure and gene flow. Thus, as a field, we are unable to confidently generalise species distributions, ranges, and phenotypes including resilience to heat stress without genotype-based analyses to adequately assess the potential for cryptic coral taxa (Textbox 3). Identifying locations with high genetic diversity that may harbour greater adaptive potential or inferring locations with high gene flow and dispersal will crucially depend on analyses that are able to detect and account for distinct taxa. Without genomic and/or experimental data on hand, a precautionary principle may be to assume that populations inhabiting distinct environments (especially depth) are likely to be evolutionarily distinct and ecologically independent.

Although observations of cryptic coral taxa are frequent, our collective knowledge regarding the evolutionary dynamics that enable closely related taxa with incomplete species boundaries to persist in sympatry remains limited. There is a vast potential to unite coral studies with the insights and approaches from studying speciation and adaptation from other fields and organisms (section 4). For example, partnerships between coral ecologists, physiologists, and population geneticists may bridge insights into microevolutionary responses to climate change, while collaborations with experts from other fields may broker novel analyses and genomic-based approaches to better understandings of speciation and hybridisation in corals.

The reality, however, is that the future for corals and coral reefs is perilous (Knowlton *et al.*, 2021). Management decisions cannot wait for perfect information. An important next step will be for evolutionary biologists to investigate how conservation and management actions can best proceed with a renewed expectation that coral species boundaries are unlikely to be well defined – a conservation challenge that ultimately afflicts many other taxa in addition to corals (Hey *et al.*, 2003; Roux *et al.*, 2016).

Textbox 1: Applying taxonomic delineation with reproductive isolation criteria

We propose three requirements for identifying and delineating coral taxa using genomics-informed ordination and model-based clustering approaches (Section 2.1). In the empirical example that follows, based on Prata *et al.* (2024, with methods detailed in their supplementary files), we outline how coral cryptic taxa were identified using these three criteria and highlight difficulties with their interpretation.

Colonies of the brooding coral, *Agaricia agaricites*, were sampled at four locations ~10-15 km apart along west Curaçao, and collections were further subset into three depths (5, 10 and 20 m) at each location. Genotyping using reduced representation sequencing of 335 colonies and 1,629 SNP-loci revealed distinct genetic groups co-occurring within four sampled sites. This study provides a clear example of cryptic taxa identified according to criteria 1-3. Furthermore, taxa occupied unique depth ranges (AA1 occurs predominantly at 20 m, whereas AA2 occurred at all depths sampled) suggesting divergence of taxa by habitat (Fig. T1A & C).




Figure T1 - *Agaricia agaricites* resolved into two distinct taxa and fulfil criteria 1, 2 and 3 for cryptic taxon delineation. A) PC1 resolves two sympatric groups at every sampling site. PC2 represents an example of geographic partitioning in AA2 (Site 1 vs Site 2 - 4) and therefore does not necessarily imply intrinsic reproductive isolation. B) Shows that the percentage of variation explaining PC1 (14.65%) is ~5x more than PC2 (3.01%). C) Analyses using ADMIXTURE for K=2 assigned individuals to

each group with high confidence ($q > 0.9$) and showed that AA1 and AA2 were sympatric at all sites. D) There is a significant drop in cross-validation error between $K=1$ and $K=2$, and greater log-likelihood, supporting the selection of $K = 2$. All three criteria are met in delineating AA1 versus AA2 as cryptic taxa within *A. agaricites*.

This example also illustrates one potential complication for interpreting differentiation among putative taxa – the presence of recent migrants. The second PC axis (also mirrored in ADMIXTURE results for $K=3$ and $K=4$) shows partitioning that largely aligns with geographic separation and would not be considered as delineating distinct taxa under our criterion of sympatry (criterion 1 in 2.1). If variation captured by the second PC axis (and $K=3$ and $K=4$) reflects geographic differentiation, then geographically mismatched genotypes likely reflect recent immigration (shown by dashes among $K=3$ and $K=4$ in Fig. T1C). Distinguishing migrants from distinct taxa may be especially difficult when sampling numbers are low. However, if gene flow is high (> 1 migrant per generation) and there are no barriers to reproduction, then the structure between populations is expected to dissipate over a few generations (Waples & Gaggiotti, 2006).

While this example highlights the utility of PCA and cluster-based modelling methods for identifying cryptic coral taxa, patterns shown in these analyses are not always straightforward to interpret. Additional subsetting and filtering steps are necessary to thoroughly scrutinise data for consistent patterns and reveal accurate groupings (see Supplementary in Prata *et al.*, 2024). To better understand the possible biases of both PCA and assignment methods, we refer readers to McVean (2009), Pritchard *et al.* (2000), and Puechmaille (2016). More detailed discussions on species delineations, especially for marine species, can be found in Pante *et al.* (2015). Once taxa are delineated, investigators can investigate signals of recent hybridisation (e.g., Anderson & Thompson, 2002) and test among demographic models of historical gene flow and divergence (as in, Roux *et al.*, 2016; De Jode *et al.*, 2023) to better understand the nature of population divergence (see section 2.5).

Textbox 2: Best practice recommendations

Future surveys of corals should be designed with the expectation that cryptic species could be encountered. This means undertaking spatially and environmentally structured and replicated sampling, reassessing field collection protocols, and testing for cryptic taxa as part of bioinformatic and population genetic analyses. Ensuring that all data and metadata are thoroughly documented ensures future investigations can re-examine published data as novel analyses emerge, thus improving the re-use and re-purposing of genetic data.

Spatial sampling at the colony level

The best evidence for discriminating cryptic taxa from population structure is when distinct taxa are observed in sympatry (or close geographic proximity) at multiple locations (Section 2.1). We acknowledge that there is an element of chance in co-sampling distinct taxa. Given that depth appears to be the most common axis of differentiation, however, researchers who are planning to sample across depth should ensure that the same depths are sampled at multiple distinct locations to enable the detection of repeated co-occurrences of distinct taxa.

Alongside structured sampling, investigators would greatly enhance their data's re-useability (and scope for future inference) by transitioning from a population sampling mindset to focusing on individual sampling and seeking to capture as much environmental context as possible at the colony level. For instance, "cryptic" species may in fact be morphologically distinguishable based on subtle characteristics (for example, *S. pistillata*, Meziere *et al.*, 2024) and therefore, could be considered "pseudo-cryptic". Taking comprehensive photographs that can be examined later (see, for example, Protocol for Coral Collection & Curation by Project Phoenix: <https://coralprojectphoenix.org/resources/#protocols>) may allow diagnostic characters to be identified post hoc. An exciting avenue for new research could be to use photographs of genotyped cryptic taxa as training datasets for machine learning approaches to rapidly identify subtle differences in their micro-morphological characters (as expert taxonomists do already) to enable non-experts to identify pseudo-cryptic taxa in the field. For a subset of samples, it would be useful to retain larger colony fragments that would be suitable as museum voucher specimens (if permits allow) and maintain genetic vouchers for future studies. Recording each colony's geolocation and depth can greatly support analyses based on depth (i.e. as a continuous rather than categorical predictor) and space, which simultaneously could provide insights on the microhabitat attributes of cryptic taxa and hybrid individuals (as in Prada & Hellberg, 2014). A particularly exciting technology that can greatly advance this colony-focused perspective is photogrammetry (Bongaerts *et al.*, 2021b). We recognise that moving the focus from coral populations to colonies will require more time, effort, and greater expense, but the insights into potential cryptic taxa and coral biology more generally will be far richer.

Datasets that link genomic genotyping with ecological context at the colony level will be immensely valuable for gaining insights into ecological and evolutionary processes relevant to conservation. To maximise this value, investigators should strive to make all facets of their data FAIR compliant (Wilkinson *et al.*, 2016), which includes linking genotypes with all recorded metadata, including metadata that might not be relevant to the original study (but that might be of value to other researchers, included with linked data files). Analytical pipelines also need to be fully reproducible by enabling consistency in bioinformatics and analytical decisions across studies such that outcomes can be confidently compared. No doubt, all this extra documentation is a substantial amount of work, and therefore should be forecasted in initial project planning. Coral biologists can take inspiration from plant population geneticists who have greatly advanced insights and impacts by sharing highly curated datasets that have been re-used to support a myriad of additional studies after their initial publication; for example, the IntraBioDiv (Meirmans *et al.*, 2011) dataset of 27 co-distributed alpine plant genotypes has supported numerous reanalyses and test cases. Additionally, the genomic (and phenotypic) datasets for lodgepole pine and spruce from the AdapTree group (<https://adaptree.forestry.ubc.ca/about/scientific-summary/>) have greatly advanced our understanding of spatial adaptive diversity in trees.

Adjusting bioinformatic pipelines and analyses

Bioinformatics and population genetic analyses also need to be sensitive to the possibility of cryptic taxa (see also Section 2.1 and Textbox 1). Missing data thresholds and other data quality filters are employed as standard practice on individuals and loci. However, the sensitivity of different missing data thresholds to test taxon assignment and hierarchy hypotheses are often omitted. The more divergent groups are, the fewer sites they will share; thus, blanket missing data thresholds on heterogeneous samples may bias outcomes especially for sites with small sample sizes. Applying different missing data filters and subsetting datasets by selecting an even representation of predetermined groups (from initial model-based clustering analyses) or isolating certain groups can help in determining if the assignment and hierarchy of groups is stable and robust to the filters selected (Pritchard *et al.*, 2000; McVean, 2009; Puechmaille, 2016). Intermediate or admixed individuals may appear as hybrid individuals, but the causes of these patterns are many, including unexplained variance due to geographic structure, under-sampled taxa, admixture with unsampled taxa, or higher levels of missing data for some individuals. Thus, we suggest formal hybrid tests be employed for clarification (e.g., NewHybrids) if datasets are suitable. Investigators should be transparent regarding how biases or decisions were handled when reporting groupings. We suggest following advice from Meirmans (2015) by always reporting multiple K values when using model-based clustering methods, as clustering analyses represent a heuristic approach that is open to interpretation for all biologically-sensible K values, even if an optimal K-value is selected by the user-defined summary statistic. Similarly, PCA results should present the percent of variation explained and include multiple axes (as there may be more than two cryptic groups and/or geographic structure within groups).

Ultimately, we hope that the guidelines presented here can be used as a framework to detect coral cryptic taxa in future population genomic investigations.

When designing experiments

While population geneticists are the primary target audience for our recommendations, any coral biologist whose data interpretations could be affected by cryptic species would do well to incorporate genotyping in their project planning or minimally keep coral tissue for future genotyping. Our research group has found that preserving tissues quickly after collection (within 30 minutes post dive with samples placed in a cool container until processing), keeping tissue in 95% EtOH and replacing the liquid with fresh EtOH several times (until liquid is clear) within the first 48 hours of preservation, and ensuring there is a large EtOH to tissue volume ratio ($\geq 10x$) are critical steps for retaining high quality DNA. Other groups have had success with DMSO (Gaither *et al.*, 2011) and ultra cold freezing (e.g., liquid nitrogen or -70 freezer) is always a good choice (although rarely feasible in field conditions). Pilot trials and consulting with geneticists who work on a particular species can point to appropriate methods. For experimental work, we propose that future studies should: 1) where possible, include larger sample sizes ($n > 30$) to screen for cryptic genetic population structure (this will ensure downstream comparisons in individual phenotypic differences are not confounded by cryptic speciation); 2) follow guidelines from 2.1 to recognise cryptic species; 3) report initial data checking methods and results (e.g., screening population structure) in publications and reports to assist the interpretation of individual- and population-level differences; and 4) clarify definitions and conventions for terms such as "cryptic species" and establish common terminology. If genotyping cannot be combined with the original study, then keeping preserved tissues and associated records will allow future genotyping. Considering and discussing how unidentified cryptic taxa might alter experimental interpretations is essential.

Textbox 3: Are coral experiments designed to detect cryptic taxa?

Overlooking cryptic taxa can bias interpretations of experimental results. To ascertain how substantial this issue might be for coral studies, we focus on experiments related to thermal tolerance as a subset of coral studies more generally. Marine heatwaves have caused extensive coral mortality events globally (Leggat *et al.*, 2019), and thus numerous coral studies have aimed to ascertain intra- and inter-specific differences in phenotypic and physiological heat stress responses using experiments (e.g., common gardens, reciprocal transplants, etc.) and natural heating events. Mounting evidence suggests that cryptic species display contrasting responses to heat stress (Gomez-Corrales & Prada, 2020; Rose *et al.*, 2021; Rivera *et al.*, 2022; Grupstra *et al.*, 2024), and so experimental results may be more accurate when considering potential cryptic taxa – identified using genomic-scale genotyping.

The Coral Research and Development Accelerator Platform (CORDAP) database (Ortiz, Humanes & Scharfenstein, 2023) represents a curated search for papers that study thermal biology of corals. We screened the database to identify records which used genome-wide data of the coral host (i.e., coral SNP data) and those that conducted either ordination or model-based clustering (as in 2.1). We evaluated the database to determine:

1. The number of studies within the database undertaking experiments that genotyped corals for multiple unlinked markers (i.e., created data that could be used for ordination or model-based clustering).
2. The proportion of these studies that performed either an ordination or model-based clustering based on individual genotypes.
3. Whether there is evidence for cryptic taxa based on applying the criteria outlined in Section 2.1.

We found that very few experimental studies genotyped coral colonies: from 562 studies, only 60 studies included any sort of host genotyping and only 17 used high-resolution genome-wide markers such as SNPs. Still fewer studies undertook either ordination or model-based clustering on their genomic data (n=8; Fig. T2). For these eight studies, it was essentially impossible to evaluate ordination and model-based clustering outputs for evidence of cryptic taxa (in line with 2.1) because the number of surveyed individuals was so low. Three studies, however, included protocols to detect or pre-select cryptic taxa (Rose *et al.*, 2017; Ruiz-Jones & Palumbi, 2017; Rose *et al.*, 2021). Given that we estimate over 50% of coral studies targeting a single species encounter cryptic taxa (2.1), it is highly likely that hundreds of experimental studies will have inadvertently sampled multiple taxa. Therefore, we would anticipate that reported variances among individuals within studies would be greater than the true variances within cryptic taxa (3.2). This could manifest as an overestimation of thermal tolerance breadth and thus may also mask differences or similarities in measured tolerance in comparative tests between morphospecies. The CORDAP database focuses on

one group of studies, but we would anticipate that similar issues arise across all coral experimental work that does not leverage genomic-level genotyping of individual colonies.




Figure T2 – Proportion of individual studies from the Coral Research and Development Accelerator Platform (CORDAP) thermal tolerance experiment database that: (1) record genotype data capable of identifying cryptic taxa via genome-wide data of hosts, and (2) used ordination-based analyses or model-based clustering analyses.

Acknowledgements

We thank M.J.H. van Oppen, K. Johannesson, R. Pears, L. DeVantier, E. Pante, and two anonymous reviewers for their comments and valuable suggestions. S. Starko generously shared photographs of *Porites* sp. Preprint version 2 of this article has been peer-reviewed and recommended by Peer Community In Evolutionary Biology (<https://doi.org/10.24072/pci.evolbiol.100783>; Pante, 2024).

Data, scripts, code, and supplementary information availability

Data and supplementary information are available, linked from EcoEvoRxiv (<https://ecoevorxiv.org/repository/view/6714/>).

Conflict of interest disclosure

The authors declare that they have no financial conflicts of interest in relation to the content of the article.

Funding

This work was supported by the Reef Restoration and Adaptation Program, funded by the partnership between the Australian Government's Reef Trust and the Great Barrier Reef Foundation. CR was supported by a guest professorship ("Marks professur") funded by Göteborgs Universitet during part of the manuscript preparation.

References

- ABBOTT, R. J., ALBACH, D., ANSELL, S., ARNTZEN, J. W., BAIRD, S. J. E., BIERNE, N., BOUGHMAN, J., BRELSFORD, A., BUERKLE, C. A., BUGGS, R., BUTLIN, R. K., DIECKMANN, U., EROUKHMANOFF, F., GRILL, A., CAHAN, S. H., HERMANSEN, J. S., HEWITT, G. M., HUDSON, A. G., JIGGINS, C., JONES, J., KELLER, B., MARCZEWSKI, T., MALLETT, J., MARTINEZ-RODRIGUEZ, P., MÖST, M., MULLEN, S., NICHOLS, R., NOLTE, A. W., PARISOD, C., PFENNIG, K., RICE, A. M., RITCHIE, M. G., SEIFERT, B., SMADJA, C. M., STELKENS, R., SZYMURA, J. M., VÄINÖLÄ, R., WOLF, J. B. W. & ZINNER, D. (2013). Hybridization and speciation. *Journal of Evolutionary Biology* **26**(2), 229-246. 10.1111/j.1420-9101.2012.02599.x
- ADEY, W. H. (2000). Coral reef ecosystems and human health: biodiversity counts! *Ecosystem health* **6**(4), 227-236. 10.1046/j.1526-0992.2000.006004227.x
- AICHELMAN, H. E. & BARSHIS, D. J. (2020). Adaptive divergence, neutral panmixia, and algal symbiont population structure in the temperate coral *Astrangia poculata* along the Mid-Atlantic United States. *PeerJ* **8**, e10201. 10.7717/peerj.10201
- ALEXANDER, D. H., NOVEMBRE, J. & LANGE, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome research* **19**(9), 1655-1664. 10.1101/gr.094052.109
- ANDERSON, E. C. & THOMPSON, E. A. (2002). A model-based method for identifying species hybrids using multilocus genetic data. *Genetics* **160**, 1217-1229. 10.1093/genetics/160.3.1217

- ANTHONY, K. R. N., HELMSTEDT, K. J., BAY, L. K., FIDELMAN, P., HUSSEY, K. E., LUNDGREN, P., MEAD, D., MCLEOD, I. M., MUMBY, P. J., NEWLANDS, M., SCHAFFELKE, B., WILSON, K. A. & HARDISTY, P. E. (2020). Interventions to help coral reefs under global change-A complex decision challenge. *PLoS ONE* **15**(8), e0236399. 10.1371/journal.pone.0236399
- AURELLE, D., PRATLONG, M., OURY, N., HAGUENAUER, A., GELIN, P., MAGALON, H., ADJEROUD, M., ROMANS, P., VIDAL-DUPIOL, J., CLAEREBOUDT, M., NOUS, C., REYNES, L., TOULZA, E., BONHOMME, F., MITTA, G. & PONTAROTTI, P. (2022). Species and population genomic differentiation in *Pocillopora* corals (Cnidaria, Hexacorallia). *Genetica* **150**(5), 247-262. 10.1007/s10709-022-00165-7
- BAKER, A. C. (2003). Flexibility and specificity in coral-algal symbiosis: diversity, ecology, and biogeography of Symbiodinium. *Annual Review of Ecology, Evolution, and Systematics* **34**(1), 661-689. 10.1146/annurev.ecolsys.34.011802.132417
- BARRACLOUGH, T. G. (2024). Does selection favour the maintenance of porous species boundaries? *Journal of Evolutionary Biology*. 10.1093/jeb/voae030
- BARTON, N. (2024). Limits to species' range: the tension between local and global adaptation. *Journal of Evolutionary Biology* **37**(6), 605-615. 10.1093/jeb/voae052
- BAY, L. K., ORTIZ, J. C., HUMANES, A., RIGINOS, C., BAUMS, I. B., SCHARFENSTEIN, H., ARANDA, M., PEIXOTO, R., NIEHAUS, A. C. & LE PORT, A. (2023). R&D technology roadmap for understanding natural adaptation and assisted evolution of corals to climate change.
- BEAUMONT, M. A. (2010). Approximate Bayesian Computation in evolution and ecology. *Annual Review of Ecology, Evolution, and Systematics* **41**(1), 379-406. 10.1146/annurev-ecolsys-102209-144621
- BIERNE, N., WELCH, J., LOIRE, E., BONHOMME, F. & DAVID, P. (2011). The coupling hypothesis: why genome scans may fail to map local adaptation genes. *Molecular Ecology* **20**(10), 2044-2072. 10.1111/j.1365-294X.2011.05080.x
- BONGAERTS, P., CARMICHAEL, M., HAY, K. B., TONK, L., FRADE, P. R. & HOEGH-GULDBERG, O. (2015). Prevalent endosymbiont zonation shapes the depth distributions of scleractinian coral species. *Royal Society Open Science* **2**(2), 140297. 10.1098/rsos.140297
- BONGAERTS, P., COOKE, I. R., YING, H., WELS, D., DEN HAAN, S., HERNANDEZ-AGREDA, A., BRUNNER, C. A., DOVE, S., ENGLEBERT, N., EYAL, G., FORET, S., GRINBLAT, M., HAY, K. B., HARI, S., HAYWARD, D. C., LIN, Y., MIHALJEVIC, M., MOYA, A., MUIR, P., SINNIGER, F., SMALLHORN-WEST, P., TORDA, G., RAGAN, M. A., VAN OPPEN, M. J. H. & HOEGH-GULDBERG, O. (2021a). Morphological stasis masks ecologically divergent coral species on tropical reefs. *Current Biology* **31**(11), 2286-2298 e8. 10.1016/j.cub.2021.03.028
- BONGAERTS, P., DUBÉ, C. E., PRATA, K. E., GIJSBERS, J. C., ACHLATHIS, M. & HERNANDEZ-AGREDA, A. (2021b). Reefscape genomics: Leveraging advances in 3D imaging to assess fine-scale patterns of genomic variation on coral reefs. *Frontiers in Marine Science* **8**, 638979. 10.3389/fmars.2021.638979
- BONGAERTS, P., FRADE, P. R., OGIER, J. J., HAY, K. B., VAN BLEIJSWIJK, J., ENGLEBERT, N., VERMEIJ, M. J. A., BAK, R. P. M., VISSER, P. M. & HOEGH-GULDBERG, O. (2013). Sharing the slope: depth partitioning of agariciid corals and associated *Symbiodinium* across shallow and mesophotic habitats (2-60 m) on a Caribbean reef. *BMC Evolutionary Biology* **13**(1), 1-1. 10.1186/1471-2148-13-205
- BONGAERTS, P., RIGINOS, C., BRUNNER, R., ENGLEBERT, N., SMITH, S. R. & HOEGH-GULDBERG, O. (2017). Deep reefs are not universal refuges: Reseeding potential varies among coral species. *Science Advances* **3**(2), e1602373. 10.1126/sciadv.1602373

- BONGAERTS, P. & SMITH, T. B. (2019). Beyond the “Deep Reef Refuge” hypothesis: a conceptual framework to characterize persistence at depth. In *Mesophotic coral ecosystems. Coral Reefs of the World* (Volume 12, eds Y. Loya, K. A. Puglise and T. C. L. Bridge), pp. 881-895. 10.1007/978-3-319-92735-0_45
- BUITRAGO-LOPEZ, C., CARDENAS, A., HUME, B. C. C., GOSSELIN, T., STAUBACH, F., ARANDA, M., BARSHIS, D. J., SAWALL, Y. & VOOLSTRA, C. R. (2023). Disparate population and holobiont structure of pocilloporid corals across the Red Sea gradient demonstrate species-specific evolutionary trajectories. *Molecular Ecology* **32**(9), 2151-2173. 10.1111/mec.16871
- CAHILL, A. E., MEGLECH, E. & CHENUIL, A. (2024). Scientific history, biogeography, and biological traits predict presence of cryptic or overlooked species. *Biological Reviews* **99**(2), 546-561. 10.1111/brv.13034
- CHAN, W. Y., PELOW, L. M., MENÉNDEZ, P., HOFFMANN, A. A. & VAN OPPEN, M. J. H. (2018). Interspecific hybridization may provide novel opportunities for coral reef restoration. *Frontiers in Marine Science* **5**, 161. 10.3389/fmars.2018.00160
- CHAN, W. Y., PELOW, L. M. & VAN OPPEN, M. J. H. (2019). Interspecific gamete compatibility and hybrid larval fitness in reef-building corals: Implications for coral reef restoration. *Scientific Reports* **9**(1), 4757. 10.1038/s41598-019-41190-5
- CHENUIL, A., CAHILL, A. E., DÉLÉMONTEY, N., DU SALLIANT DU LUC, E. & FANTON, H. (2019). Problems and questions posed by cryptic species. A framework to guide future studies. In *From Assessing to Conserving Biodiversity. History, Philosophy and Theory of the Life Sciences* (Volume 24, eds E. Casetta, J. Marques da Silva and D. Vecchi). SpringerOpen. 10.1007/978-3-030-10991-2_478
- CLEVES, P. A. (2022). A need for reverse genetics to study coral biology and inform conservation efforts. In *Coral Reef Conservation And Restoration In The ‘Omics’ Age*. (eds M. J. van Oppen and M. Aranda), pp. 167-178.
- COOKE, I., YING, H., FORÊT, S., BONGAERTS, P., STRUGNELL, J., SIMAKOV, O., ZHANG, J., FIELD, M. A., RODRIGUEZ-LANETTY, M., BELL, S. C., BOURNE, D. G., VAN OPPEN, M. J. H., RAGAN, M. A. & MILLER, D. J. (2020). Signatures of selection in the coral holobiont reveal complex adaptations to inshore environments driven by Holocene climate change. *Science Advances* **6**, eabc6318. 10.1101/2020.02.25.951905
- COYNE, J. A. & ORR, H. A. (2004). *Speciation*. Sinauer associates Sunderland, MA.
- DAVIES, S. W., GAMACHE, M. H., HOWE-KERR, L. I., KRIEFALL, N. G., BAKER, A. C., BANASZAK, A. T., BAY, L. K., BELLANTUONO, A. J., BHATTACHARYA, D., CHAN, C. X., CLAAR, D. C., COFFROTH, M. A., CUNNING, R., DAVY, S. K., DEL CAMPO, J., DIAZ-ALMEYDA, E. M., FROMMLET, J. C., FUESS, L. E., GONZALEZ-PECH, R. A., GOULET, T. L., HOADLEY, K. D., HOWELLS, E. J., HUME, B. C. C., KEMP, D. W., KENKEL, C. D., KITCHEN, S. A., LAJEUNESSE, T. C., LIN, S., MCILROY, S. E., MCMINDS, R., NITSCHKE, M. R., OAKLEY, C. A., PEIXOTO, R. S., PRADA, C., PUTNAM, H. M., QUIGLEY, K., REICH, H. G., REIMER, J. D., RODRIGUEZ-LANETTY, M., ROSALES, S. M., SAAD, O. S., SAMPAYO, E. M., SANTOS, S. R., SHOGUCHI, E., SMITH, E. G., STAT, M., STEPHENS, T. G., STRADER, M. E., SUGGETT, D. J., SWAIN, T. D., TRAN, C., TRAYLOR-KNOWLES, N., VOOLSTRA, C. R., WARNER, M. E., WEIS, V. M., WRIGHT, R. M., XIANG, T., YAMASHITA, H., ZIEGLER, M., CORREA, A. M. S. & PARKINSON, J. E. (2023). Building consensus around the assessment and interpretation of Symbiodiniaceae diversity. *PeerJ* **11**, e15023. 10.7717/peerj.15023
- DE JODE, A., LE MOAN, A., JOHANNESON, K., FARIA, R., STANKOWSKI, S., WESTRAM, A. M., BUTLIN, R. K., RAFAJLOVIC, M. & FRAISSE, C. (2023). Ten years of demographic modelling of divergence and speciation in the sea. *Evolutionary Applications* **16**(2), 542-559. 10.1111/eva.13428

- DE QUEIROZ, K. (1998). The general lineage concept of species, species criteria, and the process of speciation. In *Endless forms: species and speciation*. (eds D. J. Howard and S. H. Berlocher), pp. 57-75.
- DE QUEIROZ, K. (2005). Ernst Mayr and the modern concept of species. *Proceedings of the National Academy of Sciences of the United States of America* **102** Suppl 1(Suppl 1), 6600-7. 10.1073/pnas.0502030102
- DE-KAYNE, R., SELZ, O. M., MARQUES, D. A., FREI, D., SEEHAUSEN, O. & FEULNER, P. G. D. (2022). Genomic architecture of adaptive radiation and hybridization in Alpine whitefish. *Nature Communications* **13**(1), 4479. 10.1038/s41467-022-32181-8
- DEVANTIER, L., TURAK, E. & SZAVA-KOVATS, R. (2020). Species richness and abundance of reef-building corals in the Indo-West pacific: the local–regional relation revisited. *Frontiers in Marine Science* **7**. 10.3389/fmars.2020.00487
- DIETZEL, A., BODE, M., CONNOLLY, S. R. & HUGHES, T. P. (2021). The population sizes and global extinction risk of reef-building coral species at biogeographic scales. *Nature Ecology & Evolution* **5**(5), 663-669. 10.1038/s41559-021-01393-4
- DOBZHANSKY, T. (1937). Genetic nature of species differences. *The American Naturalist* **71**, 404-420.
- DRURY, C., PEREZ PORTELA, R., SERRANO, X. M., OLEKSIK, M. & BAKER, A. C. (2020). Fine-scale structure among mesophotic populations of the great star coral *Montastraea cavernosa* revealed by SNP genotyping. *Ecology and Evolution* **10**(12), 6009-6019. 10.1002/ece3.6340
- EDWARDS, A., GUEST, J., SHAFIR, S., FISK, D., GOMEZ, E., RINKEVICH, B., HEYWARD, A., OMORI, M., IWAO, K., DIZON, R., MORSE, A., BOCH, C., JOB, S., BONGIORNI, L., LEVY, G., SHAISH, L. & WELLS, S. (2010). *Reef Rehabilitation Manual: Coral Reef Targeted Research & Capacity Building for Management Program*. The Coral Reef Targeted Research & Capacity Building for Management Program, St. Lucia, Australia.
- FIFER, J. E., YASUDA, N., YAMAKITA, T., BOVE, C. B. & DAVIES, S. W. (2021). Genetic divergence and range expansion in a western North Pacific coral. *Science of The Total Environment* **813**, 152423. 10.1016/j.scitotenv.2021.152423
- FISHER, R., O'LEARY, R. A., LOW-CHOY, S., MENGERSEN, K., KNOWLTON, N., BRAINARD, R. E. & CALEY, M. J. (2015). Species richness on coral reefs and the pursuit of convergent global estimates. *Current Biology* **25**(4), 500-5. 10.1016/j.cub.2014.12.022
- FOGARTY, N. D. (2012). Caribbean acroporid coral hybrids are viable across life history stages. *Marine Ecology Progress Series* **446**, 145-159. 10.3354/meps09469
- FORSMAN, Z. H., KNAPP, I. S. S., TISTHAMMER, K., EATON, D. A. R., BELCAID, M. & TOONEN, R. J. (2017). Coral hybridization or phenotypic variation? Genomic data reveal gene flow between *Porites lobata* and *P. compressa*. *Molecular Phylogenetics and Evolution* **111**, 132-148. 10.1016/j.ympev.2017.03.023
- FRAÏSSE, C., POPOVIC, I., MAZOYER, C., ROMIGUIER, J., LOIRE, E., SIMON, A., GALTIER, N., DURET, L., BIERNE, N., VEKEMANS, X. & ROUX, C. (2021). DILS : Demographic inferences with linked selection. *Molecular Ecology Resources* **21**, 2629-2644. 10.1111/1755-0998.13323
- FRICHOT, E., MATHIEU, F., TROUILLON, T., BOUCHARD, G. & FRANÇOIS, O. (2014). Fast and efficient estimation of individual ancestry coefficients. *Genetics* **196**(4), 973-983. 10.1534/genetics.113.160572
- GAITHER, M. R., SZABÓ, Z., CREPEAU, M. W., BIRD, C. E. & TOONEN, R. J. (2011). Preservation of corals in salt-saturated DMSO buffer is superior to ethanol for PCR experiments. *Coral Reefs* **30**(2), 329-333. 10.1007/s00338-010-0687-1

- GISKA, I., FARELO, L., PIMENTA, J., SEIXAS, F. A., FERREIRA, M. S., MARQUES, J. P., MIRANDA, I., LETTY, J., JENNY, H., HACKLANDER, K., MAGNUSSEN, E. & MELO-FERREIRA, J. (2019). Introgression drives repeated evolution of winter coat color polymorphism in hares. *Proceedings of the National Academy of Sciences of the United States of America* **116**(48), 1-7. 10.1073/pnas.1910471116
- GOMEZ-CORRALES, M. & PRADA, C. (2020). Cryptic lineages respond differently to coral bleaching. *Molecular Ecology* **29**, 4265-4273. 10.1111/mec.15631
- GONZÁLEZ, A. M., PRADA, C. A., ÁVILA, V. & MEDINA, M. (2020). Ecological speciation in corals. In *Population Genomics: Marine Organisms. Population Genomics* (eds M. F. Oleksiak and O. P. Rajora), pp. 303-324. 10.1007/13836_2018_35
- GONZÁLEZ, I. C. O., RIVERA-VICÉNS, R. E. & SCHIZAS, N. V. (2021). Description of four *Millepora* spp. transcriptomes and their potential to delimit the Caribbean fire coral species. *Marine Genomics* **59**, 100863. 10.1016/j.margen.2021.100863
- GONZALEZ-ZAPATA, F. L., BONGAERTS, P., RAMÍREZ-PORTILLA, C., ADU-OPPONG, B., WALLJASPER, G., REYES, A. & SANCHEZ, J. A. (2018). Holobiont diversity in a reef-building coral over its entire depth range in the mesophotic zone. *Frontiers in Marine Science* **5**. 10.3389/fmars.2018.00029
- GRUPSTRA, C. G. B., GOMEZ-CORRALES, M., FIFER, J. E., AICHELMAN, H. E., MEYER-KAISER, K. S., PRADA, C. & DAVIES, S. W. (2024). Integrating cryptic diversity into coral evolution, symbiosis and conservation. *Nature Ecology & Evolution*. 10.1038/s41559-023-02319-y
- GUTENKUNST, R. N., HERNANDEZ, R. D., WILLIAMSON, S. H. & BUSTAMANTE, C. D. (2009). Inferring the joint demographic history of multiple populations from multidimensional SNP frequency data. *PLoS Genetics* **5**(10), e1000695. 10.1371/journal.pgen.1000695
- HARRISON, R. G. (1990). Hybrid zones: windows on evolutionary process. In *Oxford Surveys in Evolutionary Biology* **7**. (eds D. Futuyma and J. Antonovics), pp. 69–128. Oxford University Press, Oxford.
- HEIN, M. Y., MCLEOD, I. M., SHAVER, E. C., VARDI, T., PIOCH, S., BOSTRÖM-EINARSSON, L., AHMED, M. & GRIMSDITCH, G. (2020). Coral Reef Restoration as a strategy to improve ecosystem services A guide to coral restoration methods. United Nations Environment Program, Nairobi, Kenya.
- HEWITT, G. M. (1988). Hybrid zones-natural laboratories for evolutionary studies. *Trends in Ecology and Evolution* **3**, 158-167.
- HEY, J., WAPLES, R. S., ARNOLD, M. L., BUTLIN, R. K. & HARRISON, R. G. (2003). Understanding and confronting species uncertainty in biology and conservation. *Trends in Ecology and Evolution* **18**(11), 597-603. 10.1016/j.tree.2003.08.014
- HOBAN, S., ARCHER, F. I., BERTOLA, L. D., BRAGG, J. G., BREED, M. F., BRUFORD, M. W., COLEMAN, M. A., EKBLOM, R., FUNK, W. C., GRUEBER, C. E., HAND, B. K., JAFFE, R., JENSEN, E., JOHNSON, J. S., KERSHAW, F., LIGGINS, L., MACDONALD, A. J., MERGEAY, J., MILLER, J. M., MULLER-KARGER, F., O'BRIEN, D., PAZ-VINAS, I., POTTER, K. M., RAZGOUR, O., VERNESI, C. & HUNTER, M. E. (2022). Global genetic diversity status and trends: towards a suite of Essential Biodiversity Variables (EBVs) for genetic composition. *Biological Reviews* **97**(4), 1511-1538. 10.1111/brv.12852
- HOBBS, J.-P. A., RICHARDS, Z. T., POPOVIC, I., LEI, C., STAEUDLE, T. M., MONTANARI, S. R. & DIBATTISTA, J. D. (2021). Hybridisation and the evolution of coral reef biodiversity. *Coral Reefs*, 1-15. 10.1007/s00338-021-02193-9

- HOWELLS, E. J., ABREGO, D., MEYER, E., KIRK, N. L. & BURT, J. A. (2016). Host adaptation and unexpected symbiont partners enable reef-building corals to tolerate extreme temperatures. *Global Change Biology* **22**(8), 2702-14. 10.1111/gcb.13250
- HOWELLS, E. J., WILLIS, B. L., BAY, L. K. & VAN OPPEN, M. J. H. (2013). Spatial and temporal genetic structure of *Symbiodinium* populations within a common reef-building coral on the Great Barrier Reef. *Molecular Ecology* **22**(14), 3693-3708. 10.1111/mec.12342
- HUERTA-SÁNCHEZ, E., JIN, X., ASAN, BIANBA, Z., PETER, B. M., VINCKENBOSCH, N., LIANG, Y., YI, X., HE, M., SOMEL, M., NI, P., WANG, B., OU, X., HUASANG, LUOSANG, J., CUO, Z. X. P., LI, K., GAO, G., YIN, Y., WANG, W., ZHANG, X., XU, X., YANG, H., LI, Y., WANG, J., WANG, J. & NIELSEN, R. (2015). Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA. *Nature* **512**(7513), 194-197. 10.1038/nature13408
- HUGHES, T. P., KERRY, J. T., ÁLVAREZ-NORIEGA, M., ÁLVAREZ-ROMERO, J. G., ANDERSON, K. D., BAIRD, A. H., BABCOCK, R. C., BEGER, M., BELLWOOD, D. R., BERKELMANS, R., BRIDGE, T. C., BUTLER, I. R., BYRNE, M., CANTIN, N. E., COMEAU, S., CONNOLLY, S. R., CUMMING, G. S., DALTON, S. J., DIAZ-PULIDO, G., EAKIN, C. M., FIGUEIRA, W. F., GILMOUR, J. P., HARRISON, H. B., HERON, S. F., HOEY, A. S., HOBBS, J.-P. A., HOOGENBOOM, M. O., KENNEDY, E. V., KUO, C.-Y., LOUGH, J. M., LOWE, R. J., LIU, G., MCCULLOCH, M. T., MALCOLM, H. A., MCWILLIAM, M. J., PANDOLFI, J. M., PEARS, R. J., PRATCHETT, M. S., SCHOEPF, V., SIMPSON, T., SKIRVING, W. J., SOMMER, B., TORDA, G., WACHENFELD, D. R., WILLIS, B. L. & WILSON, S. K. (2017). Global warming and recurrent mass bleaching of corals. *Nature* **543**(7645), 373-377. 10.1038/nature21707
- ISHIDA, H., JOHN, U., MURRAY, S. A., BHATTACHARYA, D. & CHAN, C. X. (2023). Developing model systems for dinoflagellates in the post-genomic era. *Journal of Phycology* **59**(5), 799-808. 10.1111/jpy.13386
- ISHIDA, H., RIGINOS, C. & CHAN, C. X. (2024). Contaminant or goldmine? In silico assessment of Symbiodiniaceae community using coral hologenomes. *Frontiers in Protistology* **2**. 10.3389/frpro.2024.1376877
- ISOMURA, N., IWAO, K., MORITA, M. & FUKAMI, H. (2016). Spawning and fertility of F1 hybrids of the coral genus *Acropora* in the Indo-Pacific. *Coral Reefs* **35**(3), 851-855. 10.1007/s00338-016-1461-9
- KAHNG, S. E., AKKAYNAK, D., SHLESINGER, T., HOCHBERG, E. J., WIEDENMANN, J., TAMIR, R. & TCHERNOV, D. (2019). Light, temperature, photosynthesis, heterotrophy, and the lower depth limits of mesophotic coral ecosystems. In *Mesophotic coral ecosystems. Coral Reefs of the World* (Volume 12, eds Y. Loya, K. A. Puglise and T. C. L. Bridge), pp. 801-828. 10.1007/978-3-319-92735-0_45
- KAWECKI, T. J. & EBERT, D. (2004). Conceptual issues in local adaptation. *Ecology Letters* **7**(12), 1225-1241. 10.1111/j.1461-0248.2004.00684.x
- KITANOBO, S., IWAO, K., FUKAMI, H., ISOMURA, N. & MORITA, M. (2022). First evidence for backcrossing of F(1) hybrids in *Acropora* corals under sperm competition. *Scientific Reports* **12**(1), 5356. 10.1038/s41598-022-08989-1
- KITCHEN, S. A., VON KUSTER, G., KUNTZ, K. L. V., REICH, H. G., MILLER, W., GRIFFIN, S., FOGARTY, N. D. & BAUMS, I. B. (2020). STAGdb: a 30K SNP genotyping array and Science Gateway for *Acropora* corals and their dinoflagellate symbionts. *Scientific Reports* **10**(1), 12488. 10.1038/s41598-020-69101-z
- KLAUS, J. S., JANSE, I., HEIKOOP, J. M., SANFORD, R. A. & FOUKE, B. W. (2007). Coral microbial communities, zooxanthellae and mucus along gradients of seawater depth and coastal pollution. *Environmental Microbiology* **9**(5), 1291-1305. 10.1111/j.1462-2920.2007.01249.x

- KNOWLTON, N. (1993). Sibling species in the sea. *Annual Review of Ecology and Systematics*, 189-216.
- KNOWLTON, N. (2001). The future of coral reefs. *Proc Natl Acad Sci U S A* **98**(10), 5419-25. 10.1073/pnas.091092998
- KNOWLTON, N., GROTTOLI, A. G., KLEYPAS, J., OBURA, D., CORCORAN, E., DE GOEIJ, J. M., FELIS, T., HARDING, S., MAYFIELD, A., MILLER, M., OSUKA, K., PEIXOTO, R. S., RANDALL, C. J., VOOLSTRA, C. R., WELLS, S., WILD, C. & FERSE, S. (2021). Rebuilding coral reefs: A decadal grand challenge. *International coral reef society and future earth coasts* pp. 56. International Coral Reef Society and Future Earth Coasts
- KULMUNI, J., WILEY, B. & OTTO, S. P. (2024). On the fast track: hybrids adapt more rapidly than parental populations in a novel environment. *Evolution Letters* **8**(1), 128-136. 10.1093/evlett/grad002
- LAJEUNESSE, T. C., PARKINSON, J. E., GABRIELSON, P. W., JEONG, H. J., REIMER, J. D., VOOLSTRA, C. R. & SANTOS, S. R. (2018). Systematic revision of Symbiodiniaceae highlights the antiquity and diversity of coral endosymbionts. *Current Biology* **28**(16), 2570-2580 e6. 10.1016/j.cub.2018.07.008
- LAMB, A. M., PELOW, L. M., DUNGAN, A. M., FERGUSON, S. N., HARRISON, P. L., HUMPHREY, C. A., MCCUTCHAN, G. A., NITSCHKE, M. R. & VAN OPPEN, M. J. H. (2024). Interspecific hybridisation provides a low-risk option for increasing genetic diversity of reef-building corals. *Biology Open* **13**(9). 10.1242/bio.060482
- LEGGAT, W. P., CAMP, E. F., SUGGETT, D. J., HERON, S. F., FORDYCE, A. J., GARDNER, S., DEAKIN, L., TURNER, M., BEECHING, L. J., KUZHIUMPARAMBIL, U., EAKIN, C. M. & AINSWORTH, T. D. (2019). Rapid coral decay is associated with marine heatwave mortality events on reefs. *Current Biology* **29**(16), 2723-2730 e4. 10.1016/j.cub.2019.06.077
- LEICHTER, J. J., STOKES, M. D. & GENOVESE, S. J. (2008). Deep water macroalgal communities adjacent to the Florida Keys reef tract. *Marine Ecology Progress Series* **356**, 123-138. 10.3354/meps07230
- LEITWEIN, M., DURANTON, M., ROUGEMONT, Q., GAGNAIRE, P. A. & BERNATCHEZ, L. (2020). Using haplotype information for conservation genomics. *Trends in Ecology and Evolution* **35**(3), 245-258. 10.1016/j.tree.2019.10.012
- LESSER, M. P., SLATTERY, M. & LEICHTER, J. J. (2009). Ecology of mesophotic coral reefs. *Journal of Experimental Marine Biology and Ecology* **375**(1-2), 1-8. 10.1016/j.jembe.2009.05.009
- LEVITAN, D. R., FUKAMI, H., JARA, J., KLINE, D., MCGOVERN, T. M., MCGHEE, K. E., SWANSON, C. A. & KNOWLTON, N. (2004). Mechanisms of reproductive isolation among sympatric broadcast-spawning corals of the *Montastraea annularis* species complex. *Evolution* **58**(2), 308-323. 10.1111/j.0014-3820.2004.tb01647.x
- LOWRY, D. B. (2012). Ecotypes and the controversy over stages in the formation of new species. *Biological journal of the Linnean Society* **106**, 241-257.
- MACKAY, I. J., COCKRAM, J., HOWELL, P. & POWELL, W. (2021). Understanding the classics: the unifying concepts of transgressive segregation, inbreeding depression and heterosis and their central relevance for crop breeding. *Plant Biotechnol J* **19**(1), 26-34. 10.1111/pbi.13481
- MALLET, J. (1995). A species definition for the Modern Synthesis. *Trends in Ecology and Evolution* **10**, 294-299.
- MAO, Y., ECONOMO, E. P. & SATOH, N. (2018). The roles of introgression and climate change in the rise to dominance of *Acropora* corals. *Current Biology* **28**(21), 3373-3382.e5. 10.1016/j.cub.2018.08.061

- MAO, Y. & SATOH, N. (2019). A likely ancient genome duplication in the speciose reef-building coral genus, *Acropora*. *iScience* **13**, 20-32. 10.1016/j.isci.2019.02.001
- MARTIN, S. H. & JIGGINS, C. D. (2017). Interpreting the genomic landscape of introgression. *Current opinion in genetics & development* **47**, 69-74. 10.1016/j.gde.2017.08.007
- MATIAS, A. M. A., POPOVIC, I., THIA, J. A., COOKE, I. R., TORDA, G., LUKOSCHEK, V., BAY, L. K., KIM, S. W. & RIGINOS, C. (2023). Cryptic diversity and spatial genetic variation in the coral *Acropora tenuis* and its endosymbionts across the Great Barrier Reef. *Evolutionary Applications* **16**, 293-310. 10.1111/eva.13435
- MAYR, E. (1942). *Systematics and the Origin of Species*. Columbia University Press, New York.
- MCVEAN, G. (2009). A genealogical interpretation of principal components analysis. *PLoS Genetics* **5**(10), e1000686. 10.1371/journal.pgen.1000686
- MEIRMANS, P. G. (2015). Seven common mistakes in population genetics and how to avoid them. *Molecular Ecology* **24**(13), 3223-3231. 10.1111/mec.13243
- MEIRMANS, P. G., GOUDET, J., INTRABIODIV, C. & GAGGIOTTI, O. E. (2011). Ecology and life history affect different aspects of the population structure of 27 high-alpine plants. *Molecular Ecology* **20**(15), 3144-3155. 10.1111/j.1365-294X.2011.05164.x
- MEZIERE, Z., POPOVIC, I., PRATA, K., PANDOLFI, J. M. & RIGINOS, C. (2024). Exploring coral speciation: multiple *Stylophora pistillata* taxa on a divergence continuum on the Great Barrier Reef. *Evolutionary Applications* (17), e13644. 10.1111/eva.13644
- MOBERG, F. & FOLKE, C. (1999). Ecological goods and services of coral reef ecosystems. *Ecological economics* **29**(2), 215-233. 10.1016/S0921-8009(99)00009-9
- MUIR, P., WALLACE, C., BRIDGE, T. C. L. & BONGAERTS, P. (2015). Diverse staghorn coral fauna on the mesophotic reefs of north-east Australia. *PloS ONE* **10**(2), e0117933. 10.1371/journal.pone.0117933
- MUIR, P. R., OBURA, D. O., HOEKSEMA, B. W., SHEPPARD, C., PICHON, M. & RICHARDS, Z. T. (2022). Conclusions of low extinction risk for most species of reef-building corals are premature. *Nature Ecology & Evolution* **6**(4), 357-358. 10.1038/s41559-022-01659-5
- NATIONAL ACADEMIES OF SCIENCES, E., AND MEDICINE. (2018). A research review of interventions to increase the persistence and resilience of coral reefs. The National Academies Press, Washington, DC.
- NAUGLE, M. S., DENIS, H., MOCELLIN, V. J. L., LAFFY, P., POPOVIC, I., BAY, L. K. & HOWELLS, E. (2024). Environmental, host, and symbiont drivers of heat tolerance in a species complex of reef-building corals. *Communications Earth & Environment* **5**, 2662-4435. 10.1038/s43247-024-01649-4
- NOVEMBRE, J., JOHNSON, T., BRYC, K., KUTALIK, Z., BOYKO, A. R., AUTON, A., INDAP, A., KING, K. S., BERGMANN, S., NELSON, M. R., STEPHENS, M. & BUSTAMANTE, C. D. (2008). Genes mirror geography within Europe. *Nature* **456**(7218), 98-101. 10.1038/nature07331
- ORTIZ, J. C., HUMANES, A. & SCHARFENSTEIN, H. (2023). Natural adaptation and assisted evolution of corals to heat stress (ed. Australian Institute of Marine Science).
- OURY, N., NOEL, C., MONA, S., AURELLE, D. & MAGALON, H. (2023). From genomics to integrative species delimitation? The case study of the Indo-Pacific Pocillopora corals. *Molecular Phylogenetics and Evolution* **184**, 107803. 10.1016/j.ympev.2023.107803
- PANTE, E., PUILLANDRE, N., VIRICEL, A., ARNAUD-HAOND, S., AURELLE, D., CASTELIN, M., CHENUIL, A., DESTOMBE, C., FORCIOLI, D., VALERO, M., VIARD, F. & SAMADI, S. (2015). Species are hypotheses: avoid connectivity assessments based on pillars of sand. *Molecular Ecology* **24**(3), 525-544. 10.1111/mec.13048

- PATTERSON, N., PRICE, A. L. & REICH, D. E. (2006). Population structure and eigenanalysis. *PLoS Genetics* **2**(12), e190.
- PINSKY, M. L., CLARK, R. D. & BOS, J. T. (2023). Coral reef population genomics in an age of global change. *Annual Review of Genetics* (57), 87-115. 10.1146/annurev-genet-022123-102748
- PRADA, C. & HELLBERG, M. E. (2013). Long prereproductive selection and divergence by depth in a Caribbean candelabrum coral. *Proceedings of the National Academy of Sciences of the United States of America* **110**(10), 3961-3966. 10.1073/pnas.1208931110
- PRADA, C. & HELLBERG, M. E. (2014). Strong natural selection on juveniles maintains a narrow adult hybrid zone in a broadcast spawner. *The American Naturalist* **184**(6), 702-13. 10.1086/678403
- PRADA, C. & HELLBERG, M. E. (2021). Speciation-by-depth on coral reefs: Sympatric divergence with gene flow or cryptic transient isolation? *Journal of Evolutionary Biology* **34**, 128-137. 10.1111/jeb.13731
- PRATA, K. E., BONGAERTS, P., DWYER, J. M., ISHIDA, H., HOWITT, S. M., HERWARD, J. P., CRANDALL, E. D. & RIGINOS, C. (2024). Some reef-building corals only disperse metres per generation. *Proceedings of the Royal Society B* **291**(2027), 20231988. 10.1098/rspb.2023.1988
- PRATA, K. E., RIGINOS, C., GUTENKUNST, R. N., LATIJNHOUWERS, K., SÁNCHEZ, J. A., ENGLEBERT, N., HAY, K. & BONGAERTS, P. (2022). Deep connections: divergence histories with gene flow in mesophotic *Agaricia* corals. *Molecular Ecology* **9**, 2511-2527. 10.1111/mec.16391
- PRITCHARD, J. K., STEPHENS, M. & DONNELLY, P. (2000). Inference of population structure using multilocus genotype data. *Genetics* **155**, 945-959.
- PUECHMAILLE, S. J. (2016). The program structure does not reliably recover the correct population structure when sampling is uneven: subsampling and new estimators alleviate the problem. *Molecular Ecology Resources* **16**(3), 608-27. 10.1111/1755-0998.12512
- QUIGLEY, K. M., RAMSBY, B., LAFFY, P., HARRIS, J., MOCELLIN, V. J. & BAY, L. K. (2022). Symbioses are restructured by repeated mass coral bleaching. *Science Advances* **8**(49), eabq8349. 10.1126/sciadv.abq8349
- RAJ, A., STEPHENS, M. & PRITCHARD, J. K. (2014). fastSTRUCTURE: variational inference of population structure in large SNP data sets. *Genetics* **197**(2), 573-89. 10.1534/genetics.114.164350
- RAVINET, M., FARIA, R., BUTLIN, R. K., GALINDO, J., BIERNE, N., RAFAJLOVIC, M., NOOR, M. A. F., MEHLIG, B. & WESTRAM, A. M. (2017). Interpreting the genomic landscape of speciation: a road map for finding barriers to gene flow. *Journal of Evolutionary Biology* **30**(8), 1450-1477. 10.1111/jeb.13047
- REAKA-KUDLA, M. L. (1997). The global biodiversity of coral reefs: a comparison with rain forests. In *Biodiversity II: Understanding and Protecting Our Biological Resources*. (eds M. L. Reaka-Kudla, D. E. Wilson and E. O. Wilson), pp. 83–108. Joseph Henry Press, Washington, DC.
- RICHARDS, T. J., MCGUIGAN, K., AGUIRRE, J. D., HUMANES, A., BOZEC, Y.-M., MUMBY, P. J. & RIGINOS, C. (2023). Moving beyond heritability in the search for coral adaptive potential. *Global Change Biology* **29**(14), 3869-3882. 10.1111/gcb.16719
- RICHARDS, Z. T., BERRY, O. & VAN OPPEN, M. J. H. (2016). Cryptic genetic divergence within threatened species of *Acropora* coral from the Indian and Pacific Oceans. *Conservation Genetics* **17**(3), 577-591. 10.1007/s10592-015-0807-0

- RICHARDS, Z. T., VAN OPPEN, M. J. H., WALLACE, C. C., WILLIS, B. L. & MILLER, D. J. (2008). Some rare Indo-Pacific coral species are probable hybrids. *PLoS ONE* **3**(9), e3240. 10.1371/journal.pone.0003240.t001
- RICHARDSON, J. L., URBAN, M. C., BOLNICK, D. I. & SKELLY, D. K. (2014). Microgeographic adaptation and the spatial scale of evolution. *Trends in Ecology and Evolution* **29**(3), 165-76. 10.1016/j.tree.2014.01.002
- RINKEVICH, B. (1995). Restoration strategies for coral reefs damaged by recreational activities: The use of sexual and asexual recruits. *Restoration Ecology* **3**(4), 241-251. <https://doi.org/10.1111/j.1526-100X.1995.tb00091.x>
- RIPPE, J. P., DIXON, G., FULLER, Z. L., LIAO, Y. & MATZ, M. (2021). Environmental specialization and cryptic genetic divergence in two massive coral species from the Florida Keys Reef Tract. *Molecular Ecology* **30**, 3468–3484. 10.1111/mec.15931
- RIVERA, H. E., COHEN, A. L., THOMPSON, J. R., BAUMS, I. B., FOX, M. D. & MEYER-KAISER, K. S. (2022). Palau's warmest reefs harbor thermally tolerant corals that thrive across different habitats. *Communications Biology* **5**(1), 1394. 10.1038/s42003-022-04315-7
- ROSE, N. H., BAY, R. A., MORIKAWA, M. K. & PALUMBI, S. R. (2017). Polygenic evolution drives species divergence and climate adaptation in corals. *Evolution* **72**(1), 82-94. 10.1111/evo.13385
- ROSE, N. H., BAY, R. A., MORIKAWA, M. K., THOMAS, L., SHEETS, E. A. & PALUMBI, S. R. (2021). Genomic analysis of distinct bleaching tolerances among cryptic coral species. *Proceedings of the Royal Society B* **288**(1960), 20210678. 10.1098/rspb.2021.0678
- ROSSER, N. L. (2015). Asynchronous spawning in sympatric populations of a hard coral reveals cryptic species and ancient genetic lineages. *Molecular Ecology* **24**(19), 5006-5019.
- ROUGEUX, C., BERNATCHEZ, L. & GAGNAIRE, P. A. (2017). Modeling the multiple facets of speciation-with-gene-flow toward inferring the divergence history of lake whitefish species pairs (*Coregonus clupeaformis*). *Genome Biology and Evolution* **9**(8), 2057-2074. 10.1093/gbe/evx150
- ROUX, C., FRAÏSSE, C., ROMIGUIER, J., ANCIAUX, Y., GALTIER, N. & BIERNE, N. (2016). Shedding light on the grey zone of speciation along a continuum of genomic divergence. *PLoS Biology* **14**(12), e2000234-22. 10.1371/journal.pbio.2000234
- RUIZ-JONES, L. J. & PALUMBI, S. R. (2017). Tidal heat pulses on a reef trigger a fine-tuned transcriptional response in corals to maintain homeostasis. *Science Advances* **3**, e1601298.
- RUNDLE, H. D. & NOSIL, P. (2005). Ecological speciation. *Ecology Letters* **8**(3), 336-352. 10.1111/j.1461-0248.2004.00715.x
- SCHLUTER, D. (2001). Ecology and the origin of species. *Trends in Ecology and Evolution* **16**(7), 372-380.
- SCHMIDT, T. L., THIA, J. A. & HOFFMANN, A. A. (2023). How can genomics help or hinder wildlife conservation? *Annual Review of Animal Biosciences* **12**. 10.1146/annurev-animal-021022-051810
- SEEHAUSEN, O., BUTLIN, R. K., KELLER, I., WAGNER, C. E., BOUGHMAN, J. W., HOHENLOHE, P. A., PEICHEL, C. L., SAETRE, G.-P., BANK, C., BRÄNNSTRÖM, Å., BRELSFORD, A., CLARKSON, C. S., EROUKHMANOFF, F., FEDER, J. L., FISCHER, M. C., FOOTE, A. D., FRANCHINI, P., JIGGINS, C. D., JONES, F. C., LINDHOLM, A. K., LUCEK, K., MAAN, M. E., MARQUES, D. A., MARTIN, S. H., MATTHEWS, B., MEIER, J. I., MÖST, M., NACHMAN, M. W., NONAKA, E., RENNISON, D. J., SCHWARZER, J., WATSON, E. T., WESTRAM, A. M. & WIDMER, A. (2014). Genomics and the origin of species. *Nature* **15**(3), 176-192. 10.1038/nrg3644

- SHAVER, E., COURTNEY, C., WEST, J., MAYNARD, J., HEIN, M., WAGNER, C., PHILIBOTTE, J., MACGOWAN, P., MCLEOD, I. & BÖSTROM-EINARSSON, L. (2020). A manager's guide to coral reef restoration planning and design.
- SHEETS, E. A., WARNER, P. A. & PALUMBI, S. R. (2018). Accurate population genetic measurements require cryptic species identification in corals. *Coral Reefs* **37**(2), 549-563. 10.1007/s00338-018-1679-9
- SHILLING, E. N., ECKERT, R. J., STURM, A. B. & VOSS, J. D. (2023). *Porites astreoides* coral populations demonstrate high clonality and connectivity in southeast Florida. *Coral Reefs* **42**, 1131–1145. 10.1007/s00338-023-02417-0
- SOUSA, V. C. & HEY, J. (2013). Understanding the origin of species with genome-scale data: modelling gene flow. *Nature* **14**(6), 404-414. 10.1038/nrg3446
- SOUTER, D., PLANES, S., WICQUART, J., OBURO, D. & STAUB, F. (2021). Status of coral reefs of the world: 2020 report. Global Coral Reef Monitoring Network (GCRMN) and International Coral Reef Initiative (ICRI).
- STARKO, S., FIFER, J. E., CLAAR, D. C., DAVIES, S. W., CUNNING, R., BAKER, A. C. & BAUM, J. K. (2023). Marine heatwaves threaten cryptic coral diversity and erode associations among coevolving partner. *Science Advances* **9**(32), eadf0954.
- STURM, A. B., ECKERT, R. J., CARREIRO, A. M., SIMÕES, N. & VOSS, J. D. (2022). Depth-dependent genetic structuring of a depth-generalist coral and its symbiodiniaceae algal communities at Campeche Bank, Mexico. *Frontiers in Marine Science* **9**, 835789. 10.3389/fmars.2022.835789
- STURM, A. B., ECKERT, R. J., MENDEZ, J. G., GONZALEZ-DIAZ, P. & VOSS, J. D. (2020). Population genetic structure of the great star coral, *Montastraea cavernosa*, across the Cuban archipelago with comparisons between microsatellite and SNP markers. *Scientific Reports* **10**(1), 15432. 10.1038/s41598-020-72112-5
- TAYLOR, S. A. & LARSON, E. L. (2019). Insights from genomes into the evolutionary importance and prevalence of hybridization in nature. *Nature Ecology & Evolution* (3), 170-177. 10.1038/s41559-018-0777-y
- THIA, J. A. (2022). Guidelines for standardizing the application of discriminant analysis of principal components to genotype data. *Molecular Ecology Resources* **23**, 523-538. 10.1111/1755-0998.13706
- THOMAS, L., UNDERWOOD, J. N., ADAM, A. A. S., RICHARDS, Z. T., DUGAL, L., MILLER, K. J. & GILMOUR, J. P. (2019). Contrasting patterns of genetic connectivity in brooding and spawning corals across a remote atoll system in northwest Australia. *Coral Reefs* **39**(1), 55-60. 10.1007/s00338-019-01884-8
- VAN OPPEN, M. J. & GATES, R. D. (2006). Conservation genetics and the resilience of reef-building corals. *Molecular Ecology* **15**(13), 3863-83. 10.1111/j.1365-294X.2006.03026.x
- VAN OPPEN, M. J. H., BONGAERTS, P., FRADE, P. R., PELOW, L. M., BOYD, S. E., NIM, H. T. & BAY, L. K. (2018). Adaptation to reef habitats through selection on the coral animal and its associated microbiome. *Molecular Ecology* **27**(14), 2956-2971. 10.1111/mec.14763
- VAN OPPEN, M. J. H., OLIVER, J. K., PUTNAM, H. M. & GATES, R. D. (2015). Building coral reef resilience through assisted evolution. *Proceedings of the National Academy of Sciences of the United States of America* **112**(8), 2307-2313. 10.1073/pnas.1422301112
- VANWYNEN, C. M., HIGHTSHOE, M. V., FOGARTY, N. D., DAHLGREN, C. P. & GILLIAM, D. S. (2021). Should hybrids be used in coral nurseries? A case study comparing Caribbean *Acropora* spp.

- and their hybrid in the Bahamas. *Frontiers in Marine Science* **8**. 10.3389/fmars.2021.669966
- VERON, J. E. N. (1995). *Corals in Space and Time: The Biogeography and Evolution of the Scleractinia*. UNSW Press, Sydney.
- WAPLES, R. S. & GAGGIOTTI, O. E. (2006). What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology* **15**(6), 1419-1439. 10.1111/j.1365-294X.2006.02890.x
- WARNER, P. A., VAN OPPEN, M. J. H. & WILLIS, B. L. (2015). Unexpected cryptic species diversity in the widespread coral *Seriatopora hystrix* masks spatial-genetic patterns of connectivity. *Molecular Ecology* **24**, 2993–3008. 10.1111/mec.13225
- WESTRAM, A. M., RAFAJLOVIC, M., CHAUBE, P., FARIA, R., LARSSON, T., PANOVA, M., RAVINET, M., BLOMBERG, A., MEHLIG, B., JOHANNESSON, K. & BUTLIN, R. (2018). Clines on the seashore: The genomic architecture underlying rapid divergence in the face of gene flow. *Evolution Letters* **2**(4), 297-309. 10.1002/evl3.74
- WILKINSON, M. D., DUMONTIER, M., AALBERSBERG, I. J., APPLETON, G., AXTON, M., BAAK, A., BLOMBERG, N., BOITEN, J.-W., DA SILVA SANTOS, L. B., BOURNE, P. E., BOUWMAN, J., BROOKES, A. J., CLARK, T., CROSAS, M., DILLO, I., DUMON, O., EDMUNDS, S., EVELO, C. T., FINKERS, R., GONZALEZ-BELTRAN, A., GRAY, A. J. G., GROTH, P., GOBLE, C., GRETHE, J. S., HERINGA, J., T HOEN, P. A. C., HOOFT, R., KUHN, T., KOK, R., KOK, J., LUSHER, S. J., MARTONE, M. E., MONS, A., PACKER, A. L., PERSSON, B., ROCCA-SERRA, P., ROOS, M., VAN SCHAIK, R., SANSONE, S.-A., SCHULTES, E., SENGSTAG, T., SLATER, T., STRAWN, G., SWERTZ, M. A., THOMPSON, M., VAN DER LEI, J., VAN MULLIGEN, E., VELTEROP, J., WAAGMEESTER, A., WITTENBURG, P., WOLSTENCROFT, K., ZHAO, J. & MONS, B. (2016). The FAIR Guiding Principles for scientific data management and stewardship. *Scientific Data* **3**(1), e1002295-9. 10.1038/sdata.2016.18
- WILLIS, B. L., VAN OPPEN, M. J. H., MILLER, D. J., VOLLMER, S. V. & AYRE, D. J. (2006). The role of hybridization in the evolution of reef corals. *Annual Review of Ecology, Evolution, and Systematics* **37**(1), 489-517. 10.1146/annurev.ecolsys.37.091305.110136
- ZHANG, J., RICHARDS, Z. T., ADAM, A. A. S., CHAN, C. X., SHINZATO, C., GILMOUR, J., THOMAS, L., STRUGNELL, J. M., MILLER, D. J. & COOKE, I. (2023). Evolutionary responses of a reef-building coral to climate change at the end of the last glacial maximum. *Molecular Biology and Evolution* **39**, p.msac201. 10.1093/molbev/msac201
- ZHANG, J., SCHNELLER, N. M., FIELD, M. A., CHAN, C. X., MILLER, D. J., STRUGNELL, J. M., RIGINOS, C., BAY, L. & COOKE, I. (2024). Chromosomal inversions harbour excess mutational load in the coral, *Acropora kenti*, on the Great Barrier Reef. *Molecular Ecology*, e17468. 10.1111/mec.17468

Appendix

Structured review for population genomic studies

An initial search of Web of Science Core Collection was performed on 21/10/2022, using the search terms "(TI=(coral) OR TI=(scleractinia*) NOT TI=(fish)) AND (AB=(rad*) OR ALL=(snp*))". The search returned 803 studies. Titles and abstracts were filtered to exclude irrelevant studies. Studies appearing to contain population genomic data of scleractinians or octocorallians were retained for manual inspection. On 4/11/2022 the above search was repeated allowing for "coral" to be in the abstract rather than title and this yielded 853 additional studies for a total of 1620 unique papers that were reduced to 99 after skimming titles and abstracts.

Each manuscript was read and evaluated by two people independently to ensure that genomic data: i) pertained to the cnidarian coral host, ii) used many loci on a genomic scale (i.e., not microsatellites, not metabarcoding), iii) surveyed two or more sites or habitats, and iv) presented results that included ordination analysis based on individual genotypes (i.e., principal components analysis (PCA), principal coordinates analysis (PCOA) or multidimensional scaling (MDS)) and/or unsupervised model based clustering tests (such as ADMIXTURE, STRUCTURE, fastSTRUCTURE or sNMF). We did not consider papers only reporting discriminant analysis of principal components, as DAPC finds the eigenvectors that best differentiate prespecified groups. In contrast, PCA, PCOA and MDS find eigenvalues that best capture total diversity regardless of group membership (see Thia, 2022 for further discussion). If these four conditions were not met, the study was excluded. For each retained study, two evaluators independently extracted key attributes and reconciled discrepancies between their scoring through discussion. Despite attempting to undertake a rigorous and inclusive search, we noticed that several suitable manuscripts were missing and therefore on July 18, 2023 we ran an ad hoc search in Web of Science based on authors that are known to be publishing on population genomics of corals (namely Barshis, DJ; Baums, IB; Bay, LK; Bongaerts, P; Cooke, I; Matz, MV; Palumbi, SR; Richards, ZT, Underwood, JN, van Oppen, MJH) and also repeated the above search with exactly the same criteria for articles published since 21/10/2022. These new papers were evaluated as above. The ad hoc search initially identified 897 papers that were reduced to 16 papers once titles and abstracts were skimmed. In total, 41 papers were found suitable for data extraction representing a total of 51 records (unique paper by species combinations).

Heat stress studies

The CORDAP database (Ortiz *et al.*, 2023) was downloaded on 11/09/2023 and searched for any records which used genome-wide data of the coral host (i.e., coral SNP data) and those which conducted clustering analyses (as described above). First, records were filtered based on whether the database columns "Host_genotype", "Symbiodiniaceae_genotype", and

“Microbiome_genotype” were listed as “TRUE”. Second, the columns “Symbiodiniaceae_genotyping_approach”, “Host_genotyping_approach” and “Microbiome_genotyping_approach” were interrogated and only records where the genotyping method was listed as reduced representation sequencing (e.g., Restriction site Associated DNA sequencing), whole genome sequencing (WGS) or RNA sequencing (RNAseq) were kept for further checks. The titles and abstracts of the remaining records were checked, and only records for which host genotyping was performed were included. The number of cryptic species assigned in each paper by the original authors was noted, as well as the evaluators’ interpretation of the number of cryptic species based on the plots and analyses (following guidelines in 2.1).

The original database consisted of 562 records, many of which did not include host, Symbiodiniaceae, or microbiome genotyping. The initial filtering of the database for records that included some aspect of host genotyping yielded 222 results and was reduced to 17 records that included genome-scale genotyping. Of these 17 studies, nine studies included either ordination or model-based clustering analyses (e.g., PCA, ADMIXTURE). Of these, three (Rose *et al.*, 2017; Ruiz-Jones & Palumbi, 2017; Rose *et al.*, 2021) matched the criteria in Section 2.1 showing evidence for cryptic taxa. The remaining six studies were either ambiguous or showed no clear evidence for cryptic taxa.