

1 Quantifying taxon-specific habitat connectivity requirements of urban wildlife

2 using structured expert judgement

3 Stephanie K. Courtney Jones^{1,2,3,4}, Luke S. O'Loughlin¹, Danswell Starrs⁴, Jacinta E. Humphrey⁵, Stephanie A.
4 Pulsford¹, Hugh Allan⁶, Matt Beitzel¹, Kym Birgen⁷, Suzi Bond^{8,9}, Jenny Bounds¹⁰, Deborah Bower¹¹, Renee
5 Brawata¹, Ben Broadhurst⁶, Emma Carlson⁷, Simon Clulow¹², Saul Cunningham⁹, Luke Dunn⁷, Lisa Evans¹, Bruno
6 Ferronato¹, Donald B. Fletcher¹³, Arthur Georges¹², Amy-Marie Gilpin^{14,15}, Mark A. Hall¹⁵, Brian Hawkins¹, Anke
7 Maria Hoeffler¹⁶, Brett Howland¹, Damian C. Lettoof¹⁷, Mark Lintermans⁶, Michelle Littlefair⁹, Tanya Latty¹⁸,
8 Tyrone H. Lavery^{9,19}, Zohara Lucas¹, George Madani²⁰, Kim Maute^{21,22}, Richard N. C. Milner⁷, Eric J. Nordberg²³,
9 Thea O'Loughlin⁷, Woo O'Reilly⁴, Megan O'Shea²⁴, Laura Rayner⁷, Euan G. Ritchie²⁵, Natasha M. Robinson^{9,26},
10 Stephan D. Sarre¹², Manu E. Saunders²³, Ben C. Scheele⁹, Julian Seddon¹, Rob Speirs²⁷, Ricky Spencer¹⁴, Ingrid
11 Stirnemann⁷, David M. Watson²⁸, Belinda A. Wilson⁹, Peter J. Unmack⁶, Yuying Zhao⁷, Melissa A. Snape¹

12 ¹ Office of Nature Conservation, Australian Capital Territory Government, Canberra, ACT 2602, Australia

13 ² Division of Ecology and Evolution, Research School of Biology, Australian National University, Canberra, ACT 2601,
14 Australia.

15 ³ Ecosystem and Threatened Species, Northwest Branch, NSW Department of Planning and Environment, Dubbo, NSW
16 2830, Australia

17 ⁴ Office of Water, Australian Capital Territory Government, Canberra, ACT 2602, Australia

18 ⁵ Department of Environment and Genetics, Research Centre for Future Landscapes, La Trobe University, Melbourne, VIC
19 3086, Australia

20 ⁶ Centre for Water Science, Institute of Applied Ecology, University of Canberra, Bruce, ACT 6201, Australia

21 ⁷ Parks and Conservation Service, Australian Capital Territory Government, Canberra, Australian Capital Territory, Australia

22 ⁸ Australian National Insect Collection, GPO Box 1700, Canberra, ACT 2601, Australia

23 ⁹ Fenner School of Environment and Society, The Australian National University, Building 141, Linnaeus Way, Canberra, ACT
24 2601, Australia

25 ¹⁰ Canberra Ornithologist Group, Woodland Bird Monitoring Project, Canberra, ACT 2601, Australia

26 ¹¹ UNE Zoology Discipline, School of Environmental and Rural Science, University of New England, Armidale, NSW 2351,
27 Australia

28 ¹² Centre for Conservation Ecology and Genomics, Institute of Applied Ecology, University of Canberra, Bruce, ACT 6201,
29 Australia

30 ¹³ National Parks Association of the ACT Inc., General Post Office Box 544, Canberra, ACT 2601, Australia

31 ¹⁴ School of Science, Western Sydney University, Penrith, New South Wales, Australia

32 ¹⁵ Hawkesbury Institute for the Environment, Western Sydney University, Locked Bag 1797, Penrith, NSW 2751, Australia

33 ¹⁶ ACT and Region Frogwatch, Ginninderra Catchment Group, Canberra, ACT, 2615, Australia

34 ¹⁷ Centre for Environment and Life Sciences, Commonwealth Scientific and Industrial Research Organisation, Floreat, WA
35 6014, Australia

36 ¹⁸ School of Life and Environmental Science, Sydney Institute of Agriculture, The University of Sydney, Sydney, NSW 2006,
37 Australia

38 ¹⁹ School of BioSciences, The University of Melbourne, Melbourne, VIC 3010, Australia

39 ²⁰ Conservation Science Research Group, School of Environmental and Life Sciences, Biology, Building, University of
40 Newcastle, University Drive, Callaghan, NSW, 2308, Australia

41 ²¹ School of Earth, Atmospheric and Life Sciences, University of Wollongong, Wollongong, NSW 2522, Australia

42 ²² Environmental Futures Research Centre, University of Wollongong, Wollongong, NSW 2522, Australia

43 ²³ Ecosystem Management, School of Environmental and Rural Science, University of New England, Armidale, NSW 2351,
44 Australia

45 ²⁴ College of Sport, Health & Engineering, Victoria University, Melbourne, VIC 3000, Australia

46 ²⁵ School of Life and Environmental Sciences, Deakin University, Burwood, VIC 3125, Australia

47 ²⁶ Conservation and Restoration Science Branch, Science, Economics and Insights Division, NSW Department of Planning
48 and Environment, NSW, Australia

49 ²⁷ Capital Ecology Pty Ltd. Canberra, ACT 2601, Australia

50 ²⁸ Gulbali Institute and School for Agricultural, Environmental, and Veterinary Sciences, Charles Sturt University, Albury,
51 NSW 2678, Australia

52

53 **Abstract**

54 Urban planning which enhances native biodiversity in and around cities is needed to address the
55 impacts of urbanisation and conserve urban biodiversity. The “Biodiversity Sensitive Urban Design”
56 (BSUD) framework incorporates ecological knowledge into urban planning to achieve positive
57 biodiversity outcomes through improved urban design and infrastructure development. BSUD
58 includes principles to direct strategic design and placement of connected wildlife habitat. However,
59 effective BSUD implementation requires defining and quantifying the landscape-scale habitat
60 connectivity needs of a range of taxon groups within urban contexts. The aim of our study was to
61 use expert elicitation to address these gaps in landscape-scale habitat connectivity currently limiting
62 the capacity of urban planning. We estimated habitat connectivity needs for seven representative
63 taxon groups in urban environments, including ideal habitat, habitat constraints, barriers to
64 movement, and movement thresholds that determine habitat connectivity. In using expert
65 elicitation to quantify habitat connectivity requirements for urban biodiversity, our study provides
66 insights on both the usefulness of expert elicitation to inform urban habitat connectivity planning
67 generally, and the functional habitat connectivity requirements of our focal taxon groups specifically.
68 Overall, we consider our expert-derived estimates of connected habitat to be a highly useful set of
69 baseline data for habitat and connectivity modelling and urban planning for a range of taxon groups.

70 **Introduction**

71 Urbanisation threatens biodiversity through habitat loss and fragmentation, and the modification of
72 resource availability, disturbance regimes, local climate, and species assemblages within what
73 habitat remains (McKinney 2008; McDonald *et al.* 2008, 2020; Seto *et al.* 2012; Garrard *et al.* 2018,
74 Selinske *et al.* 2022). However, the urban environment is important for biodiversity conservation,
75 with many native species (including rare and threatened species) having population strongholds
76 (Maclagan *et al.* 2018) or persisting entirely within urban landscapes (Ives *et al.* 2016; Garrard *et al.*
77 2018; Soanes and Lentini 2019). Urban planning which aims to minimise the impacts of urbanisation
78 and enhance native biodiversity in and around cities is therefore urgently needed (Garrard *et al.*
79 2018; Scheele *et al.* 2018; Huang *et al.* 2018). ‘Biodiversity Sensitive Urban Design’ (BSUD) presents a
80 framework for better incorporating ecological knowledge into urban planning to promote
81 biodiversity and mitigate the impacts of urbanisation through improved urban design and
82 infrastructure development (Garrard *et al.* 2018).

83 The BSUD framework sets out five principles: (1) maintain and introduce habitat, (2) facilitate
84 dispersal, (3) minimise threats and anthropogenic disturbances, (4) facilitate natural ecological
85 processes, and (5) improve potential for positive human–nature interactions (Garrard *et al.* 2018).
86 The first two principles of BSUD intend, among other things, to direct more strategic design and
87 placement of connected wildlife habitat in urban landscapes (Garrard *et al.* 2018). However, Kirk *et al.*
88 (2018) identified two key factors that currently limit the capacity of urban design to achieve
89 habitat connectivity outcomes: (1) the assumption that connected habitat defined by structural
90 elements (e.g., patch dimensions, vegetation composition, and spatial continuity) provides
91 appropriately for target wildlife in the absence of defining functional constraints (e.g., physical,
92 physiological, or behavioural barriers to successful use, movement, or dispersal), and (2) a lack of

93 empirical information to describe taxon-specific ideal habitat requirements and constraints at the
94 relevant spatial scale to inform evidence-based urban design for target wildlife.

95 Addressing these limitations to effective BSUD implementation requires defining and quantifying the
96 landscape-scale connectivity requirements for a range of taxon groups within urban contexts. The
97 'City Biodiversity Index' – a Convention on Biological Diversity endorsed tool to monitor urban
98 biodiversity – measures ecological connectivity as the relationship between the total area of habitat
99 available and the degree to which it is functionally (dis)connected, either by distance (e.g., small
100 birds will be unable to disperse where distance between tree cover exceeds their movement
101 capacity (Tremblay and St. Clair 2009)) or by physical or behavioural barriers to movement (Chan *et al.*
102 *et al.* 2014; Deslauriers *et al.* 2017; Kirk *et al.* 2018, 2023). While recent studies have highlighted the
103 value of using this approach for spatially mapping and measuring habitat connectivity in BSUD (e.g.,
104 Kirk *et al.* 2018, 2021), the input data often remains coarse in terms of what constitutes habitat (e.g.,
105 presence of trees only without consideration of preferred spacing and composition), and taxon-
106 specific movement thresholds and movement barriers (Kirk *et al.* 2023). Applying BSUD to achieve
107 ecological connectivity outcomes requires a greater taxon-specific understanding of what
108 constitutes functional connected habitat to underpin these connectivity maps, models, and
109 measures.

110 Robust empirical data on the functional connectivity requirements of most species within urban
111 environments are severely lacking. Expert judgement is increasingly used to inform decisions where
112 empirical data are insufficient or unobtainable due to funding limitations for systematic ecological
113 surveys and monitoring (Legge *et al.* 2018). A range of methods have been developed to minimise
114 inherent bias and uncertainty, and to account for wide variances in knowledge (Martin *et al.* 2012).
115 One such method is the 'IDEA' protocol (standing for 'Investigate', 'Discuss', 'Estimate', and
116 'Aggregate') which is a structured elicitation approach designed to improve the accuracy and
117 quantitative rigor of expert judgements (Hanea *et al.* 2017; Hemming *et al.* 2018). The IDEA protocol
118 is routinely used in government policy settings (e.g., forecasting changes in biosecurity risk
119 (Wittmann *et al.* 2015)) and in ecological and conservation contexts (e.g., Geyle *et al.* 2020; Camac *et al.*
120 *et al.* 2021). However, to our knowledge, this form of structured expert elicitation has not yet been
121 used to address data gaps in taxon-specific habitat connectivity requirements in urban
122 environments.

123 The aim of our study was to use the IDEA protocol of expert elicitation to address gaps in landscape-
124 scale habitat connectivity data which limit the capacity of urban planning to adopt the BSUD
125 principles of "maintain and introduce habitat" and "facilitate dispersal". We used the city of
126 Canberra in the Australian Capital Territory (ACT) as a case study to quantify habitat connectivity
127 needs for seven taxon groups—invertebrate and vertebrate species spanning terrestrial, arboreal,
128 aquatic, and aerial habitats— of representative fauna present in that urban environment. Taxon-
129 specific experts quantitatively estimated ideal habitat, habitat constraints, barriers to movement,
130 and movement thresholds that determine habitat connectivity. In using expert elicitation to quantify
131 habitat connectivity requirements for urban biodiversity, our study provides insights on both the
132 usefulness of the IDEA protocol to inform urban habitat connectivity planning generally, and the
133 functional habitat connectivity requirements of our focal taxon groups specifically.

134 **Methods**

135 **Study area**

136 Our study was conducted for Canberra, ACT, an inland city in temperate south-eastern Australia.
137 Canberra has a population of 455,900 which has been growing at a rate of 2.3% per year since 2011,
138 faster than any other Australian city during that time (Alexandra *et al.* 2017; Alexandra and Norman
139 2020; ABS 2022). While the total urban area of Canberra is approximately 800 km², the developed
140 urbanised footprint is only around half of this, with the remaining area consisting of urban green
141 spaces and an extensive urban reserve network of remnant native vegetation (ACT Government
142 2018). As a result, the city is colloquially known as the 'Bush Capital' and has the second lowest
143 population density of any major Australian city (~1000 people per km² (ACT Government 2018; ABS
144 2022)). Canberra population densities are already increasing under a planning strategy that seeks to
145 limit urban spread through prioritising development within the existing urban footprint, however
146 new urban growth areas are also being established (ACT Government 2018). The planning strategy
147 seeks to grow Canberra in a way that protects and maintains the biodiversity values of the city.

148 Canberra is built in an area of the ecologically diverse Southern Tablelands region west of the Great
149 Dividing Range that was once dominated by box-gum grassy woodlands and natural temperate
150 grasslands. The Ngunnawal people are the Traditional Custodians of the land and waters of the ACT,
151 and for tens of thousands of years actively manipulated the woodlands, grasslands, and waterways
152 in the region, shaping the structure and function of these ecosystems. Some large intact remnants of
153 critically endangered woodland and grassland remain in and around Canberra, but most have been
154 substantially modified by land clearing, urbanisation, livestock grazing, invasion by weeds and feral
155 animals, and the loss of Indigenous management following European colonisation. Many natural
156 creeks, tributaries and associated riparian vegetation that were present throughout Canberra are
157 now highly modified, with most of these areas now existing as concreted drains of little ecological
158 value. Urbanisation presents an ongoing threat to the extent, condition, and connectedness of these
159 ecosystems in the region, and greater understanding of the habitat connectivity needs of the native
160 wildlife that rely on these areas within the city is crucial for sustainable urban policy, planning, and
161 management (Ikin *et al.* 2015; Rayner *et al.* 2014; Hale *et al.* 2015).

162 **Selection of representative taxon groups**

163 We selected seven taxon groups for which to quantify the landscape-scale habitat connectivity
164 requirements of fauna within urban Canberra. We decided to use a taxon group approach which
165 considers species that have relative ecological similarities and share broad dispersal abilities and
166 habitat requirements (as opposed to an individual species approach) (e.g., Kirk *et al.* 2018). We
167 included seven taxon groups to best capture the breadth of ecosystem associations, habitat needs,
168 and movement abilities of most fauna in urban Canberra, particularly ACT threatened species. These
169 groups of species were: (1) grassland reptiles, (2) native bees, (3) small–medium terrestrial mammals
170 (hereafter small–medium mammals), (4) small woodland birds (hereafter woodland birds), (5)
171 riparian reptiles and mammals, (6) amphibians, and (7) small freshwater fish (see Table 1 for taxon
172 group definitions, justification, and final list of species considered). While there are other taxon

173 groups that could have been considered (e.g., arboreal mammals, water birds, tree-hollow using
174 fauna, soil-dwelling fauna), we considered the selected fauna as broadly informative for taxa not
175 explicitly assessed. For example, we selected four taxon groups that are associated with box-gum
176 grassy woodlands that vary widely in their dispersal capacity and specific habitat requirements (i.e.,
177 native bees, small–medium mammals, woodland birds, and amphibians), presuming that these
178 adequately captured the variability in connected habitat needed for other non-assessed woodland-
179 associated species (e.g., native bees broadly represent other insect pollinators). Four taxon groups
180 were associated with natural temperate grasslands (i.e., grassland reptiles, native bees, small–
181 medium mammals, and amphibians), and three taxon groups were associated with aquatic zones
182 and riparian vegetation (i.e., riparian reptiles and mammals, amphibians, and small freshwater fish).

183 We refined our considered species within each taxon group to a final agreed list prior to quantifying
184 their habitat connectivity requirements (Table 1). Initial broad species lists for each taxon group
185 were established based on existing systematic lists relevant to the ACT (e.g., small woodland birds as
186 identified by Fraser *et al.* 2019; amphibians as identified by Westgate *et al.* 2015; all other groups as
187 described on the citizen-science platform Canberra Nature Map
188 <https://canberra.naturemapr.org/>). During expert elicitation workshops, we then discussed the
189 relative value of including or excluding particular species from each taxon group for our assessment.
190 Native species were included where they were considered strongly representative of the group in
191 urban areas and were (a) common but potentially threatened by increased urbanisation, (b) present
192 but listed as vulnerable in the ACT, (c) established following translocation to the ACT, or (d) absent
193 or rare in the ACT urban areas but could potentially re-establish in the future (e.g., through
194 reintroductions or assisted migration; Buckmaster *et al.* 2010). Species were excluded if they were
195 considered not representative of the group because of (a) unique habitat requirements or dispersal
196 capacities, (b) having a natural or predicted distribution which did not include the urban extent of
197 the ACT, (c) requiring direct management interventions for persistence, or (d) were absent or rare in
198 the ACT with re-establishment deemed extremely unlikely.

199 **Table 1.** Definition, species list, and justification (reasons for inclusion) for the seven taxon groups
 200 assessed for connected habitat requirements through expert elicitation in Canberra, Australian
 201 Capital Territory (ACT). Bolded species are either #endangered or critically endangered, †vulnerable,
 202 ‡regionally conservation dependant, ^locally rare, or *absent from the ACT lowlands but may occur
 203 in the future via assisted or unassisted means. Species scientific names can be found in
 204 Supplementary Material.

Taxon group and definition	Species considered	Justification
Grassland reptiles: reptile species that have a strong association to grasslands.	Blue-tongued lizard Eastern brown snake Grassland earless dragon# Pink-tailed worm-lizard† Striped legless lizard† Three-toed skink	We considered here characteristic grassland species (predominantly grassland specialists), using them as a surrogate group to ensure 'Natural Temperate Grassland' structure and functionality was protected within the urban extent.
Native Bees: all native species of the clade Anthophila (Order Hymenoptera).	All native bee species occurring within the ACT (approximately 150 species).	Native bees are a major pollinators within the urban extent and so were considered broadly representative of other insect pollinating orders (Hymenoptera, Diptera, Lepidoptera, Coleoptera).
Small-medium terrestrial mammals: mammals within the critical weight range (35–5500 g) that are predominantly terrestrial (excluding arboreal mammals such as possums, and volant mammals including bats).	Agile antechinus Brush-tailed phascogale* Bush rat Common dunnart Eastern bettong†* Eastern chestnut mouse Long-nosed bandicoot New Holland mouse† Short-beaked echidna Southern brown bandicoot#* Yellow-footed antechinus	Species considered within this group were currently present (but may be absent from urban areas, e.g., Buckmaster <i>et al.</i> 2010) or likely to occur within the urban extent of the ACT (e.g., [eastern] southern brown bandicoot; eastern bettong; and brush-tailed phascogale). Spotted-tailed and eastern quolls were considered likely to benefit from similar habitat conditions but were not considered in the expert elicitation.
Small woodland birds: smaller bird species (<40 g) of the ecologically and functionally identifiable Temperate South-eastern Mainland Australia ecoregion sub-community of the <i>Australian Temperate and Subtropical Woodland Bird Community</i> (Fraser <i>et al.</i> 2019).	Brown-headed Honeyeater Brown Treecreepert† Buff-rumped Thornbill Diamond Firetail Eastern Yellow Robin Fuscous Honeyeater Grey Fantail Leaden Flycatcher Mistletoebird Painted Button-Quail Rufous Whistler Scarlet Robin† Southern Whiteface Speckled Warbler Striated Pardalote Striated Thornbill Superb Fairy-Wren Tree Martin Weebill White-browed Scrubwren White-throated Gerygone Yellow-rumped Thornbill	Smaller species in the broader woodland bird community are most vulnerable to the threatening processes of the urban landscape (e.g., harassment by noisy miners, simplification of woodland structure). We included species that were increasing and declining, using different parts of the woodland forest column, were woodland-dependent, and already occurring the urban extent of the ACT.
Riparian reptiles and mammals: semi-aquatic species	Eastern long-necked turtle Eastern water dragon	Reptile and mammal species considered within this group were currently present

which have specific riparian or aquatic habitat requirements.	Gippsland water dragon Platypus Rakali Red-bellied black snake Tiger snake	within the urban areas of the ACT and had specific riparian or aquatic habitat requirements for population persistence.
Amphibians: any native frog, froglet, or toadlet.	Bibron's toadlet* Broad-palmed rocket frog Common eastern froglet Eastern banjo frog Eastern sign-bearing froglet Green and golden bell frog †* Stony Creek frog Peron's tree frog Smooth toadlet Spotted marsh frog Striped marsh frog Sudell's frog^ Verreaux's tree frog	Species in this taxon group included those currently occurring within or near urban areas within the ACT using data generated from the citizen-science Frogwatch ACT and Region Program (Westgate <i>et al.</i> 2015). Species which were considered candidates for reintroduction to the urban area were also included.
Small freshwater fish: freshwater fish with <10 cm total length or fork length.	Australian smelt Bald carp gudgeon* Flathead gudgeon Mountain galaxias Southern pygmy perch* Western carp gudgeon	Experts considered aquatic habitat within the urban extent of the ACT to only be suitable for small species, rather than larger species (e.g., Murray cod). As a result, the species list includes smaller species found in small stream environments, and species which transit between lake and large river core habitat. Two species, bald carp gudgeon (<i>Hypseleotris</i> sp.) and southern pygmy perch (<i>Nannoperca australis</i>), were included as potential candidates for introduction to the ACT.

205 **Selection of habitat connectivity metrics**

206 The most robust measures of functional connectivity (e.g., effective mesh size for City Biodiversity
207 Index, see Deslauriers *et al.* 2018) quantify the potential of a given landscape to provide
208 unfragmented or unobstructed habitat for particular wildlife by spatially mapping habitat and
209 barriers to movement (Deslauriers *et al.* 2018; Kirk *et al.* 2023). To be informative for such measures,
210 metrics that define taxon-specific habitat connectivity need to be both ecologically meaningful and
211 translate into spatial data layers that are location-specific and readily available (Kirk *et al.* 2023). We
212 selected 30 metrics to represent landscape-scale, functional habitat connectivity for our seven taxon
213 groups (Table 2) that were ecologically important (Doerr *et al.* 2010; 2014) and had the potential to
214 provide the spatial data inputs to underpin robust measures of functional connectivity (Kirk *et al.*
215 2018; 2023). They included metrics that represented (1) ideal habitat requirements ($n = 8$), (2)
216 habitat constraints ($n = 13$), (3) barriers to movement ($n = 6$), and (4) movement thresholds ($n = 3$).

217 We selected eight ideal habitat requirement metrics to define elements of the physical environment
218 that can promote or inhibit the presence of a taxon group (e.g., preferred distance between mature
219 trees, maximum tolerable distance from a permanent waterbody, etc.). While not included explicitly
220 in previous connectivity indices (see Chan *et al.* 2014; Deslauriers *et al.* 2018; Kirk *et al.* 2023) we
221 also included 13 habitat metrics which constrained the spatial area, vegetation composition, or
222 physical environment of available habitat. We did this to better estimate minimum spatial habitat

223 requirements, environmental tolerances, and what experts deem to be unsuitable habitat (e.g., the
224 preference of grassland reptiles for native species dominance in ground-layer vegetation; Antos and
225 Williams 2015). We selected the six metrics reflecting barriers to movement to define where
226 capacity to disperse between patches would be disrupted (i.e., reduce the movement threshold of a
227 taxon group, e.g., maximum crossable extent of paved surface and tolerable traffic flow during
228 active periods, Table 2). We selected three movement thresholds to define typical movement
229 capacity in the absence of barriers to understand where distance to the next patch of suitable
230 habitat itself became the barrier to movement.

231 Not all metrics were relevant for all taxon groups (confirmed through expert elicitation, e.g.,
232 minimum water depth of core habitat was only relevant for aquatic associated taxon groups). We
233 assessed functional connectivity using a minimum of 16 metrics (applicable to woodland birds;
234 where none of our barriers to movement metrics were relevant due to the ability of these species to
235 fly) and a maximum of 27 metrics (applicable to riparian reptiles and mammals; where terrestrial
236 and aquatic habitat use meant almost all metrics were relevant) (see Table 2 for full details). Where
237 metrics were considered only relevant for some but not all species within a taxon group (e.g., not all
238 small woodland birds require specific ground-layer vegetation conditions), the metric was retained
239 to capture the needs of more specialised (and therefore at-risk) species. All metrics considered were
240 compatible with existing spatial data layers (or layers able to be compiled) to enable habitat
241 connectivity mapping from these data in the future (e.g., Kirk *et al.* 2018).

242 **Table 2.** List of ideal habitat requirements, barriers, habitat constraints and movement threshold metrics, their
 243 description, and whether they were assessed for each of the seven taxon groups (“GR” grassland reptiles; “NB”
 244 native bees; “SM” small-medium mammals; “WB” woodland birds; “RM” riparian reptiles and mammals; “AM”
 245 amphibians; “FF” small freshwater fish). Metrics were presented as questions asked throughout the expert
 246 elicitation process. The applicability of each metric varied among the seven taxon groups as either being not
 247 relevant (and therefore not assessed = blank), assessed as relevant for some species of the group (XX), and
 248 assessed as relevant to all species in the group (XX). Ideal habitat metrics only were also determined to be a
 249 more important (but not critical) habitat element for the group (XX), or an essential (critical) habitat element
 250 for the group (XX).

	Metric	Description	Assessed taxon groups
Ideal habitat requirements	Preferred distance between tree canopies (m)	Preference in terms of tree spacing and canopy density.	<u>GR</u> NB <u>SM</u> <u>WB</u> RM AM FF
	Preferred distance between mature trees (m)	Proxy for preference in terms of access to features associated with mature trees such as fallen limbs, or tree hollows.	GR NB <u>SM</u> <u>WB</u> RM AM
	Preferred distance between mid-storey canopies (m)	Preference in terms of mid-storey spacing and canopy density.	GR NB <u>SM</u> WB
	Preferred distance from ground layer vegetation (m)	Preference in terms of proximity to ground layer vegetation, spacing between vegetation patches	<u>GR</u> NB <u>SM</u> WB RM <u>AM</u>
	Minimum height of ground layer vegetation (cm)	Preference in terms of ground layer vegetation structure and management (e.g., mowing regime).	<u>GR</u> SM WB <u>RM</u> <u>AM</u>
	Maximum height of ground layer vegetation (cm)	Preference in terms of ground layer vegetation structure and management (e.g., grazing regime).	<u>GR</u> SM WB <u>RM</u> <u>AM</u>
	Preferred distance between emergent vegetation (m)	Preference, for aquatic and riparian taxa, in terms of the distance between clumps of emergent vegetation.	<u>RM</u> <u>AM</u> FF
	Maximum distance which can be travelled from permanent waterbody (m)*	Requirements in terms of access to permanent surface water. *Represents a structural habitat requirement for aquatic species.	<u>RM</u> <u>AM</u> FF
Habitat constraints	Minimum width of core habitat patch (m)	The minimum dimension of a patch of suitable size to facilitate permanent residency.	GR NB SM WB RM AM FF
	Minimum suitable core habitat depth (m)	For aquatic habitat, the minimum depth of water required to facilitate permanent residency.	<u>RM</u> AM FF
	Minimum width of movement corridor habitat (m)	The minimum dimension of a patch of suitable size to support movement between ‘core’ habitat areas, but not permanent residency.	GR NB SM WB RM AM FF
	Minimum suitable corridor habitat depth (m)	For aquatic habitat, the minimum depth of water required to facilitate movement between ‘core’ habitat areas, but not permanent residency.	RM FF
	Percentage of trees which need to be native (%)	The proportion of trees which need to be native to facilitate habitat use.	GR NB SM <u>WB</u> RM AM FF

	Percentage of native mid-storey vegetation (%)	The proportion of shrubs which need to be native to facilitate habitat use.	GR NB SM WB AM
	Percentage of native ground layer vegetation (%)	The proportion of ground layer vegetation which needs to be native to facilitate habitat use.	<u>GR</u> NB SM WB RM AM
	Percentage of native emergent vegetation (%)	The proportion of emergent vegetation, in aquatic environments, which needs to be native to facilitate habitat use.	RM AM FF
	Maximum tolerable night-time light levels (Lux)	The level of artificial light conducive to habitat use.	GR NB SM WB RM AM FF
	Maximum tolerable surface temperature (°C)	The maximum surface temperature conducive to habitat use.	<u>GR</u> NB RM AM
	Maximum tolerable ambient temperature (°C)	The maximum ambient temperature conducive to habitat use.	<u>GR</u> NB SM WB RM AM
	Maximum tolerable water temperature (°C)	The maximum water temperature conducive to habitat use.	<u>RM AM FF</u>
	Minimum tolerable water temperature (°C)	The minimum water temperature conducive to habitat use.	<u>RM AM FF</u>
Barriers to movement	Maximum crossable extent of paved surface (m)	The maximum extent of paved surface which does not represent a physical barrier to movement, including concrete drains.	<u>GR</u> SM RM AM FF
	Maximum crossable height of vertical structure (m)	The maximum height of a vertical structure (e.g., wall or fence) which can be crossed in the absence of a suitable gap.	<u>GR</u> <u>SM</u> <u>RM AM FF</u>
	Minimum passable gap dimensions (m)	The minimum gap dimensions required to facilitate movement through an otherwise impenetrable vertical barrier.	<u>GR</u> <u>SM</u> <u>RM AM FF</u>
	Maximum crossable extent of waterbody (m)	The maximum extent of a waterbody which does not represent a physical barrier to movement.	<u>GR</u> NB SM AM
	Tolerable traffic flow during active period (vehicles/hr)	The maximum tolerable level of vehicle traffic (including boats) which does not represent a physical or behavioural barrier to movement during the taxon's active period.	GR SM RM AM
	Tolerable pedestrian traffic flow during active periods (pedestrians/hr)	The maximum tolerable level of pedestrian access (including swimmers) which does not represent a physical or behavioural barrier to movement during the taxon's active period.	GR SM RM
Movement thresholds	Typical movement distance within established home range/territory (m)	The capacity for movement within a home range or territory (used to buffer known species records to determine likely occupied habitat).	GR NB SM WB RM AM FF
	Typical capacity for movement outside of suitable habitat (m)	The capacity to move from areas of suitable habitat to other nearby patches, in the absence of a physical or behavioural barrier.	GR NB SM WB RM AM FF
	Typical dispersal distance when seeking new home range/territory (m)	The landscape scale requirements for connected habitat to facilitate the full display of life history traits.	GR NB SM WB RM AM FF

252 We used the IDEA protocol for conducting structured, iterative expert elicitation to quantify each of
253 the relevant metrics for each of our seven taxon groups (see Hanea *et al.* 2017; Hemming *et al.*
254 2018; Courtney Jones *et al.* 2023). This protocol involved four main steps: (1) *INVESTIGATE*: recruit a
255 diverse group of experts for each taxon group to answer questions with initial quantitative 4-point
256 estimate responses (i.e. best estimate, lower limit and upper limit, and a measure of confidence [or
257 a degree-of-belief] in the accuracy those estimates; Spiers-Bridge *et al.* 2010); (2) *DISCUSS*: convene
258 a workshop with experts to discuss their initial estimates to the questions, clarify their meaning,
259 share reasoning and evidence behind initial estimates, and resolve differences in interpretation of
260 the application of habitat metrics; (3) *ESTIMATE*: enable experts to provide a revised and final
261 estimate to each question that considers the workshop discussion which clarified the taxon group
262 species, existing knowledge, sources of uncertainty, and encouraged cross-examination of reasoning
263 and evidence in context of habitat connectivity within the ACT (Courtney Jones *et al.* 2023); and (4)
264 *AGGREGATE*: mathematically aggregate experts' final estimates to determine the average best,
265 lower limit and upper limit for each taxon group for each metric (Table 2).

266 We recruited experts during a two-month period leading up to a series of taxon group-themed
267 workshops held online in September and October 2021. A total of 59 experts were consulted
268 throughout the study (i.e., contributed to the collective knowledge, discussions, and interpretation
269 of results) with 47 of those providing estimates ($n = 8$ for woodland birds, $n = 7$ for amphibians, $n = 5$
270 for native bees, $n = 5$ for small freshwater fish, $n = 12$ for grassland reptiles, $n = 10$ for small–medium
271 mammals, $n = 4$ for riparian reptiles and mammals [noting that four experts contributed to two
272 taxon group estimates each]. Experts were identified based on both local-based experience and
273 taxon-specific knowledge and were selected to represent a breadth of expertise for each taxon
274 group. Experts included (a) academic researchers and post-graduate students involved in ecological
275 research on relevant taxa, (b) management agency staff involved in field ecology, surveys, and
276 management on relevant taxa within the ACT, and (c) ecological consultants, citizen-scientists,
277 naturalists, or museum and zoo staff with extensive experience with the relevant taxa. We selected a
278 diverse expert panel to capture a broad base of knowledge and perspectives, so as to yield accurate
279 aggregated judgements rather than that of a single well-credentialed expert (Page 2008).

280 Each taxon group workshop ran for between 4–6 working hours, where moderators (SKCJ and MS)
281 lead experts through each metric sequentially, discussing the initial estimates and support for those
282 estimates, the interpretation of each question and relevance of the metric for the taxon group, and
283 ensured all experts were fully informed and prepared to complete their revised estimates after the
284 workshop. A later review of metrics assessed the relative relevance and importance of each metric
285 for each taxon group (Table 2). Despite the majority decisions from such discussion, in 14% of all
286 taxon-specific metrics assessed (21/149) one or more experts felt they either could not (i.e., low
287 familiarity with the metric) or should not (i.e., disagreed with the relevance of the metric) submit
288 final estimates. We presented questions in an order that followed the workflow described by Kirk *et al.*
289 (2023), starting by estimating “ideal habitat” features without defined spatial parameters (e.g.,
290 “what are the structural features of continuous, unfragmented habitat?”), and estimating the taxon-
291 specific habitat constraints, barriers to movement and movement thresholds second (e.g., “what is
292 the minimum size/composition/distance between habitat that is still considered connected?”, see
293 Supplementary Material).

294 **Summary statistics**

295 Expert-derived data can be aggregated with or without weighting (Hanea *et al.* 2017; Hemming *et al.*
296 2018; 2022). While there are some species-level habitat association data that could be used to
297 calibrate and weight expert estimates had we taken a species-level approach, no such calibration
298 data were available at the taxon group-level at which our estimates were made. Therefore, we used
299 equally weighted aggregation using arithmetic means for all data (Hemming *et al.* 2022). We
300 estimated the means of the best, lower, and upper estimate for each metric for each taxon group in
301 which it was assessed. We also calculated standardised 80% credible intervals surrounding the best
302 estimate for each assessed metric using expert-reported confidence levels (Hemming *et al.* 2018).
303 We calculated these intervals for each estimate using linear extrapolation that considered the
304 confidence reported by the experts (see Adams-Hosking *et al.* 2016 and Hemming *et al.* 2018 for
305 equations). Where experts reported 0% confidence, their individual confidence was truncated to 1%
306 to enable calculation, and all credible intervals were averaged for each taxon group by metric
307 combination (Adams-Hosking *et al.* 2016; Hemming *et al.* 2018). Using the four-step elicitation
308 method (i.e., the expert specifying their confidence) and subsequent standardisation of credible
309 intervals reduces overconfidence in expert-derived data by presenting a confidence-informed
310 measure of certainty surrounding the mean (Speirs-Bridge *et al.* 2010; Hemming *et al.* 2018). In the
311 absence of independent empirical data on which to calibrate our expert-derived estimates, no other
312 data summarisation, transformation, or analyses were undertaken. Individual estimates were
313 removed from analysis where no response was provided, or where associated written comments
314 clearly indicated an inconsistent interpretation of the metric compared to other participants. All data
315 summarisation was performed using R version 4.1.2 (R Core Team 2022).

316 **Results**

317 We used the IDEA protocol to estimate 30 metrics to represent landscape-scale, functional habitat
318 connectivity for seven taxon groups (16–27 metrics per taxon group). They included metrics
319 representing (1) ideal habitat requirements (eight metrics), (2) habitat constraints (13 metrics), (3)
320 barriers to movement (six metrics), and (4) movement thresholds (three metrics). We present
321 averaged best estimates (\pm 80% credible intervals) and lower/upper estimates for each habitat
322 connectivity metric assessed (Table 3).

323 **Grassland reptiles**

324 We estimated functional habitat connectivity requirements for grassland reptiles across 23 relevant
325 metrics. Ideal habitat comprised a largely continuous grassy understory with a preferred grass height
326 range of 10–19 cm, and with several hundreds of metres between trees or shrubs. Core habitat was
327 estimated as requiring a minimum width of 188 m (or 38 m for a movement corridor) and high
328 native ground cover (best estimate = 72%, although they could tolerate as low as 21%). As largely
329 diurnal species, grassland reptiles were considered tolerant of high night-time light levels, and high
330 temperatures assuming refugia habitat was available. Grassland reptiles were considered unlikely to
331 cross paved surfaces >5 m wide or vertical structures >0.2 m high. Many grassland reptiles were
332 estimated as having very low movement capacity outside of ideal habitat (<10 m), although larger

333 species considered as part of this group (e.g., eastern brown snake) increased the average to 33 m.
334 Movement within home ranges or dispersal to a new home range was considered low (best = 58–69
335 m).

336 **Native bees**

337 We estimated functional habitat connectivity requirements for native bees across 17 relevant
338 metrics. Ideal habitat for native bees consisted of trees, midstory and/or ground-layer vegetation,
339 generally in an open arrangement, with variable distances between each being preferred. Estimated
340 habitat was constrained to areas with a minimum width of 241 m for core habitat or 32 m for a
341 movement corridor. High nativeness of all strata was also seen as beneficial (best estimates = 64–
342 73%, although some species could tolerate as low as 8% native cover). Native bees were considered
343 tolerant of temperatures $\geq 40^{\circ}\text{C}$ where thermal refugia was available. There was low confidence in
344 whether native bees tolerated only low or moderate night-time light levels (80% credible interval of
345 best estimate = 5–21 Lux). Movement of native bees were impacted by large expanses of pavement
346 or water, but not by vertical structures or traffic. Native bees were deemed to have moderate
347 capacity for movement outside of ideal habitat (best estimate = 214 m, although upper estimate was
348 540 m), roughly equivalent to typical foraging ranges within a habitat patch (best = 200 m).

349 **Small–medium mammals**

350 We estimated functional habitat connectivity requirements for small–medium mammals across 22
351 relevant metrics. Ideal habitat was estimated as having more dense vegetation across all strata than
352 any other taxon group, with shrubs and trees being considered the more important or essential
353 habitat elements for most species considered (best estimates of 7 and 11 m for preferred distances
354 between shrubs and trees, respectively). Core habitat was estimated as being requiring a minimum
355 width of 130 m (or 55 m for a movement corridor) with high levels of nativeness being preferred for
356 all vegetation strata, particularly for trees where the best estimate was 78% native with the low
357 estimate also relatively high at 45%. Small–medium mammals were considered only tolerant of low
358 night-time light levels (best estimate = 4 Lux). All barriers to movement assessed were considered
359 relevant, with the group unlikely to cross paved surfaces >15 m, vertical structures >0.3 m, or traffic
360 areas of >8 vehicles or >10 pedestrians per hour during the taxon groups' active period. This group
361 was assessed as having a high capacity for movement within ideal habitat, including moving a best
362 estimate of 765 m when dispersing to a new territory, but were unlikely to move more than 100 m
363 through unsuitable habitat.

364 **Woodland birds**

365 We estimated functional habitat connectivity requirements for woodland birds across 16 relevant
366 metrics. Ideal habitat was estimated as having moderate tree density, with a complex mid- and/or
367 understory comprised of shrubs or long grasses (best estimates = 41 m and 37 m for preferred
368 distances between tree and midstory canopies). Minimum width requirements for core habitat was
369 the largest for any taxon group (best estimate = 328 m for core habitat, and 28 m for a movement
370 corridor). Experts agreed native vegetation would likely represent ideal habitat but exotic vegetation

371 could also be used if it provided appropriate structure (best estimates = 59–66% native vegetation).
372 Woodland birds were considered tolerant of temperatures <40°C if thermal refugia was available,
373 although prolonged heatwaves were considered likely to impact this species group particularly
374 during breeding periods. Experts considered the group to have reasonable tolerance to artificial
375 night-time light, based on the persistence of many species in urban areas. Small woodland bird
376 movement was not impacted by any barriers assessed and they were considered capable of moving
377 substantial distances across unsuitable habitat (best estimate = 977 m with an upper estimate of 9.5
378 km).

379 **Riparian reptiles and mammals**

380 We estimated functional habitat connectivity requirements for riparian reptiles and mammals across
381 27 relevant metrics. Ideal habitat was variable due to the breadth of species considered, but was
382 generally associated with the riparian zone within 38m of permanent water where combined aquatic
383 and riparian habitat supported emergent vegetation, moderately spaced trees, and ground-layer
384 vegetation with a preferred grass height of 25–50 cm. Habitat was estimated as being constrained
385 mostly by the depth (best estimate = 2.3 m) and width (best estimate = 9 m) of the associated
386 waterbody. Corridor habitat could be narrower (4 m waterbody width) and shallower (1.3 m depth).
387 Habitat was not necessarily constrained by vegetation nativeness (best estimates = 63%) but was
388 constrained by water temperatures outside of a 5–27°C best estimate range. Barriers to movement
389 included paved surfaces >16 m, vertical surfaces >0.7 m, or traffic areas of >6 vehicles or >71
390 pedestrians per hour, however since these averages reflect a diverse group, they do not reflect
391 smaller barriers identified by experts during the discussion which would impact some species (e.g.,
392 smooth vertical barriers for eastern long-necked turtles are likely <10 cm). The average capacity for
393 movement for this taxon group was high, including moving an upper estimate of 4 km when
394 dispersing to a new territory, but their capacity to move outside of suitable habitat was best
395 estimated around 225 m.

396 **Amphibians**

397 We estimated functional habitat connectivity requirements for amphibians across 26 relevant
398 metrics. Ideal habitat was estimated as being within a few hundred metres of water which contained
399 emergent vegetation (distance from water best estimate = 304 m), with moderately spaced trees
400 and ground-layer vegetation also present to varying degrees in the broader landscape (reflecting
401 divergent habitat requirements of different species within this group). Best estimates for preferred
402 grass height were 20–48 cm. Core habitat was estimated as being constrained to a minimum width
403 of 84 m (or 11 m for a movement corridor) and a minimum water depth of 0.6 m. Amphibians were
404 not necessarily constrained by vegetation nativeness (best estimates = 49–56%) but were the least
405 tolerant of high surface and ambient temperatures of any taxon group. Most barriers to movement
406 assessed were considered relevant, with the group unlikely to cross paved surfaces >29 m, vertical
407 surfaces >0.4 m, or waterbodies >31 m. Amphibians were estimated as having moderate–low
408 movement capacity outside of ideal habitat (best = 67 m), although their capacity to disperse
409 through suitable habitat was much higher (best estimate = 479 m, to <2.5 km).

410 **Small freshwater fish**

411 We estimated functional habitat connectivity requirements for small freshwater fish across 18
412 relevant metrics. Ideal habitat was confined to permanent water, with moderately spaced emergent
413 vegetation and trees in the associated riparian environment (best estimates of 13 m and 11 m for
414 preferred distances between those elements, respectively). Core habitat was estimated as being
415 constrained to a minimum width of 5 m (or 2 m for a movement corridor) and a minimum water
416 depth of 1.4 m (or 0.6 for a movement corridor). Experts reported best habitat conditions for this
417 group with estimates of 95% and 100% for native emergent vegetation and trees, respectively. Small
418 freshwater fish were estimated to have the lowest tolerance of night-time light levels of any taxon
419 group, and water temperatures outside of a 7–24°C best estimate range. High movement barriers
420 submerged paved surfaces >12m long and exposed vertical structure >0.1 m high. Their typical
421 movement within a home range or territory was estimated to be the same as their capacity to move
422 outside of suitable habitat (both best estimates ~30–40 m).

423 **Table 3.** Summary of expert-derived functional habitat connectivity requirements for seven taxon groups representative of urban ecosystems *in Canberra, Australian*
 424 *Capital Territory*. Averaged 'Best' (\pm 80% credible intervals), lower and upper (L-U) estimates are presented for all metrics, as well as the number of expert estimates (n)
 425 used to calculate statistics for each metric.

Metric		Grassland reptiles	Native bees	Small-medium mammals	Woodland birds	Riparian reptiles and mammals	Amphibians	Small freshwater fish
Ideal habitat								
Preferred distance between tree canopies (m)	Best	114 (113–123)	40 (40–46)	11 (11–11)	41 (41–43)	28 (27–28)	23 (20–39)	11 (11–31)
	L-U (n)	54 – 965 (8)	7 – 320 (5)	2 – 49 (10)	7 – 155 (8)	8 – 88 (4)	1 – 607 (7)	1 – 440 (5)
Preferred distance between mature trees (m)	Best	865 (856–878)	116 (115–124)	23 (22–23)	75 (74–77)	53 (52–54)	54 (49–111)	
	L-U (n)	83 – 2086 (7)	55 – 510 (5)	9 – 61 (10)	24 – 189 (8)	28 – 100 (4)	5 – 957 (7)	
Preferred distance between mid-storey canopies (m)	Best	792 (788–804)	44 (43–49)	7 (7–8)	37 (36–38)			
	L-U (n)	54 – 1689 (7)	9 – 300 (4)	1 – 29 (10)	8 – 113 (8)			
Preferred distance from ground layer vegetation (m)	Best	1 (0–1)	28 (28–32)	3 (3–3)	4 (4–5)	22 (22–22)	10 (9–24)	
	L-U (n)	0 – 8 (10)	0 – 160 (5)	1 – 11 (10)	0 – 42 (7)	3 – 33 (4)	1 – 739 (7)	
Minimum height of ground layer vegetation (cm)	Best	10 (10–10)		27 (27–28)	11 (11–11)	25 (25–25)	20 (16–25)	
	L-U (n)	5 – 17 (11)		10 – 52 (10)	4 – 29 (7)	15 – 40 (4)	10 – 36 (7)	
Maximum height of ground layer vegetation (cm)	Best	19 (19–19)		50 (49–51)	23 (23–24)	50 (50–51)	48 (45–63)	
	L-U (n)	13 – 33 (11)		33 – 85 (10)	12 – 52 (8)	36 – 86 (4)	30 – 76 (7)	
Preferred distance between emergent vegetation (m)	Best					13 (12–13)	11 (11–12)	13 (13–15)
	L-U (n)					6 – 25 (4)	3 – 27 (7)	2 – 84 (5)
Maximum distance which can be travelled from permanent waterbody (m)*	Best					38 (38–43)	304 (297–375)	0 (0 – 0)
	L-U (n)					8 – 383 (4)	111 – 2021 (7)	0 – 0 (5)
Habitat constraints								
Minimum width of core habitat patch (m)	Best	188 (187–190)	241 (231–251)	130 (127–176)	328 (323–359)	9 (8–9)	84 (82–88)	5 (5–5)
	L-U (n)	82 – 323 (11)	66 – 600 (5)	49 – 1273 (10)	73 – 2075 (8)	5 – 24 (4)	22 – 177 (7)	3 – 33 (5)
Minimum suitable core habitat depth (m)	Best					2.3 (2.2–2.3)	0.6 (0.6–0.7)	1.4 (1.4–1.4)
	L-U (n)					1.5 – 4.0 (4)	0.3 – 0.9 (7)	0.3 – 3.5 (5)
	Best	38 (38–39)	32 (31–36)	55 (55–56)	28 (28–29)	4 (4–4)	11 (10–13)	2 (2–2)

Minimum width of movement corridor habitat (m)	L-U (n)	11 – 141 (11)	5 – 168 (5)	18 – 171 (10)	9 – 91 (8)	4 – 13 (3)	3 – 26 (7)	1 – 26 (5)
Minimum suitable corridor habitat depth (m)	Best L-U (n)					1.3 (1.3–1.3) 0.5 – 2.2 (4)		0.6 (0.6–0.6) 0.2 – 2.0 (5)
Percentage of trees which need to be native (%)	Best L-U (n)	48 (48–48) 23 – 68 (6)	73 (72–74) 14 – 100 (5)	78 (77–79) 45 – 94 (10)	66 (65–66) 32 – 90 (8)	63 (62–63) 38 – 98 (4)	49 (44–53) 9 – 88 (7)	100 (99–100) 12 – 100 (5)
Percentage of native mid-storey vegetation (%)	Best L-U (n)	50 (48–68) 14 – 78 (5)	73 (72–74) 18 – 100 (5)	65 (64–66) 30 – 96 (10)	59 (58–59) 8 – 89 (8)			
Percentage of native ground layer vegetation (%)	Best L-U (n)	72 (71–72) 21 – 96 (11)	64 (56–72) 8 – 98 (5)	74 (72–74) 35 – 94 (10)	64 (63–65) 13 – 94 (8)	63 (62–63) 40 – 90 (4)	53 (41–54) 1 – 91 (7)	
Percentage of native emergent vegetation (%)	Best L-U (n)					53 (46–59) 26 – 93 (4)	56 (49–59) 23 – 85 (7)	95 (93–95) 20 – 100 (5)
Maximum tolerable nighttime light levels (Lux)	Best L-U (n)	21 (21–21) 2 – 718 (7)	5 (5–21) 2 – 212 (5)	4 (4–5) 2 – 21 (8)	7 (6–8) 2 – 22 (7)	0.3 (0.3–0.3) 0.1 – 0.6 (2)	4 (4–7) 0 – 80 (7)	0.2 (0.2–0.7) 0.0 – 8.2 (5)
Maximum tolerable surface temperature (°C)	Best L-U (n)	43 (43–43) 31 – 59 (11)	39 (36–53) 34 – 78 (3)			37 (37–37) 33 – 43 (3)	25 (24–26) 19 – 33 (7)	
Maximum tolerable ambient temperature (°C)	Best L-U (n)	36 (36–36) 30 – 41 (11)	41 (41–41) 36 – 48 (5)	40 (40–40) 35 – 46 (10)	37 (37–37) 31 – 43 (8)	39 (39–40) 32 – 44 (3)	30 (30–30) 21 – 36 (7)	
Maximum tolerable water temperature (°C)	Best L-U (n)					27 (27–27) 24 – 32 (4)	25 (24–27) 21 – 31 (7)	24 (24–24) 16 – 31 (5)
Minimum tolerable water temperature (°C)	Best L-U (n)					5 (5–5) 2 – 7 (4)	8 (8–8) 4 – 12 (7)	7 (7–7) 3 – 12 (5)
Barriers to movement								
Maximum crossable extent of paved surface (m)	Best L-U (n)	5 (5–5) 2 – 22 (11)	72 (70–80) 28 – 290 (5)	15 (15–15) 7– 50 (9)		16 (16–17) 4 – 31 (4)	29 (26–37) 12 – 108 (7)	12 (12–13) 0 – 55 (5)
Maximum crossable height of vertical structure (m)	Best L-U (n)	0.2 (0.2–0.2) 0.1 – 0.6 (11)		1.1 (1.1–1.2) 0.4 – 3.3 (9)		0.7 (0.7–0.7) 0.6 – 0.9 (4)	0.4 (0.4–0.4) 0.0 – 3.0 (7)	0.1 (0.1–0.1) 0.0 – 0.2 (5)
Minimum passable gap dimensions (m)	Best L-U (n)	0.1 (0.1–0.1) 0.0 – 0.1 (11)		0.3 (0.3–0.3) 0.1 – 0.7 (10)		0.3 (0.3–0.3) 0.2 – 0.3 (4)	0.1 (0.0–0.1) 0.0 – 0.1 (7)	0.2 (0.2–0.2) 0.1 – 0.4 (5)
Maximum crossable extent of waterbody (m)	Best L-U (n)	0.8 () 0.5 – 8.1 (11)	240 (236–263) 52–780 (5)	14 (14–37) 6 – 590 (9)			31 (29–40) 14 – 196 (7)	
Tolerable traffic flow during active period (vehicles/hr)	Best L-U (n)	7 (6–9) 4 – 27 (9)		8 (8–10) 3 – 28 (9)		6 (6–6) 2 – 13 (4)	13 (12–20) 4 – 43 (7)	
	Best	11 (11–14)		10 (9–13)		71 (69–71)		

Tolerable pedestrian traffic flow during active periods (pedestrians/hr)	L-U (n)	3 – 29 (11)		3 – 42 (9)			9 – 103 (4)	
Movement thresholds								
Typical movement distance within established home range/territory (m)	Best	58 (57–59)	200 (183–340)	529 (521–562)	406 (398–418)	1625 (1614–1647)	61 (55–75)	30 (30–33)
	L-U (n)	20 – 185 (9)	22 – 800 (5)	87 – 1620 (10)	158 – 813 (8)	800 – 3250 (4)	14 – 436 (7)	7 – 226 (5)
Typical capacity for movement outside of suitable habitat (m)	Best	33 (32–40)	214 (207–228)	100 (99–110)	977 (955–1129)	225 (222–237)	67 (63–81)	32 (32–37)
	L-U (n)	2 – 224 (9)	33 – 540 (5)	34 – 699 (10)	180 – 9503 (8)	75 – 700 (4)	9 – 350 (7)	13 – 340 (5)
Typical dispersal distance when seeking new home range/territory (m)	Best	69 (68–76)	110 (107–145)	765 (753–831)	825 (808–988)	1375 (1361–1414)	479 (441–720)	90 (88–112)
	L-U (n)	18 – 500 (9)	15 – 680 (5)	110 – 3730 (10)	210 – 7375 (8)	400 – 4000 (4)	76 – 2450 (7)	11 – 820 (5)

426

427 Discussion

428 We used the IDEA protocol of expert elicitation to address gaps in landscape-scale habitat
429 connectivity data that can limit the capacity of urban planning to adopt BSUD principles. Using the
430 city of Canberra in Ngunnawal Country (ACT) as a case study, we found that the IDEA protocol was
431 effective in this application – taxon-experts were able to estimate metrics describing connected
432 habitat for the taxon-groups, the estimates were ecologically meaningful and generally consistent
433 with empirical knowledge around habitat connectivity requirements from species within the groups
434 (where it existed), and the consultative process was generally useful in determining the relevancy of
435 metrics for specific groups (see examples below). However, there were also difficulties and
436 limitations of the approach. This included difficulty identifying ‘best’ estimates for individual metrics
437 at the taxon-group level where different species within the group were expected to have quite
438 different habitat requirements or movement capabilities. Overall, we consider our expert-derived
439 estimates of connected habitat to be a highly useful set of baseline data for habitat and connectivity
440 modelling and urban planning for a range of taxon groups. Below we discuss the strengths and
441 limitations of how our taxon-specific connected habitat estimates were determined for, and their
442 potential use, in urban planning and BSUD.

443 **Applicability of the IDEA protocol to estimate habitat connectivity metrics**

444 The connected habitat estimates we derived by applying the IDEA protocol for expert elicitation
445 were, in general, both ecologically meaningful and aligned with expert expectations. These estimates
446 contribute to the identified gaps in data for biodiversity-sensitive urban design - namely that the lack
447 of taxon group-level habitat connectivity data at the relevant spatial scale (Kirk *et al.* 2018) has been
448 addressed by defining habitat preferences with greater precision than is typically used in describing
449 habitat connectivity. For instance, our expert elicitation process derived a minimum and maximum
450 grass height, required percentage of native vegetation, and minimum width for core or corridor
451 habitat areas for grassland reptiles. This contrasts with the habitat description characterised simply
452 by “a grassy ground-cover free of trees” used in a similar application by Kirk *et al.* (2018). The
453 combination of these estimates also accurately described the specialised requirements of grassland
454 reptiles when compared to empirical data (Antos and Williams, 2015; Howland *et al.* 2016). Metrics
455 that we assessed also describe well the other taxon groups that are known to be more diverse and
456 adaptable in their connected habitat needs. For example, connected habitat for small–medium
457 mammals was estimated as not only including the presence of tree canopies and midstory cover, but
458 importantly, that preferred distances between those habitat elements are required to provide
459 functionally connected habitat for the majority of species considered. All taxon groups had nuance in
460 the specific spatial arrangement - for example native versus exotic composition, or tolerance of
461 particular habitat constraints - that were estimated quantitatively (e.g., tree spacing, tolerance of
462 artificial light) using the IDEA protocol. Important qualitative elements (e.g., the relative
463 heterogeneity or ‘clumped’ distribution of structural habitat elements) was also captured through
464 the ‘DISCUSS’ step of the IDEA protocol.

465 The breadth of metrics that could be collaboratively estimated through the IDEA protocol is a major
466 strength for addressing data gaps in urban planning. Habitat connectivity modelling largely relies on

467 a limited number of metrics, such as is in Kirk *et al.* (2018) where ecological connectivity was
468 determined for taxon groups from 4–5 structural metrics, 1–2 barrier metrics, and a single dispersal
469 metric. By using expert elicitation, we have generated quantitative estimates that describe taxon
470 group habitat connectivity using 16–27 metrics (mean = 21 metrics) that consider the functional
471 dimensions of connectivity by estimating up to eight ideal habitat metrics, 13 habitat constraint
472 metrics, six barriers to movement metrics, and four movement threshold metrics. Generating such a
473 breadth of data to inform connectivity metrics is particularly important for taxon groups with
474 complex and diverse habitat needs, such as amphibians that require both terrestrial and aquatic
475 environments (Becker *et al.* 2007). Further, our approach and breadth of metrics enabled
476 determination of the impact of anthropogenic processes on connectivity. For example, Kirk *et al.*
477 (2018) determined roads with greater than 5 m width as a barrier to amphibian movement, whereas
478 our approach separated two considerations of how paved roads presented a barrier to movement
479 (i.e., crossable extent of paved surface versus impact of traffic volume) and estimated amphibians
480 were able to cross much larger road (viz. “paved surfaces” best estimate = 29 m) when traffic flow
481 during active periods was low (<13 vehicles per hour during active periods). By using the IDEA
482 protocol, we have established a large collection of quantitative estimates to describe habitat
483 connectivity for a range of taxon groups in more detail and with greater context-dependency than is
484 typical in urban planning context.

485 Using the IDEA protocol to generate ecologically meaningful habitat connectivity estimates was not
486 without limitations, with some metrics proving more difficult to estimate than others. Some of the
487 difficulty that arose was due to lumping multiple species together based on broad habitat use, but
488 without being able to represent the diversity of habitat usage between individual species. This
489 constraint was most apparent for our riparian reptiles and mammals group, where the species
490 considered broadly require riparian and/or aquatic habitat elements, but vary widely on the relative
491 importance of each. For example, defining a minimum width of core habitat required consideration
492 of both aquatic habitat (more relevant for platypus and turtles) and associated terrestrial riparian
493 habitats (more relevant for water dragons and snakes). Depending on the specific subject matter
494 expertise of the experts, responses often focused on one or the other, rather than the combined
495 requirements for the full taxon group. Careful revision of expert estimates to identify variability in
496 metric interpretation by experts, coupled with more precise refinement of species comprising the
497 taxon groups themselves (e.g., adopting a process of identifying ‘dispersal guilds’ as described by
498 Lechner *et al.* 2017) could improve our methodology.

499 Wide tolerances among species within a taxon group created difficulties in providing representative
500 estimates, and contributed to broad confidence bounds for many metrics in this study. Typically, in
501 applying the IDEA protocol, the upper and lower estimates provided by experts represent ‘plausible
502 bounds’ around the ‘best’ estimate and may reflect something akin to a 95% confidence interval. In
503 this application however, the upper and lower bounds were adopted to reflect the variability
504 between, or tolerances within, species comprising the taxon group. For example, while experts
505 unanimously agreed that native-dominated vegetation was preferable in all habitats, all taxon
506 groups were considered able to tolerate non-native dominated vegetation to some extent (Threlfall
507 *et al.* 2016; 2017). As such, in many instances this meant the lower and upper estimates for ‘percent
508 native’ vegetation metrics were close to the full 0-100% range across different taxon groups.
509 Providing a best estimate for these metrics generally reflected one of three values: (a) the mid-point

510 of the full breath of tolerance within a taxon group (e.g., amphibians), (b) the maximum value
511 indicating that 100% native vegetation will always be 'best' (e.g., small freshwater fish), or (c) a
512 native-skewed estimate indicating native vegetation was likely better than exotic within the full
513 breath of compositional tolerance (e.g., all other groups). The way in which estimates were provided
514 as 'best', 'upper', and 'lower' in this study was based on our acknowledgement that estimating the
515 single 'true' value for metrics at the taxon group-level (i.e., across a range of species) would be less
516 ecologically meaningful than representing the within-group variability. To prevent overly broad
517 metric estimates in future, researchers could select species groupings which share greater ecological
518 dependencies (such as association with a vegetation community). Additionally, deciding whether to
519 use the upper and lower estimates to capture variability among species (as we did in estimating
520 tolerance bounds) or to capture the plausible range of the true value should be carefully considered.

521 Using the IDEA protocol enabled us to estimate metrics for which there is almost no research (e.g.,
522 tolerable levels of artificial light, or traffic volumes) with a similar level of confidence to metrics with
523 considerably more knowledge (e.g., those related to structural habitat requirements). For instance,
524 the credible interval around metrics with ACT-specific empirical studies (e.g., minimum grass height
525 for grassland reptiles, Howland *et al.* 2016) were comparable to metrics where there were no
526 species- or taxon-specific literature available (e.g., tolerable levels of artificial light). However, our
527 application of the IDEA protocol did not resolve issues around metric relevance for some taxon
528 groups, which resulted in some experts not contributing estimates, thereby decreasing our sample
529 size for some metric-taxon group combinations. This was most evident for the grassland reptile
530 metrics related to preferred distances between tree canopies, mature trees, and midstory canopies.
531 All experts agreed that the presence of trees and shrubs would inhibit these grassland specialists
532 (Antos and Williams, 2015; Howland *et al.* 2016), however some experts contributed estimates for
533 large distances between trees or shrubs to represent a sufficiently 'treeless' landscapes, while others
534 provided no response, deeming tree spacing to be irrelevant for the group. The exclusion of 'no
535 response' data may have artificially reduced the confidence limits around metrics where collectively
536 there was greater uncertainty. Previous studies have adopted the confidence score to reflect
537 experts' confidence that their 'best' estimate falls within their upper and lower bounds (as opposed
538 to how confident they are that their estimate is correct) which may be a way to encourage expert
539 responses in future studies. Since we adopted upper and lower estimates to reflect the breadth of
540 suitable habitats in this study, such an approach was not appropriate here. This example highlights
541 the importance of ensuring a consistent interpretation around individual metrics within the expert
542 group, either prior to experts providing initial estimates, or during the 'DISCUSS' step. Clarifying the
543 relative value of including or excluding metrics will avoid the need for subsequent qualitative
544 descriptions of expert intent.

545 **Capacity of estimated ecological connectivity metrics to inform spatial urban planning**

546 We investigated whether using the IDEA protocol could generate data inputs that could be used to
547 directly describe or model habitat connectivity to support urban planning and BSUD. Given the
548 strengths and minimal limitations we have identified for generating ecologically sensible estimates,
549 we consider our data is most useful in extending and refining what defines ecological connectivity in
550 an urban setting, thereby enabling for more precise and taxon-specific connectivity modelling and
551 mapping in the future.

552 We have estimated habitat connectivity over a broader set of metrics than is typically considered in
553 habitat connectivity assessments. However, a smaller set of metrics in previous studies may reflect
554 limited access to accompanying spatial modelling inputs at a suitable resolution, rather than authors
555 not considering other metrics to be important. For example, connected habitat models may consider
556 the presence of trees only without consideration of preferred spacing and composition because that
557 information is not available (Kirk et al, 2018, 2023). This means many of our estimated metrics may
558 only be useful as descriptions for urban planning (e.g., ACT Government, 2023), rather than
559 contributing directly to spatial modelling. Whereas Kirk et al. (2018) presents small bird connectivity
560 in an urban environment based on presence-absence data for four vegetation metrics with
561 accompanying spatial data, we present small bird connectivity as elicited quantitative threshold data
562 for 11 vegetation metrics, alongside minimum width of core and movement corridor habitat patch.
563 These additional metrics will be useful for wildlife managers to conceptualise and advise on
564 connected habitat, and will ideally contribute to predictive habitat and fragmentation mapping
565 where associated spatial layers are available. Where possible however, using the IDEA protocol to
566 increase the number of metrics considered will limit overestimates of connected habitat (through
567 greater incorporation of limiting aspects like urban heat or light) and also underestimates (through
568 incorporating more nuance in important elements like the interaction of road width and traffic
569 volume), thereby providing more representative connected habitat model outputs overall.

570 A final strength of the IDEA protocol is that in estimating lower and upper bounds for metrics, there
571 is flexibility to explore different scenarios and contexts in habitat connectivity modelling and
572 mapping (Hanea *et al.* 2017; Hemming *et al.* 2018). This contrasts with the classical approach of
573 obtaining a single data input through behavioural aggregation of experts (O'Hagan *et al.* 2005;
574 Hanea *et al.* 2017), where habitat would be considered connected or disconnected based on the
575 'best' value only for any particular habitat metric. For example, connectivity for woodland birds in
576 Kirk *et al.* (2018) was modelled using a median dispersal distance of 1.5 km. Our best estimates for
577 typical movement within a territory (1.6 km) or typical dispersal distance when seeking a new
578 territory (1.4 km) for the same taxon group meant the results from our expert elicitation were not
579 dissimilar to those used in Kirk *et al.* (2018). However, the upper bounds provided by experts in our
580 study determined that some small woodland birds are potentially capable of moving up to three-
581 times further than the distance described as the best estimate, meaning connectivity or the
582 minimum requirements for dispersal for some species in the group is likely to be underestimated by
583 adopting only the 'best' reported value in habitat connectivity models.

584 **Conclusion**

585 Maintenance of habitat connectivity through the conservation of habitat and wildlife corridors
586 across urban landscapes is important for promoting biodiversity, including for many threatened
587 species which occur within urban extents (Ives *et al.* 2016; Garrard *et al.* 2018; Soanes and Lentini
588 2019). To identify, retain, and restore habitat and wildlife corridors to facilitate dispersal within
589 urban landscapes requires species- or taxon-specific knowledge of their ecological connectivity
590 requirements including movement abilities, habitat preferences, and potential barriers to dispersal
591 (Kirk *et al.*, 2018). Using the habitat connectivity estimates we quantified through an expert-
592 elicitation process, there is a clear opportunity to identify congruency among taxon group

593 requirements to establish urban planning and BSUD approaches that have positive effects for a
594 range of taxa (ACT Government 2023). For example, multiple species groups shared a preferred tree
595 spacing of 11–41 m, and hence the conservation of such structural elements within core habitats
596 (≥ 328 m wide) or corridors (≥ 39 m wide) will support habitat connectivity for all terrestrial groups
597 except grassland reptiles. The lack of congruency between grassland reptile habitat and that of other
598 taxon groups in this study highlights the importance of identifying taxon group-level dependencies
599 where differing ecosystems overlap or co-occur. Specific to this case study in Canberra, this will
600 involve understanding the requirements of aquatic and riparian associated fauna (i.e., amphibians,
601 riparian reptile and mammals, and freshwater fish), woodland associated fauna (i.e., native bees,
602 small–medium mammals, woodland birds, and amphibians), and grassland-associated fauna (i.e.,
603 native bees, grassland reptiles, small–medium mammals, and amphibians) and identifying a spatially
604 explicit conservation network which adequately provides for the protection and restoration of
605 connected habitat to meet the needs of all. Applying these results and BSUD in future urban
606 planning offers an opportunity to validate estimates through targeted monitoring of the taxon
607 groups. Using our approach, expert estimates can harness congruency among taxon groups to
608 maximise co-benefits and identify where additional conservation measures are required to conserve
609 habitats which are not shared by multiple species assemblages (Gordon *et al* 2009).

610 The IDEA protocol provided quantitative information on taxon-specific habitat requirements and
611 constraints in data-deficient contexts and enabled robust consideration of functional constraint data
612 (e.g., behavioural barriers) in our definitions of connected habitat. This enabled us to address the
613 two limitations of applying BSUD identified by Kirk *et al.* (2018; 2021; 2023). Through reviewing the
614 applicability of the IDEA protocol and assessing expert estimates, we identified that taxon-group
615 variability and an occasional lack of consistency around metric relevance or interpretation limited
616 the clarity around how to best interpret and apply estimates for habitat connectivity. We have
617 discussed how these limitations can be addressed in future uses of expert elicitation in similar
618 contexts. Applying these data to the calculation of connectivity indices (e.g., the City Biodiversity
619 Index) would benefit from further investigation and validation of scenario-based assumptions
620 through field-based assessments of species distribution (Kirk *et al.* 2018), as well as the creation of
621 relevant spatial layers. The application of the IDEA protocol to provide greater detail around habitat
622 connectivity metrics in this study is anticipated to represent broad benefits for urban planning and
623 developing BSUD frameworks in cities into the future.

624 **Acknowledgements**

625 We acknowledge the Ngunnawal people as Canberra’s first inhabitants and scientists, and the
626 Traditional Custodians of the land where we undertook our study, and we recognise any other
627 peoples or families with connection to the lands of the ACT region. This work included contributions
628 from people residing and having knowledge from the lands of the Anaiwan, Awabakal,
629 Boonwurrung, Dharug, Eora, Gulidjan, Kuring-Gai, Ngarigo, Tharawal, Wajuk, Wathaurong, Waveroo,
630 Whadjuk Noongar, Woiworung, and Wurundjeri People of the Kulin Nations. We acknowledge and
631 pay our respects to Elders and Traditional Custodians of these lands – both past and present – and
632 acknowledge their continuing cultures and connection to Country. We thank H. Kirk, C. Threlfall and
633 K. Soanes for their positive and enthusiastic engagement with the ACT Government for applying the

634 Linking Nature in the City protocol within the urban extent of the Australian Capital Territory. We
635 thank T. Armstrong, D. Chapple, D. Coleman, M. Mulvaney, K. Smith, and M. Wong for their valuable
636 species-specific knowledge contributing to expert elicitation and T. Armstrong, P. Arnold, J. Camac,
637 A. Hanea, C. Malam, N. McLean, A. Nicotra, M. Stewart for their helpful suggestions on the expert
638 elicitation and earlier versions of this manuscript.

639 Author Contributions

640 **Stephanie K. Courtney Jones:** Conceptualization; data curation; formal analysis; investigation;
641 methodology; project administration; supervision; validation; visualization; writing – original draft;
642 writing – reviewing and editing. **Luke S. O’Loughlin:** Conceptualization; data curation; investigation;
643 methodology; project administration; visualization; writing – original draft; writing – reviewing and
644 editing. **Danswell Starrs:** Conceptualization; data curation; formal analysis; investigation;
645 methodology; visualisation; writing – reviewing and editing. **Jacinta E. Humphrey:** Conceptualisation;
646 investigation; validation; writing – original draft; writing – reviewing and editing. **Stephanie A.**
647 **Pulsford:** Conceptualization; data curation; formal analysis; investigation; writing – reviewing and
648 editing. **Melissa Snape:** Conceptualization; data curation; formal analysis; investigation;
649 methodology; project administration; supervision; validation; visualization; writing – reviewing and
650 editing. **All other authors:** Conceptualization; investigation; writing – reviewing and editing.

651 References

- 652 ABS (2022) Australia bureau of statistics. Regional population 2021 – Australian Capital Territory.
653 Available at: [https://www.abs.gov.au/statistics/people/population/regional-](https://www.abs.gov.au/statistics/people/population/regional-population/2021#australian-capital-territory)
654 [population/2021#australian-capital-territory](https://www.abs.gov.au/statistics/people/population/regional-population/2021#australian-capital-territory) (accessed 11 January 2023)
- 655 ACT Government (2018) ACT Planning Strategy 2018. Report available at:
656 [https://www.planning.act.gov.au/_data/assets/pdf_file/0007/1285972/2018-ACT-](https://www.planning.act.gov.au/_data/assets/pdf_file/0007/1285972/2018-ACT-Planning-Strategy.pdf)
657 [Planning-Strategy.pdf](https://www.planning.act.gov.au/_data/assets/pdf_file/0007/1285972/2018-ACT-Planning-Strategy.pdf)
- 658 ACT Government (2023) ACT Biodiversity Sensitive Urban Design Guide. Available at:
659 [https://www.planning.act.gov.au/_data/assets/pdf_file/0008/2279996/ACT-Biodiversity-](https://www.planning.act.gov.au/_data/assets/pdf_file/0008/2279996/ACT-Biodiversity-Sensitive-Urban-Design-Guide.pdf)
660 [Sensitive-Urban-Design-Guide.pdf](https://www.planning.act.gov.au/_data/assets/pdf_file/0008/2279996/ACT-Biodiversity-Sensitive-Urban-Design-Guide.pdf)
- 661 Adams-Hosking, C., McBride, M.F., Baxter, G., Burgman, M., De Villiers, D., Kavanagh, R., Lawler, I.,
662 Lunney, D., Melzer, A., Menkhorst, P. and Molsher, R. (2016). Use of expert knowledge to elicit
663 population trends for the koala (*Phascolarctos cinereus*). *Diversity and Distributions*, 22(3), 249-262.
- 664 Alexandra, J., Norman, B., Steffen, W., & Maher, W. (2017). Planning and Implementing Living
665 Infrastructure in the Australian Capital Territory–Final Report. Canberra: Canberra urban and
666 regional futures, University of Canberra.
- 667 Alexandra, J., & Norman, B. (2020). The city as forest-integrating living infrastructure, climate
668 conditioning and urban forestry in Canberra, Australia. *Sustainable Earth*, 3(1), 1-11.

669 Antos M and Williams NSG 2015. The wildlife of our grassy landscapes, in Land of sweeping plains:
670 managing and restoring the native grasslands of south-eastern Australia, eds NSG Williams, A
671 Marshall, JW Morgan (CSIRO Publishing, Clayton South): pp. 87–114.

672 Becker, C. G., Fonseca, C. R., Haddad, C. F. B., Batista, R. F., & Prado, P. I. (2007). Habitat split and the
673 global decline of amphibians. *Science*, 318(5857), 1775-1777.

674 Buckmaster, A. J., Osborne, W. S., & Webb, N. (2010). The loss of native terrestrial small mammals in
675 large urban reserves in the Australian Capital Territory. *Pacific Conservation Biology*, 16(1), 36-45.

676 Camac, J. S., Umbers, K. D. L., Morgan, J. W., Geange, S. R., Hanea, A., Slatyer, R. A., McDougall, K. L.,
677 Venn, S. E., Vesk, P. A., Hoffmann, A. A., & Nicotra, A. B. (2021). Predicting species and community
678 responses to global change using structured expert judgement: An Australian mountain ecosystems
679 case study. *Global Change Biology*, 27, 4420– 4434.

680 Chan, L., Hillel, O., Elmqvist, T., Werner, P., Holman, N., Mader, A., & Calcaterra, E. (2014). User's
681 manual on the Singapore index on cities' biodiversity (also known as the City Biodiversity Index).
682 Singapore: National Parks Board, Singapore.

683 Courtney Jones, S.K., Geange, S., Hanea, A., Camac, J., Hemming, V., Doobov, B., Leigh, A., Nicotra, A.
684 (2023) IDEAcology: an interface to streamline and facilitate efficient, rigorous expert elicitation in
685 ecology. *Methods in Ecology and Evolution*, 14(8), 2019-2028

686 Deslauriers, M. R., Asgary, A., Nazarnia, N., & Jaeger, J. A. (2018). Implementing the connectivity of
687 natural areas in cities as an indicator in the City Biodiversity Index (CBI). *Ecological Indicators*, 94, 99-
688 113.

689 Doerr, V. A. J., Doerr, E. D., & Davies, M. J. (2010). Does structural connectivity facilitate dispersal of
690 native species in Australia's fragmented terrestrial landscapes. *CEE Rev*, 8, 70.

691 Doerr, E. D., Doerr, V. A., Davies, M. J., & McGinness, H. M. (2014). Does structural connectivity
692 facilitate movement of native species in Australia's fragmented landscapes?: a systematic review
693 protocol. *Environmental Evidence*, 3(1), 1-8.

694 Dugdale, T. M., Hunt, T. D., & Clements, D. (2013). Aquatic weeds in Victoria: Where and why are
695 they a problem, and how are they being controlled? *Plant Protection Quarterly*, 28(2), 35–41.

696 Fraser, H, Simmonds, JS, Kutt, AS, Maron, M. Systematic definition of threatened fauna communities
697 is critical to their conservation. *Divers Distrib*. 2019; 25: 462– 477.

698 Garrard, G. E., Williams, N. S., Mata, L., Thomas, J., & Bekessy, S. A. (2018). Biodiversity sensitive
699 urban design. *Conservation Letters*, 11(2), e12411.

700 Gaston, K. J., Davies, T. W., Bennie, J., & Hopkins, J. (2012). Reducing the ecological consequences of
701 night-time light pollution: options and developments. *Journal of Applied Ecology*, 49(6), 1256-1266.

702 Geyle H M., Tingley R, Amey A P., Cogger H, Couper P J., Cowan M, Craig M D., Doughty P, Driscoll D
703 A., Ellis R J., Emery J-P, Fenner A, Gardner M G., Garnett S T., Gillespie G R., Greenlees M J., Hoskin C
704 J., Keogh J. S, Lloyd R, Melville J, McDonald P J., Michael D R., Mitchell N J., Sanderson C, Shea G M.,
705 Sumner J, Wapstra E, Woinarski J. C. Z., Chapple D G. (2021) Reptiles on the brink: identifying the
706 Australian terrestrial snake and lizard species most at risk of extinction. *Pacific Conservation Biology*
707 27, 3-12.

708 Gordon, A., Simondson, D., White, M., Moilanen, A., & Bekessy, S. A. (2009). Integrating
709 conservation planning and landuse planning in urban landscapes. *Landscape and urban planning*,
710 91(4), 183-194.

711 Hale, R., Coleman, R., Pettigrove, V., & Swearer, S. E. (2015). Identifying, preventing and
712 mitigating ecological traps to improve the management of urban aquatic ecosystems. *Journal of*
713 *Applied Ecology*, 52(4), 928-939.

714 Hanea, A.M., McBride, M.F., Burgman, M.A., Wintle, B.C., Fidler, F., Flander, L., Twardy, C.R.,
715 Manning, B. and Mascaro, S., 2017. Investigate Discuss Estimate Aggregate for structured expert
716 judgement. *International Journal of Forecasting*, 33(1), pp.267-279.

717 Hemming, V., Burgman, M.A., Hanea, A.M., McBride, M.F. and Wintle, B.C., 2018. A practical guide
718 to structured expert elicitation using the IDEA protocol. *Methods in Ecology and Evolution*, 9(1),
719 pp.169-180.

720 Hemming, V., Hanea, A. M., Walshe, T., & Burgman, M. A. (2020). Weighting and aggregating expert
721 ecological judgments. *Ecological Applications*, 30(4), e02075.

722 Hemming, V., Hanea, A. M., & Burgman, M. A. (2022). What is a good calibration question?. *Risk*
723 *Analysis*, 42(2), 264-278.

724 Howland, B. W., Stojanovic, D., Gordon, I. J., Fletcher, D., Snape, M., Stirnemann, I. A., &
725 Lindenmayer, D. B. (2016). Habitat preference of the striped legless lizard: implications of grazing by
726 native herbivores and livestock for conservation of grassland biota. *Austral Ecology*, 41(4), 455-464.

727 Huang, C. W., McDonald, R. I., & Seto, K. C. (2018). The importance of land governance for
728 biodiversity conservation in an era of global urban expansion. *Landscape and Urban Planning*, 173,
729 44-50.

730 Ikin, K., Le Roux, D.S., Rayner, L., Villaseñor, N.R., Eyles, K., Gibbons, P., Manning, A.D. and
731 Lindenmayer, D.B. (2015), Key lessons for achieving biodiversity-sensitive cities and towns. *Ecol*
732 *Manag Restor*, 16: 206-214.

733 Ives, C.D., Lentini, P.E., Threlfall, C.G., Ikin, K., Shanahan, D.F., Garrard, G.E., Bekessy, S.A., Fuller,
734 R.A., Mumaw, L., Rayner, L. and Rowe, R., 2016. Cities are hotspots for threatened species. *Global*
735 *Ecology and Biogeography*, 25(1), pp.117-126.

736 Kirk, H., Threlfall, C., Soanes, K., Estima Ramalho, C., Parris, K., Amati, M., Bekessy, S., & Mata, L.
737 (2018). Linking Nature in the city: a framework for improving ecological connectivity across the City
738 of Melbourne. Report prepared by the CAUL hub for the City of Melbourne Urban Sustainability
739 Branch. National Environmental Science Programme.

740 Kirk H., Threlfall C., Soanes K. & Parris, K. (2020) Linking Nature in the City Part Two: Applying the
741 Connectivity Index. Report prepared for the City of Melbourne Urban Sustainability Branch. National
742 Environmental Science Programme.

743 Kirk, H., Garrard, G.E., Croeser, T., Backstrom, A., Berthon, K., Furlong, C., Hurley, J., Thomas, F.,
744 Webb, A. and Bekessy, S.A., 2021. Building biodiversity into the urban fabric: A case study in applying
745 Biodiversity Sensitive Urban Design (BSUD). *Urban Forestry & Urban Greening*, 62, p.127176.

746 Kirk, H., Soanes, K., Amati, M., Bekessy, S., Harrison, L., Parris, K., Ramalho, C., van de Ree, R. &
747 Threlfall, C. (2023). Ecological connectivity as a planning tool for the conservation of wildlife in cities.
748 *MethodsX*, 101989.

749 Lechner, A. M., Harris, R. M., Doerr, V., Doerr, E., Drielsma, M., & Lefroy, E. C. (2015). From static
750 connectivity modelling to scenario-based planning at local and regional scales. *Journal for Nature*
751 *Conservation*, 28, 78-88.

752 Lechner, A. M., Sprod, D., Carter, O., & Lefroy, E. C. (2017). Characterising landscape connectivity for
753 conservation planning using a dispersal guild approach. *Landscape Ecology*, 32, 99-113.

754 S. Legge, D. B. Lindenmayer, N. M. Robinson, B. C. Scheele, D. M. Southwell, & B. A. Wintle (Eds.).
755 (2018). *Monitoring threatened species and ecological communities*. Melbourne, Australia: CSIRO.

756 Lindenmayer, D. B., Lane, P. W., Westgate, M. J., Crane, M., Michael, D., Okada, S., & Barton, P. S.
757 (2014). An empirical assessment of the focal species hypothesis. *Conservation Biology*, 28(6), 1594-
758 1603.

759 Maclagan, S. J., Coates, T., & Ritchie, E. G. (2018). Don't judge habitat on its novelty: Assessing the
760 value of novel habitats for an endangered mammal in a peri-urban landscape. *Biological*
761 *Conservation*, 223, 11–18.

762 Martin, T. G., Burgman, M. A., Fidler, F., Kuhnert, P. M., Low-Choy, S., McBride, M., & Mengersen, K.
763 (2012). Eliciting expert knowledge in conservation science. *Conservation Biology*, 26, 29–38

764 McDonald, R. I., Kareiva, P., & Forman, R. T. (2008). The implications of current and future
765 urbanization for global protected areas and biodiversity conservation. *Biological conservation*,
766 141(6), 1695-1703.

767 McDonald, R.I., Mansur, A.V., Ascensão, F., Crossman, K., Elmqvist, T., Gonzalez, A., Güneralp, B.,
768 Haase, D., Hamann, M., Hillel, O. and Huang, K., 2020. Research gaps in knowledge of the impact of
769 urban growth on biodiversity. *Nature Sustainability*, 3(1), pp.16-24.

770 McKinney, M. L. (2008). Effects of urbanization on species richness: a review of plants and animals.
771 *Urban ecosystems*, 11(2), 161-176.

772 O'Hagan, A., Buck, C. E., Daneshkhah, A., Eiser, J. R., Garthwaite, P. H., Jenkinson, D. J., Oakley, J. and
773 Rakow, T. (2006). Uncertain judgements: eliciting experts' probabilities.

774 Page, S. E. (2008). *The Difference: How the power of diversity creates better groups, firms, schools,*
775 *and societies*. Princeton University Press. doi:10.5860/choice.45-1534

776 Peden, L., Skinner, S., Johnston, L., Frawley, K., Grant, F., and Evans, L. 2011. *Survey of Vegetation*
777 *and Habitat in Key Riparian Zones in Tributaries of the Murrumbidgee River in the ACT: Cotter,*
778 *Molonglo, Gudgenby, Naas and Paddys Rivers*. Technical Report 23. Environment and Sustainable
779 *Development Directorate, Canberra*.

780 Page, S. E. (2008). *The difference: How the power of diversity creates better groups, firms, schools,*
781 *and societies*. New Jersey: Princeton University Press.

782 R Core Team, 2022. *R: A language and environment for statistical computing*. Vienna.
783 <https://www.R-project.org/>.

784 Rayner, L., Ikin, K., Evans, M.J., Gibbons, P., Lindenmayer, D.B. and Manning, A.D. (2015), Avifauna
785 and urban encroachment in time and space. *Diversity Distrib.*, 21: 428-440.

786 Scheele, B.C., Legge, S., Armstrong, D.P., Copley, P., Robinson, N., Southwell, D., Westgate, M.J. and
787 Lindenmayer, D.B., 2018. How to improve threatened species management: An Australian
788 perspective. *Journal of Environmental Management*, 223, pp.668-675.

789 Selinske, M. J., Bekessy, S. A., Geary, W. L., Faulkner, R., Hames, F., Fletcher, C., Squires, Z. E., &
790 Garrard, G. E. Projecting biodiversity benefits of conservation behavior-change programs.
791 *Conservation Biology*, 2022; 36:e13845

792 Seto, K. C., Güneralp, B., & Hutyrá, L. R. (2012). Global forecasts of urban expansion to 2030 and
793 direct impacts on biodiversity and carbon pools. *Proceedings of the National Academy of Sciences*,
794 109(40), 16083-16088.

795 Shearer, A. W. (2005). Approaching scenario-based studies: three perceptions about the future and
796 considerations for landscape planning. *Environment and planning B: Planning and Design*, 32(1), 67-
797 87.

798 Soanes, K., & Lentini, P. E. (2019). When cities are the last chance for saving species. *Frontiers in
799 Ecology and the Environment*, 17(4), 225-231.

800 Speirs-Bridge, A., Fidler, F., McBride, M., Flander, L., Cumming, G., & Burgman, M. (2010). Reducing
801 overconfidence in the interval judgments of experts. *Risk Analysis: An International Journal*, 30(3),
802 512-523.

803 Theron, K.J., Pryke, J.S. & Samways, M.J. (2022). Maintaining functional connectivity in grassland
804 corridors between plantation forests promotes high-quality habitat and conserves range restricted
805 grasshoppers. *Landsc Ecol* 37, 2081–2097.

806 Threlfall, C. G., Ossola, A., Hahs, A. K., Williams, N. S., Wilson, L., & Livesley, S. J. (2016). Variation in
807 vegetation structure and composition across urban green space types. *Frontiers in Ecology and
808 Evolution*, 4, 66.

809 Threlfall, C. G., Mata, L., Mackie, J. A., Hahs, A. K., Stork, N. E., Williams, N. S., & Livesley, S. J. (2017).
810 Increasing biodiversity in urban green spaces through simple vegetation interventions. *Journal of
811 applied ecology*, 54(6), 1874-1883.

812 Tremblay, M.A. and St. Clair, C.C. (2009), Factors affecting the permeability of transportation and
813 riparian corridors to the movements of songbirds in an urban landscape. *Journal of Applied Ecology*,
814 46: 1314-1322

815 Wang, K., Wang, T., & Liu, X. (2018). A review: Individual tree species classification using integrated
816 airborne LiDAR and optical imagery with a focus on the urban environment. *Forests*, 10(1), 1.

817 Watson DM, Doerr VA, Banks SC, Driscoll DA, van der Ree R, Doerr ED, Sunnucks P. (2017)
818 Monitoring ecological consequences of efforts to restore landscape scale connectivity. *Biological
819 Conservation* 206: 201-209.

820 Westgate, M. J., B. C. Scheele, K. Ikin, A. M. Hoefler, R. M. Beaty, M. Evans, W. Osborne, D. Hunter, L.
821 Rayner, and D. A. Driscoll. 2015. Citizen science program shows urban areas have lower occurrence
822 of frog species, but not accelerated declines. *PLoS ONE* 10:e0140973

- 823 Wittmann, M. E., Cooke, R. M., Rothlisberger, J. D., Rutherford, E. S., Zhang, H., Mason, D. M., &
824 Lodge, D. M. (2015). Use of structured expert judgment to forecast invasions by bighead and silver
825 carp in Lake Erie. *Conservation Biology*, 29, 187–197.
- 826 Yates, C., S. Hopper, and R. Taplin. 2005. Native insect flower visitor diversity and feral honeybees on
827 jarrah (*Eucalyptus marginata*) in Kings Park, an urban bushland remnant. *Journal of the Royal Society*
828 *of Western Australia* 88: 147– 153.