

1 **Incomplete lineage sorting and hybridization underlie tree discordance in *Petunia* and related genera**
2 **(*Petunieae*, *Solanaceae*)**

3
4 Pedro H. Pezzi^{1,*}, Lucas C. Wheeler², Loreta B. Freitas¹, Stacey D. Smith²

5
6 ¹Department of Genetics, Universidade Federal do Rio Grande do Sul, Porto Alegre, Brazil;

7 ²Department of Ecology and Evolutionary Biology, University of Colorado, Boulder, USA

8
9 * pedrohenriquepezzi@gmail.com

10
11 **Abstract**

12 Despite the overarching history of species divergence, phylogenetic studies often reveal distinct
13 topologies across regions of the genome. The sources of these gene tree discordances are
14 variable, but incomplete lineage sorting (ILS) and hybridization are among those with the most
15 biological importance. *Petunia* serves as a classic system for studying hybridization in the wild.
16 While field studies suggest that hybridization is frequent, the extent of reticulation within
17 *Petunia* and its closely related genera has never been examined from a phylogenetic perspective.
18 In this study, we used transcriptomic data from 11 *Petunia*, 16 *Calibrachoa*, and 10 *Fabiana*
19 species to illuminate the relationships between these species and investigate whether
20 hybridization played a significant role in the diversification of the clade. We inferred that gene
21 tree discordance within genera is linked to hybridization events along with high levels of ILS due
22 to their rapid diversification. Moreover, network analyses estimated deeper hybridization events
23 between *Petunia* and *Calibrachoa*, genera that have different chromosome numbers. Although
24 these genera cannot hybridize at the present time, ancestral hybridization could have played a
25 role in their parallel radiations, as they share the same habitat and life history.

26
27 **Introduction**

28 Over the last two decades, systematic biology has experienced significant advances, managing to
29 work with vast volumes of data and constructing robust phylogenies to elucidate species'
30 relationships and evolutionary histories. Nevertheless, using different methods and datasets (i.e.,
31 genetic markers and sampling schemes) often results in conflicting tree topologies. These
32 discrepancies may stem from errors in model specifications, data processing, or evolutionary
33 processes such as incomplete lineage sorting (ILS) and hybridization (Galtier and Daubin, 2008).
34 Coalescent-based methods are commonly employed to mitigate conflicts in trees caused by ILS,
35 such as anomaly zones, where the topology of 'anomalous gene trees' with short branch lengths
36 differs from the species tree topology (Degnan and Rosenberg, 2006). However, these methods
37 are unreliable in situations involving gene flow among lineages (Solís-Lemus et al., 2016).
38 Despite the advances in phylogenomic methods that account for ILS and gene flow (Hibbins and
39 Hahn, 2022), detecting and distinguishing between such events remains a complex task that
40 heavily depends on the extent to which they occur (Kong and Kubatko, 2021).

41 Botanists already recognize that plant evolution likely follows a web-like pattern due to
42 the numerous examples of plant hybridization (Stull et al., 2023). However, the potential
43 outcomes of such events are highly variable (Abbott et al., 2016; Soltis and Soltis, 2009).
44 Hybridization can facilitate speciation through novel trait combinations or polyploidization
45 (Abbott et al., 2013), lead to extinction through genetic swamping (Todesco et al., 2016), or
46 introgress adaptive alleles (Suarez-Gonzalez et al., 2018). Regardless of the outcomes,

47 hybridization is a frequent evolutionary phenomenon at both shallow (Nevado et al., 2018) and
48 deep timescales (Rothfels et al., 2015) with extensive impacts on plant diversification and
49 evolution (Goulet et al., 2017; Whitney et al., 2010).

50 The *Petunia-Calibrachoa-Fabiana* Solanaceae clade presents a unique opportunity to
51 investigate the evolutionary dynamics of plant diversification in southern South America.
52 Notably, *Petunia* has become an important taxon to understand the role of hybridization in the
53 region due to the multiple hybrid zones documented (e.g., Binaghi et al., 2023; Caballero-
54 Villalobos et al., 2021; Giudicelli et al., 2019). Lineages in the *Petunia-Calibrachoa-Fabiana*
55 clade have experienced rapid diversification over the past ca. 20 million years (Zuntini et al.,
56 2024), with approximately 65 species exhibiting a diverse range of pollination syndromes and
57 inhabiting distinct biomes, from rainforests to savannahs and deserts (e.g., Alaria et al., 2022,
58 Mäder and Freitas, 2019; Reck-Kortmann et al., 2014). While *Petunia* and *Calibrachoa* are very
59 similar in morphology and ecological conditions, *Fabiana* stands out due to its drastic
60 differences from related genera: its species are xerophytic, have small flowers, and display
61 reduced or even absent leaves (Alaria et al., 2022). The three genera are classified within the
62 well-established tribe Petunieae, forming a strongly supported clade (Särkinen et al., 2013;
63 Wheeler et al., 2022). However, the internal relationships within this clade remain controversial.
64 Phylogenetic studies using distinct genetic markers, sampling schemes, and analytical
65 approaches have produced conflicting topologies (Fig. 1). Some place *Petunia* as a sister to the
66 remaining genera (Alaria et al., 2022; Olmstead et al., 2008; Wheeler et al., 2022), whereas
67 others place it as a sister genus to *Calibrachoa* (Särkinen et al., 2013) or to *Fabiana* (Mäder and
68 Freitas, 2019; Reck-Kortmann et al., 2015). These incongruences might be attributed to the
69 substantial levels of ILS due to their rapid diversification (Särkinen et al., 2013; Wheeler et al.,
70 2022), or potentially from instances of ancient hybridization.

71 The likelihood of hybridization depends on how effective reproductive barriers are at
72 preventing gene flow, and plants typically rely on a combination of barriers to achieve complete
73 reproductive isolation (Baack et al., 2015; Christie et al., 2022). In *Petunia*, gene flow is
74 primarily prevented by prezygotic barriers, including geographic and floral isolation, with
75 postzygotic barriers playing a negligible role (Dell'Olivo et al., 2011). These barriers have been
76 extensively studied in *Petunia*, which established this genus as a model in plant hybridization
77 and pollination studies (Binaghi et al., 2023; Gübitz et al., 2009; Rodrigues et al., 2018;
78 Turchetto et al., 2019). However, a comprehensive investigation into *Calibrachoa* and *Fabiana*,
79 as well as the possibility of hybridization causing tree discordance among genera and congeneric
80 species has yet to be examined under a solid phylogenetic framework.




Figure 1. Distribution map of *Petunia* (blue outline), *Calibrachoa* (pink shaded area), and *Fabiana* (green shaded area) in South America. Map dots represent sampled localities, colored by genus, except for the species collected at greenhouses. Representatives of each genus are shown in circles: *Petunia altiplana*, *Calibrachoa eglandulata*, and *Fabiana bryoides* (photos: Lucas C. Wheeler). Phylogenetic relationships of the genera are presented on the right. Olmstead et al., (2008) used two plastid markers; Särkinen et al., (2013) used two nuclear and five plastid markers; Reck-Kortmann et al., (2015) used one nuclear and two plastid markers; Mäder et al., (2019) used eight nuclear and eight plastid markers; and Alaria et al., (2022) used one nuclear and three plastid markers.

81 Even though *Petunia* and *Calibrachoa* are similar in flower morphology, ecology, and
 82 geographic distribution (Fig. 1), they have been split into two different genera due to their
 83 chromosome numbers: *Petunia* has seven chromosome pairs ($2n = 14$), whereas *Calibrachoa* has
 84 nine ($2n = 18$) (Stehmann et al., 2009; Wijsman and De Jong, 1985). The persistence in nature of
 85 hybrids between species that have different chromosome numbers is unlikely as it leads to
 86 meiotic mispairing—unless it involves polyploidization (Alix et al., 2017; Hegarty and Hiscock,

87 2008). To date, polyploidization has never been observed in *Petunia* or *Calibrachoa*. Hence, the
88 occurrence of hybrids between *Petunia* and *Calibrachoa* in the wild seems unlikely, even though
89 some species occur in sympatry. While intergeneric hybrids known as “Petchoa” have been
90 developed and are available commercially, these hybrids are sterile, and their creation requires
91 significant human intervention (Shaw, 2007). In contrast, while *Calibrachoa* and *Fabiana* share
92 the same chromosome count, which theoretically would allow successful meiosis in the hybrid,
93 their disjunct geographical distribution and distinct life histories serve as strong present-day
94 barriers that prevent gene flow.

95 In this study, we used floral transcriptome data from *Petunia*, *Calibrachoa*, and *Fabiana*
96 species to investigate the sources of discordance among phylogenetic trees. Specifically, we
97 aimed to evaluate the influence of ILS and reticulate evolution on the diversification of these
98 genera. We hypothesized that hybridization occurs frequently within genera, both in recent times
99 and throughout their evolutionary history, contributing to the observed phylogenetic discordance
100 within genera. Moreover, we tested whether intergeneric hybridization could have played a role
101 in the diversification of the clade. We predicted that intergeneric hybridization is unlikely due to
102 robust reproductive barriers, including chromosome number differences and geographic
103 isolation.

104 **Material and Methods**

105 *Taxa sampling and transcriptome data processing*

106 We used the raw RNA-seq data from Wheeler et al., (2022, 2023), focusing on the *Petunia*-
107 *Calibrachoa*-*Fabiana* clade and incorporating six outgroup species. In total, we employed 107
108 individuals, encompassing 11 *Petunia* species, 16 *Calibrachoa* species, 10 *Fabiana* species, and
109 six outgroups (Table S1). Here, we expanded the Wheeler et al., (2022) dataset by including
110 additional individuals for most sampled species derived from Wheeler et al., (2023), resulting in
111 three individuals per species collected at the same time and location (hereafter referred to as
112 replicates). We corrected the raw RNA-seq reads using Rcorrector (Song and Florea, 2015) and
113 removed adapters using Trimmomatic (Bolger et al., 2014). Subsequently, we mapped the reads
114 against the 3,672 protein-coding genes from conspecific transcriptomes, which were assembled
115 for replicate 1 by Wheeler et al., (2022) using BWA (Li and Durbin, 2010). Consensus fasta
116 sequences were generated through samtools 1.16 (Li et al., 2009) by calling the most frequent
117 base (-m simple) and then aligned with MACSE 2.06 (Ranwez et al., 2018).

118 *Phylogenetic analyses and evaluation of tree discordance*

119 We employed three distinct approaches to elucidate the phylogenetic relationships among species
120 within the *Petunia*-*Calibrachoa*-*Fabiana* clade. Firstly, we estimated the maximum likelihood
121 (ML) gene trees using the GTR+ Γ model along with 1,000 bootstrap replicates in RAxML
122 (Stamatakis, 2014) and estimated the species tree—both with and without assigning individuals
123 to species—using ASTRAL III 5.7.8 (Rabiee et al., 2019; Zhang et al., 2018). Secondly, we
124 constructed a supermatrix by concatenating the fasta alignments with the *SuperMatrix* function
125 of the evobiR R-package (Jonika et al., 2023). This supermatrix was then used to generate a
126 maximum likelihood species tree using IQTree 1.6.12 (Nguyen et al., 2015) setting the GTR+ Γ
127 model to each partition with 1,000 bootstrap replicates. Lastly, we estimated a species tree using
128 SVDQuartets, a coalescent method originally designed for SNP data but also effective with
129 multi-locus alignments (Chifman and Kubatko, 2014), implemented in PAUP* 4a (Swofford,
130 2003), which infers relationships among quartets and subsequently summarizes these

133 relationships into a species tree. We set the outgroups, assigned individuals to respective species,
134 and assessed all quartets (*evalq=all*) using 200 multi-locus bootstrap replicates.

135 We used phyparts (Smith et al., 2015) to evaluate the number of concordant and
136 conflicting bipartitions among gene trees in comparison to the inferred ASTRAL species tree
137 setting support level of at least 50% for the corresponding node (*-s 0.5*). Due to computational
138 limitations and the observed clustering of conspecific individuals (see Results), we pruned
139 replicates 2 and 3 for all species using Newick utilities (Junier and Zdobnov, 2010). Since
140 phyparts requires rooted trees as input, we set *Bouchetia erecta* as the root, which led to a dataset
141 of 3,471 gene trees where the outgroup was present. For the ML phylogenetic tree, we evaluated
142 genealogical concordance with gene concordance factor (gCF) and site concordance factor (sCF)
143 with 100 randomly sampled quartets (*-scf*), where gCF measures how often a specific branch in
144 the species tree is supported by “decisive” gene trees, while sCF measures the percentage of sites
145 that support a branch in the tree (Minh et al., 2020).

146

147 *Detection of hybridization*

148 We used HyDe (Blischak et al., 2018) to search for hybridization signals. HyDe relies on
149 phylogenetic invariants to estimate admixture (γ), where a γ value of 0.5 signifies an equal
150 genetic contribution from each parental species, and values approaching 0 or 1 indicate a greater
151 genetic contribution of one of the parental species. We used a concatenated matrix of alignments
152 and trimmed sites with trimAl (Capella-Gutiérrez et al., 2009) with options *-gt 0.5* (fraction of
153 gaps allowed) and *-cons 60* (minimum percentage of positions to keep from the original
154 alignment), which yielded 5,209,834 sites. We assigned individuals to species and set the six
155 outgroup species as outgroups, which resulted in an evaluation of 23,310 triplets. As a second
156 approach, we employed QuIBL (Edelman et al., 2019) which relies on branch lengths of gene
157 trees to assess whether hybridization provides a more plausible explanation for the divergence
158 patterns compared to ILS alone. Because QuIBL requires that all taxa be present in every gene
159 tree, we created a dataset with no missing loci for all ingroup species and *B. erecta*. All trees
160 were rooted in *B. erecta* and pruned to contain only one individual of each species with Newick
161 Utilities, which resulted in a final dataset of 826 gene trees.

162

163 *Reticulate evolution and network reconstruction*

164 Considering the possibility of a non-bifurcating evolutionary history of the *Petunia-Calibrachoa-*
165 *Fabiana*, we inferred phylogenetic networks that account for both ILS and gene flow among
166 taxa. Due to computational limitations, we constructed a reduced dataset comprising 18 ingroup
167 taxa and a single outgroup species (Table S1), not allowing for missing loci, which resulted in a
168 dataset of 1,215 loci. We estimated gene trees with RAxML and the species tree with ASTRAL
169 as described in the previous section.

170 As our first approach, we estimated a phylogenetic network with the maximum
171 pseudolikelihood method SNaQ implemented in the Julia package PhyloNetworks 0.16.2 (Solís-
172 Lemus et al., 2017; Solís-Lemus and Ané, 2016). We searched for up to five hybridization events
173 ($h = 5$) and used the ASTRAL phylogeny as the starting tree. For the following steps, we used
174 the network from the previous estimation as the starting network. The best number of
175 hybridization events was selected based on where we could detect a steep log-pseudolikelihood
176 improvement. After selecting the best number of hybridization events, we ran 100 bootstrap
177 replicates using the 1,000 bootstrap ML gene trees inferred by RAxML for each of the 1,215 loci,
178 employing default settings.

179 As a second approach, we estimated a network with the command “InferNetwork_MPL”
180 in PhyloNet 3.8.2 (Than et al., 2008), also searching for up to five hybridization events and 10
181 runs for each search. To select the best-scored network, we used the “CalGTProb” function in
182 PhyloNet (Yu et al., 2012) to get network likelihoods. We compared the networks with model
183 selection using the Akaike information criterion (AIC; Akaike 1973), the bias-corrected Akaike
184 information criterion (AICc; Sugiura 1978), and the Bayesian information criterion (BIC;
185 Schwarz 1978). We set the number of parameters to the number of estimated branch lengths and
186 hybridization probabilities, correcting for finite sample size with the number of gene trees used.


187 Considering the potential occurrence of intergeneric hybridization (see Results), we used
188 Twisst (Martin and Van Belleghem, 2017) on the reduced dataset of 19 species and 1,215 loci.
189 We categorized species according to their respective genera and designated *B. erecta* as the
190 outgroup, resulting in three potential topologies. We computed the topology weight and
191 determined the frequency of specific topologies within the gene tree set, that is, we counted the
192 number of trees supporting one of the three possible topologies. Subsequently, we conducted a
193 chi-square test to compare the occurrences of the two minor topologies (Owens et al., 2023;
194 Suvorov et al., 2022). Under the null hypothesis, i.e., without intergeneric hybridization, we
195 expect the two minor topologies to occur with similar frequency (Baum, 2007).

196

197 **Results**

198 *Phylogenetic relationships and tree discordance within the Petunia-Calibrachoa-Fabiana clade*

199 The gene count for each replicate ranged from 2,937 to 3,573 (Table S1), and the final
200 concatenated matrix consisted of 5,687,285 base pairs. The resulting phylogenetic trees
201 constructed using multiple methods consistently positioned *Petunia* as sister to *Calibrachoa* +
202 *Fabiana* (Olmstead et al., 2008; Alaria et al., 2022; Fig. 1), while revealing discordant
203 intragenetic topologies. Both the supertree (ASTRAL) and the supermatrix strategies (IQTree)
204 exhibited strong support for most branches (LPP = 0.95–1 in ASTRAL and bootstrap = 100 in
205 IQTree; Fig. 2). Nonetheless, the two methods estimated different relationships for multiple
206 branches within *Petunia* and within *Calibrachoa*, which might be expected given the high
207 proportion of conflict among gene trees apparent from the phyparts analysis (Fig. 2). We did,
208 however, find that the replicates from a single species consistently group together in the
209 ASTRAL analysis with robust support (Fig. S1), supporting assignment of individuals to species.
210 As might be expected given the differences between the ASTRAL and supermatrix trees (Fig. 2),
211 SVDQuartets displayed high support for deeper nodes, but weaker support for shallower nodes
212 within *Calibrachoa* and *Petunia* (Fig. S2), indicating extensive ILS and possibly intragenetic
213 hybridization. IQTree Concordance Factor results indicated that the gCF values were notably low
214 for shallow nodes, whereas sCF values offered greater support for these relationships than gCF,
215 suggesting that genetic sites were more consistent in inferring evolutionary relationships at these
216 shallower nodes than the genes themselves (Fig. 2).




217
 218 **Figure 2.** Phylogenetic trees of the *Petunia-Calibrachoa-Fabiana* clade inferred from ASTRAL
 219 (A) and IQTree (B). The subgenera of *Calibrachoa* are annotated on their branches on ASTRAL
 220 tree: Ca for *Calibrachoa*, and St for *Stimomphis*. All nodes are strongly supported (local
 221 posterior probability=1 for ASTRAL/bootstrap=100 for maximum likelihood tree), except when
 222 otherwise noted by an asterisk (*). Dashed lines represent species with differing positions in the
 223 two trees. Pie charts on the ASTRAL tree depict gene support based on 3,471 gene trees:
 224 corroborating (green), conflicting (orange), or uninformative (gray; < 50% bootstrap scores or
 225 missing loci) relationships. The piecharts and numbers above the branches on the maximum
 226 likelihood tree inferred from IQTree indicate gene and site Concordance Factors (gCF and sCF,
 227 respectively). Outgroup species are not shown for simplicity.

228

229 *Reticulate evolution and species networks*

230 The search for hybrids resulted in several significant hybrid triplets, both in QuIBL and HyDe
 231 (Fig. 3). HyDe resulted in 3,352 significant triplets, even between intergeneric species (Table
 232 S2). However, when we only considered events with $0.2 < \gamma < 0.8$ (to detect more recent
 233 hybridization events, where we can detect greater parental contribution from both species, and
 234 discard spurious results with low contribution from either parent), these hybridization events
 235 were constrained within genera (Fig. 3). QuIBL showed several minor topologies that could not
 236 be explained by ILS alone, although the percentage of discordant loci explained by introgression
 237 were lower than 10% in all cases (Fig. S3; Table S3).




238
239
240
241
242
243

Figure 3. Detected hybridization events using QuIBL (top) and HyDe (bottom), using ASTRAL phylogeny as the reference species tree. The HyDe graph displays hybridization events with $0.2 < \gamma < 0.8$; refer to Table S2 for complete results. QuIBL identified introgression events based on branch lengths in 826 gene trees with single individuals from each taxon; refer to Table S3 for complete results.

244
245
246
247
248
249
250
251
252

Although network inferences yielded different optimal numbers of reticulations (one in SNaQ, two in PhyloNet; Fig. 4), both agreed on an ancient hybridization edge from *Petunia* to *Calibrachoa* subgenus *Stimomphis*. However, the inheritance probabilities for this introgression were low in both analyses (less than 1% in SNaQ and 3.4% in PhyloNet). The bootstrap analyses for SNaQ showed high support for the species network nodes, but low support for the hybrid edge. The placement of minor and major edges was not consistent, with low consistency for both the origin and the source of hybridization. The addition of more hybridization events in SNaQ usually led to the impossibility of rooting the tree in the outgroup (supplemental material online), which suggests incorrect placement of that hybridization edge.




253
 254 **Figure 4.** Inferred species networks using SNaQ (left) and PhyloNet (right) based on a reduced
 255 dataset comprising 18 ingroup species and 1,125 genes, rooted in *B. erecta*. The SNaQ tree
 256 identified the optimal network with one hybridization event. All branches received 100%
 257 bootstrap support from 100 replicates, except those signed with gray circles. The dashed line
 258 represents the minor edge, displaying the inheritance probability of the best network. The
 259 bootstrap values for minor (origin) and major (source) edges of alternative networks are colored
 260 green and orange, respectively. The PhyloNet network indicated the optimal network with two
 261 hybridization events. The dashed lines also indicate the minor edges with corresponding
 262 inheritance probabilities.

263
 264 The model selection for PhyloNet revealed the network with two hybridization edges as
 265 the optimal network and highlighted that any species network is better fitting than the bifurcating
 266 species tree (Table 1). In addition to the intergeneric hybridization edge, PhyloNet also
 267 suggested a hybridization within *Fabiana* as a second hybridization event. In this case, it showed
 268 a high inheritance probability of 0.46 from *F. patagonica* to *F. australis*.

269
 270 **Table 1.** Network likelihoods derived from the reduced dataset using PhyloNet. The number of
 271 parameters (k) represents the number of estimated branch lengths and admixture probabilities.
 272 The optimal network is in bold.

Topology	Maximum number of reticulations	Number of inferred reticulations	Total log probability	lnL	k	AIC	ΔAIC	AICc	BIC
Astral	0	NA	-	-30833.49	37	61740.98	4676.06	61743.37	61781.11
Network 0	0	NA	-389607.21	-29310.04	37	58694.08	1629.16	58620.08	58734.21
Network 1	1	1	-388824.25	-28520.99	39	57119.98	55.06	57041.98	57162.28
Network 2	2	2	-388777.78	-28491.46	41	57064.92	0.00	56982.92	57109.39
Network 3	3	3	-388791.27	-28511.48	43	57108.96	44.04	57022.96	57155.60
Network 4	4	3	-388805.06	-28524.33	43	57138.66	73.74	57048.66	57181.30
Network 5	5	3	-388827.51	-28543.19	43	57180.38	115.46	57086.38	57219.02

273 Results from Twisst revealed that, between the two minor topologies, the topology
 274 positioning *Fabiana* as the sister group to *Petunia* and *Calibrachoa* exhibited a marginally
 275 greater frequency (Fig. 5), although this difference was not statistically significant (chi-square =
 276 1.15, P value = 0.28). Thus, the difference in the number of topologies of gene trees is primarily
 277 attributed to ILS and not gene flow.



278 **Figure 5.** Total count of topologies by Twisst for the subset of 1,215 gene trees. The P value
 279 indicates the lack of significance for the chi-square test between the two minor topologies that
 280 places *Calibrachoa* or *Fabiana* as the outgroup.
 281

282
 283 **Discussion**
 284

285 *Extensive ILS and ancient hybridization are the sources of tree discordance in the Petunia-*
 286 *Calibrachoa-Fabiana clade*

287 Historically, phylogenies based on Sanger sequences often yielded conflicting species trees
 288 among *Petunia*, *Calibrachoa*, and *Fabiana*. However, our results consistently placed *Petunia* as
 289 a sister group to *Calibrachoa* + *Fabiana*, mirroring previous findings by Olmstead et al., (2008),
 290 Alaria et al., (2022), and Wheeler et al., (2022). These results are surprising due to the
 291 morphological and ecological similarities between *Petunia* and *Calibrachoa* and suggest that
 292 bee-pollinated herbs (such as most extant *Calibrachoa* and *Petunia* species) represent the
 293 ancestral state with the extreme xerophyte traits found in *Fabiana* (tiny flowers and reduced
 294 leaves) being derived features. Moreover, the more arid and temperate range of the clade likely
 295 represents a southward expansion from the shared distribution of *Petunia* and *Calibrachoa*, both
 296 of which is inferred to have originated in the lowland grasslands of southern Brazil, Uruguay,

297 and northeast Argentina (Reck-Kortmann et al., 2014; 2015; Mäder and Freitas, 2019).
298 Nonetheless, the relationships within each genus remained inconsistent, with high levels of
299 conflict among all reconstructed trees. Moreover, we found extensive gene flow within genera,
300 as supported by both QuIBL and HyDe (Fig. 3) analyses.

301 Our results corroborated the subdivision of *Petunia* into two main clades, the long corolla
302 tube and the short corolla tube clades (Reck-Kortmann et al., 2014). The former is characterized
303 by a wide range of flower colors and pollinators, represented here by *P. axillaris* subsp. *parodii*
304 (white, hawkmoth-pollinated), *P. exserta* (red, hummingbird-pollinated), and *P. secreta* (purple,
305 bee-pollinated). In contrast, the latter consists of species with purple flowers primarily pollinated
306 by bees, represented here by the remaining seven *Petunia* species. The long corolla tube clade is
307 noteworthy for its documented history of extensive hybridization (e.g., Caballero-Villalobos et
308 al., 2021; Giudicelli et al., 2019), whereas records of interspecific hybridization within the short
309 corolla tube clade are rare and, until now, limited to *P. interior* and *P. inflata* (Pezzi et al., 2022).

310 The short corolla tube clade displayed a higher number of discordant gene trees and
311 shorter branch lengths compared to the long corolla tube clade, indicative of rapid radiation.
312 These species are often rare and endemic (Souza et al., 2022), occurring primarily in southern
313 Brazil. Geographic isolation serves as the primary reproductive barrier among these species, and
314 pollinators do not impose any reproductive barriers because they are shared among species
315 (Stehmann et al., 2009). Despite their geographic isolation due to microhabitat adaptation, many
316 of these species have overlapping distributions, and all are self-incompatible (Stehmann et al.,
317 2009). This scenario presents an opportunity for interspecific gene flow. Whereas field
318 observations have documented only a few hybrids, the substantial level of polymorphism shared
319 between these species could be attributed to high levels of ILS (Lorenz-Lemke et al., 2010) or
320 ongoing and recent hybridization events (Fig. 3).

321 *Calibrachoa* is classified into two subgenera: *Calibrachoa* and *Stimomphis*. The
322 subgenus *Calibrachoa* comprises just two species that exhibit significant differences in
323 reproductive biology and habitat compared to species in the subgenus *Stimomphis* (Fregonezi et
324 al., 2013). Intriguingly, no instances of hybridization have been observed between *Calibrachoa*
325 and *Stimomphis* species, underscoring the presence of robust reproductive barriers between
326 subgenera. *Stimomphis* showed a similar evolutionary history to the one observed in the short
327 corolla *Petunia* clade: rapid radiation, high levels of conflicting gene trees, and extensive ILS.
328 The topology recovered here exhibits minimal congruence with prior studies (Fregonezi et al.,
329 2012, 2013; Mäder and Freitas, 2019). Notably, the highland clade identified by Mäder and
330 Freitas (2019), represented here by *C. elegans*, *C. eglandulata*, *C. sendtneriana*, and *C. linoides*,
331 did not emerge in any of our phylogenetic reconstructions. Such high levels of conflict among
332 phylogenetic methods are expected with extensive ILS and hybridization. Moreover, it is worth
333 noting that, except for *C. parviflora*, all these species are self-incompatible (Fregonezi et al.,
334 2013), facilitating the potential for hybridization as these species cannot prevent heterospecific
335 gene flow through autonomous selfing (Brys et al., 2015).

336 It has been demonstrated that ML concatenation methods are often inadequate for
337 accurately recovering species trees when extensive ILS is involved (Kubatko and Degnan, 2007;
338 Mendes and Hahn, 2018). Therefore, ASTRAL is a more suitable alternative in scenarios such as
339 the short corolla *Petunia* clade and *Calibrachoa* subgenus *Stimomphis*. These clades display
340 extremely short branch lengths, suggesting they went through a rapid diversification process, not
341 allowing for genes to coalesce. Species that underwent rapid radiation tend to fall in the
342 “anomaly zone”, where the most frequent gene trees do not align with the species tree (Degnan

343 and Rosenberg, 2006; Linkem et al., 2016) which could explain the high levels of tree
344 discordance observed here. To address this challenge, one potential strategy is to sample multiple
345 individuals from the same species (Degnan and Rosenberg, 2006), as was done here. However, it
346 is important to note that our individuals were sampled on the same site (Table S1), and they
347 might not fully represent the species' genetic diversity. Thus, they may fail to provide sufficient
348 resolution for phylogenetic inferences. However, when investigating clades harboring numerous
349 rare and endemic species, sampling from the same locality is often unavoidable, but it still
350 provides valuable biological insights.

351 Among the three genera, *Fabiana* exhibited a lower level of tree discordance and greater
352 consistency among phylogenies. The previously available *Fabiana* phylogeny included only
353 eight species but agreed on the close relationship between *F. imbricata* and *F. australis* (Alaria et
354 al., 2022). Here, *F. viscosa* displayed varied phylogenetic placement and the highest level of
355 introgression in QuIBL analyses, indicating significant gene flow, particularly with *F. australis*
356 (Fig. 3). This implies that, in addition to ILS, hybridization plays a central role in causing tree
357 discordance for these species. Intriguingly, these two species do not currently occur in sympatry.

358 It is crucial to recognize that methods that rely on summary statistics of triplets or
359 quartets, such as HyDe, are highly sensitive to substitution rate variation across lineages and
360 genes (Frankel and Ané, 2023), resulting in a high rate of false positives. In addition, such
361 methods can fail to identify hybridization involving ghost or unsampled taxa (Bjorner et al.,
362 2022), and they often struggle to discern gene flow between sister species, where allele sharing
363 could be interpreted as ancestral polymorphism (Mallet et al., 2016). Moreover, transcriptomic
364 data is inherently more prone to natural selection because it comprises coding regions (Liu et al.,
365 2015). This can potentially influence rate variation between genes and the lengths of branches in
366 phylogenetic trees (Edwards et al., 2016), ultimately impacting the power detection of
367 hybridization analyses (Frankel and Ané, 2023). Hybridization methods that rely on branch
368 length are also susceptible to the influence of rate variation. For instance, QuIBL has
369 demonstrated a propensity for producing false positives when using shorter alignments compared
370 to alignments longer than 1,000 bp (Koppetsch et al., 2023). Notably, our QuIBL dataset
371 exhibited a median length of 494 bp, with 10% of the alignments being longer than 900 bp. The
372 *Stimomphis* species and *Petunia* short corolla clade diverged recently and are probably less prone
373 from high-rate variation across lineages. Thus, the intrageneric hybridization events detected by
374 both HyDe and QuIBL are likely authentic. However, we should interpret these results with
375 caution as there is still potential for the rate variation assumption to be violated.

376
377 *Would it be possible for Petunia and Calibrachoa to hybridize?*

378 Our network analyses suggested an introgression event from *Petunia* to *Calibrachoa* subgenus
379 *Stimomphis*, which, given recent dating estimates, would have occurred roughly between 8 mya
380 (Särkinen et al., 2013) and 10 mya (Lisa De-Silva et al., 2017). However, recent estimates
381 indicate the divergence of *Calibrachoa* and *Fabiana* around 20 mya (Zuntini et al., 2024),
382 suggesting that interspecific hybridization would have to be at least this old. These two genera
383 differ in their chromosome number, with *Petunia* having a haploid chromosome number of seven
384 (Stehmann et al., 2009) and *Calibrachoa* having nine (Wijsman and De Jong, 1985), as in
385 *Fabiana* (Acosta et al., 2006). Such differences in chromosome numbers typically impose a
386 strong postzygotic barrier against hybridization, either preventing it entirely or resulting in
387 hybrid sterility (Levin, 2002). Nevertheless, instances of hybridization and introgression between
388 plant species with different ploidy numbers have been documented (Chapman and Abbott, 2010),

389 and both *Petunia* and *Calibrachoa* subgenus *Stimomphis* share similar geographic distribution,
390 morphology, habitat, and potential group of pollinators (Stemann et al., 2009). One possible
391 explanation for our result is that this change in chromosome number occurred in the ancestral
392 lineage of *Petunia* after the admixture event, such as in the scenario posited by PhyloNet, where
393 the admixture event is from the common ancestor of all *Petunia* species. However, SNaQ
394 contradicts this hypothesis, as the introgression event is inferred to have occurred after a
395 reduction in chromosome number in the common ancestor of *Petunia* (Fig. 4). Alternatively,
396 sufficient chromosomal homology may have enabled meiotic pairing during diversification.
397 Artificial crosses between *Calibrachoa* and *Petunia* demonstrated some success in embryo
398 formation but failed in germination (Olschowski et al., 2013). However, Milicia et al., (2021)
399 crossed *P. inflata* with *C. hybrida*, and despite a significantly lower percentage of viable pollen
400 granules than intrageneric crosses, the hybrids produced 5% of viable pollen, highlighting plant
401 species' flexibility in chromosome rearrangement to allow successful meiosis. Thus,
402 hybridization between these genera may not be out of the question. Additionally, the low chance
403 of current hybridization does not exclude the possibility of ancient hybridization.

404 Despite the inference of intergeneric hybridization from SNaQ and PhyloNet, we did not
405 detect any support for such an event from our Twisst analysis. Instead, the discordance appears
406 best explained by ILS as the two minor topologies are present in nearly equal frequencies (Fig.
407 5) We note that inheritance probabilities from SNaQ and PhyloNet were very low (1 to 3%, Fig.
408 4), and thus, this reticulation event, if it occurred, might be at the boundary of detection. The
409 absence—or very low levels—of gene flow between these two genera highlights how important
410 chromosome number difference was to prevent hybridization, which allowed *Petunia* and
411 *Calibrachoa* to undergo parallel radiation despite their many ecological similarities and
412 geographic overlaps. Regardless, hybridization between the two genera merits future
413 investigation when full genomes become available for these genera (Bombarely et al., 2010).
414 Introgression of even a small fraction of the genome could potentially carry a large phenotypic
415 effect (Clarkson et al., 2014) and facilitate rapid radiations (Meier et al., 2017). However,
416 detecting such events with confidence is challenging as it involves identifying introgressed
417 genomic regions and linking them to adaptations (Taylor & Larson, 2019; Suarez-Gonzalez et
418 al., 2018).

419 **Conclusions**

421 Here, we investigated the origins of tree discordance in the *Petunia-Calibrachoa-Fabiana*
422 Solanaceae clade using a comprehensive genome-scale dataset encompassing multiple species.
423 Our results confirm *Petunia* as the sister genus to *Calibrachoa* + *Fabiana*. However, the
424 relationships among species within these genera remain unsolved. The discordance in tree
425 topologies within the short corolla tube *Petunia* clade and *Calibrachoa* subgenus *Stimomphis*
426 arises from a combination of ILS due to their rapid diversification and past and ongoing
427 hybridization events. Instances of high ILS and extensive hybridization are not uncommon in the
428 evolutionary history of plants (e.g., Kleinkopf et al., 2019; Morales-Briones et al., 2021; McLay
429 et al., 2023), but pinpointing the specific taxa involved in the hybridization events is still a
430 daunting task, and one of the reasons why different methods often yield conflicting results (Gates
431 et al., 2023). These introgression events likely contributed to the species' genetic diversity, aiding
432 their adaptation during their radiation. Additionally, our network reconstructions indicated
433 potential intergeneric hybridization between *Calibrachoa* and *Petunia*, two genera characterized
434 by distinct chromosome numbers. Considering the weak hybridization signals observed in

435 network analyses, the lack of support from gene tree topology weights, and the known current
436 barriers due to differing base chromosome numbers, it leads us to believe that such a
437 hybridization event did not occur. However, both of our network analyses indicated intergeneric
438 gene flow, suggesting there is still a remote possibility that this could have occurred and may
439 have been facilitated by strong selection despite the barrier imposed by differing chromosome
440 base numbers between the two genera. Whole genome analyses could solve the intergeneric
441 hybridization puzzle and contribute to ascertaining which genomic regions may have been
442 involved in the *Petunia-Calibrachoa* introgression. Overall, our study sheds light on the complex
443 evolutionary history of this charismatic South American clade, providing crucial insights into its
444 adaptation and diversification.

445

446 **Acknowledgments**

447 This work used the RMACC Alpine supercomputer, funded by the University of Colorado
448 Boulder, the University of Colorado Anschutz, Colorado State University, and the National
449 Science Foundation (award 2201538). SDS was supported by NSF-DEB (award 1553114). PHP
450 was supported by a CAPES/PRINT fellowship.

451

452 **Declaration of competing interest**

453 The authors declare that they have no competing interests.

454

455 **Data availability**

456 All scripts and processed data files are available at [https://github.com/pedrohpezzi/Petunia-](https://github.com/pedrohpezzi/Petunia-Calibrachoa-Fabiana_TreeDiscordance.git)
457 [Calibrachoa-Fabiana_TreeDiscordance.git](https://github.com/pedrohpezzi/Petunia-Calibrachoa-Fabiana_TreeDiscordance.git) and <https://figshare.com/s/c3f6e7305660e03031ec>.
458 The raw RNA-seq data files are available in SRA under the BioProject accession number
459 PRJNA746328.

460

461 **CRedit author contributions**

462 **Pedro H. Pezzi:** Conceptualization, Methodology, Software, Formal analysis, Investigation,
463 Writing - Original Draft, Visualization. **Lucas C. Wheeler:** Conceptualization, Data Curation,
464 Methodology, Software, Investigation, Writing - Review and Editing. **Loreta B. Freitas:**
465 Conceptualization, Writing - Review and Editing, Investigation, Supervision. **Stacey D. Smith:**
466 Conceptualization, Resources, Investigation, Writing - Review and Editing, Supervision, Project
467 administration, Funding acquisition.

468 **References**

- 469 Abbott, R., Albach, D., Ansell, S., Arntzen, J.W., Baird, S.J.E., Bierne, N., Boughman, J.,
470 Brelsford, A., Buerkle, C.A., Buggs, R., Butlin, R.K., Dieckmann, U., Eroukhmanoff, F.,
471 Grill, A., Cahan, S.H., Hermansen, J.S., Hewitt, G., Hudson, A.G., Jiggins, C., Jones, J.,
472 Keller, B., Marczewski, T., Mallet, J., Martinez-Rodriguez, P., Möst, M., Mullen, S.,
473 Nichols, R., Nolte, A.W., Parisod, C., Pfennig, K., Rice, A.M., Ritchie, M.G., Seifert, B.,
474 Smadja, C.M., Stelkens, R., Szymura, J.M., Väinölä, R., Wolf, J.B.W., Zinner, D., 2013.
475 Hybridization and speciation. *J. Evol. Biol.* 26, 229–246. [https://doi.org/10.1111/j.1420-](https://doi.org/10.1111/j.1420-9101.2012.02599.x)
476 [9101.2012.02599.x](https://doi.org/10.1111/j.1420-9101.2012.02599.x)
- 477 Abbott, R.J., Barton, N.H., Good, J.M., 2016. Genomics of hybridization and its evolutionary
478 consequences. *Mol. Ecol.* 25, 2325–2332. <https://doi.org/10.1111/mec.13685>
- 479 Acosta, M.C., Ordóñez, A.V., Cocucci, A.A., Moscone, E.A. 2006. Chromosome reports in South
480 American Nicotianeae (Solanaceae), with particular reference to *Nierembergia*. *Ann. Mo.*
481 *Bot. Gard.* 93, 634–646. [https://doi.org/10.3417/0026-](https://doi.org/10.3417/0026-6493(2006)93[634:CRISAN]2.0.CO;2)
482 [6493\(2006\)93\[634:CRISAN\]2.0.CO;2](https://doi.org/10.3417/0026-6493(2006)93[634:CRISAN]2.0.CO;2)
- 483 Akaike, H., 1973. Information Theory and an Extension of the Maximum Likelihood Principle.,
484 in: *International Symposium on Information Theory*. pp. 267–281.
- 485 Alaria, A., Chau, J.H., Olmstead, R.G., Peralta, I.E., 2022. Relationships among *Calibrachoa*,
486 *Fabiana* and *Petunia* (Petunieae tribe, Solanaceae) and a new generic placement of
487 Argentinean endemic *Petunia patagonica*. *PhytoKeys* 194, 75–93.
488 <https://doi.org/10.3897/phytokeys.194.68404>
- 489 Alix, K., Gérard, P.R., Schwarzacher, T., Heslop-Harrison, J.S. (Pat), 2017. Polyploidy and
490 interspecific hybridization: partners for adaptation, speciation and evolution in plants.
491 *Ann. Bot.* 120, 183–194. <https://doi.org/10.1093/aob/mcx079>
- 492 Baack, E., Melo, M.C., Rieseberg, L.H., Ortiz-Barrientos, D., 2015. The origins of reproductive
493 isolation in plants. *New Phytol.* 207, 968–984. <https://doi.org/10.1111/nph.13424>
- 494 Baum, D.A. 2007. Concordance trees, concordance factors, and the exploration of reticulate
495 genealogy. *Taxon*, 56, 417–426. <https://doi.org/10.1002/tax.562013>
- 496 Binaghi, M., Esfeld, K., Mandel, T., Freitas, L.B., Roesti, M., Kuhlemeier, C., 2023. Genetic
497 architecture of a pollinator shift and its fate in secondary hybrid zones of two *Petunia*
498 species. *BMC Biol.* 21, 58. <https://doi.org/10.1186/s12915-023-01561-x>
- 499 Bjorner, M., Molloy, E.K., Dewey, C.N., Solís-Lemus, C., 2022. Detectability of Varied
500 Hybridization Scenarios using Genome-Scale Hybrid Detection Methods.
501 <https://doi.org/10.48550/ARXIV.2211.00712>
- 502 Blischak, P.D., Chifman, J., Wolfe, A.D., Kubatko, L.S., 2018. HyDe: A Python Package for
503 Genome-Scale Hybridization Detection. *Syst. Biol.* 67, 821–829.
504 <https://doi.org/10.1093/sysbio/syy023>
- 505 Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina
506 sequence data. *Bioinformatics* 30, 2114–2120.
507 <https://doi.org/10.1093/bioinformatics/btu170>
- 508 Bombarely, A., Menda, N., Tecle, I.Y., Buels, R.M., Strickler, S., Fischer-York, T., Pujar, A.,
509 Leto, J., Gosselin, J., Mueller, L. A. 2010. The Sol Genomics Network (solgenomics.
510 net): growing tomatoes using Perl. *Nucleic Acids Res.* 39, 1149–1155.
511 <https://doi.org/10.1093/nar/gkq866>
- 512 Brys, R., Van Cauwenberghe, J., Jacquemyn, H. 2016. The importance of autonomous selfing in
513 preventing hybridization in three closely related plant species. *J. Ecol.* 104, 601–610.

514 <https://doi.org/10.1111/1365-2745.12524>

515 Caballero-Villalobos, L., Silva-Arias, G.A., Turchetto, C., Giudicelli, G.C., Petzold, E.,
516 Bombarely, A., Freitas, L.B., 2021. Neutral and adaptive genomic variation in hybrid
517 zones of two ecologically diverged *Petunia* species (Solanaceae). *Bot. J. Linn. Soc.* 196,
518 100–122. <https://doi.org/10.1093/botlinnean/boaa091>

519 Capella-Gutiérrez, S., Silla-Martínez, J.M., Gabaldón, T., 2009. trimAl: a tool for automated
520 alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25, 1972–1973.
521 <https://doi.org/10.1093/bioinformatics/btp348>

522 Chapman, M.A., Abbott, R.J., 2010. Introgression of fitness genes across a ploidy barrier:
523 Research review. *New Phytol.* 186, 63–71. <https://doi.org/10.1111/j.1469-8137.2009.03091.x>

525 Chifman, J., Kubatko, L., 2014. Quartet Inference from SNP Data Under the Coalescent Model.
526 *Bioinformatics* 30, 3317–3324. <https://doi.org/10.1093/bioinformatics/btu530>

527 Christie, K., Fraser, L.S., Lowry, D.B., 2022. The strength of reproductive isolating barriers in
528 seed plants: Insights from studies quantifying premating and postmating reproductive
529 barriers over the past 15 years. *Evolution* 76, 2228–2243.
530 <https://doi.org/10.1111/evo.14565>

531 Clarkson, C.S., Weetman, D., Essandoh, J., Yawson, A.E., Maslen, G., Manske, M., Field, S.G.,
532 Webster, M., Antão, T., MacInnis, B., Kwiatkowski, D., Donnelly, M.J. 2014. Adaptive
533 introgression between *Anopheles* sibling species eliminates a major genomic island but
534 not reproductive isolation. *Nat. Commun.* 5, 4248. <https://doi.org/10.1038/ncomms5248>

535 Degnan, J.H., Rosenberg, N.A., 2006. Discordance of Species Trees with Their Most Likely
536 Gene Trees. *PLoS Genet.* 2, e68. <https://doi.org/10.1371/journal.pgen.0020068>

537 Dell’Olive, A., Hoballah, M.E., Gübitz, T., Kuhlemeier, C., 2011. Isolation barriers between
538 *Petunia axillaris* and *Petunia integrifolia* (Solanaceae). *Evolution* 65, 1979–1991.
539 <https://doi.org/10.1111/j.1558-5646.2011.01279.x>

540 Edelman, N.B., Frandsen, P.B., Miyagi, M., Clavijo, B., Davey, J., Dikow, R.B., García-
541 Accinelli, G., Van Belleghem, S.M., Patterson, N., Neafsey, D.E., Challis, R., Kumar, S.,
542 Moreira, G.R.P., Salazar, C., Chouteau, M., Counterman, B.A., Papa, R., Blaxter, M.,
543 Reed, R.D., Dasmahapatra, K.K., Kronforst, M., Joron, M., Jiggins, C.D., McMillan,
544 W.O., Di Palma, F., Blumberg, A.J., Wakeley, J., Jaffe, D., Mallet, J. 2019. Genomic
545 architecture and introgression shape a butterfly radiation. *Science* 366, 594–599.
546 <https://doi.org/10.1126/science.aaw2090>

547 Edwards, S.V., Xi, Z., Janke, A., Faircloth, B.C., McCormack, J.E., Glenn, T.C., Zhong, B., Wu,
548 S., Lemmon, E.M., Lemmon, A.R., Leaché, A.D., Liu, L., Davis, C.C. 2016.
549 Implementing and testing the multispecies coalescent model: a valuable paradigm for
550 phylogenomics. *Mol. Phylogenetics Evol.* 94, 447–462.
551 <https://doi.org/10.1016/j.ympev.2015.10.027>

552 Frankel, L.E., Ané, C., 2023. Summary tests of introgression are highly sensitive to rate variation
553 across lineages. *Syst. Biol.* 72, syad056. <https://doi.org/10.1093/sysbio/syad056>

554 Fregonezi, J.N., De Freitas, L.B., Bonatto, S.L., Semir, J., Stehmann, J.R., 2012. Infrageneric
555 classification of *Calibrachoa* (Solanaceae) based on morphological and molecular
556 evidence. *TAXON* 61, 120–130. <https://doi.org/10.1002/tax.611009>

557 Fregonezi, J.N., Turchetto, C., Bonatto, S.L., Freitas, L.B., 2013. Biogeographical history and
558 diversification of *Petunia* and *Calibrachoa* (Solanaceae) in the Neotropical Pampas
559 grassland: Species Diversification in the Pampas. *Bot. J. Linn. Soc.* 171, 140–153.

560 <https://doi.org/10.1111/j.1095-8339.2012.01292.x>

561 Galtier, N., Daubin, V., 2008. Dealing with incongruence in phylogenomic analyses. *Philos.*
562 *Trans. R. Soc. B Biol. Sci.* 363, 4023–4029. <https://doi.org/10.1098/rstb.2008.0144>

563 Gates, D.J., Pilson, D., Smith, S.D., 2023. Hybridization in *Iochroma*, in: *Species Tree Inference:*
564 *A Guide to Methods and Applications.* Princeton University Press, Princeton, NJ, USA.

565 Giudicelli, G.C., Turchetto, C., Teixeira, M.C., Freitas, L.B., 2019. Morphological and genetic
566 characterization in putative hybrid zones of *Petunia axillaris* subsp. *axillaris* and subsp.
567 *parodii* (Solanaceae). *Bot. J. Linn. Soc.* 191, 353–364.
568 <https://doi.org/10.1093/botlinnean/boz057>

569 Goulet, B.E., Roda, F., Hopkins, R., 2017. Hybridization in Plants: Old Ideas, New Techniques.
570 *Plant Physiol.* 173, 65–78. <https://doi.org/10.1104/pp.16.01340>

571 Gübitz, T., Hoballah, M.E., Dell’Olivo, A., Kuhlemeier, C., 2009. *Petunia* as a Model System for
572 the Genetics and Evolution of Pollination Syndromes, in: Gerats, T., Strommer, J. (Eds.),
573 *Petunia.* Springer New York, New York, NY, pp. 29–49. [https://doi.org/10.1007/978-0-](https://doi.org/10.1007/978-0-387-84796-2_2)
574 [387-84796-2_2](https://doi.org/10.1007/978-0-387-84796-2_2)

575 Hegarty, M.J., Hiscock, S.J., 2008. Genomic Clues to the Evolutionary Success of Polyploid
576 Plants. *Curr. Biol.* 18, R435–R444. <https://doi.org/10.1016/j.cub.2008.03.043>

577 Hibbins, M.S., Hahn, M.W., 2022. Phylogenomic approaches to detecting and characterizing
578 introgression. *Genetics* 220, iyab173. <https://doi.org/10.1093/genetics/iyab173>

579 Jonika, M.M., Maximos Chin, Anderson, N.W., Adams, R.H., Demuth, J.P., Blackmon, H., 2023.
580 *coleoguy/evobir: EvobiR version 2.1.* <https://doi.org/10.5281/ZENODO.8033504>

581 Junier, T., Zdobnov, E.M., 2010. The Newick utilities: high-throughput phylogenetic tree
582 processing in the Unix shell. *Bioinformatics* 26, 1669–1670.
583 <https://doi.org/10.1093/bioinformatics/btq243>

584 Kleinkopf, J.A., Roberts, W.R., Wagner, W.L., Roalson, E.H. 2019. Diversification of Hawaiian
585 *Cyrtandra* (Gesneriaceae) under the influence of incomplete lineage sorting and
586 hybridization. *J. Syst. Evol.*, 57, 561–578. <https://doi.org/10.1111/jse.12519>

587 Kong, S., Kubatko, L.S., 2021. Comparative Performance of Popular Methods for Hybrid
588 Detection using Genomic Data. *Syst. Biol.* 70, 891–907.
589 <https://doi.org/10.1093/sysbio/syaa092>

590 Koppetsch, T., Malinsky, M., Matschiner, M. 2023. Towards reliable detection of introgression in
591 the presence of among-species rate variation. *bioRxiv*, 2023–05.
592 <https://doi.org/10.1101/2023.05.21.541635>

593 Kubatko, L.S., Degnan, J.H., 2007. Inconsistency of Phylogenetic Estimates from Concatenated
594 Data under Coalescence. *Syst. Biol.* 56, 17–24.
595 <https://doi.org/10.1080/10635150601146041>

596 Levin, D.A., 2002. *The role of chromosomal change in plant evolution.* Oxford University Press,
597 New York, NY, USA.

598 Li, H., Durbin, R., 2010. Fast and accurate long-read alignment with Burrows–Wheeler
599 transform. *Bioinformatics* 26, 589–595. <https://doi.org/10.1093/bioinformatics/btp698>

600 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
601 Durbin, R., 1000 Genome Project Data Processing Subgroup, 2009. The Sequence
602 Alignment/Map format and SAMtools. *Bioinformatics* 25, 2078–2079.
603 <https://doi.org/10.1093/bioinformatics/btp352>

604 Linkem, C.W., Minin, V.N., Leaché, A.D., 2016. Detecting the Anomaly Zone in Species Trees
605 and Evidence for a Misleading Signal in Higher-Level Skink Phylogeny (Squamata:

606 Scincidae). *Syst. Biol.* 65, 465–477. <https://doi.org/10.1093/sysbio/syw001>

607 Lisa De-Silva, D., Mota, L.L., Chazot, N., Mallarino, R., Silva-Brandão, K.L., Piñerez, L.M.G.,
608 Freitas, A.V., Lamas, G., Joron, M., Mallet, J., Giraldo, C.E., 2017. North Andean origin
609 and diversification of the largest ithomiine butterfly genus. *Sci. Rep.* 7, p.45966.
610 <https://doi.org/10.1038/srep45966>

611 Liu, L., Xi, Z., Wu, S., Davis, C.C., Edwards, S.V. 2015. Estimating phylogenetic trees from
612 genome-scale data. *Ann. N. Y. Acad. Sci.* 1360, 36–53.
613 <https://doi.org/10.1111/nyas.12747>

614 Lorenz-Lemke, A.P., Togni, P.D., Mäder, G., Kriedt, R.A., Stehmann, J.R., Salzano, F.M.,
615 Bonatto, S.L., Freitas, L.B., 2010. Diversification of plant species in a subtropical region
616 of eastern South American highlands: a phylogeographic perspective on native *Petunia*
617 (Solanaceae). *Mol. Ecol.* 19, 5240–5251. <https://doi.org/10.1111/j.1365-294X.2010.04871.x>

618

619 Mäder, G., Freitas, L.B., 2019. Biogeographical, ecological, and phylogenetic analyses clarifying
620 the evolutionary history of *Calibrachoa* in South American grasslands. *Mol. Phylogenet.*
621 *Evol.* 141, 106614. <https://doi.org/10.1016/j.ympev.2019.106614>

622 Mallet, J., Besansky, N., Hahn, M.W., 2016. How reticulated are species? *BioEssays* 38, 140–
623 149. <https://doi.org/10.1002/bies.201500149>

624 Martin, S.H., Van Belleghem, S.M., 2017. Exploring Evolutionary Relationships Across the
625 Genome Using Topology Weighting. *Genetics* 206, 429–438.
626 <https://doi.org/10.1534/genetics.116.194720>

627 McLay, T.G., Fowler, R.M., Fahey, P.S., Murphy, D.J., Udovicic, F., Cantrill, D.J., Bayly, M.J.
628 2023. Phylogenomics reveals extreme gene tree discordance in a lineage of dominant
629 trees: Hybridization, introgression, and incomplete lineage sorting blur deep evolutionary
630 relationships despite clear species groupings in *Eucalyptus* subgenus *Eudesmia*. *Mol.*
631 *Phylogenetics Evol.* 187, 107869. <https://doi.org/10.1016/j.ympev.2023.107869>

632 Meier, J.I., Marques, D.A., Mwaiko, S., Wagner, C.E., Excoffier, L., Seehausen, O. 2017.
633 Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nat. Commun.* 8,
634 14363. <https://doi.org/10.1038/ncomms14363>

635 Mendes, F.K., Hahn, M.W., 2018. Why Concatenation Fails Near the Anomaly Zone. *Syst. Biol.*
636 67, 158–169. <https://doi.org/10.1093/sysbio/syx063>

637 Milicia, V.J., Rodríguez, M.F., Hagiwara, J.C., Soto, M.S., 2021. Utilización de plantas puente en
638 el mejoramiento del complejo *Calibrachoa-Petunia*. *Chil. J. Agric. Anim. Sci.*, 37, 136–
639 150. <https://doi.org/10.29393/CHJAAS37-16UPVM40016>

640 Minh, B.Q., Hahn, M.W., Lanfear, R., 2020. New Methods to Calculate Concordance Factors for
641 Phylogenomic Datasets. *Mol. Biol. Evol.* 37, 2727–2733.
642 <https://doi.org/10.1093/molbev/msaa106>

643 Morales-Briones, D.F., Kadereit, G., Tefarikis, D.T., Moore, M.J., Smith, S.A., Brockington,
644 S.F., Timoneda, A., Yim, W.C., Cushman, J.C., Yang, Y., 2021. Disentangling Sources of
645 Gene Tree Discordance in Phylogenomic Data Sets: Testing Ancient Hybridizations in
646 *Amaranthaceae* s.l. *Syst. Biol.* 70, 219–235. <https://doi.org/10.1093/sysbio/syaa066>

647 Nevado, B., Contreras-Ortiz, N., Hughes, C., Filatov, D.A. 2018. Pleistocene glacial cycles drive
648 isolation, gene flow and speciation in the high-elevation Andes. *New Phytol.*, 219, 779–
649 793. <https://doi.org/10.1111/nph.15243>

650 Nguyen, L.-T., Schmidt, H.A., Von Haeseler, A., Minh, B.Q., 2015. IQ-TREE: A Fast and
651 Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol.*

652 Biol. Evol. 32, 268–274. <https://doi.org/10.1093/molbev/msu300>

653 Olmstead, R.G., Bohs, L., Migid, H.A., Santiago-Valentin, E., Garcia, V.F., Collier, S.M., 2008.

654 A molecular phylogeny of the Solanaceae. TAXON 57, 1159–1181.

655 <https://doi.org/10.1002/tax.574010>

656 Olschowski, S., Meyer, L., Mußmann, V., Serek, M. 2013. Investigation of Crossing Barriers

657 between *Calibrachoa* and *Petunia* species. Acta Hort. 1007, 765–769.

658 <https://doi.org/10.17660/ActaHortic.2013.1007.90>

659 Owens, G.L., Huang, K., Todesco, M., Rieseberg, L.H., 2023. Re-evaluating Homoploid

660 Reticulate Evolution in *Helianthus* Sunflowers. Mol. Biol. Evol. 40, msad013.

661 <https://doi.org/10.1093/molbev/msad013>

662 Pezzi, P.H., Guzmán-Rodríguez, S., Giudicelli, G.C., Turchetto, C., Bombarely, A., Freitas, L.B.,

663 2022. A convoluted tale of hybridization between two *Petunia* species from a transitional

664 zone in South America. Perspect. Plant Ecol. Evol. Syst. 56, 125688.

665 <https://doi.org/10.1016/j.ppees.2022.125688>

666 Rabiee, M., Sayyari, E., Mirarab, S., 2019. Multi-allele species reconstruction using ASTRAL.

667 Mol. Phylogenet. Evol. 130, 286–296. <https://doi.org/10.1016/j.ympev.2018.10.033>

668 Ranwez, V., Douzery, E.J.P., Cambon, C., Chantret, N., Delsuc, F., 2018. MACSE v2: Toolkit for

669 the Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. Mol.

670 Biol. Evol. 35, 2582–2584. <https://doi.org/10.1093/molbev/msy159>

671 Reck-Kortmann, M., Silva-Arias, G.A., Segatto, A.L.A., Mäder, G., Bonatto, S.L., Freitas, L.B.

672 2014. Multilocus phylogeny reconstruction: new insights into the evolutionary history of

673 the genus *Petunia*. Mol. Phylogenetics Evol., 81, 19–28.

674 <https://doi.org/10.1016/j.ympev.2014.08.022>

675 Reck-Kortmann, M., Silva-Arias, G.A., Stehmann, J.R., Greppi, J.A., Freitas, L.B., 2015.

676 Phylogenetic relationships of *Petunia patagonica* (Solanaceae) revealed by molecular

677 and biogeographical evidence. Phytotaxa 222, 17.

678 <https://doi.org/10.11646/phytotaxa.222.1.2>

679 Rodrigues, D.M., Caballero-Villalobos, L., Turchetto, C., Assis Jacques, R., Kuhlemeier, C.,

680 Freitas, L.B., 2018. Do we truly understand pollination syndromes in *Petunia* as much as

681 we suppose? AoB PLANTS. <https://doi.org/10.1093/aobpla/ply057>

682 Rothfels, C.J., Johnson, A.K., Hovenkamp, P.H., Swofford, D.L., Roskam, H.C., Fraser-Jenkins,

683 C.R., Windham, M.D., Pryer, K. M. 2015. Natural hybridization between genera that

684 diverged from each other approximately 60 million years ago. Am. Nat., 185, 433–442.

685 <https://doi.org/10.1086/679662>

686 Särkinen, T., Bohs, L., Olmstead, R.G., Knapp, S., 2013. A phylogenetic framework for

687 evolutionary study of the nightshades (Solanaceae): a dated 1000-tip tree. BMC Evol.

688 Biol. 13, 214. <https://doi.org/10.1186/1471-2148-13-214>

689 Schwarz, G., 1978. Estimating the Dimension of a Model. Ann. Stat. 6, 461–464.

690 Shaw, J.M.H., 2007. A new hybrid genus for *Calibrachoa* × *Petunia* (Solanaceae). Hanburyana

691 2, 50–51.

692 Smith, S.A., Moore, M.J., Brown, J.W., Yang, Y., 2015. Analysis of phylogenomic datasets

693 reveals conflict, concordance, and gene duplications with examples from animals and

694 plants. BMC Evol. Biol. 15, 150. <https://doi.org/10.1186/s12862-015-0423-0>

695 Solís-Lemus, C., Ané, C., 2016. Inferring Phylogenetic Networks with Maximum

696 Pseudolikelihood under Incomplete Lineage Sorting. PLOS Genet. 12, e1005896.

697 <https://doi.org/10.1371/journal.pgen.1005896>

- 698 Solís-Lemus, C., Bastide, P., Ané, C., 2017. PhyloNetworks: A Package for Phylogenetic
699 Networks. *Mol. Biol. Evol.* 34, 3292–3298. <https://doi.org/10.1093/molbev/msx235>
- 700 Solís-Lemus, C., Yang, M., Ané, C., 2016. Inconsistency of Species Tree Methods under Gene
701 Flow. *Syst. Biol.* 65, 843–851. <https://doi.org/10.1093/sysbio/syw030>
- 702 Soltis, P.S., Soltis, D.E., 2009. The Role of Hybridization in Plant Speciation. *Annu. Rev. Plant*
703 *Biol.* 60, 561–588. <https://doi.org/10.1146/annurev.arplant.043008.092039>
- 704 Song, L., Florea, L., 2015. Rcorrector: efficient and accurate error correction for Illumina RNA-
705 seq reads. *GigaScience* 4, 48. <https://doi.org/10.1186/s13742-015-0089-y>
- 706 Souza, A., Giudicelli, G.C., Teixeira, M.C., Turchetto, C., Bonatto, S.L., Freitas, L.B., 2022.
707 Genetic diversity in micro-endemic plants from highland grasslands in southern Brazil.
708 *Bot. J. Linn. Soc.* 199, 235–251. <https://doi.org/10.1093/botlinnean/boab057>
- 709 Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
710 large phylogenies. *Bioinformatics* 30, 1312–1313.
711 <https://doi.org/10.1093/bioinformatics/btu033>
- 712 Stehmann, J.R., Lorenz-Lemke, A.P., Freitas, L.B., Semir, J., 2009. The Genus *Petunia*, in:
713 Gerats, T., Strommer, J. (Eds.), *Petunia*. Springer New York, New York, NY, pp. 1–28.
714 https://doi.org/10.1007/978-0-387-84796-2_1
- 715 Stull, G.W., Pham, K.K., Soltis, P.S., Soltis, D.E., 2023. Deep reticulation: the long legacy of
716 hybridization in vascular plant evolution. *Plant J.* 114, 743–766.
717 <https://doi.org/10.1111/tpj.16142>
- 718 Suarez-Gonzalez, A., Lexer, C., Cronk, Q.C.B., 2018. Adaptive introgression: a plant
719 perspective. *Biol. Lett.* 14, 20170688. <https://doi.org/10.1098/rsbl.2017.0688>
- 720 Sugiura, N., 1978. Further analysis of the data by Akaike's information criterion and the finite
721 corrections: Further analysis of the data by Akaike's. *Commun. Stat. - Theory Methods* 7,
722 13–26. <https://doi.org/10.1080/03610927808827599>
- 723 Suvorov, A., Kim, B.Y., Wang, J., Armstrong, E.E., Peede, D., D'Agostino, E.R.R., Price, D.K.,
724 Waddell, P.J., Lang, M., Courtier-Orgogozo, V., David, J.R., Petrov, D., Matute, D.R.,
725 Schridder, D.R., Comeault, A.A., 2022. Widespread introgression across a phylogeny of
726 155 *Drosophila* genomes. *Curr. Biol.* 32, 111-123.e5.
727 <https://doi.org/10.1016/j.cub.2021.10.052>
- 728 Swofford, D.L., 2003. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods).
729 Version 4. Sinauer Associates, Sunderland, Massachusetts. USA.
- 730 Taylor, S.A., Larson, E.L. 2019. Insights from genomes into the evolutionary importance and
731 prevalence of hybridization in nature. *Nat. Ecol. Evol.* 3, 170–177.
732 <https://doi.org/10.1038/s41559-018-0777-y>
- 733 Than, C., Ruths, D., Nakhleh, L., 2008. PhyloNet: a software package for analyzing and
734 reconstructing reticulate evolutionary relationships. *BMC Bioinformatics* 9, 322.
735 <https://doi.org/10.1186/1471-2105-9-322>
- 736 Todesco, M., Pascual, M.A., Owens, G.L., Ostevik, K.L., Moyers, B.T., Hübner, S., Heredia,
737 S.M., Hahn, M.A., Caseys, C., Bock, D.G., Rieseberg, L.H., 2016. Hybridization and
738 extinction. *Evol. Appl.* 9, 892–908. <https://doi.org/10.1111/eva.12367>
- 739 Turchetto, C., Segatto, A.L.A., Silva-Arias, G.A., Beduschi, J., Kuhlemeier, C., Bonatto, S.L.,
740 Freitas, L.B., 2019. Contact zones and their consequences: hybridization between two
741 ecologically isolated wild *Petunia* species. *Bot. J. Linn. Soc.* boz022.
742 <https://doi.org/10.1093/botlinnean/boz022>
- 743 Wheeler, L.C., Dunbar-Wallis, A., Schutz, K., Smith, S.D., 2023. Evolutionary walks through

744 flower colour space driven by gene expression in *Petunia* and allies (Petunieae). Proc. R.
745 Soc. B Biol. Sci. 290, 20230275. <https://doi.org/10.1098/rspb.2023.0275>

746 Wheeler, L.C., Walker, J.F., Ng, J., Deanna, R., Dunbar-Wallis, A., Backes, A., Pezzi, P.H.,
747 Palchetti, M.V., Robertson, H.M., Monaghan, A., De Freitas, L.B., Barboza, G.E.,
748 Moyroud, E., Smith, S.D., 2022. Transcription Factors Evolve Faster Than Their
749 Structural Gene Targets in the Flavonoid Pigment Pathway. Mol. Biol. Evol. 39,
750 msac044. <https://doi.org/10.1093/molbev/msac044>

751 Whitney, K.D., Ahern, J.R., Campbell, L.G., Albert, L.P., King, M.S., 2010. Patterns of
752 hybridization in plants. Perspect. Plant Ecol. Evol. Syst. 12, 175–182.
753 <https://doi.org/10.1016/j.ppees.2010.02.002>

754 Wijsman, H.J.W., De Jong, J.H., 1985. On the interrelationships of certain species of *Petunia* IV.
755 Hybridization between *P. linearis* and *P. calycina* and nomenclatorial consequences in the
756 *Petunia* group. Acta Bot. Neerlandica 34, 337–349. <https://doi.org/10.1111/j.1438-8677.1985.tb01925.x>


757

758 Yu, Y., Degnan, J.H., Nakhleh, L., 2012. The Probability of a Gene Tree Topology within a
759 Phylogenetic Network with Applications to Hybridization Detection. PLoS Genet. 8,
760 e1002660. <https://doi.org/10.1371/journal.pgen.1002660>


761 Zhang, C., Rabiee, M., Sayyari, E., Mirarab, S., 2018. ASTRAL-III: polynomial time species
762 tree reconstruction from partially resolved gene trees. BMC Bioinformatics 19, 153.
763 <https://doi.org/10.1186/s12859-018-2129-y>

764 Zuntini, A.R., Carruthers, T., Maurin, O., Bailey, P.C., Leempoel, K., Brewer, G.E., Epitawalage,
765 N., Françaço, E., Gallego-Paramo, B., McGinnie, C., Negrão, R., Roy, S.R., Simpson, L.,
766 Toledo Romero, E., Barber, V.M.A., Botigué, L., Clarkson, J.J., Cowan, R.S., Dodsworth,
767 S., Johnson, M.G., Kim, J.T., Pokorny, L., Wickett, N.J., Antar, G.M., DeBolt, L.,
768 Gutierrez, K., Hendriks, K.P., Hoewener, A., Hu, A.-Q., Joyce, E.M., Kikuchi, I.A.B.S.,
769 Larridon, I., Larson, D.A., De Lirio, E.J., Liu, J.-X., Malakasi, P., Przelomska, N.A.S.,
770 Shah, T., Viruel, J., Allnutt, T.R., Ameka, G.K., Andrew, R.L., Appelhans, M.S., Arista, M.,
771 Ariza, M.J., Arroyo, J., Arthan, W., Bachelier, J.B., Bailey, C.D., Barnes, H.F., Barrett,
772 M.D., Barrett, R.L., Bayer, R.J., Bayly, M.J., Biffin, E., Biggs, N., Birch, J.L., Bogarín, D.,
773 Borosova, R., Bowles, A.M.C., Boyce, P.C., Bramley, G.L.C., Briggs, M., Broadhurst, L.,
774 Brown, G.K., Bruhl, J.J., Bruneau, A., Buerki, S., Burns, E., Byrne, M., Cable, S.,
775 Calladine, A., Callmander, M.W., Cano, Á., Cantrill, D.J., Cardinal-McTeague, W.M.,
776 Carlsen, M.M., Carruthers, A.J.A., De Castro Mateo, A., Chase, M.W., Chatrou, L.W.,
777 Cheek, M., Chen, S., Christenhusz, M.J.M., Christin, P.-A., Clements, M.A., Coffey, S.C.,
778 Conran, J.G., Cornejo, X., Couvreur, T.L.P., Cowie, I.D., Csiba, L., Darbyshire, I.,
779 Davidse, G., Davies, N.M.J., Davis, A.P., Van Dijk, K., Downie, S.R., Duretto, M.F.,
780 Duvall, M.R., Edwards, S.L., Egglí, U., Erkens, R.H.J., Escudero, M., De La Estrella, M.,
781 Fabriani, F., Fay, M.F., Ferreira, P.D.L., Ficinski, S.Z., Fowler, R.M., Frisby, S., Fu, L.,
782 Fulcher, T., Galbany-Casals, M., Gardner, E.M., German, D.A., Giaretta, A., Gibernau, M.,
783 Gillespie, L.J., González, C.C., Goyder, D.J., Graham, S.W., Grall, A., Green, L., Gunn,
784 B.F., Gutiérrez, D.G., Hackel, J., Haevermans, T., Haigh, A., Hall, J.C., Hall, T., Harrison,
785 M.J., Hatt, S.A., Hidalgo, O., Hodgkinson, T.R., Holmes, G.D., Hopkins, H.C.F., Jackson,
786 C.J., James, S.A., Jobson, R.W., Kadereit, G., Kahandawala, I.M., Kainulainen, K., Kato,
787 M., Kellogg, E.A., King, G.J., Klejevska, B., Klitgaard, B.B., Klopper, R.R., Knapp, S.,
788 Koch, M.A., Leebens-Mack, J.H., Lens, F., Leon, C.J., Lévillé-Bourret, É., Lewis, G.P.,
789 Li, D.-Z., Li, L., Liede-Schumann, S., Livshultz, T., Lorence, D., Lu, M., Lu-Irving, P.,

790 Luber, J., Lucas, E.J., Luján, M., Lum, M., Macfarlane, T.D., Magdalena, C., Mansano,
791 V.F., Masters, L.E., Mayo, S.J., McColl, K., McDonnell, A.J., McDougall, A.E., McLay,
792 T.G.B., McPherson, H., Meneses, R.I., Merckx, V.S.F.T., Michelangeli, F.A., Mitchell,
793 J.D., Monro, A.K., Moore, M.J., Mueller, T.L., Mummenhoff, K., Munzinger, J., Muriel, P.,
794 Murphy, D.J., Nargar, K., Nauheimer, L., Nge, F.J., Nyffeler, R., Orejuela, A., Ortiz, E.M.,
795 Palazzesi, L., Peixoto, A.L., Pell, S.K., Pellicer, J., Penneys, D.S., Perez-Escobar, O.A.,
796 Persson, C., Pignal, M., Pillon, Y., Pirani, J.R., Plunkett, G.M., Powell, R.F., Prance, G.T.,
797 Puglisi, C., Qin, M., Rabeler, R.K., Rees, P.E.J., Renner, M., Roalson, E.H., Rodda, M.,
798 Rogers, Z.S., Rokni, S., Rutishauser, R., De Salas, M.F., Schaefer, H., Schley, R.J.,
799 Schmidt-Lebuhn, A., Shapcott, A., Al-Shehbaz, I., Shepherd, K.A., Simmons, M.P.,
800 Simões, A.O., Simões, A.R.G., Siros, M., Smidt, E.C., Smith, J.F., Snow, N., Soltis, D.E.,
801 Soltis, P.S., Soreng, R.J., Sothers, C.A., Starr, J.R., Stevens, P.F., Straub, S.C.K., Struwe,
802 L., Taylor, J.M., Telford, I.R.H., Thornhill, A.H., Tooth, I., Trias-Blasi, A., Udovicic, F.,
803 Utteridge, T.M.A., Del Valle, J.C., Verboom, G.A., Vonow, H.P., Vorontsova, M.S., De Vos,
804 J.M., Al-Wattar, N., Waycott, M., Welker, C.A.D., White, A.J., Wieringa, J.J., Williamson,
805 L.T., Wilson, T.C., Wong, S.Y., Woods, L.A., Woods, R., Worboys, S., Xanthos, M., Yang,
806 Y., Zhang, Y.-X., Zhou, M.-Y., Zmarzty, S., Zuloaga, F.O., Antonelli, A., Bellot, S., Crayn,
807 D.M., Grace, O.M., Kersey, P.J., Leitch, I.J., Sauquet, H., Smith, S.A., Eiserhardt, W.L.,
808 Forest, F., Baker, W.J., 2024. Phylogenomics and the rise of the angiosperms. *Nature* 629,
809 843–850. <https://doi.org/10.1038/s41586-024-07324-0>
810



812
 813 **Figure S1.** ASTRAL phylogenetic tree depicting individuals without species assignments.
 814 Numerical values on branches indicate local posterior probabilities that are below 100. The scale
 815 bar denotes branch length in coalescent units.



817


818 **Figure S2.** Phylogenetic tree depicting the relationship within the *Petunia-Calibrachoa-Fabiana*

819 clade and the outgroups. The tree was constructed using quartets of taxa and the coalescent

820 model with SVDQuartets. Branch numbers indicate nonparametric bootstrap values, derived


821 from 200 bootstrap replicates.

822



823
824
825
826
827

Figure S3. Summary Graphs of QuIBL Results. The upper graph illustrates the proportion of topologies not solely explained by Incomplete Lineage Sorting (ILS), while the lower graph depicts the proportion of gene trees indicating introgression events.



828
829 **Figure S4.** Log-likelihood results from the SNaQ analysis. The steepest change in log-likelihood
830 signifies the optimal network with a single reticulation.
831

832 **Supplemental tables are available on GitHub.**