

European wild honeybee populations are endangered

Patrick L. Kohl^{1,2*} & Benjamin Rutschmann^{3,4*}

¹ Department of Livestock Population Genomics, University of Hohenheim, Garbenstrasse 17, 70599 Stuttgart, Germany

² Center for Biodiversity and Integrative Taxonomy (KomBioTa), University of Hohenheim and Stuttgart State Museum of Natural History, Wollgrasweg 23, 70599 Stuttgart, Germany

³ Department of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg, Am Hubland, 97074 Würzburg, Germany

⁴ Agroecology and Environment, Agroscope, Reckenholzstrasse 191, 8046, Zurich, Switzerland

*Correspondence: patrick.kohl@uni-hohenheim.de; benjamin.rutschmann@agroscope.admin.ch

ORCID: PLK: 0000-0001-9278-978X; BR: 0000-0001-6589-6408

Keywords: pollinator decline, IUCN, Red List, monitoring, conservation status, beekeeping, native bees

Abstract

The population trends of wild western honeybees (*Apis mellifera*) have been neglected by conservationists because the species has been considered to consist of managed colonies only. New data suggest that wild honeybee colonies (still) make up one sixth to one fifth of the overall European honeybee population. The population trends of wild cohorts can be evaluated like those of any other native wild species, albeit with some methodological adjustments to account for the bias introduced by swarms emigrating from managed cohorts. We used data on wild colony survival rates from seven European countries to model their autonomous population changes over ten-year periods, the time frame considered for population evaluation by the International Union for the Conservation of Nature (IUCN). Populations of wild honeybee colonies currently represent demographic sinks in six out of seven countries. With an estimated population decline of 56% per decade, the honeybee

should be considered “Endangered” in the wild in Europe. We believe that the formal recognition and study of honeybee populations beyond apiculture can have far-reaching consequences for the perception of this unique bee species and pollinator conservation in general.

Introduction

The western honeybee, *Apis mellifera*, is among the most abundant flower visitors across its cosmopolitan geographic range (Hung et al., 2018), so its persistence on the species level seems to be of little concern in conservation science. It is rarely appreciated, however, that the existence of honeybees in regions with temperate climates is rather remarkable. The tribe of honeybees (Apini) only comprises about a dozen species, most of which are confined to small distribution areas in (sub-)tropical Asia (Smith, 2021). Only the western honeybee (in Europe) and, to a lesser degree, the eastern honeybee *Apis cerana* (in Asia) have evolved the peculiar ability to withstand long winters (Ruttner, 1988). Forming perennial colonies, they are unique components of their respective temperate-climate bee communities. Besides its colonisation of the temperate zone, *Apis mellifera* is special because it is the only extant honeybee native to Africa, Western Asia and Europe. This shows that *Apis mellifera*, while common as a species, is uncommon from both a functional and a phylogenetic perspective. Preserving its intra-specific genetic diversity and potential to evolve naturally outside of apiculture can be seen as important goals in insect conservation (Fontana et al., 2018; Requier et al., 2019).

Interestingly, bee conservationists have not evaluated the conservation status of the western honeybee, likely because it is considered an entirely managed, or even domesticated, species, and thus, part of agriculture or the livestock sector (Geldmann & González-Varo, 2018).

Unfortunately, such view neglects that there have always been both wild and managed honeybee colonies (Kohl & Rutschmann, 2018; Visick & Ratnieks, 2023; Niklasson et al., 2024). The mating of honeybee queens and drones takes place in free flight and is not usually controlled by beekeepers, and swarms from wild colonies have traditionally been a source of beekeeping operations, while, in turn, swarm emigration from apiaries is never completely prevented. Therefore, there has always been a genetic and demographic continuum between cohorts of wild and managed colonies and there is no scientific basis for a general biological distinction between managed and wild honeybees (Johnson, 2023).

What is typically confounded are the questions whether honeybees are native versus allochthonous on the one hand or living wild versus managed on the other hand. Certain subspecies of honeybee, mainly Italian (*A. m. ligustica*) and Carnolian bees (*A. m. carnica*)

are preferred in modern apiculture. They have been introduced to countries outside their native ranges, and, due to a general lack of mating control, have hybridized with the native populations (De la Rúa et al., 2009). However, these commercially desired subspecies and their hybrids are, evolutionary speaking, not generally less wild than other honeybee populations. Furthermore, in most of the introduced regions, the respective native genetic backgrounds have not been completely replaced (Requier et al., 2019). Therefore, where *Apis mellifera* is native as a species, not considering the conservation of wild hybrid populations would mean applying a different standard compared to other taxa that are assessed on the species level. (Of course, considering wild honeybees on the species level, as we do here, does not preclude promoting the conservation of wild honeybees on the subspecies level, where this is applicable (e.g., Valentine et al., 2025).)

Apart from the question whether the attribute “wild” is appropriate for free-living colonies of honeybee within their native range, a simple reason for their neglect in conservation is the widespread belief that they have gone extinct. Due to habitat loss and/or the introduction of the invasive ectoparasitic mite *Varroa destructor* (starting around 50 years ago), the latter being a main driver of colony mortality in apiculture (Traynor et al., 2020), there would simply be no wild subpopulations left to protect. Unfortunately, there is no data documenting such alleged extinction in the wild (Kohl & Rutschmann, 2018), which is why *Apis mellifera* was listed as “Data deficient” in the “European Red List of Bees” by the International Union for the Conservation of Nature (IUCN) as of 2014 (Nieto et al., 2014). Throughout the last decade, however, motivated by the insight that temperate-adapted honeybees still form viable wild populations in their introduced range in the northeastern United States (reviewed by Seeley, 2019), researchers have started searching for, and (re-)discovered, wild honeybee colonies in Europe (Oleksa, Gawroński & Tofilski, 2013; Kohl & Rutschmann, 2018; Requier et al., 2020; Browne et al., 2021; Moro et al., 2021; Lang, Albouy & Zewen, 2022; Rutschmann et al., 2022; Albouy, 2024; Rutschmann, Remter & Roth, 2024; Visick & Ratnieks, 2024). The average colony density was estimated to be 0.26 colonies per km², equivalent to about 5.5 million colonies, 17.8% of the overall European honeybee population (which comprises ca. 25.4 million managed hives) (Visick & Ratnieks, 2023).

These figures suggest that wild honeybees are still ecologically and evolutionary relevant. In the context of ongoing updates to the European Red List of Bees (Ghisbain et al., 2023), the (re-)discovery of wild honeybees provides the opportunity to make the first informed assessment of their conservation status in Europe. Determining which IUCN Red List

category currently applies to a species is an important formal step since it will help to objectively decide whether strategies to promote the respective populations are needed. The rationale is that wild species are periodically assigned a category of threat ranging from “Least Concern” to “Extinct in the Wild” based on, for example, observed or estimated changes in population size within the last ten years (IUCN 2024).

Given that a distinction between wild and managed honeybees can generally be based only on the current mode of living of their colonies rather than on morphology or genetics, the practical problem arises of how to specifically assess the population trends of wild subpopulations. Even if actual nesting sites of wild honeybees are known, directly assessing temporal changes in the number of colonies inhabiting an area still does not suffice to estimate the wild honeybee population trend: managed colonies can revert to the wild by leaving their apiaries as swarms and establishing natural nests in cavities of their choice during the reproductive season, masking the potential autonomous population trend of the existing wild subpopulation. For example, approximately 10% of trees with black woodpecker (*Dryocopus martius*) cavities are occupied by honeybees in managed forests in Germany each summer, suggesting the existence of a stable population of wild colonies. However, studying the fate of many individual colonies revealed that only a few survive to the next spring; their relatively high abundance in summer could only be explained by the massive annual immigration of swarms from apiaries (Kohl, Rutschmann & Steffan-Dewenter, 2022).

In awareness of the problems that arise when dealing with species that contain managed populations, the IUCN (2024) has determined the following threshold to decide whether a population should be considered “wild” in the first place and thus be assessed: “*Subpopulations dependent on direct intervention are not considered wild, if they would go extinct within 10 years without ‘intensive’ management such as [...] regularly supplementing the population from captive stock to prevent imminent extinction*”. The emigration of honeybee swarms from apiaries, though a natural process, can be regarded as a regular supplementation of the wild population from captive stock. Therefore, the relevant question for determining whether a cohort of free-living colonies deserves the attribute “wild subpopulation” and whether its conservation should be assessed is, “How would the population of wild colonies change over time if there was no immigration of swarms from managed colonies?”. Understanding this requires the conceptual distinction between wild and managed cohorts and thinking of them as a metapopulation (Kohl, 2023; Rutschmann, Remter & Roth, 2024). The autonomous change of the wild subpopulation is then expressed by a

statistic called the net reproductive rate (R_0), which in turn can be derived from the average survival and reproductive rates of its members. Here, we use available data on wild colony survival rates from seven European countries to model how their populations would change autonomously over ten-year periods. Based on these projected population trends, we suggest an informed IUCN Red List category for the species in Europe.

Methods

Framework for studying the demography of wild honeybee populations

There is generally no physical or genetic barrier between managed (Figure 1a) and wild honeybee colonies (Figure 1b) and, contrary to common misconception, no stable “breeds” of domesticated honeybees exist (Seeley, 2019). We therefore define wild *Apis mellifera* colonies based on their mode of living as colonies that are ownerless and unmanaged and live in their natural comb nests, at sites they have occupied themselves (Rutschmann, Remter & Roth, 2024) (see supplementary information for further information).

To study the population demography of wild honeybee colonies, it is practical to consider regional honeybee populations as metapopulations consisting of cohorts (“subpopulations”) of managed and wild colonies (Figure 1 c) (Kohl, 2023; Rutschmann, Remter & Roth, 2024). The size of the wild cohort is affected by immigration, emigration, survival, and natality. Immigration occurs when managed colonies swarm, disperse from their home apiary and occupy a cavity on their own, thus becoming members of the wild subpopulation. Emigration occurs when beekeepers capture swarms of wild colonies, be it directly or by luring swarms into bait hives. (We assume that this process is negligible in most regions since beekeepers usually obtain new colonies by splitting their own stock or trading with other beekeepers.) To understand how the size of the wild subpopulation would change intrinsically, the survival (s) and the natality (b) (i.e., reproduction) of the colonies that are already members of that cohort need to be considered. Given the annual survival and natality rates, the net reproductive rate (R_0) describes how the population would change from year to year if no immigration occurred, with values < 1 denoting population decline and values > 1 denoting population stability or increase. Since temperate honeybee colonies typically start reproducing after completing their first year (i.e., from the age of 1 year) and colonies are hermaphrodites (all colonies can, in principle, produce swarms with queens [female colony reproduction] besides

drones [male colony reproduction]), the net reproductive rate of a wild honeybee subpopulation is (Kohl, Rutschmann & Steffan-Dewenter, 2022):

$$R_0 = s + s * b.$$

This index is the basis of our wild honeybee population trend projections.




Figure 1. *A) Apiary with honeybee colonies managed in movable-frame hives. B) Examples of typical wild honeybee nesting sites: tree cavity (left) and hollow space in a wall (right). C) Metapopulation model of managed and wild honeybee colonies. Managed and wild colonies belong to one biological population; there is a genetic exchange through the random mating of queens and drones, and colonies can migrate between managed and wild cohorts (“subpopulations”). Selection pressure under apicultural management (natural and artificial selection) and under wild conditions (natural selection) are expected to differ. Figure modified after (Kohl, 2023).*

Estimating current trends of populations of wild honeybee colonies

The annual survival rate in a population of wild honeybee colonies can be empirically studied by making repeated surveys of known nest sites (Seeley, 2017). Nine monitoring studies were available that provided information on the occupation histories of a total of 698 nesting sites of wild honeybee colonies in seven European countries gathered over several years (Figure 2, Table 1): Poland (Oleksa et al. 2013; A. Oleksa, pers. communication 2024), Germany (three studies) (Lang, Albouy & Zewen, 2022; Kohl, Rutschmann & Steffan-Dewenter, 2022; Rutschmann, Remter & Roth, 2024), France (Albouy, 2024), Spain (Rutschmann et al. 2022; Rutschmann & Kohl, unpublished data), England (Visick & Ratnieks, 2024; O. Visick, pers. communication 2025), Switzerland (Cordillot, 2024) and Luxembourg (J. Park & R. Dammé, pers. communication 2025). For each of the studied wild populations and for each country, we derived a point estimate of the average annual colony survival rate (s_{est}) from the available data (see supplementary information for details). The natality rate of wild honeybee colonies (the number of swarms produced per colony per year) is difficult to determine in the field. We therefore assumed that wild colonies produce, on average, $n = 2$ swarms per year based on studies that examined the reproductive behaviour of unmanaged honeybee colonies living in hives with limited volumes (see Rutschmann, Kohl & Steffan-Dewenter, 2024 and references therein). However, it is well known that the swarming rate can vary strongly between years due to differences in weather, with zero to four swarms per colony being typical. Although no direct data on between-year variation in colony-level swarming rate was available, we had information on the annual variation in cavity re-occupations by wild colonies in Germany (Kohl, Rutschmann & Steffan-Dewenter, 2022) and Spain (Rutschmann et al. 2022; Rutschmann & Kohl, unpublished data). Assuming that such rates of nest re-occupation closely correlate with the average colony-level swarming rate in the respective year and region (since new nests are founded by swarms), we used the observed coefficient of variation in the annual nest re-occupation rate (12.5%), a dimensionless number, to derive a realistic level of annual variation in the natality rate. Specifically, we simulated environmental stochasticity in the annual natality rate (b_{sim}) by drawing from a normal distribution truncated between 0 and 4 (the minimum and maximum number of swarms per colony expected), with a mean of 2 and a coefficient of variation of 0.125 (standard deviation: 0.25). Using estimated survival and simulated natality rates, we then simulated annual net reproductive rate for each studied wild population and country as:

$$R_{0_{sim}} = s_{est} + s_{est} \times b_{sim}$$

The relative wild honeybee population trend over a hypothetical decade without migration is given as $R_{0_sim}^x$, with x denoting the year from 0 to 10, and with R_{0_sim} randomly varying among years. Accordingly, we estimated the rate of population change per 10 years as percentage by calculating $(R_{0_sim}^{10} - 1) * 100$. These steps were repeated 10000 times for each of the eight studied populations to account for random variation in population fate due to stochasticity in the swarming rate. We present the mean and the 90% confidence interval (CI) of the population trend projections for each studied population, for each of the seven countries, and for the whole of Europe (using the median of the seven country-level estimates of the survival rate). In accordance with IUCN guidelines (IUCN 2024), we then used the mean projected population change per decade to classify a population as “Least concern” (reduction: < 30%), “Vulnerable” (reduction: $\geq 30\%$), “Endangered” (reduction: $\geq 50\%$), “Critically Endangered” (reduction: $\geq 80\%$), or “Extinct in the Wild” (reduction: 100%) (see supplementary information for details).




Figure 2. Map of Europe highlighting seven countries with data on wild honeybee colony survival rates (grey) and the respective study regions (dots, except for Luxembourg). One

study used data from three different forest regions (“German forest”; locations marked as “a”, “b”, “c”).

Table 1. Overview of eight studies on wild honeybee population demography that provide wild colony survival data. The annual colony survival rates (s_{est}) are the values used in this study (corrected by us for potential silent colony turnovers, if necessary; see supplementary information).

Country	Region	Information provided	Number of nest sites monitored	Study period	s_{est}	Reference
Germany	Three managed forest regions in southern Germany	Annual survival rate	77	2017–2021	0.106	(Kohl, Rutschmann & Steffan-Dewenter, 2022)
Germany	City of Dortmund	Annual survival rate	30	2018–2021	0.121	(Lang, Albouy & Zewen, 2022, Table 4 therein)
Germany	City of Munich	Summer, winter and spring survival rates	107	2016–2023	0.133	(Rutschmann, Remter & Roth, 2024, p. 14 therein)
Poland	Northeastern Poland	Proportion of colonies remaining after one and two years	67	2013–2015	0.307	(Oleksa, Gawroński & Tofilski, 2013), A. Oleksa, pers. communication 2024
Spain	Comarca de la Limia, Galicia	Annual survival rate	37	2019–2023	0.293	(Rutschmann et al., 2022) Rutschmann & Kohl, unpublished data
France	County of Saintonge	Summer, winter and spring survival rates	140	2018–2021	0.318	(Albouy, 2024, p. 623 therein)
Switzerland	Switzerland Regions north of the Alps	Annual survival rate	106	2020–2023	0.096	(Cordillot, 2024, p. 107 therein)
England	Southeast England	Annual survival rate	61	2021–2024	0.394	(Visick & Ratnieks, 2024; O. Visick, pers. communication 2025)
Luxembourg	Luxembourg	Summer, winter and spring survival rates	73	2019–2025	0.316	(J. Park & R. Dammé, pers. communication 2025)

Results

In six out of seven European countries, populations of wild honeybees were projected to be in decline (Figure 3). In Germany (where data from three independent studies are in close agreement) and in Switzerland, wild honeybee populations must be assumed to disappear

within 10 years without immigration (projected decline per decade [upper, lower 90% CI]: -100% [-100%, -100%]). In Spain and Poland, wild honeybee populations were projected to be declining by -78% [-68%, -87%] and -56% [-35%, -73%] per decade, meaning that they are “Endangered” according to IUCN red list criteria. The wild populations monitored in Luxembourg and France appear to be in moderate decline (-41% [-13%, -64%] and -38% [-7%, -61%] per decade); they can be classified as “Vulnerable”. Only the wild honeybee population in South England appears to have the intrinsic capacity to increase in size (projected increase per decade: +432% [+692%, +230%]) and therefore is categorised as “Least Concern”. The median population change per decade is -56% [-34%, -73%], meaning the overall wild honeybee population in Europe can be classified as “Endangered”.




Figure 3. *A) Projected wild honeybee population trends over hypothetical ten-year periods for nine studied populations and for the whole of Europe. Relative changes in population size are based on net reproductive rates, which in turn rely on observed colony survival rates and an assumed average natality rate of two swarms per colony per year. Lines describe the mean trends based on 10000 simulations for each population. The grey shaded area describes the 90% confidence interval for Europe. B) Wild honeybee population remaining (means and 90% confidence intervals based on 10000 simulations) and their mean relative change (percentages) after hypothetical ten-year periods of intrinsic development (without migration) for seven European countries and the whole of Europe. The second y-axis shows the associated IUCN conservation categories: LC = Least Concern, VU = Vulnerable, EN = Endangered, CR = Critically Endangered. (The simulations for Germany were run using the*

average of the colony survival rates of the three studies. Confidence intervals for England, Germany and Switzerland are not visible because they are outside the y-axis range.)

Discussion

Using data on wild colony survival rates gathered in seven countries during the last decade, we modelled how the sizes of wild honeybee subpopulations would have intrinsically changed over a hypothetical period of ten years. While our estimate of a median population size reduction of -56% suggests that the overall wild honeybee subpopulation represents a demographic sink, a key insight is that it would clearly not have gone completely extinct within a decade. Therefore, on the European level, the cohort of wild *Apis mellifera* can be formally considered a “wild subpopulation” according to IUCN guidelines, and the assessment of its population trend for the Red List of bees is justified.

What holds for the whole of Europe, however, is not true for each of the individual populations studied. In fact, our second key insight is that there are remarkable differences in wild population demographics among regions. On the low end of the spectrum, there are the populations studied in Germany and in Switzerland north of the Alps (Lang, Albouy & Zewen, 2022; Kohl, Rutschmann & Steffan-Dewenter, 2022; Cordillot, 2024; Rutschmann, Remter & Roth, 2024). Strikingly, these are clearly expected to go extinct within a hypothetical ten-year time window, given their low net reproductive rates. Therefore, conditional upon the discovery of populations with higher average colony survival rates, the honeybee can be considered “Extinct in the Wild” in these countries. On the other end of the spectrum, there is the population studied in southeast England (Visick & Ratnieks, 2024), which seems to be stable on its own. Here, the projected excess population means that it can act as a demographic source for other populations (including the managed subpopulation), and that it has strong evolutionary potential. Between these extremes are the populations studied in France (Albouy, 2024), Luxembourg (J. Park & R. Dammé, pers. communication 2025), Poland (Oleksa, Gawroński & Tofilski, 2013), and Spain (Rutschmann et al., 2022), which apparently fare much better than the ones in Germany and Switzerland, but also represent demographic sinks.

When referring to the individual populations by their countries of origin, it needs to be highlighted that each is represented by one or a few studies only. Depending on the spatiotemporal scopes, studies will be better or worse in representing the average

demographics in the respective wider regions, the delineation of which is arbitrary. It also needs to be considered that wild subpopulations will be structured spatially, and since they can occupy more or less favourable habitats, there will be source-sink dynamics among such patches (Dias, 1996). For example, the study of the wild honeybee subpopulation in Galicia, Spain, was conducted in a landscape dominated by intensive agriculture, not representative of the province. Since colony winter survival is significantly lower in agricultural than in adjacent semi-natural habitats (Rutschmann et al., 2022), we might have overestimated the rate of decline for that population. Despite the uncertainty in the reliability of the country-level estimates, we believe that the combination of data from nine studies and seven countries lead to an informative estimate of the overall wild honeybee population trend. Based on the best data available, we must assume that the overall population of wild honeybees is in decline in Europe.

One might argue that this intrinsic decline is not a problem insofar as the wild subpopulation could be (partly) replenished annually by feral swarms from the managed subpopulation. This view makes sense when considering that wild and managed honeybees are intrinsically the same. However, it lacks appreciation for the potential of significant genetic differences to evolve between managed and wild populations due to the accumulation of minor allele frequency changes at many loci. Such variation can reflect adaptations to contrasting selection pressures or different demographic histories. For example, a lack of medical treatment can select for colonies resisting *Varroa* mites and/or their transmitted viruses in the wild population (Mikheyev et al., 2015), while frequent trade of managed colonies (or queen bees) across countries can lead to a higher proportion of non-native alleles in the managed compared to the wild subpopulation. In fact, beekeepers have largely altered managed honeybee populations through the importation of non-native subspecies and subsequent introgressive hybridisation in many regions (De la Rúa et al., 2009; Requier et al., 2019; Espregueira Themudo et al., 2020; Kükner, Kence & Kence, 2021). If native alleles have a selective advantage over non-native ones, promoting wild honeybee subpopulations can even lead to a progressive increase in the frequency of such alleles in the overall population (Wayne & Shaffer, 2016). In that case, conserving wild honeybee populations could be understood as a long-term means of conserving native honeybee subspecies.

The perspective suggested by the present analysis of explicitly conserving wild honeybee populations is currently not implemented. Usually, no difference is made between wild and managed honeybee colonies and the species is more often regarded as a problem, rather than a

target, of European wild bee conservation (Geldmann & González-Varo, 2018). However, wild honeybee subpopulations likely suffer from the same environmental pressures as non-*Apis* wild bees (Goulson et al., 2015; Rutschmann et al., 2022; Kohl et al., 2023). In fact, local wild honeybees can be the bees most directly affected by long-distance apiary migrations, high densities of managed hives, and the trade of queen honeybees across countries (Requier et al., 2019; Panziera et al., 2022; Martínez-López, Ruiz & De La Rúa, 2022). Given that the number of managed colonies is currently growing (see supplementary information, figure S1), there is probably an ongoing shift in the relative importance of selection pressure in the overall honeybee population, with conditions under management gaining importance over wild conditions. We do not know which evolutionary role the existence of a wild subpopulation plays in the long-term health and adaptability of the overall honeybee population, but it is likely to be positive (Requier et al., 2019; Panziera et al., 2022). Therefore, the decline of wild honeybees should not only be of concern for insect conservation but also for apiculture and crop pollination in agriculture. Furthermore, the perspective that honeybees could be a subject of conservation themselves also has an impact on the conflict between apiculture and non-*Apis* bee conservation (Geldmann & González-Varo, 2018; Henry & Rodet, 2018; Beaurepaire et al., 2025). For example, the existence of a wild honeybee subpopulation in a conservation area is an excellent argument for restricting apiculture in that area.

We suggest that wild honeybee monitoring programs should be continued and adopted in many more regions to allow for better inference of population trends (Seeley, 2017; Kohl, Rutschmann & Steffan-Dewenter, 2022; Albouy, 2024; Moro et al., 2024; Rutschmann, Remter & Roth, 2024). Furthermore, we need to know which factors limit wild honeybee colony survival (e.g. Kohl et al., 2023), and why wild subpopulations fare so much better in some regions compared to others. According to the principle that we can only conserve what we understand, we believe that recognising wild honeybee subpopulations as real and tangible subjects of population demographic studies is a key step in the conservation of this unique component of the European bee fauna.

Data Availability Statement

The data used for this work are referenced in the main text.

Author contributions

Patrick L. Kohl was involved in conceptualisation, methodology, investigation, formal analysis, writing – original draft. Benjamin Rutschmann was involved in conceptualisation, methodology, investigation, acquisition of datasets, formal analysis, visualization, writing – review & editing.

Conflict of interest declaration

We declare we have no competing interests.

Acknowledgements

The idea for the presented data analysis arose in the context of a re-assessment of *Apis mellifera* for the European Red List of Bees. We would therefore like to thank the whole team of co-assessors for the discussions on the issue. We specifically thank Oliver Visick, Andrzej Oleksa, and John Park and Roger Dammé for providing information on the results of wild honeybee monitoring studies in England, Poland and Luxembourg, and Fabrice Requier for providing valuable comments on the manuscript. We are grateful to Elena Reiriz Martínez for proofreading.

References

- Albouy V. 2024. RÉSULTATS DU SUIVI 2017-2023 DES COLONIES D’ABEILLES MELLIFÈRES À L’ÉTAT SAUVAGE DANS LE NORD DE LA NOUVELLE AQUITAINE – SECTEUR 1. *Annales de la Société des Sciences Naturelles de la Charente-Maritime* 11:613–768.
- Beaurepaire AL, Hogendoorn K, Kleijn D, Otis GW, Potts SG, Singer TL, Boff S, Pirk C, Settele J, Paxton RJ, Raine NE, Tosi S, Williams N, Klein A-M, Le Conte Y, Campbell JW, Williams GR, Marini L, Brockmann A, Sgolastra F, Boyle N, Neuditschko M, Straub L, Neumann P, Charrière J-D, Albrecht M, Dietemann V. 2025. Avenues

towards reconciling wild and managed bee proponents. *Trends in Ecology & Evolution* 40:7–10. DOI: 10.1016/j.tree.2024.11.009.

Browne KA, Hassett J, Geary M, Moore E, Henriques D, Soland-Reckeweg G, Ferrari R, Mac Loughlin E, O’Brien E, O’Driscoll S, Young P, Pinto MA, McCormack GP. 2021. Investigation of free-living honey bee colonies in Ireland. *Journal of Apicultural Research* 60:229–240. DOI: 10.1080/00218839.2020.1837530.

Cordillot F. 2024. Erste Suche nach wilden Honigbienen (*Apis mellifera* L., 1758) auf der Schweizer Alpennordseite. *Entomo Helvetica* 17:97–114.

De la Rúa P, Jaffé R, Dall’Olio R, Muñoz I, Serrano J. 2009. Biodiversity, conservation and current threats to European honeybees. *Apidologie* 40:263–284. DOI: 10.1051/apido/2009027.

Dias PC. 1996. Sources and sinks in population biology. *Trends in ecology & evolution* 11:326–330.

Espregueira Themudo G, Rey-Iglesia A, Robles Tascón L, Bruun Jensen A, Da Fonseca RR, Campos PF. 2020. Declining genetic diversity of European honeybees along the twentieth century. *Scientific Reports* 10:10520. DOI: 10.1038/s41598-020-67370-2.

Fontana P, Costa C, Prisco GD, Ruzzier E, Annoscia D, Battisti A, Caoduro G, Carpana E, Contessi A, Dal A, Dall R, Cristofaro AD, Felicioli A, Floris I, Gardi T, Lodesani M, Malagnini V, Manias L, Manino A, Marzi G, Massa B, Mutinelli F, Nazzi F, Pennacchio F, Porporato M, Stoppa G, Tormen N, Valentini M, Segrè A. 2018. Appeal for biodiversity protection of native honey bee subspecies of *Apis mellifera* in Italy (San Michele all’Adige declaration). 71:257–271.

Geldmann J, González-Varo JP. 2018. Conserving honey bees does not help wildlife. *Science* 359:392–393. DOI: 10.1126/science.aar2269.

Ghisbain G, Rosa P, Bogusch P, Flaminio S, Divelec RL, Dorchin A, Kasperek M, Kuhlmann M, Litman J, Mignot M, Müller A, Praz C, Radchenko VG, Rasmont P, Risch S,

- Roberts SPM, Smit J, Wood TJ, Michez D, Reverté S. 2023. The new annotated checklist of the wild bees of Europe (Hymenoptera: Anthophila). *Zootaxa* 5327:1–147. DOI: 10.11646/zootaxa.5327.1.1.
- Goulson D, Nicholls E, Botías C, Rotheray EL. 2015. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science* 347:1255957. DOI: 10.1126/science.1255957.
- Henry M, Rodet G. 2018. Controlling the impact of the managed honeybee on wild bees in protected areas. *Scientific Reports* 8:9308. DOI: 10.1038/s41598-018-27591-y.
- Hung K-LJ, Kingston JM, Albrecht M, Holway DA, Kohn JR. 2018. The worldwide importance of honey bees as pollinators in natural habitats. *Proceedings of the Royal Society B: Biological Sciences* 285:20172140. DOI: 10.1098/rspb.2017.2140.
- IUCN Standards and Petitions Committee. 2024. *Guidelines for Using the IUCN Red List Categories and Criteria*.
- Johnson BR. 2023. *Honey bee biology*. Princeton & Oxford: Princeton University Press.
- Kohl PL. 2023. The buzz beyond the beehive: population demography, parasite burden and limiting factors of wild-living honeybee colonies in Germany. Dissertation Thesis. University of Würzburg. DOI: 10.25972/OPUS-33032.
- Kohl PL, Rutschmann B. 2018. The neglected bee trees: European beech forests as a home for feral honey bee colonies. *PeerJ* 6:e4602. DOI: 10.7717/peerj.4602.
- Kohl PL, Rutschmann B, Sikora LG, Wimmer N, Zahner V, D’Alvise P, Hasselmann M, Steffan-Dewenter I. 2023. Parasites, depredators, and limited resources as potential drivers of winter mortality of feral honeybee colonies in German forests. *Oecologia* 202:465–480. DOI: 10.1007/s00442-023-05399-6.
- Kohl PL, Rutschmann B, Steffan-Dewenter I. 2022. Population demography of feral honeybee colonies in central European forests. *Royal Society Open Science* 9:220565. DOI: 10.1098/rsos.220565.

432 Kükreer M, Kence M, Kence A. 2021. Honey Bee Diversity Is Swayed by Migratory
 433 Beekeeping and Trade Despite Conservation Practices: Genetic Evidence for the
 434 Impact of Anthropogenic Factors on Population Structure. *Frontiers in Ecology and*
 435 *Evolution* 9:556816. DOI: 10.3389/fevo.2021.556816.

436 Lang UM, Albouy V, Zewen C. 2022. Comparative monitoring of free-living honey bee
 437 colonies in three Western European regions. *Natural Bee Husbandry Magazine* 23.

438 Martínez-López V, Ruiz C, De La Rúa P. 2022. “Migratory beekeeping and its influence on
 439 the prevalence and dispersal of pathogens to managed and wild bees.” *International*
 440 *Journal for Parasitology: Parasites and Wildlife* 18:184–193. DOI:
 441 10.1016/j.ijppaw.2022.05.004.

442 Mikheyev AS, Tin MMY, Arora J, Seeley TD. 2015. Museum samples reveal rapid evolution
 443 by wild honey bees exposed to a novel parasite. *Nature Communications* 6:7991. DOI:
 444 10.1038/ncomms8991.

445 Moro A, Albouy V, Dickey M, Kohl PL, McCormack GP, Remter F, Requier F, Rogenstein S,
 446 Rutschmann B, Thiele MJ, Visick O, Dubaić JB. 2024. A Protocol for Monitoring
 447 Populations of Free-Living Western Honey Bees in Temperate Regions. *Bee World*
 448 0:1–5. DOI: 10.1080/0005772X.2024.2402109.

449 Moro A, Beaurepaire A, Dall’Olio R, Rogenstein S, Blacquièrre T, Dahle B, De Miranda JR,
 450 Dietemann V, Locke B, Licón Luna RM, Le Conte Y, Neumann P. 2021. Using Citizen
 451 Science to Scout Honey Bee Colonies That Naturally Survive Varroa destructor
 452 Infestations. *Insects* 12:536. DOI: 10.3390/insects12060536.

453 Nieto A, Roberts SPM, Kemp J, Rasmond P, Kuhlmann M, García Criado M, Biesmeijer JC,
 454 Bogusch P, Dathe HH, De La Rúa P, De Meulemeester T, Dehon M, Dewulf A, Ortiz-
 455 Sánchez FJ, Lhomme P, Pauly A, Potts SG, Praz C, Quaranta M, Radchenko VG,
 456 Scheuchl E, Smit J, Straka J, Terzo M, Tomozil B, Window J, Michez D. 2014.
 457 *European red list of bees*. LU: Publication Office of the European Union.

458 Niklasson M, Svensson E, Leidenberger S, Norrström N, Crawford E. 2024. Free-living
 459 colonies of native honey bees (*Apis mellifera mellifera*) in 19th and early 20th century
 460 Sweden. *Journal of Insect Conservation* 28:389–400. DOI: 10.1007/s10841-023-
 461 00541-4.

462 Oleksa A, Gawroński R, Tofilski A. 2013. Rural avenues as a refuge for feral honey bee
 463 population. *Journal of Insect Conservation* 17:465–472. DOI: 10.1007/s10841-012-
 464 9528-6.

465 Panziera D, Requier F, Chantawannakul P, Pirk CWW, Blacquièrre T. 2022. The Diversity
 466 Decline in Wild and Managed Honey Bee Populations Urges for an Integrated
 467 Conservation Approach. *Frontiers in Ecology and Evolution* 10:767950. DOI:
 468 10.3389/fevo.2022.767950.

469 Requier F, Garnery L, Kohl PL, Njovu HK, Pirk CWW, Crewe RM, Steffan-Dewenter I.
 470 2019. The Conservation of Native Honey Bees Is Crucial. *Trends in Ecology &*
 471 *Evolution* 34:789–798. DOI: 10.1016/j.tree.2019.04.008.

472 Requier F, Paillet Y, Laroche F, Rutschmann B, Zhang J, Lombardi F, Svoboda M, Steffan-
 473 Dewenter I. 2020. Contribution of European forests to safeguard wild honeybee
 474 populations. *Conservation Letters* 13:e12693. DOI: 10.1111/conl.12693.

475 Rutschmann B, Kohl PL, Machado A, Steffan-Dewenter I. 2022. Semi-natural habitats
 476 promote winter survival of wild-living honeybees in an agricultural landscape.
 477 *Biological Conservation* 266:109450. DOI: 10.1016/j.biocon.2022.109450.

478 Rutschmann B, Kohl PL, Steffan-Dewenter I. 2024. Swarming rate and timing of unmanaged
 479 honeybee colonies (*Apis mellifera carnica*) in a forest environment.
 480 *bioRxiv*:2024.09.07.611535. DOI: 10.1101/2024.09.07.611535.

481 Rutschmann B, Remter F, Roth S. 2024. Monitoring free-living honeybee colonies in
 482 Germany: Insights into habitat preferences, survival rates, and Citizen Science

reliability. *bioRxiv*:2024.08.02.606354. DOI:
<https://doi.org/10.1101/2024.08.02.606354>.

Ruttner F. 1988. *Biogeography and Taxonomy of Honeybees*. Berlin, Heidelberg: Springer
 Berlin Heidelberg. DOI: 10.1007/978-3-642-72649-1.

Seeley TD. 2017. Life-history traits of wild honey bee colonies living in forests around Ithaca,
 NY, USA. *Apidologie* 48:743–754. DOI: 10.1007/s13592-017-0519-1.

Seeley TD. 2019. *The Lives of Bees: The Untold Story of the Honey Bee in the Wild*. Princeton
 University Press.

Smith DR. 2021. Biogeography of Honey Bees. *Encyclopedia of Social Insects*.

Traynor KS, Mondet F, De Miranda JR, Techer M, Kowallik V, Oddie MAY, Chantawannakul
 P, McAfee A. 2020. Varroa destructor: A Complex Parasite, Crippling Honey Bees
 Worldwide. *Trends in Parasitology* 36:592–606. DOI: 10.1016/j.pt.2020.04.004.

Valentine A, Moro ,Arrigo, Briggs ,Ethan, Collier ,Brandon, Sandoval ,Kenneth, Binetti
 ,Chiara, Richardson ,Matthew, Wragg ,David, Browne ,Keith A., Barnett ,Mark, and
 McCormack GP. 2025. Introgressive hybridisation puts the distinctive population of
 Apis mellifera mellifera in Ireland at risk: evidence from a multidisciplinary approach.
Journal of Apicultural Research 64:202–216. DOI: 10.1080/00218839.2024.2404297.

Visick OD, Ratnieks FLW. 2023. Density of wild honey bee, Apis mellifera, colonies
 worldwide. *Ecology and Evolution* 13:e10609. DOI: 10.1002/ece3.10609.

Visick OD, Ratnieks FLW. 2024. Ancient, veteran and other listed trees as nest sites for wild-
 living honey bee, Apis mellifera, colonies. *Journal of Insect Conservation* 28:153–
 163. DOI: 10.1007/s10841-023-00530-7.

Wayne RK, Shaffer HB. 2016. Hybridization and endangered species protection in the
 molecular era. *Molecular Ecology* 25:2680–2689. DOI: 10.1111/mec.13642.