

1 **Advancing Amazonian botanical knowledge: a detailed ecological characterization of an**  
2 **open ombrophile forest, southwest Amazonian Brazil**

3 Bruno Umbelino<sup>1,2</sup>, Juliana Stropp<sup>3</sup>, Ingrid Mendes-Silva<sup>4</sup>, Ricardo Aleixo Correia<sup>5,6,7</sup>,

4 Antônio L. P. Silveira<sup>4</sup> and Ana C. M. Malhado<sup>2</sup>

5  
6 <sup>1</sup>Departamento de Ecologia, Instituto de Biologia, Universidade Federal do Rio de Janeiro, Rio  
7 de Janeiro, RJ, Brazil

8  
9 <sup>2</sup>Institute of Biological Sciences and Health, Laboratório de Conservação do Século XXI,  
10 Universidade Federal de Alagoas, Campus A. C. Simões, Av. Lourival Melo Mota, s/n  
11 Tabuleiro dos Martins, Maceió, AL, Brazil

12  
13 <sup>3</sup>Museo Nacional de Ciencias Naturales (MNCN- CSIC), Madrid, Spain

14  
15 <sup>4</sup>Herbarium João Geraldo Kuhlmann, Departamento de Biologia, Universidade Federal de  
16 Rondônia, Br-364 Km 9 s/ Rio Branco, Campus José Ribeiro Filho, Porto Velho, RO, Brazil

17  
18 <sup>5</sup>Department of Geosciences and Geography, University of Helsinki, FI-00014 Helsinki,  
19 Finland

20  
21 <sup>6</sup>Helsinki Institute of Sustainability Science, University of Helsinki, FI-00014 Helsinki, Finland

22  
23 <sup>7</sup>Biodiversity Unit, University of Turku, 20014 Turku, Finland

24  
25 Corresponding author: [brunosilvasantos@gmail.com](mailto:brunosilvasantos@gmail.com)

26  
27 **ORCID: BU** 0000-0002-7602-8400 / **JS** 0000-0002—2831-4066 / **IMS** 0000-0003-0053-  
28 7206 / **RAC** 0000-0001-7359-9091 / **ACMM** 0000-0003-3621-779X

29 **ABSTRACT**

30 Biodiversity inventories present excellent opportunities for ecological investigations and the  
31 classification of different threats to the community, nonetheless these applications are not  
32 frequently employed. Our main objective was to determine the tree and palm community within  
33 a one-hectare areas, also exploring the association between functional attributes and the  
34 projected threat of category for the future (by 2050) in a conservation unit situated within the  
35 Brazilian Amazon Deforestation Arc. We established a rectangular plot (10x1000m) to assess  
36 the community structure. Information on seed and fruit size attributes was obtained from the  
37 literature, along with data on the projected threat category. Overall, two taxa (Rubiaceae sp. and  
38 Rhizophoraceae sp.) were characterized only at the family level, 106 at the genera level  
39 (morpho-species) and 124 until binomial name. We found information about seed size for 55  
40 genera. Medium-sized seeds were the most frequent, occurring in 22 genera, followed by large  
41 seeds (16), small seeds (6), and very small seeds (5). As for the projected threat status for 2050,  
42 we found that 28 species were classified as vulnerable and 16 species as endangered. In our plot  
43 we founded a few numbers of species with many individuals. We conclude that floristic studies  
44 associated with ecological approach carried out in the southwestern region of the Brazilian  
45 Amazon are rare and our study, provides a significant contribution to biodiversity knowledge.

46 **Keywords:** plant inventories, tropical biodiversity, Brazilian Amazon, traits attribute, threat  
47 status

49 The Amazon region is renowned for its extraordinary biodiversity, making it one the most  
50 ecologically diverse regions in the world. Moreover, the acknowledgment of its essential  
51 contribution to critical ecosystem services has propelled the Amazon to attain global  
52 conservation eminence, commanding the attention of national and international scholars alike  
53 (Malhado et al. 2014; dos Santos et al. 2015). Regardless this premise, substantial portions of  
54 this biome remain unexplored. The vastness and isolation of the Amazon present formidable  
55 challenges for conducting systematic and regular investigations within the region, even when  
56 employing appropriate technological advancements (Ladle and Whittaker 2014). It is estimated  
57 that 40% of the Amazonian domain remains untouched by scientific inquiry (Schulman,  
58 Toivonen, and Ruokolainen 2007), indicating a significant research shortfall. Accessibility,  
59 primarily limited by poor riverine and terrestrial transportations routes, as well as the scarcity  
60 of scientific and technological facilities and infrastructure projects (*e.g.*, hydroelectric  
61 installations), restrict research efforts in the Amazon basin (dos Santos et al. 2015; Correia et  
62 al. 2016). Consequently, this situation engenders uncertainties regarding the spatial distribution  
63 of numerous species owing to prevalent sampling biases (Hortal et al. 2015).

64 Nevertheless, the Amazon confronts formidable obstacles characterized by alarming rates of  
65 habitat degradation and species extinctions. In the southern and southeastern regions, the  
66 presence of gaps is attributed to a long history of deforestation for pastureland, commonly  
67 referred to as the “Deforestation Arc”, which has hindered specimen collections in this area  
68 (Stropp et al. 2020). This region has experienced accumulative higher taxa of deforestation  
69 through the years, resulting in loss of numerous species, some of which may have become  
70 extinct even before being scientifically documented, rendering previously collected records  
71 obsolete (Stropp et al. 2020). Long-term floristic inventories play a crucial role in advancing  
72 our understanding of diversity patterns, the distribution of flora in the Amazon region and their

73 dynamics. Some studies of tree species in the Amazonian region, currently estimated to range  
74 from approximately 7000 to 10 000 species (Cardoso et al. 2017; ter Steege et al. 2020),  
75 remains far from complete, with the potential for as many as 5000 new species trees yet to be  
76 discovered (Ter Steege et al. 2016). It is widely acknowledged that there is a substantial gap in  
77 our knowledge of Amazonia biodiversity, which can be attributed to three primary factors: (i)  
78 deforestation for land-use purposes, (ii) limited accessibility and (iii) a combination of both  
79 factors (Stropp et al. 2020).

80 Even with the availability of diverse estimation methods of species surveys, it is essential to  
81 recognize that the data are susceptible to temporal degradation. Consequently, ensuring their  
82 appropriate utilization in both present and future contexts hold paramount significance  
83 (Tessarolo et al. 2017). Temporal losses of data can arise from the inherent dynamism of natural  
84 systems, encompassing local extinctions, immigration patterns, and biological invasions  
85 (Tessarolo et al. 2017) Additionally, alterations in the taxonomic relationship of biodiversity  
86 categories (Ladle and Hortal 2013) and natural shifts in species distribution and composition  
87 (Dornelas et al. 2012) can contribute to such temporal variations.

88 The literature extensively addresses the knowledge gaps surrounding species occurrences,  
89 commonly referred to as the Wallacean shortfall (Brown and Lomolino 2006; Ladle and  
90 Whittaker 2014; Hortal et al. 2015). This term refers to the insufficiency of existing knowledge  
91 about species distribution and geographical dynamics. From the standpoint of conservation and  
92 policy, it becomes imperative to prioritize extensive data collection efforts in regions  
93 characterized by high conservation value or facing imminent risks of habitat destruction,  
94 particularly where sampling completeness remains low. The Raunkiaer shortfall refers to the  
95 lack of knowledge regarding ecologically relevant species traits, encompassing both intra-  
96 specific and inter-specific trait variations (Hortal et al. 2015). Furthermore, this shortfall

97 extends to comprehending the ecological functions associated with each trait, as well as  
98 elucidating how these functions are affected by such interactions. In addition, it aims to identify  
99 the specific combination of traits that collectively contribute to the performance of distinct  
100 ecosystem functions, thus emphasizing the significance of trait bundling (Díaz et al. 2013).  
101 Significant progress has been made in stablishing a shared repertoires of valuable traits  
102 applicable to various taxa, particularly in plants, alongside the implementation of standardized  
103 sampling protocols (Hortal et al. 2015). Nevertheless, it is important to acknowledge the  
104 existence of substantial shortfall in both taxonomic coverage and geographic representation,  
105 highlighting the need for further research and exploration.

106 Local and descriptive studies highlight a foundational role in establishing our current  
107 understanding of species distributions. Much of this knowledge is derived from forest and  
108 floristic inventories that document the composition and structure of biological communities  
109 within predefined plots. These inventories yield are crucial data that enrich our comprehension  
110 of biodiversity and species distributions. Moreover, when compiled across expansive regions,  
111 forest inventories enable macroecological and biogeographical investigations that would  
112 otherwise be unattainable (ter Steege et al. 2013). However, it is worth noting that such data  
113 often falls short of elucidating the ecological mechanisms that underlie the dynamics and  
114 establishment of species within the study site. This information assumes particular significance  
115 in regions characterized by elevated levels of biodiversity, yet where local knowledge of species  
116 remains limited, and conservation planning is primarily delegated to local government entities  
117 (*e.g.*, state or municipality).

118 Our study aimed to: (i) characterize the structure and composition of a one-hectare open  
119 ombrophile forest area within the Jamari National Forest, located in the southwestern Brazilian  
120 Amazon, within the Deforestation Arc; (ii) analyse the literature about seed size of genera

121 (Malhado et al 2015); (iii) investigate the biogeographical distribution of vegetation types  
122 associated with the species present in our plot; and (iv) assess the future projected threat status  
123 of the species for the year 2050.

## 124 **MATERIAL AND METHODS**

### 125 **Plant inventory plot and botanical identification**

126 We set up a one-hectare plot for plant inventory in September of 2014. We establish one  
127 hundred consecutive sample units measuring 10x10m along a 1,000m transect within the Jamari  
128 National Forest (Floresta Nacional do Jamari – Flona do Jamari). In each sample unit, we  
129 measured the diameter of all trees and palms  $\geq 10$ cm at breast height (approximately at 1.3m  
130 high). The trees were identified in the field with aid of parataxonomists, and material was sent  
131 to specialists for identification for confirmation. Herbarium voucher numbers of all species  
132 sampled were collected and incorporated into the Herbarium Rondoniense João Geraldo  
133 Kuhlmann of the Universidade Federal de Rondônia. The dataset is available in the open  
134 database SpeciesLink (<https://specieslink.net/search/>) and can be accessed using the following  
135 search query [(coll\_groups:(botanical)) AND (coll\_ids:(((764) OR (765) OR (257)))) AND  
136 (county:(Itapuã do Oeste)) AND (locality.normal:(Módulo Potosí))]. We tagged trees of all  
137 morpho-species (fertile and/or sterile) and collected at least one sample per morpho-species.  
138 Species names follow the standard nomenclature provided by Flora do Brasil (BFG, 2021).

### 139 *Analysis of forest structure and plants species diversity*

140 We characterized forest structure by estimating the following parameters for families and  
141 species of plot: absolute density (AD), relative density (RD), absolute frequency (AF), relative  
142 frequency (RF), absolute dominance (ADo), relative dominance (RDo) and importance value  
143 (VI). Patterns of tree spatial aggregation were explored using Payandeh index (Payandeh 1970)  
144 and Shanonn Weaver ( $H'$ ) index and Pielou equability ( $J'$ ). All analysis concerned with forest

145 structure were carried out on Fitopac 2.1 software by Shepherd (2010). The Payandeh, Shannon  
146 Weaver and Pielou equability indices were calculated with Microsoft Excel 2019.

#### 147 *Seed size*

148 We obtained seed size categories for all genera in our plot from Malhado et al. (2015) and  
149 classified them into four levels: very small (<0.5 cm), small (0.5–0.99 cm), medium (1.0–1.99  
150 cm), and large ( $\geq 2.0$  cm).

#### 151 *Distribution and threat status*

152 For all taxa identified at species level, we collected information about their occurrence in each  
153 Brazilian phytogeographic domain and vegetation type according to the Flora do Brasil (BFG,  
154 2021). Moreover, we retrieved for all taxa identified at species level the Brazilian state in which  
155 it has been previously recorded using the Flora do Brasil (BFG, 2021) and the platform  
156 SpeciesLink system (CRIA, 2019). Additionally, we checked whether these taxa were present  
157 in the checklist of Brazilian Amazon tree taxa (ter Steege et al. 2015).

#### 158 *Threat status*

159 We retrieved for all taxa identified at species level the current threat status according to IUCN  
160 (2019) using the Flora do Brasil and the list provided by Ministério do Meio Ambiente (MMA)  
161 (PORTARIA MMA N° 148, DE 7 DE JUNHO DE 2022). Additionally, we incorporated the  
162 projected threat status for 2050 based on Ter Steege et al. (2015). We used the R package ‘flora’  
163 (Carvalho 2017) to retrieve information from the Brazilian Flora Checklist.

## 164 **RESULTS**

### 165 **Forest structure and plants species diversity**

166 We sampled 465 trees and palms belonging to 39 families, 100 genera and 220 species. Overall,  
167 two taxa (Rubiaceae sp. and Rhizophoraceae sp.) were characterized only at the family level,  
168 106 at the genera level (morpho-species) and 124 until binomial name. The richest genera were

169 *Protium* (52), *Eschweilera* (33), *Licania* (31), *Pouteria* (17) and *Iryanthera* (16). One hundred  
 170 and thirty-three species (60.7%) presented absolute density equal to one. Shannon ( $H'$ ) index  
 171 to species was  $5.11 \text{ nats.ind}^{-1}$  and Pielou equability index was 0.94 (Table 1). Fabaceae  
 172 represented approximately 18,58% of the importance value while another 24 families accounted  
 173 for a little more than 10%. We found that, 13 families were monospecific and no species  
 174 presented high relative values for any of the phytosociological parameters. All species studied  
 175 exhibited a value Payandeh index value below 1, there by indicating random spatial distribution.  
 176

177 **Table 1** Phytosociological parameters in decreasing of importance value for tree species recorded in one hectare  
 178 of open ombrophylous forest of Flona do Jamari, Itapuã do Oeste, Rondônia state - 2019. Absolute density (AD),  
 179 relative density (RD), absolute frequency (AF), relative frequency (RF), absolute dominance (ADo), relative  
 180 dominance (RDo) and importance value (IV).

Species	AD	RD	AF	RF	ADo	RDo	IV
<i>Licania macrophylla</i> Benth.	9	1.88	9	1.96	1.11	4.99	2.94
<i>Protium cf. gallosum</i> Daly	13	2.71	12	2.61	0.49	2.18	2.5
<i>Eschweilera bracteosa</i> (Poepp. ex O.Berg) Miers	13	2.71	13	2.83	0.42	1.87	2.47
<i>Erisma bicolor</i> Ducke	6	1.25	6	1.3	1.06	4.74	2.43
<i>Copaifera multijuga</i> Hayne	9	1.88	8	1.74	0.72	3.23	2.28
<i>Goupia glabra</i> Aubl.	2	0.42	2	0.43	1.2	5.37	2.07
<i>Qualea paraensis</i> Ducke	4	0.83	3	0.65	0.8	3.58	1.69
<i>Vatairea</i> sp3	9	1.88	9	1.96	0.22	0.98	1.61
<i>Protium grandifolium</i> Engl.	10	2.08	9	1.96	0.17	0.77	1.6
<i>Euterpe precatória</i> Mart.	10	2.08	9	1.96	0.12	0.55	1.53
<i>Licania rodriguesii</i> Prance	7	1.46	6	1.3	0.32	1.43	1.4



<b>Species</b>	<b>AD</b>	<b>RD</b>	<b>AF</b>	<b>RF</b>	<b>ADo</b>	<b>RDo</b>	<b>IV</b>
<i>Iryanthera lancifolia</i> Ducke	8	1.67	7	1.52	0.15	0.68	1.29
<i>Cespedesia spathulata</i> (Ruiz & Pav.) G.Planch.	9	1.88	7	1.52	0.09	0.41	1.27
<i>Oenocarpus bataua</i> Mart.	7	1.46	6	1.3	0.21	0.93	1.23
<i>Lueheopsis</i> sp2	1	0.21	1	0.22	0.71	3.2	1.21
<i>Protium polybotryum</i> subsp. <i>blackii</i> (Swart) Daly	7	1.46	7	1.52	0.14	0.61	1.2
<i>Licania micrantha</i> Miq.	5	1.04	4	0.87	0.36	1.63	1.18
<i>Eugenia</i> sp.	4	0.83	4	0.87	0.36	1.63	1.11
<i>Helicostylis tomentosa</i> (Poepp. & Endl.) Rusby	6	1.25	6	1.3	0.14	0.61	1.05
<i>Aspidosperma</i> sp3	2	0.42	2	0.43	0.42	1.86	0.9
<i>Cedrelinga cateniformis</i> (Ducke) Ducke	6	1.25	4	0.87	0.12	0.55	0.89
<i>Aspidosperma carapanauba</i> Pichon	2	0.42	2	0.43	0.39	1.73	0.86
<i>Eschweilera</i> sp1	4	0.83	4	0.87	0.2	0.89	0.86
<i>Alchorneopsis cf. floribunda</i> (Benth.) Müll.Arg.	3	0.63	3	0.65	0.29	1.28	0.85
<i>Astronium lecointei</i> Ducke	1	0.21	1	0.22	0.47	2.12	0.85
<i>Theobroma cf. subincanum</i>	5	1.04	5	1.09	0.1	0.43	0.85
<i>Brosimum rubescens</i> Taub.	2	0.42	2	0.43	0.37	1.66	0.84
<i>Eschweilera micrantha</i> (O.Berg) Miers	3	0.63	3	0.65	0.26	1.17	0.82
<i>Virola calophylla</i> Warb.	5	1.04	5	1.09	0.07	0.32	0.82
<i>Caryocar glabrum</i> (Aubl.) Pers.	2	0.42	2	0.43	0.34	1.54	0.8
<i>Peltogyne paniculata</i> Benth.	2	0.42	2	0.43	0.31	1.39	0.75
<i>Swartzia</i> sp4	2	0.42	2	0.43	0.3	1.33	0.73
<i>Virola</i> sp2	2	0.42	2	0.43	0.29	1.28	0.71

<b>Species</b>	<b>AD</b>	<b>RD</b>	<b>AF</b>	<b>RF</b>	<b>ADo</b>	<b>RD<sub>o</sub></b>	<b>IV</b>
<i>Endopleura</i> sp.	1	0.21	1	0.22	0.38	1.68	0.7
<i>Iryanthera sagotiana</i> (Benth.) Warb.	3	0.63	3	0.65	0.18	0.82	0.7
<i>Aniba</i> sp2	2	0.42	2	0.43	0.28	1.23	0.69
<i>Zygia racemosa</i> (Ducke) Barneby & J.W.Grimes	4	0.83	4	0.87	0.07	0.33	0.68
<i>Bowdichia</i> sp1	3	0.63	3	0.65	0.16	0.71	0.66
<i>Protium paniculatum</i> var. <i>riedelianum</i> (Engl.) Daly	4	0.83	4	0.87	0.05	0.22	0.64
<i>Bowdichia</i> sp2	3	0.63	3	0.65	0.14	0.61	0.63
<i>Helicostylis scabra</i> (J.F.Macbr.) C.C.Berg	4	0.83	4	0.87	0.04	0.18	0.63
<i>Tetragastris panamensis</i> (Engl.) Kuntze	4	0.83	3	0.65	0.09	0.4	0.63
<i>Naucleopsis macrophylla</i> Miq.	4	0.83	3	0.65	0.08	0.37	0.62
<i>Brosimum guianense</i> (Aubl.) Huber	3	0.63	3	0.65	0.12	0.54	0.61
<i>Andira</i> sp.	2	0.42	2	0.43	0.21	0.95	0.6
<i>Pseudolmedia laevis</i> (Ruiz & Pav.) J.F.Macbr.	3	0.63	3	0.65	0.1	0.46	0.58
<i>Eschweilera tessmannii</i> R.Knuth	3	0.63	3	0.65	0.1	0.43	0.57
<i>Vatairea</i> sp1	3	0.63	3	0.65	0.1	0.43	0.57
<i>Pouteria caimito</i> (Ruiz & Pav.) Radlk.	3	0.63	3	0.65	0.07	0.33	0.54
<i>Pouteria</i> cf. <i>torta</i> (Mart.) Radlk.	3	0.63	3	0.65	0.08	0.34	0.54
<i>Virola</i> sp1	3	0.63	3	0.65	0.07	0.32	0.53
<i>Heisteria densifrons</i> Engl.	3	0.63	3	0.65	0.06	0.28	0.52
<i>Pouteria</i> cf. <i>guianensis</i> Aubl.	2	0.42	2	0.43	0.16	0.7	0.52
<i>Pseudolmedia</i> sp.	3	0.63	3	0.65	0.06	0.29	0.52
<i>Aspidosperma spruceanum</i> Benth. ex Müll.Arg.	3	0.63	3	0.65	0.05	0.25	0.51

<b>Species</b>	<b>AD</b>	<b>RD</b>	<b>AF</b>	<b>RF</b>	<b>ADo</b>	<b>RDo</b>	<b>IV</b>
<i>Swartzia</i> sp2	3	0.63	3	0.65	0.05	0.24	0.51
<i>Abarema jupunba</i> (Willd.) Britton & Killip	2	0.42	2	0.43	0.15	0.66	0.5
<i>Tachigali paniculata</i> Aubl.	3	0.63	3	0.65	0.05	0.22	0.5
<i>Vochysia</i> sp4	2	0.42	2	0.43	0.14	0.64	0.5
<i>Tachigali</i> sp2	3	0.63	3	0.65	0.05	0.2	0.49
<i>Iryanthera carinata</i>	3	0.63	3	0.65	0.04	0.16	0.48
<i>Iryanthera juruensis</i> Warb.	3	0.63	3	0.65	0.03	0.13	0.47
<i>Lueheopsis</i> sp3	2	0.42	2	0.43	0.12	0.53	0.46
<i>Swartzia</i> sp6	1	0.21	1	0.22	0.2	0.88	0.44
<i>Attalea butifariz</i>	2	0.42	2	0.43	0.1	0.43	0.43
<i>Hymenaea courbaril</i>	1	0.21	1	0.22	0.19	0.87	0.43
Indet spl	2	0.42	2	0.43	0.1	0.43	0.43
<i>Pouteria</i> sp2	2	0.42	2	0.43	0.1	0.43	0.43
<i>Sterigmapetalum cf. obovatum</i> Kuhlm.	2	0.42	2	0.43	0.1	0.45	0.43
<i>Aniba</i> sp3	2	0.42	2	0.43	0.08	0.37	0.41
<i>Attalea cf. maripa</i> (Aubl.) Mart.	2	0.42	2	0.43	0.08	0.36	0.4
<i>Batesia floribunda</i> Benth.	2	0.42	2	0.43	0.08	0.35	0.4
<i>Chrysophyllum</i> sp1	2	0.42	2	0.43	0.08	0.34	0.4
<i>Eschweilera grandiflora</i> (Aubl.) Sandwith	2	0.42	2	0.43	0.07	0.33	0.39
<i>Pouteria</i> sp1	2	0.42	2	0.43	0.07	0.31	0.39
<i>Rhizophoraceae</i> sp.	1	0.21	1	0.22	0.16	0.73	0.39
<i>Erisma</i> sp.	1	0.21	1	0.22	0.16	0.72	0.38

<b>Species</b>	<b>AD</b>	<b>RD</b>	<b>AF</b>	<b>RF</b>	<b>ADo</b>	<b>RDo</b>	<b>IV</b>
<i>Eschweilera andina</i> (Rusby) J.F.Macbr.	2	0.42	2	0.43	0.06	0.28	0.38
<i>Protium</i> sp1	2	0.42	2	0.43	0.06	0.28	0.38
<i>Vitex cymosa</i> Bertero ex Spreng.	2	0.42	2	0.43	0.06	0.29	0.38
<i>Brosimum parinarioides</i> Ducke	2	0.42	2	0.43	0.06	0.27	0.37
<i>Couepia</i> sp.	2	0.42	1	0.22	0.1	0.46	0.37
<i>Abarema adenophora</i> (Ducke) Barneby & J.W.Grimes	1	0.21	1	0.22	0.14	0.65	0.36
<i>Eschweilera truncata</i> A.C.Sm.	2	0.42	2	0.43	0.05	0.24	0.36
<i>Machaerium</i> sp.	2	0.42	2	0.43	0.05	0.22	0.36
<i>Pseudolmedia macrophylla</i> Trécul	2	0.42	2	0.43	0.05	0.22	0.36
<i>Unonopsis</i> sp.	2	0.42	2	0.43	0.05	0.22	0.36
<i>Neea floribunda</i> Poepp. & Endl.	2	0.42	2	0.43	0.04	0.19	0.35
<i>Pterocarpus officinalis</i> Jacq.	2	0.42	2	0.43	0.05	0.21	0.35
<i>Oxandra</i> sp.	2	0.42	2	0.43	0.04	0.16	0.34
<i>Protium crassipetalum</i> Cuatrec.	2	0.42	2	0.43	0.03	0.16	0.34
<i>Protium</i> sp2	2	0.42	2	0.43	0.04	0.17	0.34
<i>Swartzia</i> sp1	2	0.42	2	0.43	0.04	0.18	0.34
<i>Erisma bracteosum</i> Ducke	2	0.42	2	0.43	0.03	0.15	0.33
<i>Lecythis</i> sp2	1	0.21	1	0.22	0.12	0.55	0.33
<i>Licania canescens</i> Benoist	2	0.42	2	0.43	0.03	0.14	0.33
<i>Vouacapoua americana</i> Aubl.	2	0.42	2	0.43	0.03	0.13	0.33
<i>Clarisia racemosa</i> Ruiz & Pav.	2	0.42	2	0.43	0.02	0.1	0.32
<i>Licania lata</i> J.F.Macbr.	2	0.42	2	0.43	0.02	0.11	0.32

<b>Species</b>	<b>AD</b>	<b>RD</b>	<b>AF</b>	<b>RF</b>	<b>ADo</b>	<b>RD<sub>o</sub></b>	<b>IV</b>
<i>Picramnia</i> sp.	2	0.42	2	0.43	0.02	0.1	0.32
<i>Pouteria filipes</i> Eyma	2	0.42	1	0.22	0.07	0.32	0.32
<i>Protium ferrugineum</i> (Engl.) Engl.	2	0.42	2	0.43	0.02	0.11	0.32
<i>Protium</i> sp5	2	0.42	2	0.43	0.02	0.11	0.32
<i>Pseudopiptadenia</i> sp.	1	0.21	1	0.22	0.12	0.52	0.32
<i>Psychotria</i> sp.	2	0.42	2	0.43	0.02	0.11	0.32
<i>Swartzia</i> sp5	2	0.42	2	0.43	0.02	0.1	0.32
<i>Virola</i> sp3	2	0.42	2	0.43	0.02	0.1	0.32
<i>Micropholis cylindrocarpa</i> (Poepp.) Pierre	2	0.42	2	0.43	0.02	0.09	0.31
<i>Micropholis guyanensis</i> (A.DC.) Pierre	2	0.42	2	0.43	0.02	0.09	0.31
<i>Peltogyne excelsa</i> Ducke	2	0.42	2	0.43	0.02	0.09	0.31
<i>Mouriri</i> sp.	1	0.21	1	0.22	0.1	0.46	0.3
<i>Lueheopsis</i> sp1	1	0.21	1	0.22	0.08	0.37	0.27
<i>Tachigali</i> sp4	1	0.21	1	0.22	0.08	0.38	0.27
<i>Aspidosperma</i> sp1	1	0.21	1	0.22	0.07	0.31	0.25
<i>Brosimum</i> sp2	1	0.21	1	0.22	0.07	0.32	0.25
<i>Ocotea</i> sp6	1	0.21	1	0.22	0.07	0.31	0.25
<i>Peltogyne cf. venosa</i> subsp. <i>densiflora</i> (Spruce ex Benth.) M.F.Silva	1	0.21	1	0.22	0.07	0.31	0.25
<i>Tapirira guianensis</i> Aubl.	1	0.21	1	0.22	0.07	0.33	0.25
<i>Vatairea</i> sp2	1	0.21	1	0.22	0.07	0.31	0.25
<i>Vochysia</i> sp2	1	0.21	1	0.22	0.07	0.33	0.25
<i>Aniba</i> sp4	1	0.21	1	0.22	0.07	0.3	0.24

<b>Species</b>	<b>AD</b>	<b>RD</b>	<b>AF</b>	<b>RF</b>	<b>ADo</b>	<b>RD<sub>o</sub></b>	<b>IV</b>
<i>Indet</i> sp1	1	0.21	1	0.22	0.07	0.3	0.24
<i>Licania gracilipes</i> Taub.	1	0.21	1	0.22	0.07	0.3	0.24
<i>Nectandra</i> sp2	1	0.21	1	0.22	0.06	0.28	0.24
<i>Ocotea</i> sp4	1	0.21	1	0.22	0.06	0.26	0.23
<i>Licania</i> sp2	1	0.21	1	0.22	0.05	0.23	0.22
<i>Couepia guianensis</i> Aubl.	1	0.21	1	0.22	0.05	0.21	0.21
<i>Ecclinusa</i> sp.	1	0.21	1	0.22	0.04	0.2	0.21
<i>Lecythis</i> sp3	1	0.21	1	0.22	0.05	0.21	0.21
<i>Pouteria anomala</i> (Pires) T.D.Penn.	1	0.21	1	0.22	0.05	0.21	0.21
<i>Pseudopiptadenia psilostachya</i> (DC.) G.P.Lewis & M.P.Lima	1	0.21	1	0.22	0.04	0.2	0.21
<i>Qualea</i> sp1	1	0.21	1	0.22	0.04	0.19	0.21
<i>Apeiba</i> sp2	1	0.21	1	0.22	0.04	0.18	0.2
<i>Eschweilera coriacea</i> (DC.) S.A.Mori	1	0.21	1	0.22	0.04	0.17	0.2
<i>Eschweilera</i> sp2	1	0.21	1	0.22	0.04	0.17	0.2
<i>Minuartia guianensis</i> Aubl.	1	0.21	1	0.22	0.04	0.16	0.2
<i>Platymiscium</i> sp.	1	0.21	1	0.22	0.04	0.17	0.2
<i>Alchornea</i> sp1	1	0.21	1	0.22	0.03	0.14	0.19
<i>Anacardium cf. giganteum</i> W.Hancock ex Engl.	1	0.21	1	0.22	0.03	0.14	0.19
<i>Astronium</i> sp.	1	0.21	1	0.22	0.03	0.14	0.19
<i>Glycydendron amazonicum</i> Ducke	1	0.21	1	0.22	0.03	0.13	0.19
<i>Lecythis</i> sp1	1	0.21	1	0.22	0.03	0.14	0.19
<i>Licania sothersiae</i> Prance	1	0.21	1	0.22	0.03	0.14	0.19

<b>Species</b>	<b>AD</b>	<b>RD</b>	<b>AF</b>	<b>RF</b>	<b>ADo</b>	<b>RD<sub>o</sub></b>	<b>IV</b>
<i>Micropholis trunciflora</i> Ducke	1	0.21	1	0.22	0.03	0.13	0.19
<i>Myrcia</i> sp.	1	0.21	1	0.22	0.03	0.14	0.19
<i>Nectandra pulverulenta</i> Nees	1	0.21	1	0.22	0.03	0.13	0.19
<i>Ocotea</i> sp2	1	0.21	1	0.22	0.03	0.13	0.19
<i>Ocotea</i> sp3	1	0.21	1	0.22	0.03	0.14	0.19
<i>Pouteria campanulata</i> Baehni	1	0.21	1	0.22	0.03	0.13	0.19
<i>Protium</i> sp3	1	0.21	1	0.22	0.03	0.13	0.19
<i>Ruizterania cassiquiarensis</i> (Spruce ex Warm.) Marc.-Berti	1	0.21	1	0.22	0.03	0.14	0.19
<i>Tachigali</i> sp1	1	0.21	1	0.22	0.03	0.15	0.19
<i>Vochysia</i> sp1	1	0.21	1	0.22	0.03	0.13	0.19
<i>Alchornea</i> sp2	1	0.21	1	0.22	0.02	0.1	0.18
<i>Apeiba</i> sp1	1	0.21	1	0.22	0.02	0.11	0.18
<i>Casearia javitensis</i> Kunth	1	0.21	1	0.22	0.02	0.1	0.18
<i>Chimarrhis duckeana</i> Delprete	1	0.21	1	0.22	0.02	0.1	0.18
<i>Dalbergia</i> sp.	1	0.21	1	0.22	0.03	0.12	0.18
<i>Dialium guianense</i> (Aubl.) Sandwith	1	0.21	1	0.22	0.02	0.1	0.18
Indet sp.	1	0.21	1	0.22	0.02	0.11	0.18
<i>Inga</i> sp1	1	0.21	1	0.22	0.02	0.11	0.18
<i>Leonia glycyarpa</i> Ruiz & Pav.	1	0.21	1	0.22	0.02	0.11	0.18
<i>Machaerium multifoliolatum</i> Ducke	1	0.21	1	0.22	0.03	0.12	0.18
<i>Pradosia decipiens</i> Ducke	1	0.21	1	0.22	0.02	0.1	0.18
<i>Siparuna</i> sp.	1	0.21	1	0.22	0.03	0.12	0.18

<b>Species</b>	<b>AD</b>	<b>RD</b>	<b>AF</b>	<b>RF</b>	<b>ADo</b>	<b>RD<sub>o</sub></b>	<b>IV</b>
<i>Tachigali</i> sp5	1	0.21	1	0.22	0.02	0.1	0.18
<i>Trattinnickia glaziovii</i> Swart	1	0.21	1	0.22	0.03	0.11	0.18
<i>Virola mollissima</i> (A.DC.) Warb.	1	0.21	1	0.22	0.02	0.11	0.18
<i>Xylopia nitida</i> Dunal	1	0.21	1	0.22	0.02	0.11	0.18
<i>Caryodendron</i> sp.	1	0.21	1	0.22	0.02	0.07	0.17
<i>Couma utilis</i> (Mart.) Müll.Arg.	1	0.21	1	0.22	0.01	0.07	0.17
<i>Eschweilera</i> sp3	1	0.21	1	0.22	0.01	0.07	0.17
<i>Hymenolobium</i> sp.	1	0.21	1	0.22	0.02	0.07	0.17
<i>Iryanthera</i> sp2	1	0.21	1	0.22	0.02	0.08	0.17
<i>Licania caudata</i> Prance	1	0.21	1	0.22	0.01	0.07	0.17
<i>Naucleopsis caloneura</i> (Huber) Ducke	1	0.21	1	0.22	0.02	0.07	0.17
<i>Protium altsonii</i> Sandwith	1	0.21	1	0.22	0.02	0.07	0.17
<i>Qualea</i> sp2	1	0.21	1	0.22	0.02	0.08	0.17
<i>Tachigali</i> sp3	1	0.21	1	0.22	0.01	0.07	0.17
<i>Vochysia citrifolia</i> Poir.	1	0.21	1	0.22	0.02	0.07	0.17
<i>Abuta dwyeriana</i> Krukoff & Barneby	1	0.21	1	0.22	0.01	0.04	0.16
<i>Aniba</i> sp1	1	0.21	1	0.22	0.01	0.05	0.16
<i>Aparisthium cordatum</i> (A.Juss.) Baill.	1	0.21	1	0.22	0.01	0.05	0.16
<i>Aspidosperma</i> sp3	1	0.21	1	0.22	0.01	0.06	0.16
<i>Brosimum lactescens</i> (S.Moore) C.C.Berg	1	0.21	1	0.22	0.01	0.04	0.16
<i>Brosimum</i> sp1	1	0.21	1	0.22	0.01	0.04	0.16
<i>Cariniana micrantha</i> Ducke	1	0.21	1	0.22	0.01	0.05	0.16



<b>Species</b>	<b>AD</b>	<b>RD</b>	<b>AF</b>	<b>RF</b>	<b>ADo</b>	<b>RD<sub>o</sub></b>	<b>IV</b>
<i>Cecropia cf. polystachya</i> Trécul	1	0.21	1	0.22	0.01	0.04	0.16
<i>Combretum assimile</i> Eichler	1	0.21	1	0.22	0.01	0.04	0.16
<i>Copaifera glycyarpa</i> Ducke	1	0.21	1	0.22	0.01	0.04	0.16
<i>Cybianthus</i> sp.	1	0.21	1	0.22	0.01	0.06	0.16
<i>Dinizia excelsa</i> Ducke	1	0.21	1	0.22	0.01	0.06	0.16
<i>Erismia floribundum</i> Rudge	1	0.21	1	0.22	0.01	0.04	0.16
<i>Eschweilera ovalifolia</i> (DC.) Nied.	1	0.21	1	0.22	0.01	0.05	0.16
<i>Guarea</i> sp.	1	0.21	1	0.22	0.01	0.04	0.16
<i>Heisteria</i> sp.	1	0.21	1	0.22	0.01	0.05	0.16
<i>Heisteria spruceana</i> Engl.	1	0.21	1	0.22	0.01	0.06	0.16
<i>Hirtella rodriguesii</i> Prance	1	0.21	1	0.22	0.01	0.05	0.16
<i>Hymenolobium pulcherrimum</i> Ducke	1	0.21	1	0.22	0.01	0.04	0.16
<i>Inga leiocalycina</i> Benth.	1	0.21	1	0.22	0.01	0.05	0.16
<i>Inga rubiginosa</i> (Rich.) DC.	1	0.21	1	0.22	0.01	0.04	0.16
<i>Inga</i> sp2	1	0.21	1	0.22	0.01	0.04	0.16
<i>Iryanthera</i> sp1	1	0.21	1	0.22	0.01	0.05	0.16
<i>Lecythis</i> sp4	1	0.21	1	0.22	0.01	0.04	0.16
<i>Licania</i> sp1	1	0.21	1	0.22	0.01	0.06	0.16
<i>Machaerium aureiflorum</i> Ducke	1	0.21	1	0.22	0.01	0.04	0.16
<i>Miconia</i> sp1	1	0.21	1	0.22	0.01	0.06	0.16
<i>Miconia</i> sp2	1	0.21	1	0.22	0.01	0.04	0.16
<i>Micropholis</i> sp.	1	0.21	1	0.22	0.01	0.04	0.16

<b>Species</b>	<b>AD</b>	<b>RD</b>	<b>AF</b>	<b>RF</b>	<b>ADo</b>	<b>RDo</b>	<b>IV</b>
<i>Naucleopsis glabra</i> Spruce ex Pittier	1	0.21	1	0.22	0.01	0.06	0.16
<i>Naucleopsis stipularis</i>	1	0.21	1	0.22	0.01	0.04	0.16
<i>Nectandra</i> sp1	1	0.21	1	0.22	0.01	0.05	0.16
<i>Nectandra</i> sp3	1	0.21	1	0.22	0.01	0.06	0.16
<i>Ocotea rhodophylla</i> Vicent.	1	0.21	1	0.22	0.01	0.04	0.16
<i>Ocotea</i> sp1	1	0.21	1	0.22	0.01	0.04	0.16
<i>Ocotea</i> sp5	1	0.21	1	0.22	0.01	0.04	0.16
<i>Ouratea cf. densiflora</i> Pilg.	1	0.21	1	0.22	0.01	0.04	0.16
<i>Oxandra xylopioides</i> Diels	1	0.21	1	0.22	0.01	0.04	0.16
<i>Perebea</i> sp.	1	0.21	1	0.22	0.01	0.05	0.16
<i>Protium</i> sp4	1	0.21	1	0.22	0.01	0.05	0.16
<i>Protium tenuifolium</i> (Engl.) Engl	1	0.21	1	0.22	0.01	0.05	0.16
<i>Quararibea guianensis</i> Aubl.	1	0.21	1	0.22	0.01	0.04	0.16
<i>Quiina florida</i> Tul.	1	0.21	1	0.22	0.01	0.04	0.16
<i>Rubiaceae</i> sp.	1	0.21	1	0.22	0.01	0.06	0.16
<i>Salacia</i> sp.	1	0.21	1	0.22	0.01	0.06	0.16
<i>Simarouba amara</i> Aubl.	1	0.21	1	0.22	0.01	0.05	0.16
<i>Swartzia</i> sp3	1	0.21	1	0.22	0.01	0.06	0.16
<i>Symphonia</i> sp.	1	0.21	1	0.22	0.01	0.05	0.16
<i>Taralea</i> sp.	1	0.21	1	0.22	0.01	0.04	0.16
<i>Theobroma</i> sp.	1	0.21	1	0.22	0.01	0.05	0.16
<i>Toulicia pulvinata</i> Radlk.	1	0.21	1	0.22	0.01	0.04	0.16

Species	AD	RD	AF	RF	ADo	RD <sub>o</sub>	IV
<i>Vochysia</i> sp3	1	0.21	1	0.22	0.01	0.05	0.16
<i>Xylopia</i> sp.	1	0.21	1	0.22	0.01	0.04	0.16

181

182 **Seed size**

183 We found information about seed size for 55 genera. Medium-sized seeds were the most  
 184 frequent, occurring in 22 genera, followed by large seeds (16), small seeds (6), and very small  
 185 seeds (5). Additionally, six genera had seeds that varied between small/medium (3) and  
 186 medium/large (3). The most richness genera in our plot, such as *Brosimum* (3 species),  
 187 *Iryanthera* (2 species), and *Pouteria* (2 species), were characterized by medium-sized seeds.  
 188 Genera like *Licania* (8 species), *Eschweilera* and *Iryanthera* (3 species each), *Pouteria* (4  
 189 species), and *Aspidosperma* (2 species) exhibited the largest seeds, whereas *Protium* (5 species)  
 190 and *Virola* (1 species) had the smallest seeds (Table 2).

191

192 **Table 2** Seed size and seed size category of tree species belonging to the 11 richest genera recorded at  
 193 one hectare of open ombrophylous forest of Flona do Jamari, Itapuã do Oeste, Rondônia. Seed Size  
 194 category are given for the genus and follows Malhado et al. (2015): very small (<0.5 cm), small (0.5–  
 195 0.99 cm), medium (1.0–1.99 cm), large (≥2.0 cm).

Family	Genera	Seed size
Fabaceae	Abarema	Small
Euphorbiaceae	Alchornea	Very Small
Anacardiaceae	Anacardium	Large
Lauraceae	Aniba	Large
Malvaceae	Apeiba	Very Small
Arecaceae	Attalea	Large
Moraceae	Brosimum	Medium
Caryocaraceae	Caryocar	Large

<b>Family</b>	<b>Genera</b>	<b>Seed size</b>
Salicaceae	Casearia	VerySmall
Urticaceae	Cecropia	VerySmall
Moraceae	Clarisia	Large
Fabaceae	Couepia	Large
Chrysobalanaceae	Cybianthus	Small
Myrsinaceae	Dialium	Small
Fabaceae	Endlicheria	Large
Lauraceae	Eschweilera	Large
Lecythidaceae	Eugenia	Medium/Large
Myrtaceae	Faramea	Small
Rubiaceae	Glycydendron	Large
Euphorbiaceae	Goupia	Large
Goupiaceae	Guarea	Medium
Meliaceae	Heisteria	Medium
Olacaceae	Helicostylis	Medium
Moraceae	Hirtella	Medium
Chrysobalanaceae	Inga	Small/Medium
Fabaceae	Iryanthera	Large
Myristicaceae	Leonia	Medium
Violaceae	Licania	Large
Chrysobalanaceae	Miconia	VerySmall
Melastomataceae	Minquartia	Large
Olacaceae	Mouriri	Medium
Melastomataceae	Naucleopsis	Medium
Moraceae	Nectandra	Medium
Lauraceae	Neea	Small/Medium
Nyctaginaceae	Ocotea	Medium
Lauraceae	Oenocarpus	Medium/Large
Arecaceae	Ouratea	Medium
Ochnaceae	Oxandra	Medium
Annonaceae	Pouteria	Medium/Large
Sapotaceae	Protium	Small/Medium
Burseraceae	Pseudolmedia	Small
Moraceae	Quararibea	Medium
Malvaceae	Quiina	Medium
Quiinaceae	Salacia	Large
Celastraceae	Simarouba	Medium

Family	Genera	Seed size
Simaroubaceae	Swartzia	Medium
Fabaceae	Symphonia	Large
Clusiaceae	Tetragastris	Medium
Burseraceae	Theobroma	Large
Malvaceae	Trattinnickia	Small
Burseraceae	Unonopsis	Medium
Annonaceae	Virola	Medium
Myristicaceae	Vitex	Medium
Lamiaceae	Xylopia	Medium

196

### 197 **Distribution by vegetation type**

198 We observed that five predominant vegetation types these species occur across the biomes are  
 199 Floresta de Terra Firme (106), Floresta Ombrófila (53), Campinarana (36), Floresta Ciliar (20)  
 200 and Floresta de Várzea (17) (Sup. Mat. 1). *Protium panamense* has not been reported to occur  
 201 in the Amazonian region according to Flora do Brasil, but there are occurrences reported on  
 202 Amazonas state reported on SpeciesLink database and ter Steege et al. 2016. Besides that,  
 203 *Protium picramnioides* has not occur on the region according to SpeciesLink system platform  
 204 and ter Steege et al. (2016).

### 205 **Threat status**

206 Regarding the current threat status, we identified just two species from the *Fabaceae* family as  
 207 threatened: *Peltogyne excelsa* Ducke, classified as Vulnerable (VU), and *Vouacapoua*  
 208 *americana* Aubl., classified as Endangered (EN), among the 124 species recorded in our plot.  
 209 *Vouacapoua americana* was listed as threatened in both 2014 and 2022 by MMA. For the  
 210 projected threat status in 2050, based on ter Steege et al. (2015), we found 28 species classified  
 211 as Vulnerable and 16 as Endangered by 2050 (Sup. Mat. 1).

212           **DISCUSSION**

213   The high number of rare species inventoried in our plot highlights the importance of field data  
214   in understanding species dynamics and occurrences. Moreover, the significant number of  
215   species that have not yet been assigned a threat status by the IUCN underscores the need for  
216   accurate assessments of the conservation status of tree species, especially considering the  
217   pressures faced by forests across the region. In the entire Brazilian Amazon region, knowledge  
218   partners are spatially clustered, predominantly situated near major roads or large rivers (Stropp  
219   et al. 2020). The historical speaks in botanical sampling align with prominent research  
220   initiatives, particularly the ‘Flora projects’, which emphasize comprehensive field surveys and  
221   precise species documentation (Stropp et al. 2020). This approach is of great significance in  
222   regions marked by elevated deforestation rates and ecotonal vegetation, such as the  
223   southwestern Brazilian Amazon. However, one question remains: Are protected areas suitable  
224   for implementing these projects in such areas? It is likely that they are, particularly when  
225   involving international and national research centers, policymakers (*e.g.*, municipal and state  
226   governments), indigenous leaders, and securing adequate funding. Such collaborations have the  
227   potential to reduce bureaucratic barriers and facilitate the advancement of botanical knowledge  
228   in this geographic scale.

229   The lack of knowledge regarding the threat status for the majority of species recorded in our  
230   plot suggests the necessity for a more comprehensive assessment to accurately evaluate the real  
231   threats to these species. Our study highlights the importance of improving regional floristic  
232   knowledge to address existing shortfalls in understanding about species distribution, ecology  
233   interactions, like dispersal characteristics, and threat status. The high number of species with  
234   low density and possible new occurrences highlights the importance of gaining knowledge  
235   about ecological dynamics. The substantial presence of species exhibiting relatively low  
236   population densities, alongside the potential identification of new occurrences, serves to

237 underscore the paramount importance of acquiring comprehensive knowledge concerning the  
238 flora, encompassing its composition and distribution, within this conspicuously diverse yet  
239 understudied ecological region. Our investigation effectively highlights the indispensable role  
240 played by both floristic and phytosociological studies in elucidating the intricate structure and  
241 unique floristic aspects of Amazonian forests. Our endeavours, a considerable expanse of the  
242 region's floristic composition remains unexplored, necessitating further in-depth inquiries to  
243 gain a comprehensive understanding of species distribution. Positioned in an ecotonal area,  
244 Rondônia holds significant promise as a contributor in this vital undertaking.

### 245 *Structure*

246 The number of species recorded in our inventory of the Jamari National Forest was the highest  
247 recorded so far in studies carried out in Rondônia state. Woody plants families with the highest  
248 densities also showed greater specific richness. In the present study, Fabaceae accounted for  
249 18% of the total number of woody plants, confirming its high representativeness in tropical  
250 forests along with Burseraceae, Lecythidaceae and Arecaceae (ter Steege et al. 2016; Cardoso  
251 et al. 2017; Beech et al., 2017). The Equability Index was 0.94, suggesting uniformity in the  
252 relationship between the number of individuals per species within the plant community. Similar  
253 values were found by Oliveira et al. (2008) in one hectare of 'terra firme' forest in Central  
254 Amazon, where the diversity index was ( $H' = 5.1 \text{ nat. individual}^{-1}$ ) and the equability ( $e' = 0.92$ ).

255 In Amazonian terra firme forests, it is common to find a large number of species with few  
256 individuals and few species with a larger number of individuals (ter Steege et al. 2013). This  
257 pattern, however, was not found in our plot: *Licania macrophyla* showed highest VI (2.94) with  
258 nine individuals, followed by *Protium cf. gallosum* and *Eschueilera bracteosa*, besides that the  
259 large number of singletons (133) and doubletons (54) registered. Oliveira and Nelson (2001)  
260 made study with a 31 tree inventories from 12 sites in the Brazilian Amazon, one site in the

261 Bolivian Amazon and one in the northeast Brazilian Atlantic coastal forest, and they concluded  
262 that the western Amazon regions has greater species diversity than inventories in the east.  
263 However, inventories of terra firme forest in central Amazon present similar the tree species  
264 diversity to those found in the inventories of the western Amazon (Valencia et al 1994; Oliveira  
265 and Amaral 2004; Hubbell et al. 2008). The random pattern of species distribution and this  
266 pattern it is similar to reports by Oliveira (1997). This work emphasized that areas of terra firme  
267 in the Amazon generally show high diversity owing to the high frequency of low-density  
268 species. The importance of the Fabaceae family in our study is in agreement with the work  
269 carried out in terra firme forests across several regions of the Amazon (*e.g.*, Carim et al. 2013).  
270 In addition to Fabaceae, we identified Burseraceae, Chrysobalanaceae, Lecythidaceae and  
271 Moraceae as the five most important families in our study area. All of them are families widely  
272 represented in the structure of the Amazonian mainland forests.

### 273 *Traits attributes*

274 Our finding underscores a higher abundance of genera with medium-sized seeds compared to  
275 larger seeds. Amazonian tree genera with smaller seeds are prevalent in the southwestern and  
276 western margins of Amazonia (northern Bolivia and southeast Peru), while genera with larger  
277 seeds are more commonly found in central and northern Amazonia (Malhado et al. 2015).  
278 According to those authors, there is a significant correlation between seed size, seasonality,  
279 temperature and larger seeds are associated with warmer and less seasonal climatic conditions.  
280 These findings are consistent with the observation that forests in the southwestern of  
281 Amazonian, inhabit a climatological and ecological region, exhibiting lower temperatures,  
282 higher seasonality, and relatively more open canopies (Sombroek 2001; Coe et al. 2013; Saatchi  
283 et al. 2012).



284 *Spatial distributions*

285 According to the Flora do Brasil (BFG, 2021), seven species occurring in our plot are common  
286 across Amazon, Atlantic Forest, Cerrado and Caatinga biomes. However, this number increases  
287 to fifteen according to information from SpeciesLink system (CRIA, 2016). This pattern  
288 supports the findings of Carvalho and Almeida (2016) who suggest that most species have a  
289 restricted geographic distribution with a large number of individuals occurring in small areas  
290 and few occurring in many regions. Alternatively, a study highlighted the presence of species  
291 shared between biomes, focusing on trees and shrubs (Méio et al. 2003). The authors found that  
292 41.1% of the analysed species were exclusive to the Cerrado, indicating endemism, while  
293 58.9% were also found in the Atlantic and/or Amazonian Forest. Although the study was  
294 conducted in the Cerrado, our results demonstrated a similar pattern of species associations with  
295 species founded on our plot.

296 *Threat status*

297 However, it is important to note that our research was limited to consulting the IUCN list  
298 evaluation criteria due to the absence of a local threat list. Nineteen percent of the inventoried  
299 species represent new occurrences in the Rondônia state, indicating that the local flora may still  
300 be poorly known. Even species with a wide geographical distribution in the region are not  
301 exempt from threats.

302 We must emphasize that our study underscores the necessity for meticulous investigations,  
303 where the accurate taxonomic identification of species occupies a central position in research  
304 methodologies. The revelation of 220 distinct species within a single hectare raises a pivotal  
305 query that warrants consideration: could the estimates of diversity and richness for this  
306 particular section of the Amazon be underestimated? What is the impact of unknown species  
307 and inaccuracy of determining them on richness and diversity estimates for the region?  
308 Questions like that must be made by scientists for accessible areas in proximity to research

309 bases cannot be overstated. Instead, there is a pressing need to extend research efforts into  
310 regions that are susceptible to the impacts of climate change or deforestation (Carvalho et al.  
311 2023). We conclude that floristic studies carried out in the southwestern region of the Brazilian  
312 Amazon are rare and our study, though small scale, provides a significant contribution to  
313 biodiversity knowledge of the region. The need for such studies is urgent, given the continuing  
314 loss of native vegetation due to development pressures, especially on native vegetation. We  
315 emphasize the importance for scientists to conduct studies addressing the new species (Linnean  
316 shortfall), spatial under sampling (Wallacean shortfall) and ecology interactions (Raunkiaer  
317 shortfall) in relations to plant structure investigations.

#### 318 *author contributions statement*

319 **Bruno Umbelino** - Conceptualization, Formal analysis, Methodology, Visualization, Writing  
320 – original draft, Writing – review & editing; **Juliana Stropp** – Formal analysis, Methodology,  
321 Visualization, Writing – original draft, Writing – review & editing; **Ingrid Mendes-Silva** -  
322 Visualization, Writing – original draft, Writing – review & editing; **Ricardo Aleixo Correia** –  
323 Writing – review & editing; **Antônio L. P. Silveira** – Conceptualization, Supervision, Writing  
324 – review & editing; **Ana C. M. Malhado** - Conceptualization, Supervision, Writing – review  
325 & editing.

#### 326 *conflict of interest statement*

327 The authors declare that they have no known financial conflicts of interest or personal  
328 relationships that could appear to have influenced the work reported in this paper.

329 **Acknowledgements** – The ideas presented in this study it was discussed in depth with  
330 Marcus Vinicius Vieira. We thank the botanists Douglas Daly, Wendeson Castro, Herison  
331 Medeiros, Daniel Silva, the parataxonomists Edilson C. Oliveira, Adriano S. Lima, and the

332 team of Herbarium João Geraldo Kuhlmann. The first author acknowledges the support  
333 provided by the Programa Institucional de Bolsas de Iniciação Científica (PIBIC, Brazilian  
334 portuguese acronym), which granted a scholarship during the initial stages of this project.  
335 Additionally, the ongoing support through a PhD scholarship from Coordenação de  
336 Aperfeiçoamento de Pessoas de Nível Superior – Brasil (CAPES) – Financial Code 001 is also  
337 greatly appreciated.

## 338 REFERENCE

- 339 Beech, Emily, M Rivers, S Oldfield, and P P Smith. 2017. “GlobalTreeSearch: The First  
340 Complete Global Database of Tree Species and Country Distributions.” *Journal of*  
341 *Sustainable Forestry* 36 (5): 454–89.
- 342 Cardoso, Domingos, Tiina Särkinen, Sara Alexander, André M Amorim, Volker Bittrich, and  
343 Marcela Celis. 2017. “Amazon Plant Diversity Revealed by a Taxonomically Verified  
344 Species List” 114 (40). <https://doi.org/10.1073/pnas.1706756114>.
- 345 Carvalho, Raquel L., Angelica F. Resende, Jos Barlow, Filipe M. França, Mario R. Moura,  
346 Rafaella Maciel, Fernanda Alves-Martins, et al. 2023. “Report Pervasive Gaps in  
347 Amazonian Ecological Research Pervasive Gaps in Amazonian Ecological Research.”  
348 *Current Biology*, 1–9. <https://doi.org/10.1016/j.cub.2023.06.077>.
- 349 Coe, Michael T, Toby R Marthews, Marcos Heil Costa, David R Galbraith, Nora L  
350 Greenglass, Hewlley M A Imbuzeiro, Naomi M Levine, et al. 2013. “Deforestation and  
351 Climate Feedbacks Threaten the Ecological Integrity of South – Southeastern  
352 Amazonia.” *The Royal Society*.
- 353 Correia, Ricardo A., Ana C. M. Malhado, Lays Lins, Norah Costa Gamarra, Waltyane A. G.  
354 Bonfim, Anyelet Valencia-Aguilar, Chiara Bragagnolo, Paul Jepson, and Richard J.

355 Ladle. 2016. "The Scientific Value of Amazonian Protected Areas." *Biodiversity and*  
356 *Conservation*. <https://doi.org/10.1007/s10531-016-1122-x>.

357 Díaz, Sandra, Andy Purvis, Johannes H. C. Cornelissen, Georgina M. Mace, Michael J.  
358 Donoghue, Robert M. Ewers, Pedro Jordano, and William D. Pearse. 2013. "Functional  
359 Traits, the Phylogeny of Function, and Ecosystem Service Vulnerability." *Ecology and*  
360 *Evolution* 3: 2958–75. <https://doi.org/10.1002/ece3.601>.

361 Dornelas, Maria, Anne E Magurran, Stephen T Buckland, Anne Chao, Robin L Chazdon,  
362 Robert K Colwell, Tom Curtis, et al. 2012. "Quantifying Temporal Change in  
363 Biodiversity : Challenges and Opportunities." *ProceedingsThe Royal Society*.

364 Hortal, Joaquín, Francesco De Bello, José Alexandre F. Diniz-Filho, Thomas M. Lewinsohn,  
365 Jorge M. Lobo, and Richard J. Ladle. 2015. "Seven Shortfalls That Beset Large-Scale  
366 Knowledge of Biodiversity." *Annual Review of Ecology, Evolution, and Systematics* 46  
367 (October): 523–49. <https://doi.org/10.1146/annurev-ecolsys-112414-054400>.

368 Hubbell, S. P., F. He, R. Condit, L. Borda-de-Agua, J. Kellner, and H. ter Steege. 2008. "How  
369 Many Tree Species Are There in the Amazon and How Many of Them Will Go  
370 Extinct?" *Proceedings of the National Academy of Sciences* 105 (Supplement 1): 11498–  
371 504. <https://doi.org/10.1073/pnas.0801915105>.

372 Ladle, Richard, and Joaquín Hortal. 2013. "Mapping Species Distributions: Living with  
373 Uncertainty." *Frontiers of Biogeography* 5 (1). <https://doi.org/10.21425/f55112942>.

374 Malhado, Ana C. M., Rafael S. D. de Azevedo, Peter A. Todd, Ana M. C. Santos, Nídia N.  
375 Fabré, Vandick S. Batista, Leonardo J. G. Aguiar, and Richard J. Ladle. 2014.  
376 "Geographic and Temporal Trends in Amazonian Knowledge Production." *Biotropica*  
377 46 (1): 6–13.

378 Malhado, Ana C M, Jose A Oliveira-neto, Juliana Stropp, Giovanni Strona, Livia C P Dias,  
379 Luciana B Pinto, and Richard J Ladle. 2015. "Climatological Correlates of Seed Size in  
380 Amazonian Forest Trees" 26: 956–63. <https://doi.org/10.1111/jvs.12301>.

381 Méio, Beatriz B, Cristiane V Freitas, Leuseroberta Jatobá, Mario E F Silva, José F Ribeiro,  
382 and Raimundo P. B. Henriques. 2003. "Influência Da Flora Das Florestas Amazônica e  
383 Atlântica Na Vegetação Do Cerrado Sensu Stricto." *Brazilian Journal of Botany* 26:  
384 437–44.

385 Oliveira, Alexandre Adalardo de, and Bruce Walker Nelson. 2001. "Floristic Relationships of  
386 Terra Firme Forests in the Brazilian Amazon." *Forest Ecology and Management* 146:  
387 169–79.

388 Oliveira, Arlem Nascimento De, Michele Braule, Pinto Ramos, Luciana Bovino Couto, and  
389 Rosana Martiniano Sahdo. 2008. "Composição e Diversidade Florístico-Estrutural de  
390 Um Hectare de Floresta Densa de Terra Firme Na Amazônia." *Acta Amazonica* 38 (4):  
391 627–42.

392 Saatchi, Sassan, Salvi Ase, Yadvinder Malhi, Luiz E O C Aragão, and Liana O Anderson.  
393 2012. "Persistent Effects of a Severe Drought on Amazonian Forest Canopy." *PNAS* 110  
394 (565–570). <https://doi.org/10.1073/pnas.1204651110>.

395 Santos, Jhonatan Guedes dos, Ana C M Malhado, Richard J. Ladle, Ricardo A. Correia, and  
396 Marcos H. Costa. 2015. "Geographic Trends and Information Deficits in Amazonian  
397 Conservation Research." *Biodiversity and Conservation* 24 (11): 2853–63.  
398 <https://doi.org/10.1007/s10531-015-0981-x>.

399 Schulman, Leif, Tuuli Toivonen, and Kalle Ruokolainen. 2007. "Analysing Botanical  
400 Collecting Effort in Amazonia and Correcting for It in Species Range Estimation."

401 *Journal of Biogeography* 34 (8): 1388–99. <https://doi.org/10.1111/j.1365->  
402 [2699.2007.01716.x](https://doi.org/10.1111/j.1365-2699.2007.01716.x).

403 Sombroek, Wim. 2001. “Spatial and Temporal Patterns of Amazon Rainfall.” *A Journal of the*  
404 *Human Environment* 30 (7): 388–96.

405 Steege, Hans Ter, Nigel C.A. Pitman, Daniel Sabatier, Christopher Baraloto, Rafael P.  
406 Salomão, Juan Ernesto Guevara, Oliver L. Phillips, et al. 2013. “Hyperdominance in the  
407 Amazonian Tree Flora.” *Science* 342 (6156). <https://doi.org/10.1126/science.1243092>.

408 Steege, Hans ter, Paulo I. Prado, Renato A.F.de Lima, Edwin Pos, Luiz de Souza Coelho,  
409 Diogenes de Andrade Lima Filho, Rafael P. Salomão, et al. 2020. “Biased-Corrected  
410 Richness Estimates for the Amazonian Tree Flora.” *Scientific Reports* 10 (1): 1–13.  
411 <https://doi.org/10.1038/s41598-020-66686-3>.

412 Steege, Hans Ter, Rens W. Vaessen, Dairon Cárdenas-López, Daniel Sabatier, Alexandre  
413 Antonelli, Sylvia Mota De Oliveira, Nigel C.A. Pitman, Peter Møller Jørgensen, and  
414 Rafael P. Salomão. 2016. “The Discovery of the Amazonian Tree Flora with an Updated  
415 Checklist of All Known Tree Taxa.” *Scientific Reports* 6: 1–15.  
416 <https://doi.org/10.1038/srep29549>.

417 Stropp, J., B. Umbelino, R. A. Correia, J. V. Campos-Silva, R. J. Ladle, and A. C.M.  
418 Malhado. 2020. “The Ghosts of Forests Past and Future: Deforestation and Botanical  
419 Sampling in the Brazilian Amazon.” *Ecography* 43 (7): 979–89.  
420 <https://doi.org/10.1111/ecog.05026>.

421 Tessarolo, Geiziane, Richard Ladle, Thiago Rangel, and Joaquin Hortal. 2017. “Temporal  
422 Degradation of Data Limits Biodiversity Research.” *Ecology and Evolution*.  
423 <https://doi.org/10.1002/ece3.3259>.

424 Valencia, Renato, Henrik Balslev, and Guilherme Paz Y Miño C. 1994. "High Tree Alpha-  
425 Diversity in Amazonian Ecuador." *Biodiversity and Conservation* 28: 21–28.

426