

Subterranean environments contribute to three-quarters of classified ecosystem services

Stefano Mammola^{1,2,3,‡,*}, David Brankovits^{1,‡}, Tiziana Di Lorenzo^{3,4,5,6}, Isabel R. Amorim⁷, Raluca I. Bancila^{5,8}, Adrià Bellvert¹, Enrico Bernard⁹, Anna S. Blomberg², Paulo A.V. Borges^{7,10,11}, Martina Cappelletti^{12,13}, Rodrigo Ferreira¹⁴, Rosalina Gabriel⁷, Diana M. P. Galassi¹⁵, Laura Garzoli¹, Vasilis Gerovasileiou^{16,17}, Grant C Hose¹⁸, Kathryn L Korbel¹⁸, Youness Mabrouki¹⁹, Simone Martino²⁰, Ana Z Miller^{21, 22}, Nataša Mori²³, Veronica Nanni¹, Giuseppe Nicolosi²⁴, Mattia Saccò^{25,26,27}, Troy S. Sakihara²⁸, Marconi Souza Silva¹⁴, Anne E. Tamalavage²⁹, Maja Zagmajster³⁰, Efraín Chávez^{31,32}, Christian Griebler^{33,^}, Pedro Cardoso^{2,6,^}, Alejandro Martínez^{1,^}

‡ Shared first

^ Shared last

* Corresponding: stefano.mammola@cnr.it

1. Molecular Ecology Group (MEG), Water Research Institute, National Research Council of Italy (CNR-IRSA), Verbania Pallanza, Italy
2. Finnish Museum of Natural History, University of Helsinki, Helsinki, Finland
3. NBFC, National Biodiversity Future Center, Palermo, Italy
4. Research Institute on Terrestrial Ecosystems, National Research Council (CNR-IRET), Via Madonna del Piano 10, 50019 Sesto Fiorentino, Florence, Italy
5. “Emil Racoviță” Institute of Speleology, Cluj-Napoca Department, Str. Clinicilor, Nr. 5-7, 400006 Cluj-Napoca, Romania
6. Centre for Ecology, Evolution and Environmental Changes & CHANGE – Global Change and Sustainability Institute, and Departamento de Biologia Animal, Faculdade de Ciências, Universidade de Lisboa, Campo Grande 1749-016, Lisbon, Portugal
7. University of Azores, CE3C—Centre for Ecology, Evolution and Environmental Changes, Azorean Biodiversity Group, CHANGE —Global Change and Sustainability Institute, Rua Capitão João d’Ávila, Pico da Urze, 9700-042 Angra do Heroísmo, Azores, Portugal
8. “Emil Racoviță” Institute of Speleology, Calea 13 Septembrie, Nr. 13, 050711 Bucharest, Romania
9. Laboratório de Ciência Aplicada à Conservação da Biodiversidade, Departamento de Zoologia, Instituto de Biociências, Universidade Federal de Pernambuco, Recife PE, Brazil
10. IUCN SSC Atlantic Islands Invertebrate Specialist Group, 9700-042 Angra do Heroísmo, Azores, Portugal
11. IUCN SSC Species Monitoring Specialist Group, 9700-042 Angra do Heroísmo, Azores, Portugal
12. Department of Pharmacy and Biotechnology, University of Bologna, Bologna 40126, Italy
13. La Venta Geographic Explorations Association, Treviso 31100, Italy
14. Center for Studies in Subterranean Biology, Department of Ecology and Conservation, Institute of Natural Sciences, Federal University of Lavras, Lavras, Minas Gerais, Brazil
15. Department of Life, Health & Environmental Sciences, University of L’Aquila, L’Aquila, Italy
16. Department of Environment, Faculty of Environment, Ionian University, Zakynthos, Greece
17. Institute of Marine Biology, Biotechnology and Aquaculture (IMBBC), Hellenic Centre for Marine Research (HCMR), Heraklion, Greece
18. School of Natural Sciences, Macquarie University, NSW 2109, Australia
19. Biotechnology, Conservation and Development of Bioresources Laboratory, Faculty of Sciences Dhar El Mehraz, University Sidi Mohamed Ben Abdellah, Fez, Morocco.
20. The James Hutton Institute, Craigiebuckler, Aberdeen, AB15 8QH, Scotland

21. BIOGEOCOM - Instituto de Recursos Naturales y Agrobiología de Sevilla (IRNAS-CSIC), Avenida Reina Mercedes 10, 41012 Seville, Spain
22. HERCULES Lab, University of Évora, Largo Marquês de Marialva 8, 7000-809 Évora, Portugal.
23. Department of Organisms and Ecosystem Research, National institute of Biology, Večna pot 121, 1000, Ljubljana, Slovenia
24. Department of Biological, Geological and Environmental Sciences, Section Animal Biology, University of Catania, 95124 Catania, Italy
25. Subterranean Research and Groundwater Ecology (SuRGE) Group, Trace and Environmental DNA (TrEnD) Lab, School of Molecular and Life Sciences, Curtin University, Perth, WA 6102, Australia
26. Department of Chemistry, Life Sciences and Environmental Sustainability, University of Parma, Parco Area delle Scienze 11/A, 43124 Parma, Italy;
27. Laboratoire de Biologie des Organismes et des Écosystèmes Aquatiques-BOREA. Muséum national d'Histoire naturelle, SU, CNRS, IRD, UA, F-75005 Paris, France
28. Department of Land and Natural Resources, Division of Aquatic Resources, Hilo, Hawai'i, USA
29. Smithsonian Environmental Research Center, Edgewater, Maryland USA
30. University of Ljubljana, Biotechnical Faculty, Department of Biology, Subterranean Biology Lab (SubBioLab), Jamnikarjeva 101, 1000, Ljubljana, Slovenia
31. Unidad Multidisciplinaria de Docencia e Investigación, Facultad de Ciencias, Universidad Nacional Autónoma de México, Sisal, Yucatán, México
32. Secretaria de Ciencia, Humanidades, Tecnología e Innovación, Mexico City, Mexico.
33. Department for Functional and Evolutionary Ecology, University of Vienna, Vienna, Austria

Abstract

Beneath the Earth's surface lies a network of interconnected caves, voids, and systems of fissures forming in rocks of sedimentary, igneous, or metamorphic origin. Though largely inaccessible to humans, this hidden realm supports and regulates services critical to ecological health and human well-being. Subterranean ecosystems are integral to major biogeochemical cycles, sustain diverse surface habitats, and serve as the primary source of irrigation and drinking water. They also offer non-material benefits, including scientific discovery, education, and cultural practices. Yet, these contributions often go unrecognized, partly due to the lack of a unified synthesis of ecosystem services across terrestrial, freshwater, and marine subterranean compartments. This gap limits effective communication of their value to scientists, practitioners, and the public. Through a systematic expert-based review, we show that subterranean ecosystems contribute to up to 75% of classified ecosystem services. Notably, many of these contributions are described only qualitatively, lacking numerical or economic quantification. Next, we provide examples of the main services to offer a global overview of their multifaceted value and vulnerability to environmental change. We believe this synthesis provides researchers and practitioners with concrete examples and targeted metaphors to more effectively communicate the importance of subterranean ecosystems to diverse audiences.

Keywords: Groundwater; Hypogean; Nature value; Drinking water; Food production; Biotechnology; Geothermal energy; Sustainability; Ecotourism; Cultural heritage

Introduction

Whether engaging in high-stakes discussions with policymakers or navigating casual conversations at social gatherings, scientists studying subterranean biodiversity may find themselves in the uncomfortable position of defending the very essence of their work. Questions like, “*Why waste your time in a muddy cave to count tiny beetles?*”, “*Are we really worried about some blind shrimp no one's ever seen?*” or “*What’s next —national parks for glow-in-the-dark worms?*” are all too common. They reflect a deep misunderstanding of the hidden world beneath our feet, the fragile ecosystems it sustains, and the profound influence it has on the surface environments where humans live.

Studying “unremarkable” species thriving beneath the Earth surface might seem like an indulgent pursuit, far removed from the pressing concerns of modern life. After all, how could the presence of a whitish shrimp in a remote cave pond possibly contribute to global challenges such as economic growth, public health, or technological development? Far from trivial, these discussions reflect a broader struggle to spotlight the invisible services provided by nature. The challenge, then, is not merely defending one’s research but broadening collective understanding of biodiversity’s essential functions—its intrinsic value and its critical role in maintaining a healthy, habitable planet. The public cannot grasp what is at risk if scientists fail to communicate these values.

When the concept of ecosystem services gained momentum after 1997, it offered biodiversity scientists a powerful framework to articulate the societal relevance of their work. Ecosystem services encompass all the functions and products of ecosystems that benefit humans and contribute to societal welfare. Initially conceived as a metaphor, the concept quickly evolved into a robust research agenda focused on cataloging, quantifying, and mapping humanity’s reliance on nature^{1–3}. For example, ecosystem services are frequently categorized into: provisioning services (e.g., food, water), regulation and maintenance services (e.g., climate regulation, pollination, air and water quality), and cultural services (e.g., recreational, traditional practices and spiritual well-being). Notwithstanding the

inherent risk of putting a price tag on nature¹⁰, many of these services are also often measured economically. This reflects the need to highlight the value of services that are, in part, subjective and difficult to perceive outside of academic contexts^{11,12}.

While the quantification of ecosystem services has occupied the research community for decades, knowledge remains incomplete for subterranean ecosystems. Despite their hidden nature, subterranean ecosystems (Box 1) provide and regulate services that are as critical to human well-being and ecological health as those in surface ecosystems (Figure 1). The benefits derived from subterranean ecosystems are remarkably diverse, with direct and indirect links to essential functions such as freshwater provisioning, food production, and the regulation of diverse biogeochemical and physical processes^{13–16}. Subterranean ecosystems also contribute to essential “non-material” values, including scientific research and inspiration^{17–19}, ecotourism^{20,21}, aesthetic appreciation^{22,23}, and cultural practices^{24,25}.

The questions driving this review are straightforward yet important: What services do subterranean ecosystems provide, and how many of these have been quantified to some extent? Answering these questions is urgent because, paraphrasing the common adage, “you can’t manage what you can’t see and measure”. Currently, information on the benefits provided by subterranean ecosystems is scattered across numerous sources, many of which remain inaccessible to the public. While a handful of reviews have compiled subsets of these services for specific subterranean ecosystems^{16,26–30} or species³¹, a comprehensive scheme that unifies all services across terrestrial, freshwater, and marine subterranean compartments is still lacking. Moreover, integrating quantitative rigor into this mapping exercise could enhance the perceived importance of these services and help establish connections to the eco-evolutionary processes that sustain them. Such an understanding could shift the narrative—from viewing subterranean ecosystems merely as sources of water, geothermal energy, and minerals to recognizing their broader ecological value. This, in turn, would reinforce the importance of even partial data in designing conservation strategies that prioritize ecosystem functions over isolated species or habitats³².



156 **Figure 1.** A visual summary of the main services associated with subterranean ecosystems. Original
 157 illustration by Jagoba Malumbres-Olarte.

Box 1. What do we mean by “subterranean ecosystem”? Subterranean ecosystems are globally distributed and vary widely in extent and type of matrix. Following the function-based classification of Earth’s ecosystems^{33,34}, we considered ecosystems belonging to these biomes in terrestrial, freshwater, and marine domains: ‘Subterranean’ (S) [including the ‘Subterranean lithic’ (S1) and ‘Anthropogenic subterranean voids’ (S2) biomes], ‘Subterranean-freshwater’ (SF) [including the ‘Subterranean freshwater’ (SF1) and ‘Anthropogenic subterranean freshwater’ (SF2) biomes], and ‘Subterranean tidal’ (SM1). These include various types of caves (e.g., aerobic caves, lava tubes, volcanic pits, anchialine caves, sea caves) and other voids (e.g., fissure systems, deep scree strata), groundwater ecosystems and their ecotones (e.g., aquifers, underground streams, ponds, lakes, subterranean estuaries, anchialine pools, sinkholes, cenotes, blueholes, springs, hyporheic systems), as well as anthropogenic subterranean voids (e.g., mines, underground bunkers and tunnels, water pipes, subterranean canals, wells). Conversely, excluded ‘Endolithic systems’ (S1.2)³³ or, in other words, the deep biosphere—microbial communities occurring kilometers beneath the Earth’s surface—due to the limited ecological data available and the uncertainty regarding their connectivity to shallower ecosystems and ecosystem service provision^{35,36}.

Subterranean ecosystems services in numbers

To map subterranean ecosystem services, we used the Common International Classification of Ecosystem Services (CICES Version 5.1). CICES is a classification scheme designed to measure, account for, and assess ecosystem services⁹. The services are categorized into three main “Sections” (Provisioning, Regulation & Maintenance, and Cultural services) and two broad types within each section (biotic and abiotic), with further breakdowns into levels of Division, Group, and Class. Conveniently, CICES is interoperable with other ecosystem service classification systems by providing equivalency across various schemes. CICES lists 90 primary services—63 biotic and 27 abiotic. Using this backbone classification, we assessed whether subterranean ecosystems contribute to the various ecosystem services listed in CICES based on our expert knowledge and the literature. To strengthen our evaluation, we conducted a literature review for each service to assess quantitative estimates of the services provided by subterranean ecosystems.

According to our mapping exercise (see [Data availability statement](#)), subterranean ecosystems contribute to up to 75% (68 out of 90) of the ecosystem services classified by CICES. This contribution is higher than the estimations for ecosystem services provided by grasslands (36%)³⁷, urban water bodies (43%)³⁸, mangrove ecosystems (33% of the biotic services)³⁹, or vineyards (64%)⁴⁰. When considering specific systems, terrestrial, freshwater, and saltwater subterranean compartments match 48%, 57%, and 54% of the services classified by CICES, respectively.

Of all the 68 matching services, between one third and a half have been quantified (i.e., measured numerically) ([Figure 2](#)), primarily by local case studies. Most of the identified services benefit society at large, although specific services appear to be most important for specific economic sectors ([Figure 3](#)). Groundwaters, particularly freshwater systems, dominate in the percentage of measured ecosystem services. This is likely both due to their accessibility and measurability compared to terrestrial and marine systems and the crucial importance of groundwater for drinking and irrigation. Indeed, human settlements are often located where there is access to

206 aquifers, through springs, caves, wells and boreholes. These features allow for direct
207 sampling and regular monitoring. In comparison, terrestrial and marine subterranean
208 ecosystems are less accessible, often requiring specialized and costly technologies
209 for exploration^{41–43}.




Figure 2. The number of Provisioning, Regulation & Maintenance, and Cultural services provided by terrestrial, freshwater, and saltwater subterranean ecosystems (colored bars) compared to the total services mapped by the Common International Classification of Ecosystem Services (white bars). Darker shades indicate the fraction of subterranean services that have been quantitatively assessed in at least one study.

Provisioning services

Provisioning ecosystem services are the tangible goods and resources that ecosystems provide to humans⁹. These services are the direct products we obtain from nature, such as freshwater, food, raw materials, medicinal resources, and energy. Subterranean ecosystems contribute to as many as 63% of the provisioning ecosystem services classified by CICES ([Figure 3](#)).

Water supply

Groundwater, the largest unfrozen continental reserve of freshwater globally^{44,45}, is a prime source of water for drinking, irrigation, and industrial use¹³ ([Figure 4A, 4B](#)). It is estimated that groundwater constitutes approximately 95% of the planet's accessible liquid freshwater resources, including drinking water. Over a quarter of the global population relies on this resource, either partially or entirely⁴⁶. Current human groundwater use is estimated to exceed the capacity of aquifers by about 3.5 times and groundwater decline is accelerating at the global scale⁴⁷. About 43% of irrigation water and 49% for domestic use is sourced from groundwater⁴⁸, and this figure is likely to become even more pronounced due to continuous population growth and increasing frequency of droughts and extreme events connected with climate change intensity^{48,49}.

Still, there are large uncertainties in global estimations of the total volume of groundwater^{44,45}, where it is distributed⁴⁵, its depth⁵⁰, recharge rates⁵¹, and patterns of human extractions⁵². Besides quantity, water quality is very relevant, which primarily depends on geochemical processes and anthropogenic impacts but also, at least partially, on the presence of subterranean organisms (see section "Regulation & Maintenance Services").




Figure 3. Importance of ecosystem services provided by terrestrial, freshwater, and marine subterranean ecosystems to socio-economic sectors (primary: resource extraction; secondary: manufacturing; tertiary: services; quaternary: knowledge-based activities). “All society” represents services with transversal benefits, not limited to a single sector.

Energy production

Subterranean ecosystems are increasingly used for heating, cooling, and direct energy production. Geothermal heat pump systems, which use heat from shallow underground sources, are the fastest-growing segment of geothermal technology and one of the fastest-growing renewable energy options in the world. Other direct uses, such as heating buildings, bathing, swimming, industrial processes, farming (especially greenhouses), and fish farming, are generally based on deep hydrothermal resources⁵³. Deep geothermal energy plants produce hot water, directly used for heating purposes (e.g., via district heating networks) or turn the heat into electrical power. The global geothermal energy production of 95 TWh yr⁻¹ represents about 10% of the sustainable electricity generated annually. Among renewables, geothermal power has the highest potential in the future accounting for about 67%. Solar power, wind power, power from biomass, and hydropower account for 21%, 8%, 3.5%, and 0.1%, respectively^{53–55}.

When considering subterranean ecosystems, it is the use of shallow geothermal energy that requires the most attention. In geology, the boundary between 'shallow' and 'deep' is typically set at a depth of 400–500 meters, which corresponds, with only a few exceptions, to the deepest known occurrence of subterranean fauna⁵⁶. Most geothermal heat pump systems operate with relatively shallow closed-loop borehole heat exchangers, often complemented by open, groundwater-based systems. The shallow subsurface is warmer in winter and cooler in summer compared to the outside air. By using geothermal heat pump systems, this temperature difference can provide heating in winter and cooling in summer. However, it is important to note that extracting heat or cold from the subsurface can alter thermal conditions in ways that may be harmful from an ecological perspective. It can also induce temperature fluctuations resembling surface seasonality, though with much smaller temperature differences. Among these effects, warming is the main factor that puts pressure on subterranean communities⁵⁷. Warming also accelerates the metabolism of both microbes and fauna, leading to faster consumption of dissolved oxygen and potentially resulting in hypoxic or anoxic conditions. These oxygen-depleted conditions can cause the disappearance of fauna

and are followed by a decline in water quality⁵⁸. Thus, energy production through geothermal use can be in conflict with the health of subterranean ecosystems— alongside other global driver of subsurface warming such as climate change⁵⁹ and urbanisation^{60,61}.

Food production

Groundwater is critical for global food security, supplying over 40% of the water used for irrigation and supporting approximately 13% of total food production⁶² (Figure 4B). Groundwater enables both large- and small-scale farmers to enhance agricultural output, particularly in regions where rainfall is insufficient to meet crop water demands⁶³. Even though the negative effect of irrigation can be mitigated^{64,65}, groundwater resources are increasingly being overexploited, especially in major agricultural regions such as California's Central Valley, the High Plains Aquifer in the U.S. Midwest, the Middle East, the Indus and Ganges Basins, and the North China Plain⁶⁶. Currently, India is the world's largest consumer of groundwater, supplying approximately 60% of its irrigation needs⁶⁷. Among internationally traded crops, rice is the most groundwater-intensive, accounting for 29% of global usage, followed by wheat (12%), cotton (11%), maize (4%), and soybeans (3%). Citrus and sugar crops also account for approximately 5% of groundwater use for irrigation each⁶⁸.

Beyond these agricultural trends, subterranean environments have been central to food production and foraging practices for millennia. Shepherds have historically used caves and caverns as shelters to protect livestock from harsh weather conditions^{69,70}. Additionally, caves were integral to traditional food preservation and production, particularly in cheese and wine-making, as well as mushroom cultivation, where their stable temperatures and humidity make them natural analogs to cellars^{71,72}. A case-in-point is *Penicillium roqueforti*, a fungus discovered in the limestone caves above Roquefort, France, where the mold accidentally transformed cheese into a flavorful delicacy, now renowned as Roquefort cheese.

Subterranean ecosystems and their ecotones also contribute to the service of food production by enabling aquaculture or as habitat for commercially or culturally

important species. For example, anchialine pools have been used to keep fish for fresh consumption or even to cultivate fish bait used in traditional mackerel fisheries⁷³, as in the case of the red shrimp *Halocaridina rubra* (‘ōpae‘ula) in Hawai‘i⁷⁴. The nests and eggs of cave swiftlets (*Collocalia linchi*) and Cory’s shearwaters (*Calonectris diomedea*) are harvested for their nutritional ⁷⁵ or traditional medical value⁷⁶. Oilbirds (*Steatornis caripensis*) are exploited in South America for their flesh and fat (oil), used for cooking and lighting⁷⁷. Bats are hunted as a meat source in Asia and Africa⁷⁸. Depending on species and locations, bats are either considered a delicacy or an affordable source of protein during times of food scarcity^{79,80}. However, such practices may threaten endangered species and their habitats⁸¹.




Figure 4. Global mapping of major subterranean ecosystem services based on proxy variables.

A) Groundwater depth and human population density, illustrating potential hotspots where there will be pressure in terms of groundwater extraction. Pink areas indicate regions where groundwater is deeply underground and difficult to access, with low population densities. Dark orange areas represent regions where groundwater is also difficult to access but have high population densities. B) Groundwater depth and irrigation intensity, illustrating potential hotspots where there will be pressure in terms of groundwater extraction. Dark blue areas indicate regions that are highly irrigated and face greater difficulty accessing groundwater. In both A and B, groundwater availability is measured as the depth from the land surface to the point where groundwater begins (source: ref. ⁸²). A shallow depth means groundwater is close to the surface, whereas a greater depth indicates it is further underground. Dark orange and dark blue areas highlight regions where accessing groundwater is particularly challenging, especially in the absence of shallow water resources. C) Global mining pressure, illustrating potential hotspots where mining activities may reduce subterranean habitat availability. Mining intensity is calculated based on the percentage of each country's area occupied by mines (source: ref. ⁸³) and the total extracted tonnage of target minerals (source: World Mineral Statistics contributed by permission of the British Geological Survey). D) Annual number of show cave visitors per country and associated income, estimated based on cave entrance fees (in dollars) (source: ref. ²⁰). Note that political borders in the maps are based on open sources, and do not reflect the views of the coauthors or the institutions in which the work.

Raw materials

Rock, mineral, and materials extracted from subterranean ecosystems account for a major part of the global economy (Figure 4C). The effects of mining, including rock or mineral extraction itself and all the infrastructure involved, potentially influences 50 million km² of the planet's surface⁸⁴. In 2025, the global production of minerals is expected to reach 15 billion tons⁸⁵, with a value exceeding 7 trillion USD in 2024 and constituting an important part of the national GDP in many countries⁸⁶.

Many of the mining areas coincide with protected, key biodiversity and wilderness areas. Hence, mining activities impact subterranean ecosystems, either directly (e.g., loss of habitat) or indirectly (aquifer contamination)^{87,88}. For example, iron ore production in Brazil accounts for approximately 1.6% of the country's GDP, generating around 31 billion US\$ in 2022. With thousands of caves associated with iron ore landscapes, mining activities severely threaten these unique subterranean ecosystems, which are recognized for their significant diversity of cave-restricted species^{89,90}.

A special case of mining involves bat and bird guano, which can be locally abundant—millions of bats gathering in cave colonies can produce guano piles as high as 10 meters⁹¹. Guano is widely used as a fertilizer due to its high nitrogen and phosphorus content⁹², or as a source of chitin and chitosan for cosmetics, pharmaceuticals, and textiles^{93,94}. Bat guano fertilizer typically costs US\$ 2.50–24.00 per 1 kg⁹².

Biomolecular resources and emerging technologies

Subterranean ecosystems are a promising source of molecules and compounds with biotechnological applications, though bioprospective activities are still in their early stages. Subterranean microbial biofilms often influence mineral precipitation and dissolution⁹⁵, particularly through polymeric substances that are produced and secreted by microbes (mediating microbial adhesion on surfaces) and may serve as nucleation sites for mineral precipitation, promoting the growth of cave formations (speleothems)^{96–98}. Secondary metabolites produced by microbes within those

biofilms may have biotechnological and pharmaceutical applications as well, including use as enzymes, biosurfactants, or as antitumoral, immunosuppressive, and immunostimulatory agents^{98–105}. Some subterranean microorganisms with extracellular hydrolytic activity and antimicrobial compound production may be relevant against multidrug-resistant pathogens^{99,106,107}. For example, extracts of bacterial isolates from lava tubes of Lanzarote (Canary Islands) showed antimicrobial activity against the pathogenic strains *Staphylococcus aureus*, *Escherichia coli*, and *Pseudomonas aeruginosa*, and exhibited antiproliferative activity against human breast cancer cells¹⁰⁴.

Beyond microbes, larger subterranean organisms have also been explored for their biomolecular potential. For example, many sessile invertebrates in marine caves (e.g., sponges, anthozoans, bryozoans, and tunicates) contain or secrete compounds with significant application potential^{108–112}. This biotechnological potential may also arise from more subtle interactions between microscopic and macroscopic organisms. For instance, animal excrement in caves, which often harbors pathogenic viruses, may stimulate microorganisms to produce antiviral substances¹¹³.

Finally, the unique biological adaptations of several subterranean species hold promise for biomimicry, particularly in developing sensors, biomaterials, adhesives, and biologically inspired robotic movement¹⁹. In recent years, medical applications inspired by subterranean adaptations have also gained attention, ranging from potential treatments for diabetes¹¹⁴ and autism¹¹⁵ to innovations in blindness research¹¹⁶. Despite these possibilities, this potential remains largely untapped, with most studies still far from yielding concrete applications.

Regulation & Maintenance services

Regulation & Maintenance services provide the abiotic and biotic processes and environmental conditions that benefit living organisms, including humans⁹. Hence, these services offer stability, safety, and resilience to both ecosystems and human societies, and subterranean ecosystems contribute to as many as 82% of these (Figure 3).

Regulation of physico-chemical conditions

Subterranean ecosystems are central to global water and (bio)geochemical cycles, including carbon, nitrogen, and other key elements (e.g., phosphorus, sulphur, and iron)^{13,36,117}. Given their role in maintaining freshwater, seawater, and atmospheric balance, subterranean ecosystems are increasingly recognized as vital to global sustainability efforts. In particular, subterranean environments may be integral to Earth System governance frameworks such as the planetary boundaries, where groundwater has already been proposed as a key component¹³. The planetary boundaries define a set of critical biogeophysical processes that collectively regulate the stability and resilience of the Earth System^{118,119}.

Hotspots for these biogeochemical processes are typically located along environmental gradients, redox interfaces, ecotones, and other transition zones in both terrestrial (e.g., subsurface-surface atmosphere, sediment/rock-atmosphere interfaces) and aquatic settings (e.g., land-sea, sediment-water, and water-atmosphere interfaces)¹²⁰. These environmental gradients span micro (<mm) to regional scales (>km), and their role in regulating chemical fluxes and ecosystem functioning is often disproportionately large relative to their size¹²¹. Some of these processes may also be mediated within the so-called “deep biosphere” (Box 1)—following the recognition that bacteria and archaea can occur kilometers deep in the Earth's crust³⁵. Yet, major gaps remain in our understanding of their extent, function, and role in global biogeochemical cycling³⁶.

Biogeochemical processes associated with subterranean ecosystems primarily regulate the chemical conditions of freshwater and marine habitats. Natural and anthropogenic inputs of nutrients and organic matter from the surface into the groundwater increase dissolved organic carbon (DOC) and nitrate concentrations—important indicators of water quality—that are then attenuated through microbial activity¹⁶. For instance, redox-driven microbial processes under aerobic or anaerobic conditions (e.g., denitrification and iron reduction) can substantially consume nitrate and reduce or transform DOC as groundwater migrates through subterranean freshwater environments¹⁵ or discharges into the sea¹²². Marine caves and cavities in

tropical regions are also areas of heterotrophic DOC consumption¹²³, which depletes dissolved oxygen¹²⁴.

Fresh groundwater discharge only accounts for a minor portion (~0.6%) of the total freshwater input to the world's oceans¹²⁵, but it can be critical locally for coastal ecosystem functioning due to its high solute and nutrient loads¹²⁶. At the land-sea interface, the region of a coastal aquifer where seawater and groundwater mix—the subterranean estuary—is a biogeochemical reaction zone that modulates nutrient and carbon fluxes from rocky, sandy, and muddy coastlines to marine ecosystems and fisheries^{122,127}. For example, microbial activity reduces nitrate and methane concentrations in groundwater discharged from sandy coasts^{128,129}, and methane and DOC in groundwater discharging from karstic coastlines^{130,131}. Moreover, sinkholes along karst coastlines are hotspots for carbon burial¹³², highlighting their potential for inclusion in blue carbon stocks. Given that approximately 40% of the world's population lives within 100 km of the coast, understanding these dynamics is of growing global importance¹³³.

Beyond biogeochemical cycles, subterranean ecosystems regulate key physical conditions in the environment. For example, hydrogeological conditions in aquifers control land subsidence, a phenomenon mainly driven by excessive groundwater extraction and aquifer compaction, a process that occurs when an aquifer's volume and storage capacity decreases due to the removal of water, often resulting in land subsidence. Globally, land subsidence leads to the loss of aquifer storage (~17 km³/year) and affects mainly cropland and urban areas (73%)¹³⁴. Consequences include damage to infrastructure, increased flood hazards, and substantial economic and human impacts^{135,136}.

Groundwater also supports ecosystem services provided by groundwater-dependent ecosystems¹³⁷, such as water storage, purification, and flood control. In turn, groundwater-fed vegetation controls erosion rates¹³⁸, buffers and attenuates mass movement, and regulates the overall hydrological cycle and water flow, and contributes to flood control and coastal protection. For example, tidal marsh vegetation stabilizes sediment and prevents shoreline erosion¹³⁹, while riparian trees like willows (*Salix* spp.) reduce erosion along riverbanks during heavy flow¹⁴⁰. The value of groundwater-fed vegetation in flood control has been estimated at about €16

billion in the EU alone¹⁴¹. Moreover, groundwater provides essential baseflow to rivers, particularly during dry seasons, sustaining river flow and influencing nutrient cycling and contaminant transport. Globally, baseflow is estimated to account for 59% ± 7% of river flow¹⁴².

Regulation of biological conditions

Subterranean ecosystems largely support surface vegetation^{137,143} and marine habitats^{122,144}. Approximately 37% of the world's vegetation depends on groundwater to some extent^{145,146}. The quality and availability of groundwater influence the distribution, diversity, functioning, and resilience of these plant communities¹⁴³. This dependency is particularly pronounced in drought-prone regions, where threshold levels of groundwater availability serve as indicators of potential drought refugia¹⁴⁷. Groundwater discharged into the marine environment delivers nutrients and affects water quality in estuaries, coral reefs, lagoons, mangroves, and saltmarshes^{122,148}.

Subterranean ecosystems also act as temporary, daily, or seasonal habitats for many surface animals and plants, all of which are integral to interconnected subterranean-surface food webs¹⁷. Surface vertebrates shelter or nest in cave entrances^{149–151}, while bats mate near entrances, but breed and hibernate in deeper sections¹⁵². Different vertebrates and invertebrates move in and out of terrestrial caves, often guided by circadian rhythms or seasonal cues^{151,153–155}. Aquatic insects, crustaceans, and fish seek refuge in the hyporheic zone of rivers during droughts¹⁵⁶. Groundwater inputs also heavily influence freshwater fish behavior, migration, spawning, and distribution¹⁵⁶. Similarly, marine caves host diverse sessile invertebrates (e.g., sponges, corals, bryozoans and brachiopods), fishes, and crustaceans, including many economically and ecologically valuable species such as the precious red coral *Corallium rubrum*¹⁵⁷. As climate become more unpredictable, these subterranean refugia are expected to grow in importance because of their environmental stability⁵⁷.

Arguably, cave-dwelling bats represent the best-studied example of biological regulation by subterranean ecosystems. Bats provide critical pollination and seed dispersal services for economically important plants, including figs, durian, mango,

and agave^{31,158}. For instance, the pollination services of *Eonycteris spelaea* to durian farmers in Sulawesi, Indonesia, were valued at US\$117 per hectare during each fruiting season¹⁵⁹. Another notable example is the mutualistic relationship between bats and agave. The pollination of agave relies on bats, particularly the cave-dwelling *Leptonycteris nivalis*, which, in turn, depend on agave during their seasonal migrations¹⁶⁰. Agave holds cultural and economic significance in Mexico as a source of food, spirits (tequila and mezcal), and fiber.

Insectivorous bats are also key biological controllers due to their hunting efficiency. For example, the cave-dwelling species *Pteronotus gymnonotus* and *P. personatus* consume 5–28% of their body weight in insects each night¹⁶¹. At least 81 species of insectivorous bats, including several obligate or facultative cave-dwellers, prey on over 760 species of insect pests that affect economically important crops such as corn, coffee, cotton, rice, apples, macadamia nuts, cocoa, and grapes¹⁶². Some of these species form massive colonies. For example, Mexican free-tailed bats (*Tadarida brasiliensis*) can form colonies of millions of individuals. During the summer, when bat populations peak in Bracken Cave, Texas, they can remove approximately 100 tons of insects per night, with the annual value of this pest suppression estimated at US\$3.42 million³¹. The economic importance of insectivorous bats in Northern America has been estimated to be as high as exceeding US\$3.7 billion per year¹⁶³.

Mitigation of pollutants

The Chemical Abstracts Service lists >200 million organic and inorganic synthesized compounds, with 20,000–30,000 new entries added daily¹⁶⁴. Many of these chemicals, especially those produced in large volumes, are released into the environment and eventually make their way underground, either passively (e.g., through percolating water) or intentionally (historically, shallow aquifers and caves were often used as waste disposal sites)^{165,166}.

Against this backdrop, a critical service is self-purification—the removal or immobilization of pollutants by natural processes²⁷. Subterranean microorganisms are key actors in this process, transforming harmful substances into more stable or

less toxic forms. For example, bacteria such as *Alcaligenes*, *Acinetobacter*, and *Pseudomonas* can immobilize heavy metals or dissolve phosphate minerals, aiding in the removal of contaminants¹⁶⁷. As for organic pollutants, microbes can degrade or mineralize compounds like petroleum hydrocarbons and halogenated solvents, particularly in point-source contamination scenarios¹⁶⁸. However, these processes are often slow, as microbial activity in subsurface environments is limited, and groundwater contamination can persist for years^{169,170}. For example, nitrates persist in groundwater for decades unless hypoxic or anoxic conditions and an appropriate electron donor (e.g., organic matter, pyrite) are present¹⁷¹.

Self-purification processes may be stimulated via amendment of electron acceptors (e.g., dissolved oxygen), electron donors (e.g., molasses), and bacterial strains (termed bioaugmentation)¹⁷². Managed aquifer recharge systems can effectively remove contaminants¹⁷³, including pharmaceuticals and antibiotics, through degradation processes that depend on the aquifer's redox state and temperature¹⁷⁴. However, biotransformation processes can sometimes produce byproducts that are recalcitrant to further degradation or more toxic than their parent compounds, highlighting the complexity of chemical regulation in groundwater systems¹⁷⁵.

Beyond microorganisms, larger subterranean fauna may also contribute to water purification through bioturbation of sediments and filtration^{176,177}. Based on consumption rates and rough density estimates of the isopod *Phreatoicus typicus* in New Zealand, it has been estimated that a population of 100 individuals can process approximately 7–28 tonnes of sediment per hectare annually and assimilate 120–650 grams of organic carbon per hectare annually¹⁷⁶. Synergistic effects with microorganisms appear to be particularly important in this context. Amphipods, isopods, and other invertebrates bioturbate and aerate sediments, creating favorable conditions for microbial communities to degrade contaminants^{176–178}. For example, the isopod *Coecidotaea tridentata* enhances both planktonic and sedimentary bacterial abundance and activity through the excretion of nitrogen, which promotes microbial growth, the disturbance of sediments, and the direct consumption of bacteria¹⁷⁹.

Potential for climate change mitigation

Subterranean ecosystems—particularly karst environments and caves—play a surprisingly important yet understated role in locally regulating atmospheric composition. Microbially-mediated formation of speleothems, such as moonmilks, sequesters and stores CO₂^{180,181}. Furthermore, aerobic caves act as net sinks for atmospheric methane (CH₄), actively consuming this greenhouse gas through microbial oxidation mediated by methane-feeding (methanotrophic) bacteria^{182–184} or through other processes¹⁸⁵. Within flooded caves of a karst subterranean estuary, it is estimated that ~1.4 tons of methane was consumed during 6 months across a ~100 km² catchment region in the Yucatán Peninsula¹³¹. It is unlikely that this magnitude of methane removal would affect global greenhouse gas budgets, but it quantifies the contribution of a critical energy source for an anchialine food web¹³⁰.

Beyond gas fluxes, subterranean ecosystems exert influence on microclimatic conditions. Their ability to buffer temperature and maintain high humidity levels creates stable environments that interact with aboveground climates, especially in regions with extensive karst topography^{186,187}. In terrestrial systems, this kind of regulation is often aided by bryophyte cushions (mosses and liverworts) developing in the entrance-zone of caves, which function as living sponges, intercepting rainfall, fog, and dew and retaining water volumes several times their dry mass. By slowly releasing this stored moisture into the substrate and underlying fissures, they buffer hydrological extremes at the subterranean-surface interface, sustain high local humidity for microbial and faunal communities, and contribute measurably to the water-storage service of groundwater-dependent ecosystems¹⁸⁸. In aquatic and marine settings, flooded caves and other subterranean environments have an important role in heat transfer through groundwater transport. Aquifers in rocky coastlines, such as karstic and volcanic platforms, are distinct from others, because the fissures and conduits enhance hydraulic transport and exchange of material with the sea through diffuse processes or submarine springs^{144,189,190}. Tidal driven oscillation of fresh groundwater discharge has been shown to transport heat to the sea from a volcanic platform¹⁹¹. On the contrary, tropical carbonate platforms may

cool the nearby sea through fresh groundwater discharge while facilitating the marine-derived saline water to import heat from the coast to inland¹⁹².

Beyond caves, groundwater-dependent ecosystems such as groundwater-fed wetlands, fens, riparian forests, and woodlands facilitate atmospheric CO₂ uptake through photosynthesis, root respiration, bicarbonate formation in soil, and the subsequent storage of carbon in groundwater or its precipitation as calcium carbonate¹⁹³. Vegetation supported by groundwater, such as the redwood forests of Northern California, grows more robustly and for longer periods compared to vegetation without groundwater access, sequestering significantly more carbon¹⁹⁴. Notably, areas with groundwater-dependent ecosystems store approximately 790 million tons of CO₂—nearly double California’s annual emissions¹⁹⁴. However, these benefits can be counterbalanced by the dewatering of groundwater-dependent ecosystems. For example, estimates suggest that wetlands could emit ~408 gigatons of CO₂ between 2021 and 2100 if degraded or drained¹⁹⁵.

Cultural services

Cultural ecosystem services are the non-material benefits people derive from ecosystems, contributing to cultural identity, spirituality, scientific endeavors, and quality of life⁹. Subterranean ecosystems contribute to all ([Figure 3](#)).

Tourism and recreation

Terrestrial and marine caves are among the most frequently visited geo- and ecotourism attractions worldwide. A recent synthesis identified 1,223 show caves across 95 countries, involving an estimated 79 million visitors in 2019²⁰. This generates around 800 million euros in entrance fees, with an even greater economic impact when considering related tourist activities—souvenir shops, restaurants, bars, and local transport ([Figure 4D](#)). Inevitably, this level of tourism comes not without impacts, including structural damage to caves, alterations to local climatic conditions,

the introduction of external organic matter and non-native fungi, bacteria, and animals, and the growth of photosynthetic organisms due to artificial lighting²¹.

Furthermore, geothermal phenomena linked to subterranean ecosystems—such as boiling lakes, mud ponds, and geysers—serve as striking natural attractions, drawing visitors to destinations that blend wonder with recreation. Some of these features also fuel the wellness sector. Thermal springs, long used by humans—and other apes¹⁹⁶—for health and wellness, are increasingly being transformed into modern hot spring resorts and water parks. Similarly, speleotherapy, particularly speleoclimatotherapy and radon therapy, offers drug-free therapeutic benefits. For example, the unique microclimate of salt caves and mines—characterized by fine aerosols of NaCl, K⁺, and Mg²⁺, high humidity, low radiation, light air ions, hypoallergenic air, and stable temperature—effectively alleviates different respiratory syndromes¹⁹⁷.

Terrestrial and aquatic caves are popular recreational sites for activities such as caving, snorkeling, scuba diving, and boat tours^{198,199}. These activities range from spontaneous experiences lasting a few hours, undertaken solo or in groups, to more structured expeditions and cave trips that require advanced speleological knowledge and skills. Often this kind of tourism brings visitors to caves that would be closed to humans otherwise, which may cause local impact to the ecosystems but also enhance scientific knowledge by citizens, amateur scientists, and speleologists.

Finally, subterranean-related ecotourism offers opportunities for wildlife enthusiasts to observe animals in their natural habitats. For instance, bat-watching is increasingly popular worldwide²⁰⁰. The nightly emergence of millions Mexican free-tailed bats from caves in the Southwestern US is estimated to attract over 240,000 visitors each year, conservatively valued at \$6.5 million annually²⁰¹. Such activities support local economies and provide unique educational experiences for the public, raising awareness about the ecological significance of subterranean ecosystems.

Aesthetic and artistic value

Subterranean landscapes inspire and support a range of artistic expressions^{22,23}. For instance, artistic practices have explored groundwater as a theme through creative

expressions of its sensory qualities—tastes, smells, sounds, textures, and movements—as well as its landscapes, cultural significance, and community connections²³. Contemporary abstract art frequently draws from the textures and patterns of speleothems, as seen in the cave-inspired works of artist Ana Teresa Barboza. Literature has frequently embraced subterranean themes, such as Jules Verne’s *Journey to the Center of the Earth* and Haruki Murakami’s *Hard-Boiled Wonderland* and the *End of the World*. Music, too, draws inspiration from subterranean acoustics, with composers like John Luther Adams creating pieces that echo the resonant and mysterious qualities of caves. Architecture has similarly demonstrated how caves and sinkholes can be reimagined into cultural and artistic venues, with spaces like *Los Jameos del Agua* in Lanzarote, shaped by César Manrique and Jesús Soto. These are just a few examples among many^{22,23}.

It has been argued that subterranean-related art may improve scientific communication and support the conservation of subterranean ecosystems^{22,23,202}. For example, projects such as the virtual reconstructions of cave art by the Chauvet Cave team not only preserve these fragile environments but also educate the public about their ecological and historical significance. Likewise, the Cenoteando initiative (<https://cenoteando.mx/>) in Mexico, has developed several educational materials that combine scientific accuracy with artistic expression to promote environmental awareness and proper stewardship of cenotes, promoting a sustainable interaction with these fragile environments. Similarly, artworks and photography that highlight the fragility of subterranean ecosystems, such as those by environmental artists like Agnes Denes and Martin Broen, can galvanize support and financial backing for conservation campaigns. Lastly, there is a practical significance to exploring aesthetics of subterranean features. For example, groundwater aesthetics—taste, odor, color, and clarity—is essential in shaping cultural perceptions and public trust in water supplies²⁰³.

Scientific research

Terrestrial caves have long been regarded as model systems for scientific research across various fields^{18,204,205}. The convergent adaptations of subterranean organisms make subterranean ecosystems a rich subject for evolutionary research, with a

lineage of studies tracing back to Charles Darwin²⁰⁶. Several cave-adapted species, such as cavefish and crustaceans, serve as established model organisms for evolutionary studies and beyond⁴². Furthermore, due to their climatic stability, low biological diversity, simple habitat structure, and often isolated nature, caves allow researchers to minimize many confounding factors that typically complicate ecological studies in surface environments¹⁷. Similarly, marine caves in the littoral zone have been described as “deep-sea mesocosms”, providing direct human access to deep-sea-like conditions²⁰⁷.

Importantly, this expanding research agenda builds upon the observations made by individuals who regularly explore subterranean environments, often driven by personal passion and a deep appreciation for nature. Speleological and cave diving clubs are typically composed of highly experienced, non-scientific explorers who possess the technical expertise necessary to access and map these underground spaces. Scientific research is also increasingly supported by dedicated subterranean research facilities, such as the Moulis Experimental Ecology Station in France and the Boulby Underground Laboratory in the UK, which provide controlled environments for ecological and evolutionary experiments⁴². Other underground laboratories, including Gran Sasso (Italy) and SNOLAB (Canada), further highlight the broader scientific value of caves, extending beyond biology to fields such as astroparticle physics.

Caves play a crucial role in archaeology and paleontology by safeguarding fossils, sediments, prehistoric artifacts, and even recently extinct species—such as certain birds known only from cave deposits in Macaronesia^{208,209}—as well as numerous human remains discovered in caves around the world^{210,211}. Stalagmites are archives for paleoclimate research, offering high-resolution records of past climatic fluctuations through isotopic and geochemical analyses²¹², while sediment deposits within cave systems record paleoenvironmental history, such as changes in sea level^{213–215}. All these archives provide clues into past ecosystems useful for reconstructing paleoenvironments and their biodiversity—yielding important implications for establishing baseline references for conservation and restoration efforts²¹⁶. For example, the analysis of speleothems has provided evidence of past

environmental changes and the anthropogenic impacts that contributed to the well-documented ecocide on Easter Island²¹⁷. Similarly, speleothems from lava tubes in the Galapagos Islands have revealed biomarkers of surface vegetation changes and human-induced pollution, emphasizing the need for robust conservation policies to mitigate the impact of anthropogenic activities²¹⁸.

The inspirational value of caves may even extend beyond Earth^{219–221}. The detection of volcanic caves on Mars and their protective properties against surface radiation, extreme temperatures, and atmospheric variability, have led researchers to explore caves on Earth from planetary science and astrobiological perspectives. A rich research agenda is shaping up, showing that these subterranean environments could serve as analogs for space exploration and planetary research^{220,222}, and offer insights into the possibility of extraterrestrial life^{223,224}. Specifically, microbial metabolism and mineral interactions in caves and lava tubes on Earth generate a variety of biosignatures^{222,225–227}, which provide reference models for potentially detecting extraterrestrial microbial life²²⁸. Moreover, deep caves offer polygons for training for astronauts (programme by European Space Agency), allowing them to practice behavior and tasks in harsh environments that resemble conditions in space.

Education

Subterranean ecosystems offer vast educational potential, especially for fostering scientific literacy and environmental awareness. Every cave provides visitors with an unforgettable experience, combining natural beauty with rich site-specific educational opportunities. Cave interpretation centers, guided tours, and interactive activities can help students and visitors appreciate the uniqueness of cave ecosystems and the importance of their conservation. Similarly, groundwater-fed springs enhance the natural beauty of their surroundings and serve as ideal settings for educational school trips. These sites allow students and teachers to observe firsthand the interactions between groundwater systems, biodiversity, and human activities²²⁹. Activities such as water quality testing, species identification, and habitat mapping

can transform these visits into living labs, offering hands-on learning experiences that reinforce classroom lessons.

This interplay between natural and cultural elements creates opportunities for educational projects that explore connections across disciplines such as biology, earth sciences, history, and even art. For example, studying speleothems can teach students about geological processes, offering a concrete visual representation of time accumulation, while analyzing the unique adaptations of cave-dwelling organisms can illustrate fundamental evolutionary principles. Importantly, these educational activities can be reinforced through citizen science initiatives. A recent citizen science project collected biological samples from over 300 municipal groundwater sites across Switzerland. This initiative bridged educational objectives with research goals, leading to the discovery of new species²³⁰ and enabling the mapping of macroecological patterns at unprecedented resolutions^{231–233}.

Cultural heritage and identity

Subterranean ecosystems often shape traditions, customs, and identities, influencing both positive and negative cultural narratives. Historically, caves were often perceived as liminal spaces—thresholds between the world of the living and the underworld. In European folklore, they often symbolize fear of the unknown and are believed to be entrances to Hell or lairs for dragons, trolls, and other sinister beings. This is illustrated in 17th-century engravings published in the monograph on the Duchy of Carniola by J. V. Valvasor, a Slovenian scientist, which depicts the beliefs of local inhabitants at the time²³⁴. Yet, caves have also held positive associations, for example by serving as places of refuge²⁵. Quintessential examples are underground cities in the Mediterranean region, such as Matera (Italy), Bulla Regia (Tunisia), and Cappadocia (Turkey), with tunnels, living quarters, and even chapels carved into the rock. Similarly, Coober Pedy, South Australia, is renowned for its man-made “dugouts”, subterranean residences bored into the hillsides of the desert. Beyond human-accessible cavities, features such as springs, anchialine pools, and oases played vital roles in community life, fostering social interaction and cohesion.

Specific organisms, such as bats, are often protagonists of these cultural narratives²³⁵. In some traditions, bats are feared as harbingers of darkness and death, a view perpetuated by Gothic literature and popular media. However, bats are also revered as symbols of luck, fertility, or protection. For example, in Chinese culture, bats are associated with happiness and prosperity, as the word for bat (*fu*) sounds like the word for good fortune. In the Americas, indigenous communities such as the Maya often incorporate bats into their mythology, viewing them as powerful guardians of the underworld.

Slovenia offers a prime example of how, even today, subterranean landscapes and their fauna can be deeply intertwined with national identity. The country is home to the renowned Postojna Cave, a UNESCO-listed site that has become a source of national pride²³⁶. This is the cave where the first scientific descriptions of exclusive cave-dwelling animals originated, beginning with the beetle *Leptodirus hochenwartii*, which marks the start of speleobiological research in 1832²³⁷. Slovenia is also the land where the discovery and scientific description of the olm (*Proteus anguinus*)—a blind, pale groundwater salamander—took place. *Proteus* has achieved iconic status, celebrated across various facets of Slovenian culture, from beer labels and public street art to the textile industry and contemporary art projects.

Spiritual and religious significance

Caves, anchialine pools, subterranean rivers, springs, and cenotes were often regarded as sacred or spiritually significant^{24,238}. For example, the caves of Crete were religious sites for the ancient Minoans, while Zeus was believed to have been born in a cave. In Greek mythology, the river Styx delineated Hades, the underworld (the prefix “stygo-” is still used today for “stygebionts”, a technical term referring to groundwater-dwelling organisms). Similarly, the cenotes of the Yucatán Peninsula were viewed by the Maya as both gateways to Xibalba, the underworld, and essential sources of life-giving water^{239,240}. Likewise, many anchialine pools in Hawai‘i

are revered as wahi pana (celebrated places), or strictly reserved for various uses, including royal baths, rituals, ceremonies, and other daily activities²⁴¹.

Countless rock-cut churches and monasteries worldwide further highlight the spiritual dimensions of subterranean sites²⁵. Likewise, groundwater provides spiritual and religious services through sacred water sites, often linked to natural features such as trees, stones, caves, and hills. These places offer a sensory connection to spiritual practices, with holy wells and springs frequently serving as focal points for rituals and supernatural engagement. While not all water sources are considered sacred, many cultures believe in offering gifts to water spirits to sustain their blessings. Springs emerging from caves hold particular significance, often seen as miraculously pure and ritually powerful, with evidence of reverence spanning from prehistoric times to contemporary cultures worldwide²³⁸. In Australia, many Aboriginal nations consider groundwater sites fundamental to their Dreamtime creation stories, in which the Rainbow Serpent is believed to have shaped landforms, springs, and river upwelling zones. Many sacred sites associated with fertility, teachings of lore, and cultural customs are linked to groundwater, holding immeasurable value for these communities²⁴².

Subterranean ecosystem disservices

Alongside their many positive contributions, ecosystems can also have effects that are perceived as harmful, unpleasant, or unwanted—termed “ecosystem disservices”²⁴³. While research on subterranean ecosystem disservices is virtually non-existent and beyond the scope of this assessment, it is important to briefly mention the potential human health and infrastructural risks associated with these environments. For instance, subterranean ecosystems can serve as reservoirs of pathogens and facilitate disease transmission. They harbor harmful microbes, fungi, and viruses, which may exist freely or be associated with specific organisms. Cave-roosting bats, in particular, are significant vectors of pathogens, including *Histoplasma* fungi found in bat guano, which can cause histoplasmosis in humans²⁴⁴. Additionally, subterranean environments can accumulate potentially toxic gases such as carbon dioxide, methane, hydrogen sulfide, and radon. These gases pose risks of

asphyxiation or poisoning, while radon may increase lung cancer risk for frequent visitors.

At the same time, the public's fascination with the underworld has often led to unfortunate accidents, particularly when individuals engage in caving or cave diving without adequate training or equipment—as in the famous Thailand cave rescue²⁴⁵ or the harrowing account of Sheck Exley in the Túnel de la Atlántida²⁴⁶. Subterranean ecosystems can also evoke some of the most common human phobias, as ranked by ref. ²⁴⁷. These environments are often dark (nyctophobia), enclosed (claustrophobia), contain deep pits or abysses (acrophobia/vertigo), and host fear-inducing organisms such as spiders (arachnophobia) and bats (chiroptophobia), potentially causing psychological distress in visitors. Beyond direct health risks, subterranean environments also pose threats to human infrastructure. Natural underground erosion, combined with human activities such as mining and groundwater extraction, can lead to cave collapses and sinkholes, damaging buildings and roads.

This discussion of disservices is far from exhaustive. Yet, it serves as a placeholder for further research in this area. Indeed, studying ecosystem disservices has been proposed as a way to better balance the benefits and drawbacks of nature, ultimately leading to a more objective evaluation of its net impact on human well-being²⁴⁸.

Box 2. The economic dimension of subterranean ecosystem services

Valuation of services provided by subterranean ecosystems is still in its early stages. A recent review of over 1,300 studies, yielding more than 9,400 monetary value estimates, found that subterranean ecosystems accounted for only 0.08% of the sample¹². Similarly, ref. ²⁴⁹ reported negligible research effort toward subterranean ecosystems when analyzing the correlation between ecological and economic assessments of 15 regulating services across 32 ecosystem types. While ecological roles such as nutrient cycling, soil formation, and groundwater provision are well-documented, their economic valuation remains limited, with groundwater being the most studied²⁴⁹.

In subterranean ecosystems, most valuation efforts focus on provisioning services, particularly groundwater. Methods include market prices, replacement costs, and production functions that measure the marginal impact of water on economic outputs like agricultural crops²⁵⁰. However, market prices often fail to capture the full social value of groundwater due to distortions like subsidies, requiring adjustments to reflect true economic value²⁵¹. Replacement cost methods, which estimate the expenses needed to restore lost services, offer an alternative approach²⁵².

Regulating services, though frequently reported for subterranean ecosystems, are rarely valued economically. For example, studies on erosion control, flood protection, and water quality regulation typically focus on surface ecosystems rather than subterranean ones^{253,254}. Similarly, cultural services like geo- and ecotourism are gaining attention, with examples including the recreational value of mining heritage and willingness-to-pay estimates for geo-guided tours^{255–258}.

Outlook: Communicating the value of subterranean ecosystems

Although still emerging, research on subterranean ecosystem services is likely to expand rapidly²⁸. We now have reasonable estimates of the global distribution and volume of certain types of subterranean ecosystems^{44,45}, a growing understanding of subterranean biodiversity patterns^{259,260}, and insights into how much of these ecosystems and their biodiversity are protected^{32,261}. Increasingly available open data²⁶² and emerging technologies—from omics tools²⁶³ and environmental DNA²⁶⁴ to terrestrial laser scanning²⁶⁵ and computer simulations⁴²—enable us to map and quantify subterranean ecosystems at unprecedented resolutions. Simultaneously, state-of-the-art economic theory provides a set of approaches to quantify the socio-economic relevance of these services at meaningful scales (Box 2). If harnessed effectively, these tools could bridge critical knowledge gaps in subterranean ecosystem services research.

Yet the importance of subterranean ecological processes to support surface ecosystems and human societies often goes unnoticed. Why do we celebrate climbing the highest mountains, yet overlook the exploration of the deepest caves? Why are so many unaware of the remarkable biodiversity thriving underground? And why do we study distant galaxies while Earth's subterranean environments may hold solutions to today's ecological and societal challenges?

Considering the importance of communicating these findings to inform real-world decision-making, this review aims to equip researchers and practitioners with a comprehensive *vade mecum* of examples, concepts, and ideas for conveying the importance of subterranean ecosystems. Effective communication requires tailoring messages to specific target audiences, using the right metaphors and psychological triggers. For some, subterranean biodiversity can be framed as a form of “life insurance,” emphasizing its role in maintaining ecosystem stability and resilience². Others may respond to economic metaphors, recognizing the monetary value of services like water filtration, carbon sequestration, and raw material provision (Box 2). At the same time, indigenous cultures, which have depended on subterranean ecosystems for centuries, offer invaluable traditional ecological knowledge and biocultural values that can enrich natural resource management strategies²⁴¹ These

perspectives often tap into metaphors related to the spiritual connection with these places, whereas the aesthetic allure and sense of mystery inherent to subterranean ecosystems can captivate audiences drawn to the unknown²⁶⁶.

By integrating these diverse perspectives, we can foster a deeper appreciation for subterranean ecosystems and their role in sustaining life on Earth. Subterranean biodiversity is not just a scientific curiosity—it is a cornerstone of planetary health, a source of resilience in the face of environmental change, and a testament to the interconnectedness of all ecosystems. With this knowledge, we can transform awkward questions about subterranean ecosystems into opportunities for inspiration and advocacy. As the world rallies to address environmental change and biodiversity loss, acknowledging and valuing the vital services provided by nature is essential to driving meaningful progress toward a more sustainable future. Ultimately, ensuring that subterranean ecosystems receive the attention and protection they deserve begins with one simple act: shifting the attitude of the next listener from indifference to appreciation.

Author contribution

SM, CG, and PC conceived the main idea, with suggestions by all authors. All authors contributed to the classification of services. SM and AB analyzed the data. SM wrote the first draft. All authors contributed to the writing of specific sections, and provided suggestions and additions to the overall text.

Fundings

This research was funded by Biodiversa+, the European Biodiversity Partnership under the 2021–2022 BiodivProtect joint call for research proposals, co-funded by the European Commission (GA N°101052342) and with the funding organisations Ministry of Universities and Research (Italy), Agencia Estatal de Investigación – Fundación Biodiversidad (Spain), Fundo Regional para a Ciência e Tecnologia (Azores, Portugal), Suomen Akatemia – Ministry of the Environment (Finland), Belgian Science Policy Office (Belgium), Agence Nationale de la Recherche (France), Deutsche Forschungsgemeinschaft e.V. (Germany), Schweizerischer Nationalfonds (Grant N° 31BD30_209583, Switzerland), Fonds zur Förderung der

Wissenschaftlichen Forschung (Austria), Ministry of Higher Education, Science and Innovation (Slovenia), and the Executive Agency for Higher Education, Research, Development and Innovation Funding (Romania). Additional support is provided by the P.R.I.N. 2022 projects “DEEPCHANGE” (2022MJSYF8) and “ANCHIALOS” (2022LLNF3N), funded by the Ministry of Universities and Research (Italy), by project “FRAGGLE” (PID2021-124640NB-I00), funded by MCIN/AEI/10.13039/501100011033 and by “ERDF A way of making Europe”. S.Mam. and T.D.L. acknowledge additional support of NBFC, funded by the Italian Ministry of University and Research, P.N.R.R., Missione 4 Componente 2, “Dalla ricerca all’impresa”, Investimento 1.4, Project CN00000033. V.G. was supported by the project “SponBIODIV - Marine sponge biodiversity from genes to ecosystems: delivering knowledge and tools for sustainable management and conservation”, funded by Biodiversa+, the European Biodiversity Partnership under the 2021-2022 BiodivProtect joint call for research proposals, co-funded by the European Commission (GA N°101052342) and the funding organisation General Secretariat for Research and Innovation (GSRI) under the Hellenic Ministry of Development (ΓΓΒΙΟ-0559429). P. C. is funded by cE3c (<https://doi.org/10.54499/UIDB/00329/2020>) and CHANGE (<https://doi.org/10.54499/la/p/0121/2020>). S.Mar. acknowledges the project HJI-D5-1 “Bringing in participatory approaches to widen the scope of natural capital valuation”, RESAS SRP 2022-2027, Scottish Government. I.R.A. was funded by Portuguese national funds through FCT – Fundação para a Ciência e a Tecnologia, I.P., under the Norma Transitória – <https://doi.org/10.54499/DL57/2016/CP1375/CT0003>. The MICROCENO project (<https://doi.org/10.54499/PTDC/CTA-AMB/0608/2020>), funded by the Portuguese Foundation for Science and Technology (FCT), and the MICROLAVA project (PROYEXCEL_00185) funded by the Regional Government of Andalusia, are also acknowledged.

Conflict of Interest

None declared.

Data and code availability

Data and code to reproduce the analysis is available in Github (<https://github.com/StefanoMammola/Subterranean-ecosystem-services>).

References

1. Rieb, J. T. *et al.* When, where, and how nature matters for ecosystem services: Challenges for the next generation of ecosystem service models. *Bioscience* **67**, 820–833 (2017).
2. Loreau, M. *et al.* Biodiversity as insurance: from concept to measurement and application. *Biol. Rev.* **96**, 2333–2354 (2021).
3. Bekessy, S. A., Runge, M. C., Kusmanoff, A. M., Keith, D. A. & Wintle, B. A. Ask not what nature can do for you: A critique of ecosystem services as a communication strategy. *Biol. Conserv.* **224**, 71–74 (2018).
4. Costanza, R. *et al.* The value of the world's ecosystem services and natural capital. *Nature* **387**, 253–260 (1997).
5. Lele, S., Springate-Baginski, O., Lakerveld, R., Deb, D. & Dash, P. Ecosystem services: origins, contributions, pitfalls, and alternatives. *Conserv. Soc.* **11**, 343–358 (2013).
6. Norgaard, R. B. Ecosystem services: From eye-opening metaphor to complexity blinder. *Ecol. Econ.* **69**, 1219–1227 (2010).
7. Benra, F. *et al.* National ecosystem restoration pledges are mismatched with social-ecological enabling conditions. *Commun. Earth Environ.* **5**, 731 (2024).
8. Chaplin-Kramer, R. *et al.* Wildlife's contributions to people. *Nat. Rev. Biodivers.* **1**, 68–81 (2025).
9. Haines-Young, R. & Potschin-Young, M. B. Revision of the Common International Classification for Ecosystem Services (CICES V5.1): A Policy Brief. *One Ecosyst.* **3**, e27108 (2018).
10. Temel, J., Jones, A., Jones, N. & Balint, L. Limits of monetization in protecting ecosystem services. *Conserv. Biol.* **32**, 1048–1062 (2018).
11. Chee, Y. E. An ecological perspective on the valuation of ecosystem services. *Biol. Conserv.* **120**, 549–565 (2004).
12. Brander, L. M. *et al.* Economic values for ecosystem services: A global synthesis and way forward. *Ecosyst. Serv.* **66**, 101606 (2024).
13. Gleeson, T., Cuthbert, M., Ferguson, G. & Perrone, D. Global groundwater sustainability, resources, and systems in the Anthropocene. *Annu. Rev. Earth Planet. Sci.* **48**, 431–463 (2020).
14. Siebert, S. *et al.* Groundwater use for irrigation – a global inventory. *Hydrol. Earth Syst. Sci.* **14**, 1863–1880 (2010).
15. Chapelle, F. H. The significance of microbial processes in hydrogeology and geochemistry. *Hydrogeol. J.* **8**, 41–46 (2000).

16. Griebler, C. & Avramov, M. Groundwater ecosystem services: a review. *Freshw. Sci.* **34**, 355–367 (2015).
17. Mammola, S. Finding answers in the dark: caves as models in ecology fifty years after Poulson and White. *Ecography (Cop.)*. **42**, 1331–1351 (2019).
18. Mammola, S. *et al.* Fundamental research questions in subterranean biology. *Biol. Rev.* **95**, 1855–1872 (2020).
19. Hesselberg, T. The biomimetic potential of novel adaptations in subterranean animals. *Front. Ecol. Evol.* **11**, 1127728 (2023).
20. Chiarini, V., Duckeck, J. & De Waele, J. A Global Perspective on Sustainable Show Cave Tourism. *Geoheritage* **14**, 82 (2022).
21. Piano, E., Mammola, S., Nicolosi, G. & Isaia, M. Advancing tourism sustainability in show caves. *Cell Reports Sustain.* **1**, 100057 (2024).
22. Mammola, S. *et al.* On art, science, and the conservation of subterranean ecosystems. *Subterr. Biol.* **51**, 1–19 (2025).
23. Gleeson, T. Groundwater connected art: practicing arts-based research to enrich how hydrogeology engages people, place and other disciplines. (2024).
24. Moyes, H. *Sacred Darkness: A Global Perspective on the Ritual Use of Caves*. (University Press of Colorado, 2012).
25. Bertini, A. Underground cities, cave dwelling, cave homes: yesterday, to day, tomorrow. *Reg. Archit. Mediterr. Area* 104 (2010).
26. Herman, J. S., Culver, D. C. & Salzman, J. Groundwater ecosystems and the service of water purification. *Stanford Environ. Law J.* **20**, 479 (2001).
27. Griebler, C., Avramov, M. & Hose, G. Groundwater ecosystems and their services: current status and potential risks. *Atlas Ecosyst. Serv. drivers, risks, Soc. responses* 197–203 (2019).
28. Canedoli, C. *et al.* Integrating landscape ecology and the assessment of ecosystem services in the study of karst areas. *Landsc. Ecol.* **37**, 347–365 (2022).
29. Iliopoulos, V. G. & Damigos, D. Groundwater Ecosystem Services: Redefining and Operationalizing the Concept. *Resources* **13**, 13 (2024).
30. Charchousi, D., Goula, A. & Papadopoulou, M. P. Mapping and Assessment of Groundwater Dependent Ecosystems (GDEs) Services—An Expert-based Land Use/Land Cover Scoring Approach. *Environ. Process.* **12**, 2 (2025).
31. Medellín, R. A., Wiederholt, R. & Lopez-Hoffman, L. Conservation relevance of bat caves for biodiversity and ecosystem services. *Biol. Conserv.* **211**, 45–50 (2017).
32. Mammola, S. *et al.* Perspectives and pitfalls in preserving subterranean biodiversity through protected areas. *npj Biodivers.* **3**, 2 (2024).
33. Keith, D. A., Ferrer-Paris, J. R., Nicholson, E. & Kingsford, R. T. *The IUCN Global Ecosystem Typology 2.0: Descriptive Profiles for Biomes and Ecosystem Functional Groups*. (IUCN., Gland, Switzerland, 2020). doi:10.2305/IUCN.CH.2020.13.en.

1064 34. Keith, D. A. *et al.* A function-based typology for Earth's ecosystems. *Nature* **610**, 513–518
1065 (2022).

1066 35. Pedersen, K. Exploration of deep intraterrestrial microbial life: current perspectives. *FEMS*
1067 *Microbiol. Lett.* **185**, 9–16 (2000).

1068 36. Edwards, K. J., Becker, K. & Colwell, F. The Deep, Dark Energy Biosphere: Intraterrestrial Life
1069 on Earth. *Annu. Rev. Earth Planet. Sci.* **40**, 551–568 (2012).

1070 37. Richter, F. *et al.* A guide to assess and value ecosystem services of grasslands. *Ecosyst.*
1071 *Serv.* **52**, 101376 (2021).

1072 38. Jakubiak, M. & Chmielowski, K. Identification of urban water bodies ecosystem services. *Acta*
1073 *Sci. Pol. Form. Circumiectionis* **19**, 73–82 (2020).

1074 39. Mukherjee, N. *et al.* Ecosystem service valuations of mangrove ecosystems to inform decision
1075 making and future valuation exercises. *PLoS One* **9**, e107706 (2014).

1076 40. Winkler, K. J., Viers, J. H. & Nicholas, K. A. Assessing ecosystem services and
1077 multifunctionality for vineyard systems. *Front. Environ. Sci.* **5**, 15 (2017).

1078 41. Kreamer, D. K. *et al.* The future of groundwater science and research. in *Global Groundwater*
1079 503–517 (Elsevier, 2021).

1080 42. Mammola, S. *et al.* Collecting eco-evolutionary data in the dark: Impediments to subterranean
1081 research and how to overcome them. *Ecol. Evol.* **11**, 5911–5926 (2021).

1082 43. Navarro-Barranco, C. *et al.* Conservation of dark habitats. in *Coastal habitat conservation*
1083 147–170 (Elsevier, 2023).

1084 44. Ferguson, G. *et al.* Crustal groundwater volumes greater than previously thought. *Geophys.*
1085 *Res. Lett.* **48**, e2021GL093549 (2021).

1086 45. Gleeson, T., Befus, K. M., Jasechko, S., Luijendijk, E. & Cardenas, M. B. The global volume
1087 and distribution of modern groundwater. *Nat. Geosci.* **9**, 161–167 (2016).

1088 46. Taylor, R. G. *et al.* Ground water and climate change. *Nat. Clim. Chang.* **3**, 322–329 (2013).

1089 47. Jasechko, S. *et al.* Rapid groundwater decline and some cases of recovery in aquifers
1090 globally. *Nature* **625**, 715–721 (2024).

1091 48. Kuang, X. *et al.* The changing nature of groundwater in the global water cycle. *Science* (80-.).
1092 **383**, eadf0630 (2024).

1093 49. Wu, W.-Y. *et al.* Divergent effects of climate change on future groundwater availability in key
1094 mid-latitude aquifers. *Nat. Commun.* **11**, 3710 (2020).

1095 50. Reinecke, R. *et al.* Uncertainty in model estimates of global groundwater depth. *Environ. Res.*
1096 *Lett.* (2024).

1097 51. Reinecke, R. *et al.* Uncertainty of simulated groundwater recharge at different global warming
1098 levels: a global-scale multi-model ensemble study. *Hydrol. Earth Syst. Sci.* **25**, 787–810
1099 (2021).

1100 52. Loaiciga, H. A. & Doh, R. Groundwater for People and the Environment: A Globally
1101 Threatened Resource. *Groundwater* **62**, 332–340 (2024).

- 1102 53. Rybach, L. Global status, development and prospects of shallow and deep geothermal energy.
1103 *Int. J. Terr. Heat Flow Appl. Geotherm.* **5**, 20–25 (2022).
- 1104 54. Huttner, G. W. Geothermal power generation in the World 2015-2020 Update report. in
1105 *Proceedings World Geothermal Congress* vol. 1 (2020).
- 1106 55. Murdock, H. E. *et al.* *Renewables 2021-Global Status Report*. [https://www.ren21.net/wp-](https://www.ren21.net/wp-content/uploads/2019/05/GSR2021_Full_Report.pdf)
1107 [content/ uploads/2019/05/GSR2021_Full_Report.pdf](https://www.ren21.net/wp-content/uploads/2019/05/GSR2021_Full_Report.pdf) (2021).
- 1108 56. Fišer, C., Pipan, T. & Culver, D. C. The Vertical Extent of Groundwater Metazoans: An
1109 Ecological and Evolutionary Perspective. *Bioscience* **64**, 971–979 (2014).
- 1110 57. Vaccarelli, I. *et al.* A global meta-analysis reveals multilevel and context-dependent effects of
1111 climate change on subterranean ecosystems. *One Earth* **6**, 1510–1522 (2023).
- 1112 58. Griebler, C. *et al.* Potential impacts of geothermal energy use and storage of heat on
1113 groundwater quality, biodiversity, and ecosystem processes. *Environ. Earth Sci.* **75**, 1391
1114 (2016).
- 1115 59. Benz, S. A. *et al.* Global groundwater warming due to climate change. *Nat. Geosci.* **17**, 545–
1116 551 (2024).
- 1117 60. Bayer, P. *et al.* Extracting past atmospheric warming and urban heating effects from borehole
1118 temperature profiles. *Geothermics* **64**, 289–299 (2016).
- 1119 61. Becher, J., Englisch, C., Griebler, C. & Bayer, P. Groundwater fauna downtown – Drivers,
1120 impacts and implications for subsurface ecosystems in urban areas. *J. Contam. Hydrol.* **248**,
1121 104021 (2022).
- 1122 62. de Graaf, I. E. M., Gleeson, T., Van Beek, L. P. H., Sutanudjaja, E. H. & Bierkens, M. F. P.
1123 Environmental flow limits to global groundwater pumping. *Nature* **574**, 90–94 (2019).
- 1124 63. Davis, K. F., Rulli, M. C., Seveso, A. & D’Odorico, P. Increased food production and reduced
1125 water use through optimized crop distribution. *Nat. Geosci.* **10**, 919–924 (2017).
- 1126 64. Fišer, C. *et al.* Toward sustainable irrigation practices safeguarding groundwater biodiversity
1127 and ecosystem services. *Bioscience* b1af016 (2025) doi:10.1093/biosci/b1af016.
- 1128 65. Carlson, G. *et al.* Intensive irrigation buffers groundwater declines in key European
1129 breadbasket. *Nat. Water* (2025) doi:10.1038/s44221-025-00445-4.
- 1130 66. Famiglietti, J. S. The global groundwater crisis. *Nat. Clim. Chang.* **4**, 945–948 (2014).
- 1131 67. Rodell, M., Velicogna, I. & Famiglietti, J. S. Satellite-based estimates of groundwater depletion
1132 in India. *Nature* **460**, 999–1002 (2009).
- 1133 68. Dalin, C., Wada, Y., Kastner, T. & Puma, M. J. Groundwater depletion embedded in
1134 international food trade. *Nature* **543**, 700–704 (2017).
- 1135 69. Hernández-Marrero, J.-C. *et al.* An approach to prehistoric herding in La Gomera (Canary
1136 Islands) through the study of domestic spaces. *Quat. Int.* **414**, 337–349 (2016).
- 1137 70. Delhon, C., Martin, L. & Thiébault, S. Neolithic herders and sheepfold caves in Southern
1138 France and adjacent areas: An overview from 40 years of bioarchaeological analyses. *Quat.*
1139 *Int.* **683**, 61–75 (2024).

71. Pardo, J. M. F. & Guerrero, I. C. Subterranean wine cellars of Central-Spain (Ribera de Duero): An underground built heritage to preserve. *Tunn. Undergr. Sp. Technol.* **21**, 475–484 (2006).
72. Krishnan, K. *et al.* Effects of temperature, relative humidity, and protective netting on *Tyrophagus putrescentiae* (schrank)(sarcoptiformes: Acaridae) infestation, fungal growth, and product quality of cave-aged Cheddar cheese. *J. Stored Prod. Res.* **83**, 44–53 (2019).
73. Maly, K. & Maly, O. *Ka Hana Lawai'a a Me Nā Ko'a O Na Kai 'ewalu: A History of Fishing Practices and Marine Fisheries of the Hawaiian Islands.* (Kumu Pono Associates, 2003).
74. Keliipuleole, K. *Aia Ke Kumu Waiwai Ma Makalawena: Analysis of the Natural Resources within the Makalawena Loko Wai'ōpae Complex.* (University of Hawai'i at Manoa, 2022).
75. Yan, T. H., Babji, A. S., Lim, S. J. & Sarbini, S. R. A Systematic Review of Edible Swiftlet's Nest (ESN): Nutritional bioactive compounds, health benefits as functional food, and recent development as bioactive ESN glycopeptide hydrolysate. *Trends Food Sci. Technol.* **115**, 117–132 (2021).
76. Rath, B. S., Kumar, P. S. & Vo, D.-V. N. Critical review on hazardous pollutants in water environment: Occurrence, monitoring, fate, removal technologies and risk assessment. *Sci. Total Environ.* **797**, 149134 (2021).
77. Brinkløv, S. & Warrant, E. Oilbirds. *Curr. Biol.* **27**, R1145–R1147 (2017).
78. Mickleburgh, S., Waylen, K. & Racey, P. Bats as bushmeat: a global review. *Oryx* **43**, 217–234 (2009).
79. Jenkins, R. K. B. & Racey, P. A. Bats as bushmeat in Madagascar. *Madagascar Conserv. Dev.* **3**, (2008).
80. Kamins, A. O. *et al.* Uncovering the fruit bat bushmeat commodity chain and the true extent of fruit bat hunting in Ghana, West Africa. *Biol. Conserv.* **144**, 3000–3008 (2011).
81. Tanalgo, K. C., Sritongchuay, T., Agduma, A. R., Cruz, K. C. Dela & Hughes, A. C. Are we hunting bats to extinction? Worldwide patterns of hunting risk in bats are driven by species ecology and regional economics. *Biol. Conserv.* **279**, 109944 (2023).
82. Verkaik, J., Sutanudjaja, E. H., Oude Essink, G. H. P., Lin, H. X. & Bierkens, M. F. P. GLOBGM v1. 0: a parallel implementation of a 30 arcsec PCR-GLOBWB-MODFLOW global-scale groundwater model. *Geosci. Model Dev.* **17**, 275–300 (2024).
83. Tang, L. & Werner, T. T. Global mining footprint mapped from high-resolution satellite imagery. *Commun. Earth Environ.* **4**, 134 (2023).
84. Sonter, L. J., Dade, M. C., Watson, J. E. M. & Valenta, R. K. Renewable energy production will exacerbate mining threats to biodiversity. *Nat. Commun.* **11**, 4174 (2020).
85. Statista. Mining - Worldwide. Statista (n.d.) <https://www.statista.com/outlook/io/mining/worldwide#production> (2025).
86. Reichl, C. & Schatz, M. *World Data Mining 2024.* (2024).
87. Mammola, S. *et al.* Climate change going deep: The effects of global climatic alterations on cave ecosystems. *Anthr. Rev.* **6**, 98–116 (2019).

88. Nanni, V., Piano, E., Cardoso, P., Isaia, M. & Mammola, S. An expert-based global assessment of threats and conservation measures for subterranean ecosystems. *Biol. Conserv.* **283**, 110136 (2023).
89. Ferreira, R. L., de Oliveira, M. P. A. & Silva, M. S. Subterranean biodiversity in ferruginous landscapes. *Cave Ecol.* 435–447 (2018).
90. Ferreira, R. L. *et al.* Brazilian cave heritage under siege. *Science* (80-.). **375**, 1238–1239 (2022).
91. Cleary, D. M. *et al.* Guano stable isotopic evidence of Anthropocene climate change influence on aridity and vegetation dynamics. *Ecol. Indic.* **145**, 109710 (2022).
92. Sakoui, S. *et al.* The Life Hidden Inside Caves: Ecological and Economic Importance of Bat Guano. *Int. J. Ecol.* **2020**, 9872532 (2020).
93. Rinaudo, M. Chitin and chitosan: Properties and applications. *Prog. Polym. Sci.* **31**, 603–632 (2006).
94. Kaya, M., Seyyar, O., Baran, T. & Turkes, T. Bat guano as new and attractive chitin and chitosan source. *Front. Zool.* **11**, 59 (2014).
95. Riquelme, C. *et al.* Actinobacterial diversity in volcanic caves and associated geomicrobiological interactions. *Front. Microbiol.* **6**, 1342 (2015).
96. Miller, A. Z. *et al.* Biogenic Mn oxide minerals coating in a subsurface granite environment. *Chem. Geol.* **322**, 181–191 (2012).
97. Paar, D. *et al.* Geochemical and mineralogical characterization of speleothems from the karst of Croatia as potential sources of data for environmental researches. *J. geochemical Explor.* **167**, 20–37 (2016).
98. Sauro, F. *et al.* Microbial diversity and biosignatures of amorphous silica deposits in orthoquartzite caves. *Sci. Rep.* **8**, 17569 (2018).
99. Zada, S. *et al.* Cave microbes as a potential source of drugs development in the modern era. *Microb. Ecol.* 1–14 (2021).
100. Zhu, H.-Z. *et al.* Bacteria and metabolic potential in karst caves revealed by intensive bacterial cultivation and genome assembly. *Appl. Environ. Microbiol.* **87**, e02440-20 (2021).
101. Ghezzi, D. *et al.* Insights into the microbial life in silica-rich subterranean environments: microbial communities and ecological interactions in an orthoquartzite cave (Imawari Yeuta, Auyan Tepui, Venezuela). *Front. Microbiol.* **13**, 930302 (2022).
102. Ghezzi, D. *et al.* Ancient and remote quartzite caves as a novel source of culturable microbes with biotechnological potential. *Microbiol. Res.* **286**, 127793 (2024).
103. Pipite, A. *et al.* Isolation, antibacterial screening, and identification of bioactive cave dwelling bacteria in Fiji. *Front. Microbiol.* **13**, 1012867 (2022).
104. Gatinho, P. *et al.* From cultural and natural heritage to reservoir of biomedicine: Prospection of bioactive compounds produced by bacterial isolates from caves. *Int. Biodeterior. Biodegradation* **190**, 105773 (2024).
105. Salazar-Hamm, P. S. *et al.* Subterranean marvels: microbial communities in caves and underground mines and their promise for natural product discovery. *Nat. Prod. Rep.* (2025).

1219 106. Cheeptham, N. *et al.* Cure from the cave: Volcanic cave actinomycetes and their potential in
1220 drug discovery. *Int. J. Speleol.* **42**, 35–47 (2013).

1221 107. Kosznik-Kwaśnicka, K., Golec, P., Jaroszewicz, W., Lubomska, D. & Piechowicz, L. Into the
1222 unknown: microbial communities in caves, their role, and potential use. *Microorganisms* **10**,
1223 222 (2022).

1224 108. Uriz, M. J. *et al.* An approach to the ecological significance of chemically mediated bioactivity
1225 in Mediterranean benthic communities. *Mar. Ecol. Prog. Ser.* 175–188 (1991).

1226 109. Turon, X., Martí, R. & Uriz, M. J. Chemical bioactivity of sponges along an environmental
1227 gradient in a Mediterranean cave. *Sci. Mar.* **73**, 387–397 (2009).

1228 110. Audoin, C. *et al.* Balibalosides, an original family of glucosylated sesterterpenes produced by
1229 the Mediterranean sponge *Oscarella balibalo*. *Mar. Drugs* **11**, 1477–1489 (2013).

1230 111. Rotter, A. *et al.* The essentials of marine biotechnology. *Front. Mar. Sci.* **8**, 629629 (2021).

1231 112. Suárez-Moo, P. *et al.* Exploring the microbial community and biotechnological potential of the
1232 sponge *Xestospongia* sp. from an anchialine cave in the Yucatán Peninsula. *Ciencias Mar.* **50**,
1233 (2024).

1234 113. Gatinho, P., Salvador, C., Silva, A. M. & Caldeira, A. T. Prokaryotic communities from pristine
1235 cave environments: biotechnological potential with sustainable production. *Sustainability* **15**,
1236 7471 (2023).

1237 114. Riddle, M. R. *et al.* Insulin resistance in cavefish as an adaptation to a nutrient-limited
1238 environment. *Nature* **555**, 647 (2018).

1239 115. Yoshizawa, M. *et al.* The evolution of a series of behavioral traits is associated with autism-risk
1240 genes in cavefish. *BMC Evol. Biol.* **18**, 89 (2018).

1241 116. Gore, A. V *et al.* An epigenetic mechanism for cavefish eye degeneration. *Nat. Ecol. Evol.* **2**,
1242 1155–1160 (2018).

1243 117. Griebler, C. & Lueders, T. Microbial biodiversity in groundwater ecosystems. *Freshw. Biol.* **54**,
1244 649–677 (2009).

1245 118. Rockström, J. *et al.* Planetary boundaries: exploring the safe operating space for humanity.
1246 *Ecol. Soc.* **14**, (2009).

1247 119. Steffen, W. *et al.* Planetary boundaries: Guiding human development on a changing planet.
1248 *Science (80-.).* **347**, 1259855 (2015).

1249 120. McClain, M. E. *et al.* Biogeochemical hot spots and hot moments at the interface of terrestrial
1250 and aquatic ecosystems. *Ecosystems* 301–312 (2003).

1251 121. Schmidt, S. I., Cuthbert, M. O. & Schwientek, M. Towards an integrated understanding of how
1252 micro scale processes shape groundwater ecosystem functions. *Sci. Total Environ.* **592**, 215–
1253 227 (2017).

1254 122. Santos, I. R. *et al.* Submarine groundwater discharge impacts on coastal nutrient
1255 biogeochemistry. *Nat. Rev. Earth Environ.* **2**, 307–323 (2021).

1256 123. de Goeij, J. M., van den Berg, H., van Oostveen, M. M., Epping, E. H. G. & Van Duyl, F. C.
1257 Major bulk dissolved organic carbon (DOC) removal by encrusting coral reef cavity sponges.
1258 *Mar. Ecol. Prog. Ser.* **357**, 139–151 (2008).

124. Young, C. *et al.* Effects of short-term variations in sea level on dissolved oxygen in a coastal karst aquifer, Quintana Roo, Mexico. *Limnol. Oceanogr.* **63**, 352–362 (2018).
125. Luijendijk, E., Gleeson, T. & Moosdorf, N. Fresh groundwater discharge insignificant for the world's oceans but important for coastal ecosystems. *Nat. Commun.* **11**, 1260 (2020).
126. Slomp, C. P. & Van Cappellen, P. Nutrient inputs to the coastal ocean through submarine groundwater discharge: controls and potential impact. *J. Hydrol.* **295**, 64–86 (2004).
127. Moore, W. S. The subterranean estuary: a reaction zone of ground water and sea water. *Mar. Chem.* **65**, 111–125 (1999).
128. Santos, I. R. S. *et al.* Nutrient biogeochemistry in a Gulf of Mexico subterranean estuary and groundwater-derived fluxes to the coastal ocean. *Limnol. Oceanogr.* **53**, 705–718 (2008).
129. Schutte, C. A., Wilson, A. M., Evans, T., Moore, W. S. & Joye, S. B. Methanotrophy controls groundwater methane export from a barrier island. *Geochim. Cosmochim. Acta* **179**, 242–256 (2016).
130. Brankovits, D. *et al.* Methane-and dissolved organic carbon-fueled microbial loop supports a tropical subterranean estuary ecosystem. *Nat. Commun.* **8**, 1–12 (2017).
131. Brankovits, D. *et al.* Hydrologic controls of methane dynamics in karst subterranean estuaries. *Global Biogeochem. Cycles* **32**, 1759–1775 (2018).
132. Adame, M. F., Santini, N. S., Torres-Talamante, O. & Rogers, K. Mangrove sinkholes (cenotes) of the Yucatan Peninsula, a global hotspot of carbon sequestration. *Biol. Lett.* **17**, 20210037 (2021).
133. UNEP (United Nations Environment Programme). *Coastal Resilience*. (2021).
134. Hasan, M. F., Smith, R., Vajedian, S., Pommerenke, R. & Majumdar, S. Global land subsidence mapping reveals widespread loss of aquifer storage capacity. *Nat. Commun.* **14**, 6180 (2023).
135. Erkens, G., Bucx, T., Dam, R., De Lange, G. & Lambert, J. Sinking coastal cities. *Proc. Int. Assoc. Hydrol. Sci.* **372**, 189–198 (2015).
136. Connor, R. & Miletto, M. *The United Nations World Water Development Report 2022: Groundwater: Making the Invisible Visible*. (UNESCO Publishing, 2022).
137. Saccò, M. *et al.* Groundwater is a hidden global keystone ecosystem. *Glob. Chang. Biol.* **30**, e17066 (2024).
138. Lowry, C. S. & Loheide, S. P. Groundwater-dependent vegetation: Quantifying the groundwater subsidy. *Water Resour. Res.* **46**, (2010).
139. Temmerman, S. *et al.* Impact of vegetation on flow routing and sedimentation patterns: Three-dimensional modeling for a tidal marsh. *J. Geophys. Res. Earth Surf.* **110**, (2005).
140. Borysiak, J., Czyryca, P. & Stępniewska, M. Erosion control ecosystem service provided by *Salix acutifolia* Willd. Neophyte on the South Baltic coast: Insights from Wolin Island, Poland. *Quaest. Geogr.* **43**, 5–19 (2024).
141. Vallecillo, S. *et al.* Accounting for changes in flood control delivered by ecosystems at the EU level. *Ecosyst. Serv.* **44**, 101142 (2020).

- 1298 142. Xie, J. *et al.* Majority of global river flow sustained by groundwater. *Nat. Geosci.* **17**, 770–777
1299 (2024).
- 1300 143. Glanville, K., Sheldon, F., Butler, D. & Capon, S. Effects and significance of groundwater for
1301 vegetation: A systematic review. *Sci. Total Environ.* **875**, 162577 (2023).
- 1302 144. Moore, W. S. The effect of submarine groundwater discharge on the ocean. *Ann. Rev. Mar.*
1303 *Sci.* **2**, 59–88 (2010).
- 1304 145. Barbeta, A. & Peñuelas, J. Relative contribution of groundwater to plant transpiration
1305 estimated with stable isotopes. *Sci. Rep.* **7**, 10580 (2017).
- 1306 146. Evaristo, J. & McDonnell, J. J. Prevalence and magnitude of groundwater use by vegetation: a
1307 global stable isotope meta-analysis. *Sci. Rep.* **7**, 44110 (2017).
- 1308 147. Rohde, M. M. *et al.* Establishing ecological thresholds and targets for groundwater
1309 management. *Nat. water* **2**, 312–323 (2024).
- 1310 148. Moosdorf, N. *et al.* A State-Of-The-Art Perspective on the Characterization of Subterranean
1311 Estuaries at the Regional Scale. *Front. Earth Sci.* **Volume 9-**, (2021).
- 1312 149. Baker, G. M. Quantifying wildlife use of cave entrances using remote camera traps. *J. Cave*
1313 *Karst Stud.* **77**, 200–210 (2015).
- 1314 150. dos Santos, T., de Souza, A. M., Bondezan, F. L. & Eterovick, P. C. Going underground: What
1315 the natural history traits of cave users can tell us about cave use propensity. *J. Herpetol.* **56**,
1316 153–163 (2022).
- 1317 151. Tuniyev, B. S., Koval, A. G. & Vargovitsh, R. S. Amphibians and reptiles in the caves of the
1318 Greater Caucasus. *Nat. Conserv. Res.* **6**, 1–16 (2021).
- 1319 152. Furey, N. M. & Racey, P. A. Conservation Ecology of Cave Bats. in *Bats in the Anthropocene:*
1320 *Conservation of Bats in a Changing World* (eds. Voigt, C. C. & Kingston, T.) 463–500
1321 (Springer International Publishing, Cham, 2016). doi:10.1007/978-3-319-25220-9_15.
- 1322 153. Novak, T., Thirion, C. & Janžekovič, F. Hypogean ecophase of three hymenopteran species in
1323 Central European caves. *Ital. J. Zool.* **77**, 469–475 (2010).
- 1324 154. Mammola, S. & Isaia, M. Day-night and seasonal variations of a subterranean invertebrate
1325 community in the twilight zone. *Subterr. Biol.* **27**, 31–51 (2018).
- 1326 155. Moog, O., Christian, E. & Eis, R. Increased cave use by butterflies and moths: a response to
1327 climate warming? *Int. J. Speleol.* **50**, 15–24 (2021).
- 1328 156. Land, E. & Peters, C. N. Groundwater impacts on stream biodiversity and communities: a
1329 review. *J. Freshw. Ecol.* **38**, 2260801 (2023).
- 1330 157. Gerovasileiou, V. & Bianchi, C. N. Mediterranean marine caves: a synthesis of current
1331 knowledge. *Oceanogr. Mar. Biol. An Annu. Rev.* **59**, 1–88 (2021).
- 1332 158. Ramírez-Fráncel, L. A. *et al.* Bats and their vital ecosystem services: a global review. *Integr.*
1333 *Zool.* **17**, 2–23 (2022).
- 1334 159. Sheherazade, Ober, H. K. & Tsang, S. M. Contributions of bats to the local economy through
1335 durian pollination in Sulawesi, Indonesia. *Biotropica* **51**, 913–922 (2019).

160. Trejo-Salazar, R. *et al.* Historical, temporal, and geographic dynamism of the interaction between Agave and Leptonycteris nectar-feeding bats. *Am. J. Bot.* **110**, e16222 (2023).
161. Pimentel, N. T., da Rocha, P. A., Pedroso, M. A. & Bernard, E. Estimates of insect consumption and guano input in bat caves in Brazil. *Mammal Res.* **67**, 355–366 (2022).
162. Tuneu-Corral, C. *et al.* Pest suppression by bats and management strategies to favour it: a global review. *Biol. Rev.* **98**, 1564–1582 (2023).
163. Boyles, J. G., Cryan, P. M., McCracken, G. F. & Kunz, T. H. Economic Importance of Bats in Agriculture. *Science (80-.)*. **332**, 41–42 (2011).
164. Chemical Abstracts Service. (n.d.). *CAS Content*. (2025).
165. Misstear, B., Vargas, C. R., Lapworth, D., Ouedraogo, I. & Podgorski, J. A global perspective on assessing groundwater quality. *Hydrogeol. J.* **31**, 11–14 (2023).
166. Lapworth, D. J., Boving, T. B., Kreamer, D. K., Kebede, S. & Smedley, P. L. Groundwater quality: Global threats, opportunities and realising the potential of groundwater. *Science of the Total Environment* vol. 811 152471 at (2022).
167. Putilina, V. S. & Yuganova, T. I. The Role of Microbiological Processes in the Formation of Geochemical Barriers and Redox Zones under Conditions of Contamination of Soils and Aquifers with Metals Near MSW Disposal Sites. *Water Resour.* **49**, S83–S93 (2022).
168. Das, N. & Chandran, P. Microbial degradation of petroleum hydrocarbon contaminants: an overview. *Biotechnol. Res. Int.* **2011**, 941810 (2011).
169. Tatti, F. *et al.* Experimental and numerical evaluation of Groundwater Circulation Wells as a remediation technology for persistent, low permeability contaminant source zones. *J. Contam. Hydrol.* **222**, 89–100 (2019).
170. Li, P., Karunanidhi, D., Subramani, T. & Srinivasamoorthy, K. Sources and Consequences of Groundwater Contamination. *Arch. Environ. Contam. Toxicol.* **80**, 1–10 (2021).
171. Basu, N. B. *et al.* Managing nitrogen legacies to accelerate water quality improvement. *Nat. Geosci.* **15**, 97–105 (2022).
172. Logeshwaran, P., Megharaj, M., Chadalavada, S., Bowman, M. & Naidu, R. Petroleum hydrocarbons (PH) in groundwater aquifers: An overview of environmental fate, toxicity, microbial degradation and risk-based remediation approaches. *Environ. Technol. Innov.* **10**, 175–193 (2018).
173. Laws, B. V., Dickenson, E. R. V., Johnson, T. A., Snyder, S. A. & Drewes, J. E. Attenuation of contaminants of emerging concern during surface-spreading aquifer recharge. *Sci. Total Environ.* **409**, 1087–1094 (2011).
174. Burke, V., Duennbier, U. & Massmann, G. The effect of aeration on the removal of wastewater-derived pharmaceutical residues from groundwater—A laboratory study. *Water Sci. Technol.* **67**, 658–666 (2013).
175. Postigo, C. & Barceló, D. Synthetic organic compounds and their transformation products in groundwater: occurrence, fate and mitigation. *Sci. Total Environ.* **503**, 32–47 (2015).
176. Boulton, A. J. A., Fenwick, G., Hancock, P. J. A. & Harvey, M. S. Biodiversity, functional roles and ecosystem services of groundwater invertebrates. *Invertebr. Syst.* **22**, 103–116 (2008).

177. Hose, G. C. & Stumpp, C. Architects of the underworld: bioturbation by groundwater invertebrates influences aquifer hydraulic properties. *Aquat. Sci.* **81**, 20 (2019).
178. Malard, F. & Hervant, F. Oxygen supply and the adaptations of animals in groundwater. *Freshw. Biol.* **41**, 1–30 (1999).
179. Edler, C. & Dodds, W. K. The ecology of a subterranean isopod, *Caecidotea tridentata*. *Freshw. Biol.* **35**, 249–259 (1996).
180. Martin-Pozas, T. *et al.* Role of subterranean microbiota in the carbon cycle and greenhouse gas dynamics. *Sci. Total Environ.* **831**, 154921 (2022).
181. Ghezzi, D. *et al.* The microbiota characterizing huge carbonatic moonmilk structures and its correlation with preserved organic matter. *Environ. Microbiome* **19**, 25 (2024).
182. Waring, C. L. *et al.* Seasonal total methane depletion in limestone caves. *Sci. Rep.* **7**, 8314 (2017).
183. Webster, K. D. *et al.* Subterranean karst environments as a global sink for atmospheric methane. *Earth Planet. Sci. Lett.* **485**, 9–18 (2018).
184. Ojeda, L. *et al.* Methane sources and sinks in karst systems: The Nerja cave and its vadose environment (Spain). *Geochim. Cosmochim. Acta* **259**, 302–315 (2019).
185. Fernandez-Cortes, A. *et al.* Subterranean atmospheres may act as daily methane sinks. *Nat. Commun.* **6**, 7003 (2015).
186. Caldwell, T. G. *et al.* Spring discharge and thermal regime of a groundwater dependent ecosystem in an arid karst environment. *J. Hydrol.* **587**, 124947 (2020).
187. Goldscheider, N. *et al.* Global distribution of carbonate rocks and karst water resources. *Hydrogeol. J.* **28**, 1661–1677 (2020).
188. Cedrés-Perdomo, R. D., Polaiño-Martín, C., Jennings, L. & Gabriel, R. Seeking a hideout: Caves as refuges for various functional groups of bryophytes from Terceira Island (Azores, Portugal). *Diversity* **16**, 58 (2024).
189. Fleury, P., Bakalowicz, M. & De Marsily, G. Submarine springs and coastal karst aquifers: a review. *J. Hydrol.* **339**, 79–92 (2007).
190. Moosdorf, N. & Oehler, T. Societal use of fresh submarine groundwater discharge: An overlooked water resource. *Earth-Science Rev.* **171**, 338–348 (2017).
191. Taniguchi, M., Ishitobi, T. & Shimada, J. Dynamics of submarine groundwater discharge and freshwater-seawater interface. *J. Geophys. Res. Ocean.* **111**, (2006).
192. Beddows, P. A., Smart, P. L., Whitaker, F. F. & Smith, S. L. Decoupled fresh–saline groundwater circulation of a coastal carbonate aquifer: Spatial patterns of temperature and specific electrical conductivity. *J. Hydrol.* **346**, 18–32 (2007).
193. Singh, P., Jiroušek, M., Hájková, P., Horsák, M. & Hájek, M. The future of carbon storage in calcareous fens depends on the balance between groundwater discharge and air temperature. *CATENA* **231**, 107350 (2023).
194. Howard, J. K., Dooley, K., Brauman, K. A., Klausmeyer, K. R. & Rohde, M. M. Ecosystem services produced by groundwater dependent ecosystems: a framework and case study in California. *Front. Water* **5**, 1115416 (2023).

- 1416 195. Zou, J. *et al.* Rewetting global wetlands effectively reduces major greenhouse gas emissions. *Nat. Geosci.* **15**, 627–632 (2022).
1417
- 1418 196. Matsuzawa, T. Hot-spring bathing of wild monkeys in Shiga-Heights: origin and propagation of
1419 a cultural behavior. *Primates* **59**, 209–213 (2018).
- 1420 197. Munteanu, C. SPELEOTHERAPY-scientific relevance in the last five years (2013–2017)—A
1421 systematic review. *Balneo Res. J.* **8**, 252–254 (2017).
- 1422 198. Gunn, J. Diving in caves. *Encyclopedia of caves and karst science* 653–655 (2004).
- 1423 199. Wilson, J. M. Recreational caving. *Encyclopedia of Caves* 861–870 (2019).
- 1424 200. Kunz, T. H., Braun de Torrez, E., Bauer, D., Lobova, T. & Fleming, T. H. Ecosystem services
1425 provided by bats. *Ann. N. Y. Acad. Sci.* **1223**, 1–38 (2011).
- 1426 201. Bagstad, K. J. & Wiederholt, R. Tourism values for Mexican free-tailed bat viewing. *Hum.*
1427 *Dimens. Wildl.* **18**, 307–311 (2013).
- 1428 202. Danielopol, D. L. Conservation and protection of the biota of karst: assimilation of scientific
1429 ideas through artistic perception. *J. Cave Karst Stud.* **60**, 67 (1998).
- 1430 203. Burlingame, G. A., Dietrich, A. M., Adams, H. & Bartrand, T. A. Managing Drinking Water
1431 Aesthetics With Water Safety Planning. *Journal of American Water Work. Assoc.* **116**, 6–17
1432 (2024).
- 1433 204. Poulson, T. L. & White, W. B. The Cave Environment. *Science (80-.)*. **165**, 971–981 (1969).
- 1434 205. Martinez, A. & Mammola, S. Let research on subterranean habitats resonate! *Subterr. Biol.* **36**,
1435 63–71 (2020).
- 1436 206. Juan, C., Guzik, M. T., Jaume, D. & Cooper, S. J. B. Evolution in caves: Darwin's 'wrecks of
1437 ancient life' in the molecular era. *Mol. Ecol.* **19**, 3865–3880 (2010).
- 1438 207. Harmelin, J.-G. & Vacelet, J. Clues to deep-sea biodiversity in a nearshore cave. *Vie*
1439 *Milieu/Life Environ.* 351–354 (1997).
- 1440 208. Rando, J. C., Alcover, J. A., Olson, S. L. & Pieper, H. A new species of extinct scops owl
1441 (Aves: Strigiformes: Strigidae: Otus) from São Miguel island (Azores archipelago, north
1442 Atlantic ocean). *Zootaxa* **3647**, 343–357 (2013).
- 1443 209. Rando, J. C., Pieper, H., Olson, S. L., Pereira, F. & Alcover, J. A. A new extinct species of
1444 large bullfinch (Aves: Fringillidae: Pyrrhula) from Graciosa Island (Azores, North Atlantic
1445 Ocean). *Zootaxa* (2017).
- 1446 210. Chatters, J. C. *et al.* Late Pleistocene Human Skeleton and mtDNA Link Paleoamericans and
1447 Modern Native Americans. *Science (80-.)*. **344**, 750–754 (2014).
- 1448 211. Berger, L. R. *et al.* Homo naledi, a new species of the genus Homo from the Dinaledi
1449 Chamber, South Africa. *Elife* **4**, e09560 (2015).
- 1450 212. Fairchild, I. J. & Baker, A. *Speleothem Science: From Process to Past Environments*. (John
1451 Wiley & Sons, 2012).
- 1452 213. van Hengstum, P. J., Reinhardt, E. G., Beddows, P. A. & Gabriel, J. J. Linkages between
1453 Holocene paleoclimate and paleohydrogeology preserved in a Yucatan underwater cave.
1454 *Quat. Sci. Rev.* **29**, 2788–2798 (2010).

- 1455 214. van Hengstum, P. J., Scott, D. B., Gröcke, D. R. & Charette, M. A. Sea level controls
1456 sedimentation and environments in coastal caves and sinkholes. *Mar. Geol.* **286**, 35–50
1457 (2011).
- 1458 215. van Hengstum, P. J., Cresswell, J. N., Milne, G. A. & Iliffe, T. M. Development of anchialine
1459 cave habitats and karst subterranean estuaries since the last ice age. *Sci. Rep.* **9**, 11907
1460 (2019).
- 1461 216. Hughes, A. C. *et al.* Reconstructing cave past to manage and conserve cave present and
1462 future. *Ecol. Indic.* **155**, 111051 (2023).
- 1463 217. Miller, A. Z. *et al.* Analytical pyrolysis and stable isotope analyses reveal past environmental
1464 changes in coralloid speleothems from Easter Island (Chile). *J. Chromatogr. A* **1461**, 144–152
1465 (2016).
- 1466 218. Miller, A. Z. *et al.* Organic geochemistry and mineralogy suggest anthropogenic impact in
1467 speleothem chemistry from volcanic show caves of the Galapagos. *iScience* **25**, 104556
1468 (2022).
- 1469 219. Titus, T. N. *et al.* A roadmap for planetary caves science and exploration. *Nat. Astron.* **5**, 524–
1470 525 (2021).
- 1471 220. Sauro, F. *et al.* Speleology as an analogue to space exploration: The ESA CAVES training
1472 programme. *Acta Astronaut.* **184**, 150–166 (2021).
- 1473 221. Sauro, F. *et al.* Training astronauts for scientific exploration on planetary surfaces: The ESA
1474 PANGAEA programme. *Acta Astronaut.* **204**, 222–238 (2023).
- 1475 222. Wynne, J. J. *et al.* Fundamental science and engineering questions in planetary cave
1476 exploration. *J. Geophys. Res. Planets* **127**, e2022JE007194 (2022).
- 1477 223. Northup, D. E. *et al.* Lava Cave Microbial Communities Within Mats and Secondary Mineral
1478 Deposits: Implications for Life Detection on Other Planets. *Astrobiology* **11**, 601–618 (2011).
- 1479 224. Popa, R., Smith, A. R., Popa, R., Boone, J. & Fisk, M. Olivine-Respiring Bacteria Isolated from
1480 the Rock-Ice Interface in a Lava-Tube Cave, a Mars Analog Environment. *Astrobiology* **12**, 9–
1481 18 (2011).
- 1482 225. Westall, F. *et al.* Biosignatures on Mars: what, where, and how? Implications for the search for
1483 martian life. *Astrobiology* **15**, 998–1029 (2015).
- 1484 226. Palma, V. *et al.* Connecting molecular biomarkers, mineralogical composition, and microbial
1485 diversity from Mars analog lava tubes. *Sci. Total Environ.* **913**, 169583 (2024).
- 1486 227. Palma, V. *et al.* Decoding organic compounds in lava tube sulfates to understand potential
1487 biomarkers in the Martian subsurface. *Commun. Earth Environ.* **5**, 530 (2024).
- 1488 228. Macalady Jennifer, L. *et al.* Dominant Microbial Populations in Limestone-Corroding Stream
1489 Biofilms, Frasassi Cave System, Italy. *Appl. Environ. Microbiol.* **72**, 5596–5609 (2006).
- 1490 229. Reinfried, S., Aeschbacher, U., Kienzler, P. M. & Tempelmann, S. The model of educational
1491 reconstruction—a powerful strategy to teach for conceptual development in physical geography:
1492 the case of water springs. *Int. Res. Geogr. Environ. Educ.* **24**, 237–257 (2015).
- 1493 230. Alther, R., Bongni, N., Borko, Š., Fišer, C. & Altermatt, F. Citizen science approach reveals
1494 groundwater fauna in Switzerland and a new species of Niphargus (Amphipoda, Niphargidae).

- 1495 *Subterr. Biol.* **25**, 1–31 (2021).
- 1496 231. Schneider, A. S., Knüsel, M. & Altermatt, F. Assessment of occurrence, diversity, and biomass
1497 of macroinvertebrates in Swiss groundwater systems using citizen science data. *Subterr. Biol.*
1498 **46**, 147–164 (2023).
- 1499 232. Knüsel, M. *et al.* Systematic and highly resolved modelling of biodiversity in inherently rare
1500 groundwater amphipods. *J. Biogeogr.* **51**, 2094–2108 (2024).
- 1501 233. Knüsel, M., Alther, R. & Altermatt, F. Pronounced changes of subterranean biodiversity
1502 patterns along a Late Pleistocene glaciation gradient. *Ecography (Cop.)*. e07321 (2024).
- 1503 234. Crane, R. & Fletcher, L. *Cave: Nature and Culture*. (Reaktion Books, 2015).
- 1504 235. Sieradzki, A. & Mikkola, H. Bats in folklore and culture: a review of historical perceptions
1505 around the world. *Bats–Disease-prone But Benef.* (2022).
- 1506 236. Zagmajster, M., Polak, S. & Fišer, C. Postojna-Planina Cave System in Slovenia, a Hotspot of
1507 Subterranean Biodiversity and a Cradle of Speleobiology. *Diversity* vol. 13 at
1508 <https://doi.org/10.3390/d13060271> (2021).
- 1509 237. Schmidt, F. Beitrag zu Krain's Fauna. *Illyrisch- es Blatt* **3**, 9–10 (1832).
- 1510 238. Ray, C. Holy wells and sacred springs. in *Sacred Waters* 1–31 (Routledge, 2020).
- 1511 239. Munro, P. G. & de Lourdes Melo, M. The role of cenotes in the social history of Mexico's
1512 Yucatan Peninsula. *Environ. Hist. Camb.* **17**, 583–612 (2011).
- 1513 240. Melo Zurita, M. de L. Holes, subterranean exploration and affect in the Yucatan Peninsula.
1514 *Emot. Sp. Soc.* **32**, 100584 (2019).
- 1515 241. Gibson, V. L., Bremer, L. L., Burnett, K. M., Lui, N. K. & Smith, C. M. Biocultural values of
1516 groundwater dependent ecosystems in Kona, Hawai'i. *Ecol. Soc.* **27**, (2022).
- 1517 242. Moggridge, B. J. Aboriginal people and groundwater. *Proc. R. Soc. Queensland*, **126**, 11–27
1518 (2020).
- 1519 243. Blanco, J., Dendoncker, N., Barnaud, C. & Sirami, C. Ecosystem disservices matter: Towards
1520 their systematic integration within ecosystem service research and policy. *Ecosyst. Serv.* **36**,
1521 100913 (2019).
- 1522 244. Gugnani, H. C. & Denning, D. W. Infection of bats with *Histoplasma* species. *Med. Mycol.* **61**,
1523 myad080 (2023).
- 1524 245. Beech, H., Paddock, R. C. & Suhartono, M. Still can't believe it worked. The story of the
1525 Thailand cave rescue. *New York Times* (2018).
- 1526 246. Exley, S. *Caverns Measureless to Man*. (Cave Books, 1994).
- 1527 247. Correia, R. A. & Mammola, S. The searchscape of fear: A global analysis of internet search
1528 trends for biophobias. *People Nat.* **6**, 958–972 (2024).
- 1529 248. Schaubroeck, T. A need for equal consideration of ecosystem disservices and services when
1530 valuing nature; countering arguments against disservices. *Ecosyst. Serv.* **26**, 95–97 (2017).
- 1531 249. Kadykalo, A. N., Kelly, L. A., Berberi, A., Reid, J. L. & Findlay, C. S. Research effort devoted to
1532 regulating and supporting ecosystem services by environmental scientists and economists.
1533 *PLoS One* **16**, e0252463 (2021).

- 1534 250. Aziz, T. *et al.* Monetizing the role of water in sustaining watershed ecosystem services using a
1535 fully integrated subsurface–surface water model. *Hydrol. Earth Syst. Sci. Discuss.* **2023**, 1–34
1536 (2023).
- 1537 251. Singh, S., Kaur, P., Sachdeva, J. & Bhardwaj, S. Profitability analysis of major crops in punjab:
1538 some evidence from cost of cultivation survey data. *Indian J. Econ. Dev.* **13**, 71–78 (2017).
- 1539 252. Carrera-Hernández, J. J. & Gaskin, S. J. Water management in the Basin of Mexico: current
1540 state and alternative scenarios. (2009).
- 1541 253. Lundin-Frisk, E. *et al.* Improved assessments of subsurface projects: Systematic mapping of
1542 geosystem services and a review of their economic values. *J. Environ. Manage.* **365**, 121562
1543 (2024).
- 1544 254. Patault, E. *et al.* Analysis of off-site economic costs induced by runoff and soil erosion:
1545 Example of two areas in the northwestern European loess belt for the last two decades
1546 (Normandy, France). *Land use policy* **108**, 105541 (2021).
- 1547 255. Pérez-Álvarez, R., Torres-Ortega, S., Díaz-Simal, P., Husillos-Rodríguez, R. & De Luis-Ruiz,
1548 J. M. Economic valuation of mining heritage from a recreational approach: Application to the
1549 case of El Soplao Cave in Spain (Geosite UR004). *Sustainability* **8**, 185 (2016).
- 1550 256. Khalaf, E. E. D. A. H. Geoheritage and cultural-religious heritage of Samalute-Minia area in
1551 North Egypt. *Geoheritage* **16**, 5 (2024).
- 1552 257. Kubalíková, L. Cultural ecosystem services of geodiversity: A case study from Stranska skala
1553 (Brno, Czech Republic). *Land* **9**, 105 (2020).
- 1554 258. Cheung, L. T. O., Fok, L. & Fang, W. Understanding geopark visitors' preferences and
1555 willingness to pay for global geopark management and conservation. *J. Ecotourism* **13**, 35–51
1556 (2014).
- 1557 259. Zagmajster, M., Ferreira, R. L., Humphreys, W. F., Niemiller, M. L. & Malard, F. Patterns and
1558 determinants of richness and composition of the groundwater fauna. in *Groundwater ecology
1559 and evolution* 141–164 (Elsevier, 2023).
- 1560 260. Martínez, A. *et al.* Stygofauna Mundi: a comprehensive global biodiversity database of
1561 groundwater-related habitats across marine and freshwater realms. 69–73 at (2024).
- 1562 261. Sánchez-Fernández, D., Galassi, D. M. P., Wynne, J. J., Cardoso, P. & Mammola, S. Don't
1563 forget subterranean ecosystems in climate change agendas. *Nat. Clim. Chang.* **11**, 458–459
1564 (2021).
- 1565 262. Huggins, X. *et al.* A review of open data for studying global groundwater in social-ecological
1566 systems. *EarthArXiv* (2025) doi:10.31223/X5XB07.
- 1567 263. Pérez-Moreno, J. L., Iliffe, T. M. & Bracken-Grissom, H. D. Life in the Underworld: Anchialine
1568 cave biology in the era of speleogenomics. *Int. J. Speleol.* **49**, 149–170 (2016).
- 1569 264. Saccò, M. *et al.* eDNA in subterranean ecosystems: Applications, technical aspects, and future
1570 prospects. *Sci. Total Environ.* **820**, 153223 (2022).
- 1571 265. Idrees, M. O. & Pradhan, B. A decade of modern cave surveying with terrestrial laser
1572 scanning: A review of sensors, method and application development. *Int. J. Speleol.* **45**,
1573 71–88 (2016).

1574 266. Ryan-Davis, J. & Scalice, D. Co-Creating Ethical Practices and Approaches for Fieldwork.
1575 *AGU Adv.* **3**, e2022AV000762 (2022).