

# Flower constancy in pollinators: a bouquet of agendas shape interactions among mutualistic partners

Christoph Grüter

School of Biological Sciences, University of Bristol, 24 Tyndall Avenue, BS8 1TQ Bristol, UK

Correspondence: [c.grueter@bristol.ac.uk](mailto:c.grueter@bristol.ac.uk)

## Abstract

Plant-pollinator interactions have become a major research area due to their impact on key ecosystem services. One pollinator behaviour of particular importance is flower constancy, *i.e.* the tendency of pollinators to temporarily specialise on one flower species during a foraging trip, thereby promoting cross-pollination. The costs and benefits of flower constancy for both plants and pollinators are varied, complex and far from understood. This review aims to synthesise and interpret studies spanning the last decades, from both plant and pollinator perspectives. Flower constancy is often viewed as an epiphenomenon of pollinator cognition, but there is increasing recognition that pollinators show remarkable behavioural flexibility in their flower choice, often in response to ecological and social factors. Plants usually benefit from flower constancy, which reduces pollen loss and interspecific pollen transfer. However, in some situations, pollinator inconstancy can be advantageous due to increased visits from pollinators shared with co-flowering plants. The fitness consequences of pollinator behaviour for rare or invasive plants are intriguing yet little understood, with important implications plant conservation. Rather than seeing flower constancy as a strategy imposed on pollinators by

cognitive constraints, this review emphasises that plants and pollinators pursue varied agendas depending on their ecological context and lifestyle.

**Keywords:** Pollination, cognition, facilitation, flower constancy, behavioural flexibility

## 1. Introduction

Most flowering plants depend on animal pollinators for sexual reproduction [1]. The effectiveness of pollinators for this role depends on a range of morphological and behavioural traits. One behaviour of particular importance is flower constancy, the tendency of a pollinator to visit flowers of the same plant species during a foraging trip, sometimes for days [2,3], while bypassing rewarding alternative flowers [2,4–7]. While flower constancy has been observed and studied predominantly in bees (e.g. [2,5,8–11]), it has also been reported in butterflies [12–14], hover flies [15], beetles [16] and birds [17]. Flower constancy ensures that flowers receive pollen from the same species, facilitating cross-pollination. From a pollinator perspective, flower constancy is often viewed as a suboptimal strategy and the result of pollinators attempting to mitigate the limitations of their nervous system [2,5–7]. And yet, a growing number of studies highlight that pollinators are flexible and differ in their flower choice behaviour, e.g. in response to ecological or social factors. This review brings together traditional views and recent insights into the causes and consequences of flower constancy from both pollinator and plant perspectives with the aim to better understand the varied interests that shape interactions among plant and their pollinators.

At first glance flower constancy is a deceptively simple behaviour, a choice of a flower of one type over another type, and yet a closer look reveals a perplexing complexity that spans biological levels, from neurons to species communities, shapes plant-pollinator communities and raises a number of intertwined mechanistic and evolutionary questions. Here, flower

constancy refers to the tendency of pollinators to move between flowers of the same species when other rewarding flower types are available. Flower constancy is based on associative learning – e.g. between colour, odour or shape and floral rewards [6] – and is different from the innate dietary preferences found in specialist (oligolectic) pollinators [5,20; note that bees can be specialists when collecting pollen, but generalists when collecting nectar [18]). This definition of flower constancy is descriptive and agnostic regarding the causes and adaptive value of flower constancy (similar to Takagi & Ohashi [19], but different from Waser [5] who considered flower constancy a non-optimal foraging strategy caused by cognitive limitations).


Researchers have used two main approaches to quantify flower constancy: first, by observing movements between natural or artificial flowers, sometimes with the aim to calculate a flower constancy index (e.g. [2,5,20–23], see [5,20,24] for discussion of different indices). Second, by analysing pollen load purity of foraging bees, which reflects the sum of its foraging choices during a foraging trip (e.g. [25,26]). While the study of natural flower movements provides information about flower choice, the diversity and frequency of available options, it is often not feasible in natural habitats due to habitat and interaction network complexity or inaccessibility (e.g. the forest canopy). Analysing pollen load purity can be a convenient shortcut but findings can be misleading as pollen purity could be the result of local abundance and clustering of flowers. Pollinators could also be visiting different flower types while collecting pollen only from one [5,6]. Artificial flower setups solve many of these challenges (e.g. [8,19,22,27]), but they are feasible only for a small number of pollinators and artificial setups work best with sugar solution as a reward. Pollinators are likely to modify their strategies depending on whether they collect pollen or nectar, e.g. by being more specialised, when collecting pollen [9,18].

## 2. The plant perspective

Animal-mediated pollination involves the transfer of pollen from the male parts of the flower, the anthers, to the female parts, the stigma, with the help of a pollinator. Darwin [4] noted that flower constancy is of “great significance to the plant, as it favours cross fertilization of distinct individuals of the same species”. Flower constancy and inconstancy differentially shape competition between plant species by affecting both the quantity and quality of plant-pollinator interactions (Fig. 1) [14,28–39].

### *(a) Interspecific pollen transfer (IPT)*

Inconstant pollinators will frequently transfer pollen from one species to the stigmas of a different species. This interspecific pollen transfer (IPT) is ubiquitous [33–35,40–43] and can reduce plant fitness through a variety of mechanisms [31,35,43–46]. If a pollinator switches plant species, much of the original pollen is lost on the heterospecific flowers or it is groomed off in the process, which can reduce the seed set of conspecifics that receive less pollen as a result [29,31,35,37,47,48]. Pollen misplacement can also lower outcrossing rate, thereby reducing not only offspring quantity but also quality [48].



**Figure 1.** Pollinator behaviour affects interactions between plants. If plants from different species share the same pollinator, then this will likely affect the fitness of plants through different mechanisms, depending on whether pollinators are flower constant or not. While sharing pollinators will often have a negative impact on flowers through competition, there may be circumstances when the sharing of inconstant pollinators benefits plants through facilitation [32,34–37,49,50]. Plants visited by inconstant pollinators may still pay some costs due to interspecific pollen transfer, but these costs are outweighed by the benefits of increased visitation.

For the plant receiving heterospecific pollen, the deposition of this pollen can impair seed set [29,31,37], for example, through stigma clogging, which is the reduction of the receptive stigma surface area for conspecific pollen, stigma closure, which reduces the receptivity for conspecific pollen or active inhibition of germination and hybridisation (see [31] for review). IPT could be especially costly for rare plants co-flowering with more common species as they are likely to receive relatively more heterospecific pollen [28,31,37,46,51,52], which could increase the risk of extinction of rare plants [31,53]. Interspecific pollen transfer from invasive

to native plants has the potential to reduce the reproductive success of native plants ([31,43]; but see [40,41,54]). Flower constancy affects the strength of this effect, for example, the solitary bee *Rhodanthidium septemdentatum* switched more often from an invasive to native plants and carried more pollen between species than the highly flower constant honey bee *Apis mellifera*, with negative impacts on seed production in one of three native plant species visited by the solitary bees [55]. This highlights how different types of pollinators create different interaction pollen transfer networks (see also [56]). More flower constant pollinators reduce the risk of ITP and facilitate cross-pollination [57], but they may also increase repeat visits to flowers of the same plant, thereby increasing self-pollination (geitonogamy), which can impair plant fitness [58].

The costs of these interspecies pollen transactions can select for greater divergence in floral traits, e.g. different habitat choice, flowering times or floral morphology (e.g. morphology, colour or scent; see Section 4) and self-pollination [14,28–31,33,35,44,59]. A striking example are the closely related plants *Phlox drummondii* and *P. cuspidate*, which share the butterfly *Battus philenor* as their primary pollinator. Butterflies show little constancy when the two plants appear in their ancestral colours, leading to pollen loss and hybridisation [14]. Where the two species are sympatric, however, selection on an allele affecting flower pigment intensity causes flowers to have more distinctive colours, which increases flower constancy of their pollinator and promotes reproductive isolation in the plants [14]. This example highlights how flower constancy can drive evolutionary divergence in closely related sympatric lineages (see also [60,61]).

## (b) Facilitation

Not all studies, however, find that pollinator sharing has negative effects on plant fitness. Floral infidelity can be neutral or even beneficial for co-flowering plant species through facilitation (Fig. 1) [34,36,49,50,62], *i.e.* an increase in visitation rate due to the presence of co-flowering plants, for example through a “magnet effect”. For example, nonrewarding orchids benefit from [63] or provide benefits to [64] co-flowering rewarding plants (see also [65,66]). In high-altitude plant communities where pollinators are scarce, this facilitation could be common [34]. More generally, when plant abundance is low and pollination services are limited, sharing inconstant pollinators with co-flowering plants species could increase visitation rates due to a “mass effect”, *i.e.* the attraction of pollinators to the area (or the survival of pollinators in an area) [34,37,50,67]. Similarly, in environments experiencing widespread pollinator decline, inconstant pollinators may be of increasing value to plants.

Another potential benefit of gene flow driven by pollinator inconstancy is increased genetic variation or an exchange of adaptive traits (adaptive introgression) between non-fully reproductively isolated plant species, which may have been an important evolutionary force in many plant clades [35]. In sum, the effects of floral inconstancy on plant fitness are varied and a variation in the degree of flower constancy among pollinators, both at species and individual level, may benefit plant biodiversity (see also [68,69]).

## 3. The pollinator perspective

Pollinators are expected to maximise energy intake – either in terms of rate [70] or efficiency [71] – while making sure they also collect all required nutrients for a healthy diet [72,73]. The first requirement is primarily achieved by collecting nectar, the second is achieved by collecting different types of pollen. Flower constancy appears to be at odds with both of these

requirements: bypassing rewarding flowers risks reducing energy intake [74] due to time (increased foraging trip duration) and opportunity (missed rewards) costs [74], sub-optimal flower choice [5,75] and increased flight distances [76]. Accordingly, computer simulations of flower constant and inconstant bee colonies suggest that strict flower constancy is usually less successful in terms of net energy intake than foraging indiscriminately [74]. In addition, focusing on a particular plant species risks missing out on important nutrients (see section 4b) [77].

So why would a pollinator be flower constant? One common view is that cognitive limitations imposed by the processing, storage and retrieval of information favour flower constancy. Extracting a reward efficiently requires pollinators to learn a wealth of information about floral colours, odours, shapes, textures and motor skills, involving a range of cognitive tools [6]. Due to limitations of these cognitive tools, pollinators switching between plants would be less efficient at extracting rewards [5–7,78,79]. An alternative view is that flower constancy is the economically most successful foraging strategy given the limited information a pollinator has about its environment [7,20].

#### (a) Cognitive limitations

Darwin [4] and von Frisch [80] suggested that flower constancy is favoured due to the benefits of learning: a bee that has learned how to extract nectar from, for example, linden (*Tilia*) blossoms is more efficient than a bee unfamiliar with this flower [2,14] (an idea sometimes misleadingly been called “Darwin’s interference hypothesis” [81,82]). The flipside of this argument is that learning has costs and a pollinator switching to a new flower type would need time and energy to learn how to extract rewards efficiently [5,83–85]. However, research on the learning efficiency of pollinators suggests that pollinators learn fast: cabbage white butterflies (*Pieris rapae*) need only a couple of flower visits to learn how to efficiently reach



the nectar in two different flower types [12] and naïve *Bombus impatiens* bees collecting pollen from four plant reached a plateau in pollen collection efficiency within the first 20 visits [85]. On the other hand, *B. terrestris* collecting pollen from poppy flowers (*Papaver rhoeas*) improved their foraging efficiency over several foraging trips that included visiting hundred flower visits [84], showing that learning to exploit flowers sometimes does take time. Whether and how plant fitness is affected by pollinator learning is not well understood, but a recent study found that pollination success did not change as a result of learning how to handle flowers by *Bombus impatiens* foragers [85]. Overall, observations lead to the conclusion that bees and butterflies are quick learners, which challenges Darwin's idea that the benefits of knowing how to exploit a flower species are sufficient to favour bypassing rewarding alternatives.

Limitations in working- and long-term memory may also favour flower constancy because they cause time costs if pollinators are inconstant. Long-term memories, while robust and durable, may be costly or slow to retrieve [5,6,19]. Working memory, on the other hand, is prone to rapid decay and interference from competing information, such as new floral traits [6,12,86,87]. Inter-flower flights of inconstant bumble bees were found to be longer (~1-3 seconds) than flights between flowers of the same species [21,88]. These delays in inconstant flights could be due to a transient working memory that is stable for only a few seconds, or they could indicate that bees switching flower species need time to access long-term memory about the alternative type (if the second flower type is already familiar). These time delays in inter-floral flights and learning how to extract nectar are likely to accelerate flower visitation due to flower constancy, while also facilitating information consolidation from working- to long-term memory [21]. Overall, however, time costs due to inconstancy appear to be low. Lavery [89], for instance, found that *Bombus fervidus* experienced no time costs when switching between two types of flower with simple morphologies and only a ~1sec delay when switching between flower types with more complex morphologies, similar to the small

increases in handling time found by Raine & Chittka [21] and Goulson et al. [13]. One might argue that even small time gains can add up to a substantial amount of time saved as pollinators visit hundreds or even thousands of flowers per day. However, pollinators also accumulate time, opportunity and energy costs every time they fly past a rewarding flower. Computer simulations suggest that time costs of flower inconstancy need to be considerable (>30sec with each visit in the simulations) to make flower constancy more successful than inconstancy [74]. I am not aware of any empirical studies comparing the time costs of switching flower type – due to learning and/or memory processes – to the costs of ignoring flowers of a different type. Such studies are most likely absent because they are exceedingly difficult to perform.

The “search image” hypothesis proposes that flower constancy is the result of pollinators establishing a search image of a specific flower type in a complex visual environment [82]. Search images are thought to be useful when looking for cryptic targets, *i.e.* those that are difficult to find [90,91]. In contrast, flower constancy tends to be stronger when flowers become easier to detect or flower species become easier to differentiate (see Section 4a). Convincing evidence that “search images” cause of flower constancy is currently missing, but more research into visual background effects on flower constancy and attention priming [90,91] is needed to rule out that search “search images” affect plant-pollinator interactions.

While research suggests that multiple, non-mutually exclusive cognitive processes may contribute to the strength of flower constancy in some pollinator species, the idea that cognitive constraints predispose pollinators to flower constancy is at odds with evidence that pollinators are (i) able to process impressive amounts of information efficiently and (ii) are varied and flexible in their behaviour, often in response to ecological circumstances and their social lifestyle (Section 4). This suggests that ecological and social interactions shape the cognitive tools and behavioural strategies pollinators have at their disposal.

226

227 *(b) Informational limitations*

228 The costly information hypothesis [6,7] posits that flower constancy is the best strategy in an  
229 uncertain environment where acquiring information about better plant species would cost time  
230 and energy. Assessing the profitability of alternatives may require sampling a large number of  
231 flowers since different flowers of the same plant species offer variable rewards [92]. Flower  
232 constancy might then be the best option if the rewards currently experienced by a pollinator  
233 are above a threshold. Honey bees are indeed almost fully flower constant when the rewards  
234 they receive are above a reward threshold [22,93]. As the number of plant species in an  
235 environment increases, so do the sampling costs to obtain reliable information, thus favouring  
236 flower constancy [6,7]. While this hypothesis is intuitively appealing, computer simulations  
237 suggest that flower constancy becomes more costly as plant species diversity increases [74].  
238 This is because the time, energy and opportunity costs of bypassing flowers also increase when  
239 alternative options become relatively more numerous. Pollinators should be *less* flower  
240 constant in a habitat with more flower species. The simulation findings highlight an important  
241 point often missed in discussions about flower constancy: even if pollinators visit the most  
242 profitable flower species, flower constancy may not be the best strategy, especially when  
243 alternative options become more abundant [74]. Empirical evidence for this is mixed, however:  
244 while Gervais et al. [94] and Martínez-Bauer et al. [95] found that increasing plant diversity  
245 was indeed associated with lower flower constancy in bumble bees, Austin et al. [96] found  
246 that bumble bees became more flower constant when there are more options available. The first  
247 two studies were performed under natural conditions, whereas Austin et al. [20] used artificial  
248 flower arrays and lab conditions. More research is needed to understand how increasing plant  
249 diversity affects flower constancy in different pollinators.

250

## 4. Behavioural flexibility and species variation

If the discussion so far has given the impression that strict flower constancy is the rule among pollinators, then this would be misleading. We know little about the degree of flower constancy for the vast majority of pollinators, but numerous studies and the widespread phenomenon of heterospecific pollen transfer (see Section 2) suggest that pragmatism and flexibility guide flower choice in the pollinator world. Indeed, we would expect an optimal pollinator with multiple options to divide her time between exploiting familiar flowers and sampling alternative ones [97,98]. The impression that bees are commonly flower constant may have resulted from the focus on the Western honey bee *Apis mellifera*, a highly flower constant species [3,75,99], but even honey bees show flexibility in their floral choices in response to reward characteristics [7]. These two findings – variation between species and flexibility within species – suggest that there is ample scope for natural selection to drive flower choice strategies [7]. Below I discuss three types of drivers of that may be key to understanding behavioural flexibility: floral features, ecological factors and sociality.

### (a) Floral features

A key determinant of the strength of flower constancy is the value of the reward offered by flowers, which depends on the quality (*e.g.* sugar concentration) and quantity (amount or production rate) of the offered rewards [88,100–105]. Bumble bees preferentially foraged on more complex flowers only if it offered sucrose solution of higher concentration than simple flowers in experimental setups [102,103]. Bumble bees foraging naturally on different shrubs preferred the species with higher daily sucrose production per flower [106]. The small skipper butterfly was twice as likely to switch plant species after receiving a below-average reward quantity [13]. Even the highly flower constant honey bee *Apis mellifera* adjusted the degree of

flower constancy within seconds following a flower visit, with bees becoming less flower constant when reward quality, quantity and number were reduced [22].

The response to rewards also interacts with other floral features and flower constancy often increases as options become more dissimilar, either in a specific trait, such as flower colour and morphology or in the number of traits [5,19,27,45,79,88,99,101,107–109]. The link between flower constancy and pollinator perception is likely due to pollinators being able to learn to differentiate more efficiently among flowers with divergent traits, which, in turn, can drive evolutionary divergence among similar plant morphs (see Section 2). Bees also increase flower constancy as flower size and floral display size (*i.e.* larger number of inflorescences per plant) increase [110,111]. Since flower and display size have been shown to correlate positively with reward size [112], pollinators may use flower and display size as proxies of relative profitability of a flower species, and become less inclined to switch away from large flowers.


Reward value and extraction costs also depend on flower morphology, since the way flowers are built affects how fast pollinators can learn to extract rewards as well as the subsequent handling time costs and foraging rate (see also Section 3) [85,105,108,113,114]. For instance, visiting complex flowers is associated with handling times several times longer than those for simple flowers (up to 25 seconds *vs.* a few seconds for simple flowers) [85,113,114]. One might, therefore, predict that pollinators prefer simple flowers. Evidence, however, is mixed: while *Bombus impatiens* preferred the simpler of two artificial flower types [102], naïve *Bombus terrestris* foraging on natural flowers preferred complex types [113]. Similarly, *Bombus fervidus* were only flower constant when visiting plants with complex morphologies [89]. What could explain these counterintuitive observations? First, bees might be discouraged to switch to an alternative type due to their experience that becoming an expert forager is costly. Second, morphological complexity might discourage or exclude some pollinators while providing rewards for the expert forager [115,116]. Visiting a complex

flowers could, thus, be beneficial for pollinators experiencing intense competition, while plants might benefit if flower morphology filters out ineffective pollinators [116].

### *(b) Ecological factors: spatial distribution of flowers*

Recent studies have highlighted the importance of the spatial arrangement of flowers in influencing how pollinators balance floral fidelity *versus* behavioural flexibility [19,23]. Cape Sugarbirds (*Promerops cafer*), a rare example of a flower constant bird, adjust the degree of flower constancy in relation to the relative abundance of flower species [17]. Abundance and degree of clustering of co-flowering plants determine the distances pollinators need to fly between flowers. As distances increase, both honey bees and bumble bees become less flower constant and switching to the nearest flower type becomes more common [19,23,27,88,117–119]. Pollinators could gauge these distances based on travel costs [5] or based on floral visual angles [111]. Computer simulations similarly found that flower constancy reduces energy intake when flower density is low [74] as energy, time and opportunity costs of bypassing flowers increase when flower abundance is low. Thus, flower constancy is predicted to increase with flower abundance based on both energetics and cognitive limitations (working memory instability) arguments [6,119]. Working memory instability could be the underlying mechanism that allows bees to adjust flower constancy adaptively in relation to food source abundance.

The effects of inter-floral distances also explain why flower constancy is often higher when food sources are arranged in clusters rather than when evenly mixed (Fig. 2) [19,120]. For plants, on the other hand, the more frequent switching in more evenly mixed situations could be costly due to increased IPT.



**Figure 2.** Factors affecting behavioural flexibility and the strength of flower constancy. Flower constancy depends on floral features like reward size [22], ecological factors like the arrangement of flowers in space (e.g. [19,23,120]), flower diversity [94,95] or social factors like social information [74,121]. + and – indicate whether a particular factor is expected to increase or decrease flower constancy.

### (c) Interspecies variation and the effects of sociality

Different observers have noticed that pollinator species can vary considerably in how flower constant they are [2,5,7,39,77,122]. Bateman [123] and Waser [5], observing bees moving between flowers, found that honey bees were more flower constant than bumble bees. Furthermore, honey bees returning to their hive have mostly pure pollen loads [3], whereas those of bumble bees are commonly mixed (Fig. 3) [124]. In controlled laboratory experiments, *Bombus impatiens* foragers were also quicker to switch from a deteriorating sugar solution to an alternative one than *Apis mellifera* [125]. Even within bumble bees (*Bombus*) there appear to be considerable differences: in the South American *B. atratus* and *B. bellicosus*, 80% and

84% of pollen foragers visited just one plant during a foraging trip [9], whereas only 23% of pollen foragers showed flower constancy in the European *B. terrestris* [126]. Whether these differences indeed reflect innate interspecific differences or differences in ecological or floral factors remains to be studied. Overall, however, evidence supports the view that pollinator groups vary in the degree of flower constancy.

One factor that has been linked to flower constancy is sociality and social lifestyle. Solitary pollinators, including bees, butterflies and flies, are often less flower constant than highly social bees [42,55,123,127]. In a comparative study of pollen loads of 56 bee species in a temperate bee community, Smith et al. [122] found that social bees were more flower constant than solitary bees. Three explanations for this sociality effect have been proposed: the resource-partitioning hypothesis [6,7,100], the communication hypothesis [6,74,121] and the diet breadth hypothesis [77,128].

The resource partitioning hypothesis argues that flower constancy in social bees is a form of task partitioning that helps foragers reduce competition with nestmates by specialising on different flower species [6,100,126]. However, if a subset of nestmates specialises on a subset of flowers then they still compete within their group, while now also paying the energy, time and opportunity costs of flower constancy.





**Fig. 3.** A bumble bee forager collecting pollen on meadowsweet (*Filipendula ulmaria*). Her pollen package has two colours, showing the bee has visited two types of flowers during the same foraging trip (*Rubus* before switching to meadowsweet, photo: CG).

Sociality could favour flower constancy in bees due because many social bees communicate about food sources [74,121]. Since social bees share information selectively about high quality food sources, nestmates using social information are likely to discover more profitable food sources [129–133]. For example, honey bees use the waggle dance and some stingless bees lay pheromone trails to food sources [129,131,133–136]. Most social bees perform excitatory behaviours, such as jostling runs and trophallaxis (food sharing) inside the nest after finding particularly good food sources, which facilitate the learning of floral odours and stimulate the search for flowers with these odours [130–132,135,137,138]. This selective information sharing lowers the risk of specialising on flowers offering low-quality rewards and, therefore, reduces the benefits of sampling alternatives. Simulated bee colonies with communication and flower constancy indeed collected more energy than those without communication because selective information sharing allowed colonies to specialise on the

most profitable flower species in their environment [74]. In environments with abundant food sources and large rewards, flower constancy in combination with communication was the best strategy overall when plant diversity was low [74]. As plant diversity increased, inconstancy became the best strategy (see Section 3b). Thus, while communication about profitable flower types reduces one cost of flower constancy, specialising on sub-optimal flower species, it does not affect the time and energy costs of flower constancy.

Finally, sociality could affect flower constancy through indirect effects on diet diversity. Pollinators require a range of nutrients for a healthy diet [72,73,139]. Nectar is the main source of carbohydrates, while pollen provides most of the proteins, lipids and micronutrients [72,139]. As pollen from different plant species differ in their nutrient composition [72], collecting a small number of pollen types risks nutritional imbalances [140,141], with potentially negative fitness consequences [142–144]. In social bees, nectar and different types of pollen are often collected by different colony members as a form of division of labour [145–147]. A solitary bee, on the other hand, needs to collect both nectar and pollen by herself. Williams & Tepedino [128] found that the need to collect both nectar and pollen most likely explained why the solitary mason bee *Osmia lignaria* switched between plant species during foraging trips.


Since different colony members exploit different flower species in social species, flower constancy might not affect diet breadth or even increase it, especially with larger colony sizes [74,126,128]. Pollen analysis confirms that colonies of highly flower constant species exploit many plant species simultaneously [148–150]. However, only a small number of pollen types, usually <5, is collected in significant quantities. Thus, social bees may still face the risk nutritional deficiencies due to flower constancy, especially in environments with low plant diversity. To the best of my knowledge, controlled empirical studies on the link between diet breadth, colony size and flower constancy are lacking. However, computer simulations have

found that flower constant colonies with fewer than 50 foragers often exploit less than half as many plant species as inconstant colonies [77]. A larger colony size did increase the number of flower types visited even if colonies were flower constant, suggesting that larger colony sizes mitigate the risks of nutritional deficiencies due to flower constancy. This could help explain why bumble bees, which have smaller colony sizes than honey bees and stingless bees are less flower constant than the latter two groups [77].

## 5. Summary: a bouquet of agendas

Plants and pollinators have different agendas, *i.e.* fitness interests, and these agendas are often viewed as contrasting, with plants favouring flower constancy and pollinators favouring behavioural flexibility. However, plant and pollinator interests are varied, for instance, common and rare plants may experience different outcomes when their pollinators are flower constant. Likewise, social and solitary pollinators are likely to experience different costs and benefits due to flower constancy. As a result, interests of plants and pollinators may align in some situations, yet contrast in others (Fig. 4): for example, when a plant is abundant, flower visitation and flower constancy increase [17,152,153], which is likely to favour both plants and pollinators in terms of seed set for the former [153] and energy intake rate for the latter [74] (*Scenario 1*). Deceptive unrewarding plants and rare plants co-flowering with abundant flower species, on the other hand, may experience reduced visitation rates due to flower constancy (*Scenario 2*). These costs could outweigh the benefits of reduced interspecific pollen transfer due to flower constancy under some circumstances (Fig. 1). In plant species-rich habitats with a relatively even abundance and distribution, interspecific pollen transfer due to inconstancy is likely to be common, and flowers are likely to benefit from flower constancy. Flower constant pollinators, on the other hand, pay considerable opportunity costs in such

biodiverse habitats as most of the flowers they encounter will not be their preferred type (*Scenario 3*) [6,74]. Finally, when plants and pollinators are scarce, flower constancy could reduce pollinator visits to plants and reward rates for pollinators, thus negatively impacting the fitness of both plants and pollinators (*Scenario 4*).



**Fig 4.** Interests of both plants and their pollinators when pollinators are *flower constant*. Four Scenarios (1-4) are suggested that reflect different situations.

While some empirical and theoretical support for these scenarios exists, the interests of plants and pollinators remain far from understood. These knowledge gaps, especially when considering rare plants, remain a blind spot for plant conservation: it has been suggested that rare plants experience higher costs due to IPT [28,31,52,53], but whether and when these costs outweigh the benefits of facilitation [34,37,50] is poorly understood. Another important knowledge gap are the dietary requirements and foraging strategies, including flower constancy, of the vast majority of pollinators, especially in small-bodied pollinators and those in the tropics [18,35,39]. Research in both controlled environments and natural communities is

needed to better understand how different behavioural strategies impact pollen transfer networks, and how changes in species composition impact interactions among plants and their pollinators. Such an understanding is essential for linking mutualistic interactions to ecosystem functioning.

## 6. References

1. Ollerton J, Winfree R, Tarrant S. 2011 How many flowering plants are pollinated by animals? *Oikos* **120**, 321–326. (doi:10.1111/j.1600-0706.2010.18644.x)
2. Grant V. 1950 The flower constancy of bees. *The Botanical Review* **16**, 379–398.
3. Free JB. 1963 The flower constancy of honeybees. *Journal of Animal Ecology* **32**, 119–131.
4. Darwin C. 1876 *Cross and self fertilization in the vegetable kingdom*. London: Murray.
5. Waser NM. 1986 Flower constancy: definition, cause, and measurement. *American Naturalist* **127**, 593–603.
6. Chittka L, Thomson JD, Waser NM. 1999 Flower constancy, insect psychology, and plant evolution. *Naturwissenschaften* **86**, 361–377.
7. Grüter C, Ratnieks FLW. 2011 Flower constancy in insect pollinators: Adaptive foraging behaviour or cognitive limitation? *Communicative and Integrative Biology* **4**, 1–4.
8. Slaa EJ, Cevaál A, Sommeijer MJ. 1998 Floral constancy in *Trigona* stingless bees foraging on artificial flower patches: a comparative study. *Journal of Apicultural Research* **37**, 191–198.
9. Rossi N, Santos E, Salvarrey S, Arbulo N, Invernizzi C. 2015 Determination of flower constancy in *Bombus atratus* Franklin and *Bombus bellicosus* Smith (Hymenoptera: Apidae) through palynological analysis of nectar and corbicular pollen loads. *Neotrop Entomol* **44**, 546–552. (doi:10.1007/s13744-015-0322-5)
10. White D, Cribb BW, Heard TA. 2001 Flower constancy of the stingless bee *Trigona carbonaria* Smith (Hymenoptera: Apidae: Meliponini). *Australian Journal of Entomology* **40**, 61–64. (doi:10.1046/j.1440-6055.2001.00201.x)
11. Gaiarsa MP, Rehan S, Barbour MA, McFrederick QS. 2022 Individual dietary specialization in a generalist bee varies across populations but has no effect on the richness of associated microbial communities. *The American Naturalist* **200**, 730–737. (doi:10.1086/721023)
12. Lewis AC. 1986 Memory constraints and flower choice in *Pieris rapae*. *Science* **232**, 863–865.
13. Goulson D, Stout J, Hawson SA. 1997 Can flower constancy in nectar butterflies be explained by Darwin's interference hypothesis? *Oecologia* **112**, 225–231.

- 472 14. Hopkins R, Rausher MD. 2012 Pollinator-mediated selection on flower color allele drives  
473 reinforcement. *Science* **335**, 1090–1092. (doi:10.1126/science.1215198)
- 474 15. Goulson D, Wright NP. 1998 Flower constancy in the hoverflies *Episyrphus balteatus* (Degeer)  
475 and *Syrphus ribesii* (L.) (Syrphidae). *Behavioral Ecology* **9**, 213–219.
- 476 16. Pellmyr O. 1985 Flower constancy in individuals of an anthophilous beetle, *Byturus ochraceus*  
477 (Scriba) (Coleoptera: Byturidae). *The Coleopterists Bulletin* **39**, 341–345.
- 478 17. Schmid B, Nottebrock H, Esler KJ, Pagel J, Böhning-Gaese K, Schurr FM, Mueller T,  
479 Schleuning M. 2016 A bird pollinator shows positive frequency dependence and constancy of  
480 species choice in natural plant communities. *Ecology* **97**, 3110–3118. (doi:10.1002/ecy.1565)
- 481 18. Cane JH, Sipes S. 2006 Characterizing floral specialization by bees: analytical methods and a  
482 revised lexicon for oligolecty. In *Plant-pollinator interactions: from specialization to*  
483 *generalization* (eds NM Waser, J Ollerton), pp. 99–122. Chicago and London: The University of  
484 Chicago Press.
- 485 19. Takagi K, Ohashi K. 2025 Realized flower constancy in bumble bees: Optimal foraging strategy  
486 balancing cognitive and travel costs and its possible consequences for floral diversity. *Functional*  
487 *Ecology* **39**, 863–875. (doi:10.1111/1365-2435.70008)
- 488 20. Gegear RJ, Thomson JD. 2004 Does the flower constancy of bumble bees reflect foraging  
489 economics? *Ethology* **110**, 793–805. (doi:10.1111/j.1439-0310.2004.01010.x)
- 490 21. Raine NE, Chittka L. 2007 Flower constancy and memory dynamics in bumblebees  
491 (Hymenoptera: Apidae: *Bombus*). *Entomologia Generalis* , 179–199.  
492 (doi:10.1127/entom.gen/29/2007/179)
- 493 22. Grüter C, Moore H, Firmin N, Helanterä H, Ratnieks FLW. 2011 Flower constancy in honey bee  
494 foragers (*Apis mellifera*) depends on ecologically realistic rewards. *Journal of Experimental*  
495 *Biology* **214**, 1397–1402.
- 496 23. Bruninga-Socular B, Winfree R, Crone EE. 2022 The contribution of plant spatial arrangement to  
497 bumble bee flower constancy. *Oecologia* **198**, 471–481. (doi:10.1007/s00442-022-05114-x)
- 498 24. Chittka L, Spaethe J, Schmidt A, Hickelsberger A. 2001 Adaptation, constraint, and chance in  
499 the evolution of flower color and pollinator color vision. In *Cognitive Ecology of Pollination:*  
500 *Animal Behaviour and Floral Evolution* (eds JD Thomson, L Chittka), pp. 106–126. Cambridge:  
501 Cambridge University Press. (doi:10.1017/CBO9780511542268.007)
- 502 25. Ramalho M, Giannini TC, Malagodi-Braga KS, Imperatriz-Fonseca VL. 1994 Pollen harvest by  
503 stingless bee foragers (Hymenoptera, Apidae, Meliponinae). *Grana* **33**, 239–244.  
504 (doi:10.1080/00173139409429005)
- 505 26. Somme L, Vanderplanck M, Michez D, Lombaerde I, Moerman R, Wathelet B, Wattiez R,  
506 Lognay G, Jacquemart A-L. 2015 Pollen and nectar quality drive the major and minor floral  
507 choices of bumble bees. *Apidologie* **46**, 92–106. (doi:10.1007/s13592-014-0307-0)
- 508 27. Hill PSM, Hollis J, Wells H. 2001 Foraging decisions in nectarivores: unexpected interactions  
509 between flower constancy and energetic rewards. *Animal Behaviour* **62**, 729–737.
- 510 28. Levin DA, Anderson WW. 1970 Competition for pollinators between simultaneously flowering  
511 species. *The American Naturalist* **104**, 455–467.

- 512 29. Waser NM. 1978 Interspecific pollen transfer and competition between co-occurring plant  
513 species. *Oecologia* **36**, 223–236. (doi:10.1007/BF00349811)
- 514 30. Geber MA, Moeller DA. 2006 Pollinator responses to plant communities and implications for  
515 reproductive character evolution. In *Ecology and Evolution of Flowers* (eds LD Harder, SCH  
516 Barrett), p. 0. Oxford University Press. (doi:10.1093/oso/9780198570851.003.0006)
- 517 31. Morales CL, Traveset A. 2008 Interspecific pollen transfer: magnitude, prevalence and  
518 consequences for plant fitness. *Critical Reviews in Plant Sciences* **27**, 221–238.  
519 (doi:10.1080/07352680802205631)
- 520 32. Mitchell RJ, Flanagan RJ, Brown BJ, Waser NM, Karron JD. 2009 New frontiers in competition  
521 for pollination. *Annals of Botany* **103**, 1403–1413. (doi:10.1093/aob/mcp062)
- 522 33. Ashman T-L, Arceo-Gómez G. 2013 Toward a predictive understanding of the fitness costs of  
523 heterospecific pollen receipt and its importance in co-flowering communities. *American Journal*  
524 *of Botany* **100**, 1061–1070. (doi:10.3732/ajb.1200496)
- 525 34. Tur C, Sáez A, Traveset A, Aizen MA. 2016 Evaluating the effects of pollinator-mediated  
526 interactions using pollen transfer networks: evidence of widespread facilitation in south Andean  
527 plant communities. *Ecology Letters* **19**, 576–586. (doi:10.1111/ele.12594)
- 528 35. Moreira-Hernández JI, Muchhala N. 2019 Importance of pollinator-mediated interspecific pollen  
529 transfer for angiosperm evolution. *Annual Review of Ecology, Evolution, and Systematics* **50**,  
530 191–217. (doi:10.1146/annurev-ecolsys-110218-024804)
- 531 36. Sargent RD, Ackerly DD. 2008 Plant–pollinator interactions and the assembly of plant  
532 communities. *Trends in Ecology & Evolution* **23**, 123–130. (doi:10.1016/j.tree.2007.11.003)
- 533 37. Rathcke B. 1983 Competition and facilitation among plants for pollination. In *Pollination*  
534 *Biology* (ed L Real), pp. 305–329. Academic Press. (doi:10.1016/B978-0-12-583980-8.50019-3)
- 535 38. Brosi BJ, Briggs HM. 2013 Single pollinator species losses reduce floral fidelity and plant  
536 reproductive function. *Proceedings of the National Academy of Sciences* **110**, 13044–13048.  
537 (doi:10.1073/pnas.1307438110)
- 538 39. Bruninga-Socular B, Socolar JB, Konzmann S, Lunau K. 2023 Pollinator-mediated plant  
539 coexistence requires high levels of pollinator specialization. *Ecology and Evolution* **13**, e10349.  
540 (doi:10.1002/ece3.10349)
- 541 40. Bartomeus I, Bosch J, Vilà M. 2008 High invasive pollen transfer, yet low deposition on native  
542 stigmas in a *Carpobrotus*-invaded community. *Annals of Botany* **102**, 417–424.  
543 (doi:10.1093/aob/mcn109)
- 544 41. Montgomery BR, Rathcke BJ. 2012 Effects of floral restrictiveness and stigma size on  
545 heterospecific pollen receipt in a prairie community. *Oecologia* **168**, 449–458.  
546 (doi:10.1007/s00442-011-2094-x)
- 547 42. Fang Q, Huang S-Q. 2013 A directed network analysis of heterospecific pollen transfer in a  
548 biodiverse community. *Ecology* **94**, 1176–1185. (doi:10.1890/12-1634.1)
- 549 43. Parra-Tabla V, Alonso C, Ashman T-L, Raguso RA, Albor C, Sosenski P, Carmona D, Arceo-  
550 Gómez G. 2021 Pollen transfer networks reveal alien species as main heterospecific pollen  
551 donors with fitness consequences for natives. *Journal of Ecology* **109**, 939–951.  
552 (doi:10.1111/1365-2745.13520)

- 553 44. Waser NM. 1978 Competition for hummingbird pollination and sequential flowering in two  
554 colorado wildflowers. *Ecology* **59**, 934–944. (doi:10.2307/1938545)
- 555 45. de Jager ML, Dreyer LL, Ellis AG. 2011 Do pollinators influence the assembly of flower colours  
556 within plant communities? *Oecologia* **166**, 543–553. (doi:10.1007/s00442-010-1879-7)
- 557 46. Runquist RB, Stanton ML. 2013 Asymmetric and frequency-dependent pollinator-mediated  
558 interactions may influence competitive displacement in two vernal pool plants. *Ecology Letters*  
559 **16**, 183–190. (doi:10.1111/ele.12026)
- 560 47. Campbell DR, Motten AF. 1985 The mechanism of competition for pollination between two  
561 forest herbs. *Ecology* **66**, 554–563. (doi:10.2307/1940404)
- 562 48. Bell JM, Karron JD, Mitchell RJ. 2005 Interspecific competition for pollination lowers seed  
563 production and outcrossing in *Mimulus ringens*. *Ecology* **86**, 762–771. (doi:10.1890/04-0694)
- 564 49. Ashman T-L, Alonso C, Parra-Tabla V, Arceo-Gómez G. 2020 Pollen on stigmas as proxies of  
565 pollinator competition and facilitation: complexities, caveats and future directions. *Annals of*  
566 *Botany* **125**, 1003–1012. (doi:10.1093/aob/mcaa012)
- 567 50. Braun J, Lortie CJ. 2019 Finding the bees knees: A conceptual framework and systematic review  
568 of the mechanisms of pollinator-mediated facilitation. *Perspectives in Plant Ecology, Evolution*  
569 *and Systematics* **36**, 33–40. (doi:10.1016/j.ppees.2018.12.003)
- 570 51. de Waal C, Anderson B, Ellis AG. 2015 Relative density and dispersion pattern of two southern  
571 African Asteraceae affect fecundity through heterospecific interference and mate availability, not  
572 pollinator visitation rate. *Journal of Ecology* **103**, 513–525. (doi:10.1111/1365-2745.12358)
- 573 52. Arceo-Gómez G. 2021 Spatial variation in the intensity of interactions via heterospecific pollen  
574 transfer may contribute to local and global patterns of plant diversity. *Annals of Botany* **128**,  
575 383–394. (doi:10.1093/aob/mcab082)
- 576 53. Wolf DE, Takebayashi N, Rieseberg LH. 2001 Predicting the risk of extinction through  
577 hybridization. *Conservation Biology* **15**, 1039–1053. (doi:10.1046/j.1523-  
578 1739.2001.0150041039.x)
- 579 54. Charlebois JA, Sargent RD. 2017 No consistent pollinator-mediated impacts of alien plants on  
580 natives. *Ecology Letters* **20**, 1479–1490. (doi:10.1111/ele.12831)
- 581 55. Jakobsson A, Padrón B, Traveset A. 2008 Pollen transfer from invasive *Carpobrotus* spp. to  
582 natives – A study of pollinator behaviour and reproduction success. *Biological Conservation* **141**,  
583 136–145. (doi:10.1016/j.biocon.2007.09.005)
- 584 56. Carneiro LT, Williams JN, Barker DA, Anderson JW, Martel C, Arceo-Gomez G. 2024 Patterns  
585 and drivers of pollen co-transport network structure vary across pollinator functional groups.  
586 *Journal of Ecology* **112**, 2319–2332. (doi:10.1111/1365-2745.14397)
- 587 57. Knight TM, Ashman T-L, Bennett JM, Burns JH, Passonneau S, Steets JA. 2018 Reflections on,  
588 and visions for, the changing field of pollination ecology. *Ecology Letters* **21**, 1282–1295.  
589 (doi:10.1111/ele.13094)
- 590 58. Travis DJ, Kohn JR. 2023 Honeybees (*Apis mellifera*) decrease the fitness of plants they  
591 pollinate. *Proceedings of the Royal Society B: Biological Sciences* **290**, 20230967.  
592 (doi:10.1098/rspb.2023.0967)



- 593 59. Trunschke J, Lunau K, Pyke GH, Ren Z-X, Wang H. 2021 Flower color evolution and the  
594 evidence of pollinator-mediated selection. *Front. Plant Sci.* **12**. (doi:10.3389/fpls.2021.617851)
- 595 60. Oyama RK, Jones KN, Baum DA. 2010 Sympatric sister species of californian *Antirrhinum* and  
596 their transiently specialized pollinators. In *The American Midland Naturalist*, pp. 337–347.  
597 (doi:10.1674/0003-0031-164.2.337)
- 598 61. Wenzell KE, Neequaye M, Paaianen P, Hill L, Brett P, Byers KJRP. 2025 Within-species floral  
599 evolution reveals convergence in adaptive walks during incipient pollinator shift. *Nat Commun*  
600 **16**, 2721. (doi:10.1038/s41467-025-57639-3)
- 601 62. Feldman TS, Morris WF, Wilson WG. 2004 When can two plant species facilitate each other's  
602 pollination? *Oikos* **105**, 197–207.
- 603 63. Johnson SD, Peter CI, Nilsson LA, Ågren J. 2003 Pollination success in a deceptive orchid is  
604 enhanced by co-occurring rewarding magnet plants. *Ecology* **84**, 2919–2927.
- 605 64. Pellegrino G, Bellusci F, Palermo AM. 2016 Who helps whom? Pollination strategy of *Iris*  
606 *tuberosa* and its relationship with a sexually deceptive orchid. *J Plant Res* **129**, 1051–1059.  
607 (doi:10.1007/s10265-016-0853-9)
- 608 65. Thomson JD, Fung HF, Ogilvie JE. 2019 Effects of spatial patterning of co-flowering plant  
609 species on pollination quantity and purity. *Annals of Botany* **123**, 303–310.  
610 (doi:10.1093/aob/mcy120)
- 611 66. Ye Z-M, Dai W-K, Jin X-F, Gituru RW, Wang Q-F, Yang C-F. 2014 Competition and  
612 facilitation among plants for pollination: can pollinator abundance shift the plant–plant  
613 interactions? *Plant Ecol* **215**, 3–13. (doi:10.1007/s11258-013-0274-y)
- 614 67. Moeller DA. 2004 Facilitative interactions among plants via shared pollinators. *Ecology* **85**,  
615 3289–3301. (doi:10.1890/03-0810)
- 616 68. Valdovinos FS, Moisset de Espanés P, Flores JD, Ramos-Jiliberto R. 2013 Adaptive foraging  
617 allows the maintenance of biodiversity of pollination networks. *Oikos* **122**, 907–917.  
618 (doi:10.1111/j.1600-0706.2012.20830.x)
- 619 69. Song Z, Feldman MW. 2014 Adaptive foraging behaviour of individual pollinators and the  
620 coexistence of co-flowering plants. *Proceedings of the Royal Society B: Biological Sciences* **281**,  
621 20132437. (doi:10.1098/rspb.2013.2437)
- 622 70. Pattrick JG, Symington HA, Federle W, Glover BJ. 2023 Bumblebees negotiate a trade-off  
623 between nectar quality and floral biomechanics. *iScience* **26**, 108071.  
624 (doi:10.1016/j.isci.2023.108071)
- 625 71. Charlton NL, Houston AI. 2010 What currency do bumble bees maximize? *PLoS One* **5**, e12186.  
626 (doi:10.1371/journal.pone.0012186)
- 627 72. Brodschneider R, Crailsheim K. 2010 Nutrition and health in honey bees. *Apidologie* **41**, 278–  
628 294. (doi:10.1051/apido/2010012)
- 629 73. Vaudo AD *et al.* 2020 Pollen protein: lipid macronutrient ratios may guide broad patterns of bee  
630 species floral preferences. *Insects* **11**, 132. (doi:10.3390/insects11020132)

- 631 74. Hayes L, Grüter C. 2023 When should bees be flower constant? An agent-based model highlights  
632 the importance of social information and foraging conditions. *Journal of Animal Ecology* **92**,  
633 580–593.
- 634 75. Wells H, Wells PH. 1983 Honey bee foraging ecology: optimal diet, minimal uncertainty or  
635 individual constancy? *Journal of Animal Ecology* **52**, 829–836.
- 636 76. Grüter C, Hayes L. 2022 Sociality is a key driver of foraging ranges in bees. *Current Biology* **32**,  
637 5390–5397. (doi:10.1016/j.cub.2022.10.064)
- 638 77. Grüter C, Segers FHID, Hayes L. 2024 Extensive loss of forage diversity in social bees due to  
639 flower constancy in simulated environments. *Proceedings of the Royal Society B: Biological*  
640 *Sciences* **291**, 20241036.
- 641 78. Chittka L, Thomson JD. 1997 Sensori-motor learning and its relevance for task specialization in  
642 bumble bees. *Behavioral Ecology and Sociobiology* **41**, 385–398.
- 643 79. Gegear RJ, Lavery TM. 2005 Flower constancy in bumblebees: a test of the trait variability  
644 hypothesis. *Animal Behaviour* **69**, 939–949.
- 645 80. von Frisch K. 1923 Über die ‘Sprache’ der Bienen. *Zoologisches Jahrbuch (Zoologie und*  
646 *Physiologie)* **40**, 1–186.
- 647 81. Woodward GL, Lavery TM. 1992 Recall of flower handling skills by bumble bees: a test of  
648 Darwin’s interference hypothesis. *Animal Behaviour* **44**, 1045–1051. (doi:10.1016/S0003-  
649 3472(05)80316-1)
- 650 82. Goulson D. 2000 Are insects flower constant because they use search images to find flowers?  
651 *Oikos* **88**, 547–552.
- 652 83. Lavery TM. 1994 Bumble bee learning and flower morphology. *Animal Behaviour* **47**, 531–545.
- 653 84. Raine NE, Chittka L. 2007 Pollen foraging: learning a complex motor skill by bumblebees  
654 (*Bombus terrestris*). *Naturwissenschaften* **94**, 459–464.
- 655 85. Mayberry MM, Naumer KC, Novinger AN, McCart DM, Wilkins RV, Muse H, Ashman T-L,  
656 Russell AL. 2024 Learning to handle flowers increases pollen collection for bees but does not  
657 affect pollination success for plants. *Behavioral Ecology* **35**, arae083.  
658 (doi:10.1093/beheco/arae083)
- 659 86. Menzel R. 1999 Memory dynamics in the honeybee. *Journal of Comparative Physiology A-*  
660 *Neuroethology Sensory Neural and Behavioral Physiology* **185**, 323–340.
- 661 87. Menzel R. 2012 The honeybee as a model for understanding the basis of cognition. *Nature*  
662 *Reviews Neuroscience* **13**, 758–768.
- 663 88. Chittka L, Gumbert A, Kunze J. 1997 Foraging dynamics of bumble bees: correlates of  
664 movements within and between plant species. *Behavioral Ecology* **8**, 239–249.
- 665 89. Lavery TM. 1994 Costs to foraging bumble bees of switching plant species. *Can. J. Zool.* **72**,  
666 43–47. (doi:10.1139/z94-007)
- 667 90. Shettleworth SJ. 2010 *Cognition, Evolution, and Behavior*. Oxford: Oxford University Press.

- 668 91. Jackson RR, Cross FR. 2018 Search Image. In *Encyclopedia of Animal Cognition and Behavior*  
669 (eds J Vonk, T Shackelford), pp. 1–7. Cham: Springer International Publishing.  
670 (doi:10.1007/978-3-319-47829-6\_1538-1)
- 671 92. Willmer P. 2011 *Pollination and Floral Ecology*. Princeton and Oxford: Princeton University  
672 Press.
- 673 93. Greggers U, Menzel R. 1993 Memory dynamics and foraging strategies of honeybees.  
674 *Behavioral Ecology and Sociobiology* **32**, 17–29.
- 675 94. Gervais A, Courtois È, Fournier V, Bélisle M. 2020 Landscape composition and local floral  
676 resources influence foraging behavior but not the size of *Bombus impatiens* Cresson  
677 (Hymenoptera: Apidae) workers. *PLOS ONE* **15**, e0234498. (doi:10.1371/journal.pone.0234498)
- 678 95. Martínez-Bauer AE, Chadwick FJ, Westmoreland AJ, Lander TA. 2021 Novel pollen analogue  
679 technique shows bumblebees display low floral constancy and prefer sites with high floral  
680 diversity. *Landscape Ecol* **36**, 3231–3247. (doi:10.1007/s10980-021-01304-9)
- 681 96. Austin MW, Horack P, Dunlap AS. 2019 Choice in a floral marketplace: the role of complexity  
682 in bumble bee decision-making. *Behavioral Ecology* **30**, 500–508. (doi:10.1093/beheco/ary190)
- 683 97. Charnov EL. 1976 Optimal foraging, the marginal value theorem. *Theoretical Population*  
684 *Biology* **9**, 129–136. (doi:10.1016/0040-5809(76)90040-X)
- 685 98. Heinrich B. 1979 ‘Majoring’ and ‘Minoring’ by foraging bumblebees, *Bombus vagans*: An  
686 experimental analysis. *Ecology* **60**, 246–255. (doi:10.2307/1937652)
- 687 99. Hill PSM, Wells PH, Wells H. 1997 Spontaneous flower constancy and learning in honey bees as  
688 a function of colour. *Animal Behaviour* **54**, 615–627.
- 689 100. Wells H, Rathore RRS. 1994 Foraging ecology of the Asian hive bee, *Apis cerana indica*, within  
690 artificial flower patches. *Journal of Apicultural Research* **33**, 219–230.
- 691 101. Slaa EJ, Tack AJM, Sommeijer MJ. 2003 The effect of intrinsic and extrinsic factors on flower  
692 constancy in stingless bees. *Apidologie* **34**, 457–468.
- 693 102. Muth F, Keasar T, Dornhaus A. 2015 Trading off short-term costs for long-term gains: how do  
694 bumblebees decide to learn morphologically complex flowers? *Animal Behaviour* **101**, 191–199.  
695 (doi:10.1016/j.anbehav.2014.12.024)
- 696 103. Krishna S, Keasar T. 2021 Generalization of foraging experience biases bees toward flowers with  
697 complex morphologies. *Front. Ecol. Evol.* **9**. (doi:10.3389/fevo.2021.655086)
- 698 104. Cnaani J, Thomson JD, Papaj D. 2006 Flower choice and learning in foraging bumblebees:  
699 effects of variation in nectar volume and concentration. *Ethology* **112**, 278–285.
- 700 105. Cakmak I *et al.* 2009 Different solutions by bees to a foraging problem. *Animal Behaviour* **77**,  
701 1273–1280.
- 702 106. Rathcke B. 1988 Interactions for pollination among coflowering shrubs. *Ecology* **69**, 446–457.  
703 (doi:10.2307/1940443)
- 704 107. Ishii HS, Kadoya EZ. 2016 Legitimate visitors and nectar robbers on *Trifolium pratense* showed  
705 contrasting flower fidelity versus co-flowering plant species: could motor learning be a major

determinant of flower constancy by bumble bees? *Behav Ecol Sociobiol* **70**, 377–386.  
(doi:10.1007/s00265-016-2057-7)

108. Sanderson CE, Orozco BS, Hill PSM, Wells H. 2006 Honeybee (*Apis mellifera ligustica*)  
response to differences in handling time, rewards and flower colours. *Ethology* **112**, 937–946.

109. Kulahci IG, Dornhaus A, Papaj DR. 2008 Multimodal signals enhance decision making in  
foraging bumble-bees. *Proceedings of the Royal Society of London Series B-Biological Sciences*  
**275**, 797–802.

110. Ishii HS. 2006 Floral display size influences subsequent plant choice by bumble bees. *Functional*  
*Ecology* **20**, 233–238. (doi:10.1111/j.1365-2435.2006.01106.x)

111. Ishii HS, Masuda H. 2014 Effect of flower visual angle on flower constancy: a test of the search  
image hypothesis. *Behavioral Ecology* **25**, 933–944.

112. Wright EK, Timberlake TP, Baude M, Vaughan IP, Memmott J. 2024 Quantifying the  
production of plant pollen at the farm scale. *New Phytologist* **242**, 2888–2899.  
(doi:10.1111/nph.19763)

113. Krishna S, Keasar T. 2019 Bumblebees forage on flowers of increasingly complex morphologies  
despite low success. *Animal Behaviour* **155**, 119–130. (doi:10.1016/j.anbehav.2019.06.028)

114. Ohashi K. 2002 Consequences of floral complexity for bumblebee-mediated geitonogamous self-  
pollination in *Salvia nipponica* miq. (Labiatae). *Evolution* **56**, 2414–2423. (doi:10.1111/j.0014-  
3820.2002.tb00167.x)

115. Krishna S, Keasar T. 2018 Morphological complexity as a floral signal: from perception by  
insect pollinators to co-evolutionary implications. *Int J Mol Sci* **19**, 1681.  
(doi:10.3390/ijms19061681)

116. Heinrich B. 1979 *Bumblebee Economics*. Cambridge, Massachusetts and London, England:  
Harvard University Press.

117. Marden JH, Waddington KD. 1981 Floral choices by honey bees in relation to the relative  
distances to flowers. *Physiological Entomology* **6**, 431–435.

118. Kunin WE. 1993 Sex and the single mustard: population density and pollinator behavior effects  
on seed-set. *Ecology* **74**, 2145–2160.

119. Ishii HS. 2005 Analysis of bumblebee visitation sequences within single bouts: implication of the  
overstrike effect on short-term memory. *Behav Ecol Sociobiol* **57**, 599–610.  
(doi:10.1007/s00265-004-0889-z)

120. Katz EJ, Essenberg CJ. 2018 The effect of the dispersion of rewarding and rewardless flowers on  
visitation and constancy by bumblebees (*Bombus impatiens*). *Journal of Pollination Ecology* **23**,  
119–126. (doi:10.26786/1920-7603(2018)14)

121. Heinrich B. 1976 The foraging specializations of individual bumblebees. *Ecological*  
*Monographs* **46**, 105–128. (doi:10.2307/1942246)

122. Smith C, Weinman L, Gibbs J, Winfree R. 2019 Specialist foragers in forest bee communities are  
small, social or emerge early. *Journal of Animal Ecology* **88**, 1158–1167. (doi:10.1111/1365-  
2656.13003)

- 745 123. Bateman AJ. 1951 The taxonomic discrimination of bees. *Heredity* **5**, 271–278.  
746 (doi:10.1038/hdy.1951.24)
- 747 124. Leonhardt SD, Blüthgen N. 2012 The same, but different: pollen foraging in honeybee and  
748 bumblebee colonies. *Apidologie* **43**, 449–464. (doi:10.1007/s13592-011-0112-y)
- 749 125. Townsend-Mehler JM, Dyer FC, Maida K. 2011 Deciding when to explore and when to persist: a  
750 comparison of honeybees and bumblebees in their response to downshifts in reward. *Behavioral*  
751 *Ecology and Sociobiology* **65**, 305–312.
- 752 126. Yourstone J, Varadarajan V, Olsson O. 2023 Bumblebee flower constancy and pollen diversity  
753 over time. *Behavioral Ecology* **34**, 602–612. (doi:10.1093/beheco/arad028)
- 754 127. Alarcón R. 2010 Congruence between visitation and pollen-transport networks in a California  
755 plant–pollinator community. *Oikos* **119**, 35–44. (doi:10.1111/j.1600-0706.2009.17694.x)
- 756 128. Williams NM, Tepedino VJ. 2003 Consistent mixing of near and distant resources in foraging  
757 bouts by the solitary mason bee *Osmia lignaria*. *Behavioral Ecology* **14**, 141–149.  
758 (doi:10.1093/beheco/14.1.141)
- 759 129. von Frisch K. 1967 *The dance language and orientation of bees*. Cambridge, Massachusetts:  
760 Harvard University Press.
- 761 130. Farina WM, Grüter C, Arenas A. 2012 Olfactory information transfer during recruitment in  
762 honey bees. In *Honeybee Neurobiology and Behavior - A Tribute to Randolph Menzel* (eds CG  
763 Galizia, D Eisenhardt, M Giurfa), pp. 89–101. Heidelberg: Springer.
- 764 131. Alves DA, George EA, Kaur R, Brockmann A, Hrncir M, Grüter C. 2023 Diverse  
765 communication strategies in bees as a window into adaptations to an unpredictable world.  
766 *Proceedings of the National Academy of Sciences* **120**, e2219031120.  
767 (doi:10.1073/pnas.2219031120)
- 768 132. Hrncir M. 2009 Mobilizing the foraging force mechanical signals in stingless bee recruitment. In  
769 *Food Exploitation by Social Insects: Ecological, Behavioral, and Theoretical Approaches* (eds S  
770 Jarau, M Hrncir), Boca Raton, Florida.: CRC Press, Taylor & Francis Group.
- 771 133. Jarau S. 2009 Chemical communication during food exploitation in stingless bees. In *Food*  
772 *Exploitation by Social Insects: Ecological, Behavioral, and Theoretical Approaches* (eds S Jarau,  
773 M Hrncir), Boca Raton, Florida: CRC University Press.
- 774 134. Grüter C. 2020 *Stingless Bees: Their Behaviour, Ecology and Evolution*. Springer International  
775 Publishing. See <https://www.springer.com/gp/book/9783030600891>.
- 776 135. Lindauer M, Kerr WE. 1960 Communication between the workers of stingless bees. *Bee World*  
777 **41**, 29–71.
- 778 136. Nieh JC. 2004 Recruitment communication in stingless bees (Hymenoptera, Apidae,  
779 Meliponini). *Apidologie* **35**, 159–182.
- 780 137. Dornhaus A, Chittka L. 1999 Evolutionary origins of bee dances. *Nature* **401**, 38–38.
- 781 138. Hrncir M, Schmidt VM, Schorkopf DLP, Jarau S, Zucchi R, Barth FG. 2006 Vibrating the food  
782 receivers: a direct way of signal transmission in stingless bees (*Melipona seminigra*). *Journal of*  
783 *Comparative Physiology A* **192**, 879–887.

139. Vaudo AD, Tooker JF, Grozinger CM, Patch HM. 2015 Bee nutrition and floral resource restoration. *Current Opinion in Insect Science* **10**, 133–141. (doi:10.1016/j.cois.2015.05.008)
140. Eckhardt M, Haider M, Dorn S, Müller A. 2014 Pollen mixing in pollen generalist solitary bees: a possible strategy to complement or mitigate unfavourable pollen properties? *Journal of Animal Ecology* **83**, 588–597. (doi:10.1111/1365-2656.12168)
141. Filipiak M, Kuszewska K, Asselman M, Denisow B, Stawiarz E, Woyciechowski M, Weiner J. 2017 Ecological stoichiometry of the honeybee: Pollen diversity and adequate species composition are needed to mitigate limitations imposed on the growth and development of bees by pollen quality. *PLOS ONE* **12**, e0183236. (doi:10.1371/journal.pone.0183236)
142. Ruedenauer FA, Raubenheimer D, Kessner-Beierlein D, Grund-Mueller N, Noack L, Spaethe J, Leonhardt SD. 2020 Best be(e) on low fat: linking nutrient perception, regulation and fitness. *Ecology Letters* **23**, 545–554. (doi:10.1111/ele.13454)
143. Génissel A, Aupinel P, Bressac C, Tasei J-N, Chevrier C. 2002 Influence of pollen origin on performance of *Bombus terrestris* micro-colonies. *Entomologia Experimentalis et Applicata* **104**, 329–336.
144. Schmidt JO, Thoenes SC, Levin MD. 1987 Survival of honey bees, *Apis mellifera* (Hymenoptera: Apidae), fed various pollen sources. *Annals of the Entomological Society of America* **80**, 176–183. (doi:10.1093/aesa/80.2.176)
145. Arenas A, Lajad R, Peng T, Grüter C, Farina W. 2021 Correlation between octopaminergic signalling and foraging task specialisation in honeybees. *Genes, Brain and Behavior* **20**, e12718. (doi:https://doi.org/10.1111/gbb.12718)
146. Mateus S, Ferreira-Caliman MJ, Menezes C, Grüter C. 2019 Beyond temporal-polyethism: division of labor in the eusocial bee *Melipona marginata*. *Insectes Sociaux* **66**, 317–328.
147. Balbuena MS, Farina WM. 2020 Chemosensory reception in the stingless bee *Tetragonisca angustula*. *Journal of Insect Physiology* **125**, 104076. (doi:10.1016/j.jinsphys.2020.104076)
148. Ramalho M. 1990 Foraging by stingless bees of the genus, *Scaptotrigona* (Apidae, Meliponinae). *Journal of Apicultural Research* **29**, 61–67. (doi:10.1080/00218839.1990.11101198)
149. Brodschneider R *et al.* 2021 CSI Pollen: Diversity of honey bee collected pollen studied by citizen scientists. *Insects* **12**, 987. (doi:10.3390/insects12110987)
150. M. AJ, N. KM, Sreekumar S, Kelber A, Somanathan H. 2025 Influence of floral traits on visitation patterns in a miniature tropical stingless bee, *Tetragonula iridipennis*. *Sci Nat* **112**, 44. (doi:10.1007/s00114-025-01994-0)
151. Michener CD. 1974 *The Social Behavior of the Bees*. Cambridge, Massachusetts: Harvard University Press.
152. Bruninga-Socular B, Crone EE, Winfree R. 2016 The role of floral density in determining bee foraging behavior: a natural experiment. *naar* **36**, 392–399. (doi:10.3375/043.036.0406)
153. Dauber J *et al.* 2010 Effects of patch size and density on flower visitation and seed set of wild plants: a pan-European approach. *Journal of Ecology* **98**, 188–196. (doi:10.1111/j.1365-2745.2009.01590.x)